

Serveur Académique Lausannois SERVAL [serval.unil.ch](http://serval.unil.ch)

## Author Manuscript

Faculty of Biology and Medicine Publication

This paper has been peer-reviewed but does not include the final publisher proof-corrections or journal pagination.

Published in final edited form as:

**Title:** Low emotion-oriented coping and informal help-seeking behaviour as major predictive factors for improvement in major depression at 5-year follow-up in the adult community.

**Authors:** Rodgers S, Vandeleur CL, Strippoli MF, Castelao E, Tesic A, Glaus J, Lasserre AM, Müller M, Rössler W, Ajdacic-Gross V, Preisig M

**Journal:** Social psychiatry and psychiatric epidemiology

**Year:** 2017

**Issue:** 52

**Volume:** 9

**Pages:** 1169-1182

**DOI:** 10.1007/s00127-017-1421-x

In the absence of a copyright statement, users should assume that standard copyright protection applies, unless the article contains an explicit statement to the contrary. In case of doubt, contact the journal publisher to verify the copyright status of an article.

**Coping behaviours as a major predictive factor for improvement in major depression at 5-year follow-up in a Swiss community sample of adults**

**Short title: Predictive factors of improvement in depression after 5 years**

*S. Rodgers<sup>1,2\*</sup>, C. L. Vandeleur<sup>3</sup>, M.-P. F. Strippoli<sup>3</sup>, E. Castelao<sup>3</sup>, A. Tesic<sup>1</sup>, J. Glaus<sup>3,4</sup>, A. M. Lasserre<sup>3</sup>, M. Müller<sup>1</sup>, W. Rössler<sup>1,5,6</sup>, V. Ajdacic-Gross<sup>1,2</sup>, M. Preisig<sup>3</sup>*

<sup>1</sup>Department of Psychiatry, Psychotherapy and Psychosomatics, Psychiatric Hospital, University of Zurich, Switzerland

<sup>2</sup>Epidemiology, Biostatistics and Prevention Institute, Swiss MS Registry, University of Zurich, Switzerland

<sup>3</sup>Psychiatric Epidemiology and Psychopathology Research Centre, Department of Psychiatry, Lausanne University Hospital, Lausanne, Switzerland

<sup>4</sup>Genetic Epidemiology Research Branch, Intramural Research Program, National Institute of Mental Health, Bethesda, MD, USA

<sup>5</sup>Collegium Helveticum, University of Zurich and Swiss Federal Institute of Technology, Zurich, Switzerland

<sup>6</sup>Institute of Psychiatry, Laboratory of Neuroscience (LIM 27), University of Sao Paulo, Brazil

Word count (excluding abstract, tables/figures and references): 3711

Manuscript submitted as original paper to:  
*Social Psychiatry and Psychiatric Epidemiology*

\*Corresponding author:

Stephanie Rodgers

Department of Psychiatry, Psychotherapy and Psychosomatics

Psychiatric Hospital, University of Zurich

PO Box 1930

CH-8021 Zurich

Switzerland

Tel: 0041-44-296 74 22

Fax: 0041-44-296 73 09

E-Mail: [stephanie.rodgers@dgsp.uzh.ch](mailto:stephanie.rodgers@dgsp.uzh.ch)

**Acknowledgements**

The authors would like to express their gratitude to the inhabitants of Lausanne who volunteered to participate in the PsyCoLaus study and to the collaborators who contributed to the coordination of the study and the collection of data. We would also like to thank all the investigators of the CoLaus|PsyCoLaus study, who made the psychiatric study possible, as well as many GSK employees who contributed to the execution of this study.

**Abstract** (250 words)

**Purpose:** Given the broad range of biopsychosocial difficulties resulting from Major Depressive Disorder (MDD), reliable evidence for predictors of improved mental health is essential, particularly from unbiased prospective community samples. Consequently, potential clinical and non-clinical predictors of improved mental health, defined as an absence of current MDE at follow-up, were examined over a 5-year period in a community sample.

**Methods:** The longitudinal population-based PsyCoLaus study from the city of Lausanne, Switzerland, was used. Subjects having a lifetime MDD with a current Major Depressive Episode (MDE) at baseline assessment were selected, resulting in a subsample of 210 subjects. Logistic regressions were applied to the data.

**Results:** Coping behaviours were the most important predictive factors in the present study. More specifically, low emotion-oriented coping and informal help-seeking behaviour at baseline were associated with the absence of an MDD diagnosis at follow-up. Notably, the partitioning of depression into subtypes did not show differential associations and only a trend-level association was found between regular physical activity and the absence of MDD diagnosis at follow-up.

**Conclusions:** The paramount role of coping behaviours as predictors of improvement in depression found in the present study might be a valuable target for resource-oriented therapeutic models. On the one hand, the positive impact of low emotion-oriented coping highlights the utility of clinical interventions interrupting excessive mental ruminations during MDE. On the other hand, the importance of informal social networks raises questions regarding how to enlarge the personal network of affected subjects and on how to best support informal caregivers.

**Keywords:** *Depression, epidemiology, prospective study, population survey.*

## **Introduction**

Understanding the circumstances of improved mental health state in individuals with Major Depressive Disorder (MDD) is a crucial public health concern, given that this disorder represents a highly prevalent mental illness and is associated with high levels of individual suffering and enormous burdens on society [1-3]. A large number of studies have focused on the risk factors for poor MDD outcomes, such as relapse [e.g., 4,5-7], recurrence [e.g., 8,9-11], or, with a focus on the current episode, persistence [e.g., 12,13,14]. In contrast, there is less evidence on predictive factors of positive outcomes, such as remission states/recovery [e.g., 15,16,17], apart from the knowledge logically derived from risk factors, such as baseline severity, and comorbidity [18], which can be seen as the other side of the same coin. Yet it is important to gain more knowledge about relevant predictive factors – particularly for resource-oriented therapeutic models [19] – in order to help individuals improve their mental health state. The generalization of recent findings is, however, limited by the fact that the few existing studies focusing on remission states/recovery have used clinical samples [20-22]. Knowledge from representative community samples is rather sparse, despite its high potential external validity.

A recent community study addressed this limitation by examining non-clinical predictors on time to remission over a 12-year period [23]. The results indicate that a history of childhood physical abuse is significantly associated with longer time to remission, whereas lack of social support and presence of pain and migraines show trend-level associations. On the other hand, no single demographic variable was related to time to remission [23]. Clinical and depression-related factors (i.e., depression history, index episode severity, index comorbidity, and personality factors including low neuroticism and coping styles) were shown to have an impact on depression outcomes in clinical-based studies [13,24-26]. Of the commonly distinguished problem-oriented and emotion-oriented coping styles [27], emotion-oriented coping (EOC) was specifically correlated with depression [28,29]. While EOC is geared towards managing the negative emotions often generated by stressful circumstances, problem-oriented coping (POC) is geared towards changing the stressful situation itself [30]. Moreover, level of

education, marital status, physical activity, younger age, less childhood adversity, gender (at least in some studies), and social support were associated with remission/positive depression outcomes [26,23,31]. The issue of social support needs to be complemented by help-seeking behaviour, considering that subjective willingness to seek help and the personal feelings and beliefs about receiving aid from others is the most direct predictor of whether or not a person receives help [32]. Based on this evidence, we conclude that both clinical and non-clinical variables should be considered in community surveys.

Little is known about remission/recovery from depression subtypes to date, despite growing evidence demonstrating that MDD is highly heterogeneous comprising different clinical symptom subtypes [33]. Recent studies based on clinical samples have accounted for the remission of depression symptom subtypes by focusing on reduction of symptom severity scales after antidepressant treatment interventions [34-37]. In the recent longitudinal study of Lamers et al. [38], involving a combined clinical and community sample, baseline severity differentiated between depression subtypes, with an unfavourable course for the atypical subtype (regarding somatic health) and the melancholic subtype (regarding suicidal thoughts and anxiety) compared to the control and moderate groups, while the trajectories of the severe subtypes mostly ran parallel over the 6-year time-period. To our knowledge, no study has yet considered symptom-based depression subtypes as predictors of remission in a pure community sample of adults.

Taken together, the studies above highlight the importance of identifying factors that predict positive outcomes of individuals suffering from MDD in community samples. Therefore, we sought to investigate potential predictive factors of improved mental health over a 5-year period in a prospective, population-based sample of adults meeting criteria for lifetime MDD with a current Major Depressive Episode (MDE) at baseline. The extensive collection of data enabled us to investigate a broad spectrum of potential clinical and non-clinical predictors, including a series of childhood and disorder-related factors, behavioural and personality characteristics as well as the body mass index (BMI).

## **Material and methods**

### *Study design and participants*

The data for this article stemmed from CoLaus|PsyCoLaus, a prospective cohort study designed to study mental disorders and cardiovascular risk factors in the community and to determine their associations. The methodological features of this study have been already described in detail [39,40]. The original sample of 6734 subjects was randomly selected from the 35- to 75-year-old residents of the city of Lausanne (Switzerland) between 2003 and 2006 according to the civil register. Sixty-seven percent of the 35- to 66-year-old participants who underwent the physical exam ( $n = 5535$ ) also accepted the psychiatric evaluation, resulting in a sample of 3719 individuals in the baseline assessment ( $T_0$ ). All subjects were also invited to participate at the first follow-up ( $T_1$ ), which took place between 2009 and 2013. A total of 5064 subjects participated in the physical and 4004 subjects in the psychiatric follow-up evaluations.

The present analyses relied on those subjects who fulfilled the criteria of lifetime MDD with a current MDE at  $T_0$  ( $n = 293$ ) who also took part in the psychiatric follow-up evaluation  $T_1$  ( $n=210$ ). Moreover, analyses comprising the Problem Resolution Strategy questionnaire needed to be restricted to a smaller subsample ( $n = 144$ ), since this questionnaire was only completed by a subgroup of participants.

The study was approved by the Institutional Ethics Committee of the University of Lausanne. According to the declaration of Helsinki, all participants signed written informed consent after being informed about the goal and funding of the study.

## ***Measurements***

### *Mental disorders*

Information on mental disorders and their characteristics, treatment, and illness-related factors was collected using the semi-structured Diagnostic Interview for Genetic Studies [DIGS; 41,42]. The DIGS was developed and validated by the National Institute of Mental Health (NIMH) Molecular Genetics Initiative [42] to obtain a wide spectrum of DSM-IV [43] Axis I criteria. The French version of the DIGS was modified to include sections on anxiety disorders from the Schedule for Affective Disorders and Schizophrenia – Lifetime Version (SADS-LA) [44].

The French translation of the DIGS [45] was comprehensively tested and showed excellent inter-rater (kappa = 0.93) and a slightly lower 6-week test–retest reliability (kappa = 0.62) for MDD [41]. Inter-rater reliability was excellent for substance use disorders [46], and inter-rater and test-retest reliabilities were also satisfactory for the anxiety sections [47,48].

The DIGS section assessing depressive phenomenology records precise information on age of onset, the current/recent/last/most severe depressive episodes, information on course including duration of MDEs, and duration and chronology of comorbid conditions [40]. The criteria for the MDD *specifiers* according to the DSM-IV [49] were used to diagnose atypical and melancholic depressive episodes. Based on the anamnesis of the MDD lifetime diagnosis and the depressive phenomenology, atypical, melancholic, combined atypical-melancholic and unspecified depression subtypes were computed (see Glaus et al., [50] for more information).

Information on demographic characteristics at T<sub>0</sub> was derived from the DIGS. The Global Assessment of Functioning (GAF) score was applied at baseline. Socio-economic status (SES) was assessed using the Hollingshead scale [51].



### *Childhood-related factors at baseline*

Information on childhood-related adversities was dichotomized into any versus none if one of the following questions was confirmed:

- did your parents fight frequently amongst themselves? (interparental violence);
- did your parents ever do anything that frightened you (like lock you in a closet)? (fear of maltreatment by parents);
- did any of the following occur before your 16th birthday: .... put in foster care? (foster care);

General childhood unhappiness was rated by the participants by answering the question “overall, how would you characterize your childhood?” on a four-point Likert scale ranging from 1 = “happy” to 4 = “very unhappy”. The trauma items stemmed from the French version of the SADS-La. A cut-off of <17 years was used to define childhood trauma, which covered the events 1. accident, 2. physical assault, 3. combat and/or war, 4. witness of murder, violence or accidental death, and 5. sexual abuse.

### *Behavioural characteristics at baseline*

The behavioural characteristics regular physical activity, and regular alcohol consumption were derived from the DIGS at baseline and led to three binary coded variables (1 = “yes”; 0 = “no”). Regular smoking was grouped into the three categories “never”, “former”, and “current”. The definition of regular smoking required at least 5 days’ smoking per week and regular alcohol consumption entailed consumption at least once a week for at least 6 months. Those participants who were physically active at least 2-3 times per week were assigned as physically active.

### *Personality characteristics at baseline*

Subjective coping strategies were measured by the French translation of the self-rating Problem Resolution Strategy questionnaire [52,53]. This questionnaire consists of 17 items on possible reactions to problematic situations. Every item is rated on a four-level Likert scale, ranging from “very unusual for me” to “very usual for me”. Perrin et al. [54] demonstrated the division of the items of the French version into the three broad domains of *emotional problem resolution strategy* (9 items), *help-seeking behaviours* (4 items) and *active problem resolution strategy* (4 items). The standardized

Cronbach's alpha coefficients were 0.65 (EOC), 0.69 (help-seeking), and 0.44 (POC), respectively [54]. Item 17 referring to professional help was excluded from the help-seeking subscale for the current study to distinguish professional help from informal help (i.e., family, friends, colleagues, neighbours). The Cronbach's alpha coefficient for this informal help subscale without item 17 was 0.74.

#### *Assessment of physical features at baseline*

The BMI providing information on adiposity was calculated as weight in kilograms divided by height in meters squared. Participants were dressed in light indoor clothes without shoes while measured and weighed. These physical measurements were taken after an 8-hour fast and after refraining from any extensive physical activity for 12 hours.

#### *Definition of the outcome at the psychiatric follow-up*

Improvement in major depression at  $T_1$  was defined by the absence of a current MDE at  $T_1$ .

#### *Statistical analysis*

Logistic regressions for binary outcome variables [55] were applied to the data of the 210 depressive participants at  $T_0$ . The binary coded dependent variable contained information about the absence versus the presence of a current MDE at  $T_1$ . The following  $T_0$ -predictor variables were considered as variables of interest: marital status, living status, childhood-related factors (general unhappiness, adversities, trauma), disorder-related factors (age of onset of MDD, depression subtype at  $T_0$ , number of depression symptoms during the current MDE, duration of all MDE, duration of the current MDE, number of MDEs, current GAF score of all disorders at  $T_0$ , help from a doctor or other healthcare professional, psychopharmacological treatment (antidepressants), anxiety disorder at  $T_0$ , drug abuse at  $T_0$ , and number of comorbid diagnoses (lifetime)), behavioural characteristics (regular smoking, alcohol consumption, physical activity), personality characteristics (EOC, informal help-seeking behaviour, POC), and the somatic characteristic BMI.

Independent predictors were simultaneously tested in a final logistic regression model, adjusted for sex, age and SES, if they were significantly associated with MDE status at follow-up in the initial marginal logistic regression (cut-off level:  $p < 0.20$ ). This non-restrictive cut-off level was chosen due to the small sample sizes. In addition, this model was complemented by a model following stepwise backward selection ( $p_{in} \leq 0.05$  and  $p_{out} \geq 0.10$ ), and a model using imputed data (missing value analysis (MVA)) for the missing values in the coping variables.

The  $p$ -values were considered significant at  $< 0.05$ . All analyses were performed using SPSS version 20 for Macintosh (SPSS Inc., USA).

## Results

Among the 210 individuals meeting the criteria for lifetime MDD with a current MDE at  $T_0$  and also taking part in the psychiatric follow-up evaluation, 65.7% were women. The mean age of this sample was 50.1 years ( $y$ ) ( $SD = 8.1$ ; min. 36.3  $y$ ; max. 68.6  $y$ ) and the mean SES according to Hollingshead's index was 3.1 ( $SD = 1.3$ ) (a value of 3 represents a middle class). At  $T_1$ , 72.9% no longer fulfilled criteria for a MDE and were considered improved, whereas 27.1% still had ongoing MDE, either because of the persistence of the initial episode or due to the occurrence of a new episode.

Table 1 displays the results of the marginal logistic regression models yielding potential determinants of improvement in major depression. The following  $T_0$ -variables were significantly associated with the absence of a current MDE at  $T_1$ : depression subtype (i.e., the category combined atypical-melancholic subtype compared to the unspecified subtype), a lower number of depression symptoms during the current MDE, a higher GAF score of all disorders, and a lower EOC.

*--Insert Table 1 about here--*

Based on the cut-off criterion of  $p < 0.2$ , the following variables were simultaneously included as independent predictors in the final model, adjusted for sex, age and SES: depression subtypes, number of depression symptoms during the current MDE, current GAF score of all disorders, help from a doctor or other healthcare professional, psychopharmacological treatment (antidepressants), regular physical activity, EOC and informal help-seeking behaviour (Table 2). Low EOC and high informal help-seeking coping were significantly associated with absence of MDE at  $T_1$  in this final model. All the other predictors found in the previous marginal logistic regression models were no longer significantly associated with MDE status at  $T_1$ . Both the model following stepwise backward selection and the model using MVA confirmed that low EOC and high informal help-seeking coping were significantly associated with absence of MDE at  $T_1$  (data not shown).

*--Insert Table 2 about here--*

## **Discussion**

This longitudinal study provides important insight into predictive factors of improved mental health state in a population-based sample of depressive residents of the city of Lausanne, Switzerland. The present study examined the effects of both clinical and non-clinical variables on improved mental health state, defined as an absence of a current MDE at the end of a five-year follow-up period. The predictive effect of coping-related behaviours was demonstrated in the sense that help-seeking behaviour, which was restricted to the informal, private social environment, and decreased EOC were unequivocally associated with improvement in major depression.

### *Coping*

Low EOC scores – but not POC scores – emerged as a predictive factor with regard to an absence of MDE at follow-up. The opposite association, i.e., the link between high EOC and depression, has been demonstrated in previous research [29,28]. The problems of EOC when dealing with one's own negative emotions/distress could be related to the concept of rumination. At least certain elements of this concept are included in the EOC subscales [27] and some theorists have argued that rumination may be one type of EOC [56]. According to Nolen-Hoeksema et al. [57-59] rumination holds the potential to exacerbate and perpetuate depressed mood. In the meantime, rumination was shown to be strongly associated with the level of distress and onset, course, and duration of depression [60-62]. Ruminative thoughts also represent a core element of the influential cognitive model of Aaron T. Beck [63], introduced nearly 50 years ago. In fact, current neurobiological research focusing on an integrated cognitive-neurobiological perspective identified possible neurobiological underpinnings of Beck's model [60]. Ruminative thoughts were associated with hyperactivity in regions involved in emotional recall (such as the amygdala and hippocampus) and hypoactivity of higher cortical regions (such as the dorsolateral prefrontal cortex), adaptively regulating lower regions and therefore resulting in inhibited top-down processes [60]. Whatever the underlying biological mechanisms are, the predictive effect of low EOC is a promising target for therapy. For example, the skill of accepting/tolerating negative emotions was particularly highlighted as being beneficial to mental

health [27]. In contrast, POC, an adaptive strategy, showed no influence on improvement in major depression in the present study. This confirms earlier findings demonstrating that adaptive emotion regulation strategies show weaker associations with psychopathology than maladaptive strategies [64].

Informal – but not formal – help-seeking behaviour was a further predictor for the absence of depression 5 years after baseline. The absence of formal help-seeking could be a result of the high age-span of the PsyCoLaus sample. In a large Norwegian population, elderly people sought less professional help than did younger ones, with a distinct drop after middle age [65], suggesting a shift from formal to informal help-seeking with increasing age. Furthermore, in a large national population-based sample, subjects tended to endorse informal rather than formal help when they were asked how they would cope in the case of psychological problems [66]. The importance of informal help-seeking was also highlighted in another community study [67] and a clinical sample [68]. It is surprising then that informal help-seeking has generally been less investigated than formal help-seeking. Considering the deterrent effect of stigma on help-seeking for mental health problems [69], a possible explanation for the beneficial effect of informal compared to formal help might be less stigma-associated stress [70]. However, this view is not supported by others, who explicitly emphasise the stigmatising effect of informal help-seeking [71]. Beyond that, the health of informal caregivers, including relatives, spouses and particularly offspring, also requires special attention, since problematic consequences occur frequently and can cause distress among caregivers [72].

In sum, coping strategies showed strong predictive validity with regard to the depression outcome. This is in line with Conradi [25] who found that, apart from the number of prior MDEs, the coping potential was the strongest predictor of time to recurrence. Considering that coping-behaviour can be seen as the behavioural expression of various personality traits as well as cross-situational beliefs (so-called *generalized expectancies*) about one's ability to terminate, e.g., a negative mood state, [73-76], it represents an important target for treatment.

*Variables lacking impact on improvement in major depression*

The association between regular physical activity and the absence of a current MDE at the end of the five-year follow-up period showed only a borderline significant trend. Therefore, we could only partly confirm previous studies that highlighted the positive effect of physical activity in depressive subjects [77,78,31]. Plausible explanations for this predictive trend are the ability of physical activity to “energise” and produce more positive mood, improve body image [79], enhance the social aspects of physical activity [80] and enhance immuno-neuroendocrine stabilization in subjects with dysregulated inflammatory and stress feedback [81]. Finally, the question remains as to how much physical activity is needed to produce beneficial effects on MDD: clear guidelines with regard to the exact intensity and frequency are still lacking [82]. Moreover, a systematic review and meta-analysis confirmed the short-term effect of physical exercise in clinically depressed adults, but evidence regarding long-term effectiveness is still sparse [83].

In the present paper, the depression subtypes did not differ regarding improvement in major depression. Aside from the way we defined improvement in major depression, this could have resulted from the fact that we included depression severity in the model. Melancholic depression was shown to be more severe than unspecified depression in a prospective, longitudinal community study [84]. In another study, the lower remission rates of depressives with atypical features were no longer present after adjustment for baseline characteristics [85]. The importance of baseline severity on course trajectories was recently confirmed by the longitudinal study of Lamers et al. [38]. Considering that symptom-based depression subtypes have led to inconsistent results in predicting differential outcomes of antidepressant treatments [36,35,e.g., 85] and that clear evidence concerning psychotherapeutic treatment-distinctions is still lacking, the relevance of subtyping depression with regard to specific depression outcomes thus requires further examination.

Some variables, such as neuroticism, have shown predictive effects but were nevertheless excluded from the present analysis due to combined high correlation with EOC. With regard to the lack of associations with sociodemographic variables, the current analysis supported previous evidence that

challenged the importance of sociodemographic factors in the course of depression [13,12,23]. Finally, childhood-related factors showed surprisingly little impact on improvement in major depression. Presumably, these factors would have been more relevant for a first MDE, but as the age of the current sample was quite high, the impact of childhood-related factors was not strong enough.

### *Limitations*

There were some limitations in this investigation, which should be acknowledged. Potential bias could have resulted from our definition of improvement in major depression. The DIGS did not include enough information to distinguish between remission and recovery as defined in previous studies [86,8]. Moreover, although absence of a current MDE at follow-up is most likely linked to an improvement in mental health, we cannot ensure that this was always subjectively the case. Furthermore, we combined subjects with potentially varying depression courses without considering information on their specific individual states between  $T_0$  and  $T_1$ , such as continuous remission, one new MDE/recurrent MDD. Beyond that, the older age of our sample precluded the opportunity to examine MDD course characteristics at a younger age. On the other hand, considering that the mean age of onset of mood disorders is situated at around 30 years [87], the present study likely captured the relevant time-span. Additionally, we only accounted for diagnosis-cut-off levels, despite some evidence showing that the presence of residual symptoms after a MDE is associated with a higher risk of relapse, recurrence, chronicity, and even suicide [88]. Also, the subjective measure of coping behaviour used in the current study might have entailed several sources of potential bias. First, a balance with regard to objective measures of received social support is warranted [89]. Second, the extent to which individuals can accurately self-report on their EOC styles has been questioned by some authors [90,91]. Third, the relationship between emotion-regulation strategies might be inflated due to item overlap with depression. However, Aldao et al. [61] showed similar patterns of results when using rumination measures with and without depression symptom overlap. Fourth, beyond the components of emotion regulation considered in our article, the entire process of emotion regulation is much more complex and can also include unconscious and physiological (i.e., rapid pulse) cognitive processes [27].



*Conclusions*

Long-term evidence of predictive factors for improvement in individuals affected by MDD with current MDEs is particularly relevant for resource-oriented therapeutic approaches in clinical practice. The predictors found in the current study were primarily located at the behavioural level. Low EOC as a predictive factor underlined the importance of interventions aimed at interrupting the intensive mental preoccupation during MDE by the training of alternative coping strategies. Beyond that, the present study highlighted that differentiation between formal and informal help-seeking behaviour is warranted. These findings therefore raise questions for clinical practice on how to enlarge the social network of individuals suffering from loneliness as a characteristic feature of MDD, and on how to best support informal caregivers.

**Financial support**

The CoLaus|PsyCoLaus study was and is supported by research grants from GlaxoSmithKline, the Faculty of Biology and Medicine of Lausanne, and the Swiss National Science Foundation (grants 3200B0-105993, 3200B0-118308, 33CSCO-122661, 33CS30-139468 and 33CS30-148401).

**Conflict of interest**

None.

## References

1. Gustavsson A, Svensson M, Jacobi F, Allgulander C, Alonso J, Beghi E, Dodel R, Ekman M, Faravelli C, Fratiglioni L, Gannon B, Jones DH, Jenum P, Jordanova A, Jonsson L, Karampampa K, Knapp M, Kobelt G, Kurth T, Lieb R, Linde M, Ljungcrantz C, Maercker A, Melin B, Moscarelli M, Musayev A, Norwood F, Preisig M, Pugliatti M, Rehm J, Salvador-Carulla L, Schlehofer B, Simon R, Steinhausen HC, Stovner LJ, Vallat JM, Van den Bergh P, van Os J, Vos P, Xu W, Wittchen HU, Jonsson B, Olesen J (2011) Cost of disorders of the brain in Europe 2010. *Eur Neuropsychopharmacol* 21 (10):718-779. doi:10.1016/j.euroneuro.2011.08.008
2. Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR, Rush AJ, Walters EE, Wang PS (2003) The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *Jama* 289 (23):3095-3105. doi:10.1001/jama.289.23.3095
3. DeRubeis RJ, Siegle GJ, Hollon SD (2008) Cognitive therapy versus medication for depression: treatment outcomes and neural mechanisms. *Nat Rev Neurosci* 9 (10):788-796. doi:10.1038/nrn2345
4. Keller MB, Shapiro RW, Lavori PW, Wolfe N (1982) Relapse in major depressive disorder: analysis with the life table. *Arch Gen Psychiatry* 39 (8):911-915
5. Kanai T, Takeuchi H, Furukawa TA, Yoshimura R, Imaizumi T, Kitamura T, Takahashi K (2003) Time to recurrence after recovery from major depressive episodes and its predictors. *Psychol Med* 33 (5):839-845
6. Lin EH, Katon WJ, VonKorff M, Russo JE, Simon GE, Bush TM, Rutter CM, Walker EA, Ludman E (1998) Relapse of depression in primary care. Rate and clinical predictors. *Arch Fam Med* 7 (5):443-449
7. Judd LL, Paulus MJ, Schettler PJ, Akiskal HS, Endicott J, Leon AC, Maser JD, Mueller T, Solomon DA, Keller MB (2000) Does incomplete recovery from first lifetime major depressive episode herald a chronic course of illness? *Am J Psychiatry* 157 (9):1501-1504. doi:10.1176/appi.ajp.157.9.1501
8. Richards D (2011) Prevalence and clinical course of depression: a review. *Clin Psychol Rev* 31 (7):1117-1125. doi:10.1016/j.cpr.2011.07.004
9. Weissman MM, Kasl SV (1976) Help-seeking in depressed out-patients following maintenance therapy. *Br J Psychiatry* 129:252-260
10. Angst J (1986) The course of major depression, atypical bipolar disorder, and bipolar disorder. In: H. Hippus, G. L. Klerman & N. Matussek (Eds.), *New results in depression research* (pp. 26–35). Springer-Verlag, Berlin/Heidelberg
11. Mueller TI, Leon AC, Keller MB, Solomon DA, Endicott J, Coryell W, Warshaw M, Maser JD (1999) Recurrence after recovery from major depressive disorder during 15 years of observational follow-up. *Am J Psychiatry* 156 (7):1000-1006. doi:10.1176/ajp.156.7.1000
12. Spijker J, de Graaf R, Bijl RV, Beekman AT, Ormel J, Nolen WA (2004) Determinants of persistence of major depressive episodes in the general population. Results from the Netherlands Mental Health Survey and Incidence Study (NEMESIS). *J Affect Disord* 81 (3):231-240. doi:10.1016/j.jad.2003.08.005
13. Sargeant JK, Bruce ML, Florio LP, Weissman MM (1990) Factors Associated With 1-Year Outcome of Major Depression in the Community. *Arch Gen Psychiatry* 47 (6):519-526
14. Furukawa TA, Kitamura T, Takahashi K (2000) Time to recovery of an inception cohort with hitherto untreated unipolar major depressive episodes. *Br J Psychiatry* 177:331-335
15. De Raedt R, Koster EH (2010) Understanding vulnerability for depression from a cognitive neuroscience perspective: A reappraisal of attentional factors and a new conceptual framework. *Cogn Affect Behav Neurosci* 10 (1):50-70. doi:10.3758/CABN.10.1.50
16. Hoorelbeke K, Marchetti I, De Schryver M, Koster EH (2016) The interplay between cognitive risk and resilience factors in remitted depression: A network analysis. *J Affect Disord* 195:96-104. doi:10.1016/j.jad.2016.02.001
17. Marchetti I, Koster EH, Sonuga-Barke EJ, De Raedt R (2012) The default mode network and recurrent depression: a neurobiological model of cognitive risk factors. *Neuropsychol Rev* 22 (3):229-251. doi:10.1007/s11065-012-9199-9
18. Steinert C, Hofmann M, Kruse J, Leichsenring F (2014) The prospective long-term course of adult depression in general practice and the community. A systematic literature review. *J Affect Disord* 152-154:65-75. doi:10.1016/j.jad.2013.10.017

19. Priebe S, Omer S, Giacco D, Slade M (2014) Resource-oriented therapeutic models in psychiatry: conceptual review. *Br J Psychiatry* 204:256-261. doi:10.1192/bjp.bp.113.135038
20. Harbeck S, Kliem S, Wollburg E, Braukhaus C, Kroger C (2013) [Remission, response, and its prediction in depressive inpatients]. *Psychother Psychosom Med Psychol* 63 (7):272-279. doi:10.1055/s-0032-1327715
21. Sotsky SM, Glass DR, Shea MT, Pilkonis PA, Collins JF, Elkin I, Watkins JT, Imber SD, Leber WR, Moyer J, et al. (1991) Patient predictors of response to psychotherapy and pharmacotherapy: findings in the NIMH Treatment of Depression Collaborative Research Program. *Am J Psychiatry* 148 (8):997-1008. doi:10.1176/ajp.148.8.997
22. DeRubeis RJ, Gelfand LA, Tang TZ, Simons AD (1999) Medications versus cognitive behavior therapy for severely depressed outpatients: mega-analysis of four randomized comparisons. *Am J Psychiatry* 156 (7):1007-1013. doi:10.1176/ajp.156.7.1007
23. Fuller-Thomson E, Battiston M, Gadalla TM, Brennenstuhl S (2014) Bouncing back: remission from depression in a 12-year panel study of a representative Canadian community sample. *Soc Psychiatry Psychiatr Epidemiol* 49 (6):903-910. doi:10.1007/s00127-013-0814-8
24. Keller MB, Klerman GL, Lavori PW, Coryell W, Endicott J, Taylor J (1984) Long-term outcome of episodes of major depression. Clinical and public health significance. *Jama* 252 (6):788-792
25. Conradi HJ, de Jonge P, Ormel J (2008) Prediction of the three-year course of recurrent depression in primary care patients: different risk factors for different outcomes. *J Affect Disord* 105 (1-3):267-271. doi:10.1016/j.jad.2007.04.017
26. Rhebergen D, Lamers F, Spijker J, de Graaf R, Beekman AT, Penninx BW (2012) Course trajectories of unipolar depressive disorders identified by latent class growth analysis. *Psychol Med* 42 (7):1383-1396. doi:10.1017/S0033291711002509
27. Garnefski N, Kraaij V, Spinhoven P (2001) Negative life events, cognitive emotion regulation and emotional problems. *Personality and Individual Differences* 30:1311-1327
28. Ravindran AV, Griffiths J, Waddell C, Anisman H (1995) Stressful life events and coping styles in relation to dysthymia and major depressive disorder: variations associated with alleviation of symptoms following pharmacotherapy. *Prog Neuropsychopharmacol Biol Psychiatry* 19 (4):637-653
29. Endler NS, Parker JDA (1990) State and trait anxiety, depression and coping styles. *Australian Journal of Psychology* 42 (2):207-220
30. O'Brien TB, DeLongis A (1996) The interactional context of problem-, emotion-, and relationship-focused coping: the role of the big five personality factors. *J Pers* 64 (4):775-813
31. Rosenbaum S, Tiedemann A, Sherrington C, Curtis J, Ward PB (2014) Physical activity interventions for people with mental illness: a systematic review and meta-analysis. *J Clin Psychiatry* 75 (9):964-974. doi:10.4088/JCP.13r08765
32. Kaniasty K, Norris FH (2000) Help-seeking comfort and receiving social support: the role of ethnicity and context of need. *Am J Community Psychol* 28 (4):545-581
33. Baumeister H, Parker G (2012) Meta-review of depressive subtyping models. *Journal of affective disorders* 139:126-140
34. Ulbricht CM, Rothschild AJ, Lapane KL (2015) The association between latent depression subtypes and remission after treatment with citalopram: A latent class analysis with distal outcome. *J Affect Disord* 188:270-277. doi:10.1016/j.jad.2015.08.039
35. Arnow BA, Blasey C, Williams LM, Palmer DM, Rekschan W, Schatzberg AF, Etkin A, Kulkarni J, Luther JF, Rush AJ (2015) Depression Subtypes in Predicting Antidepressant Response: A Report From the iSPOT-D Trial. *Am J Psychiatry* 172 (8):743-750. doi:10.1176/appi.ajp.2015.14020181
36. Uher R, Dernovsek MZ, Mors O, Hauser J, Souery D, Zobel A, Maier W, Henigsberg N, Kalember P, Rietschel M, Placentino A, Mendlewicz J, Aitchison KJ, McGuffin P, Farmer A (2011) Melancholic, atypical and anxious depression subtypes and outcome of treatment with escitalopram and nortriptyline. *J Affect Disord* 132 (1-2):112-120. doi:10.1016/j.jad.2011.02.014
37. Gili M, Roca M, Armengol S, Asensio D, Garcia-Campayo J, Parker G (2012) Clinical patterns and treatment outcome in patients with melancholic, atypical and non-melancholic depressions. *PLoS One* 7 (10):e48200. doi:10.1371/journal.pone.0048200
38. Lamers F, Beekman AT, van Hemert AM, Schoevers RA, Penninx BW (2016) Six-year longitudinal course and outcomes of subtypes of depression. *Br J Psychiatry* 208 (1):62-68. doi:10.1192/bjp.bp.114.153098

39. Firmann M, Mayor V, Vidal PM, Bochud M, Pecoud A, Hayoz D, Paccaud F, Preisig M, Song KS, Yuan X, Danoff TM, Stirnadel HA, Waterworth D, Mooser V, Waeber G, Vollenweider P (2008) The CoLaus study: a population-based study to investigate the epidemiology and genetic determinants of cardiovascular risk factors and metabolic syndrome. *BMC Cardiovasc Disord* 8:6. doi:10.1186/1471-2261-8-6
40. Preisig M, Waeber G, Vollenweider P, Bovet P, Rothen S, Vandeleur C, Guex P, Middleton L, Waterworth D, Mooser V, Tozzi F, Muglia P (2009) The PsyCoLaus study: methodology and characteristics of the sample of a population-based survey on psychiatric disorders and their association with genetic and cardiovascular risk factors. *BMC Psychiatry* 9:9. doi:10.1186/1471-244X-9-9
41. Preisig M, Fenton BT, Matthey ML, Berney A, Ferrero F (1999) Diagnostic interview for genetic studies (DIGS): inter-rater and test-retest reliability of the French version. *Eur Arch Psychiatry Clin Neurosci* 249 (4):174-179
42. Nurnberger JI, Jr., Blehar MC, Kaufmann CA, York-Cooler C, Simpson SG, Harkavy-Friedman J, Severe JB, Malaspina D, Reich T (1994) Diagnostic interview for genetic studies. Rationale, unique features, and training. NIMH Genetics Initiative. *Arch Gen Psychiatry* 51 (11):849-859; discussion 863-844
43. APA (2000) *Diagnostic and Statistical Manual of Mental Disorders*. 4th ed, text revision. American Psychiatric Association, Washington, DC
44. Endicott J, Spitzer RL (1978) A diagnostic interview: the schedule for affective disorders and schizophrenia. *Arch Gen Psychiatry* 35 (7):837-844
45. Leboyer M, Barbe B, Gorwood P, Teherani M, Allilaire JF, Preisig M, Matthey ML, Poyetton V, Ferrero F (1995) *Interview Diagnostique pour les Etudes Génétiques Paris*: INSERM.
46. Berney A, Preisig M, Matthey ML, Ferrero F, Fenton BT (2002) Diagnostic interview for genetic studies (DIGS): inter-rater and test-retest reliability of alcohol and drug diagnoses. *Drug Alcohol Depend* 65 (2):149-158
47. Rougemont-Buecking A, Rothen S, Jeanpretre N, Lustenberger Y, Vandeleur CL, Ferrero F, Preisig M (2008) Inter-informant agreement on diagnoses and prevalence estimates of anxiety disorders: direct interview versus family history method. *Psychiatry Res* 157 (1-3):211-223. doi:10.1016/j.psychres.2006.04.022
48. Leboyer M, Maier W, Teherani M, Lichtermann D, D'Amato T, Franke P, Lepine JP, Minges J, McGuffin P (1991) The reliability of the SADS-LA in a family study setting. *Eur Arch Psychiatry Clin Neurosci* 241 (3):165-169
49. APA (2000) *Practice guideline for the treatment of patients with major depressive disorder (revision)*. *Am J Psychiatry* 157:1-45
50. Glaus J, Vandeleur C, Gholam-Rezaee M, Castelao E, Perrin M, Rothen S, Bovet P, Marques-Vidal P, von Kanel R, Merikangas K, Mooser V, Waterworth DM, Waeber G, Vollenweider P, Preisig M (2013) Atypical depression and alcohol misuse are related to the cardiovascular risk in the general population. *Acta Psychiatr Scand* 128 (4):282-293. doi:10.1111/acps.12057
51. Hollingshead AB (1975) *Four factor Index of Social Status*. Yale University Press, New Haven, CT
52. Grob A, Bodmer N, Flammer A (1993) *Living Conditions and the Development of Adolescents in Europe: The Case of Switzerland*. Research Report (Nr. 5). Institute of Psychology. University of Berne, Switzerland
53. Bodmer NM, Grob A (1996) Bien-être et contraintes d'adolescents: une comparaison entre adolescents de familles monoparentales et de familles biparentales. *International Journal of Psychology* 31:39-48
54. Perrin M, Vandeleur CL, Castelao E, Rothen S, Glaus J, Vollenweider P, Preisig M (2014) Determinants of the development of post-traumatic stress disorder, in the general population. *Soc Psychiatry Psychiatr Epidemiol* 49 (3):447-457. doi:10.1007/s00127-013-0762-3
55. Nelder JA, Wedderburn RWM (1972) Generalized Linear Models. *Journal of the Royal Statistical Society Series A (General)* 135 (3):370-384
56. Smith JM, Alloy LB (2009) A roadmap to rumination: a review of the definition, assessment, and conceptualization of this multifaceted construct. *Clin Psychol Rev* 29 (2):116-128. doi:10.1016/j.cpr.2008.10.003

57. Nolen-Hoeksema S, Morrow J (1993) Effects of rumination and distraction on naturally occurring depressed mood. *Cognition and Emotion* 7:561–570
58. Nolen-Hoeksema S (2000) The role of rumination in depressive disorders and mixed anxiety/depressive symptoms. *J Abnorm Psychol* 109 (3):504-511
59. Lyubomirsky S, Nolen-Hoeksema S (1993) Self-perpetuating properties of dysphoric rumination. *J Pers Soc Psychol* 65 (2):339-349
60. Disner SG, Beevers CG, Haigh EA, Beck AT (2011) Neural mechanisms of the cognitive model of depression. *Nat Rev Neurosci* 12 (8):467-477. doi:10.1038/nrn3027
61. Aldao A, Nolen-Hoeksema S, Schweizer S (2010) Emotion-regulation strategies across psychopathology: A meta-analytic review. *Clin Psychol Rev* 30 (2):217-237. doi:10.1016/j.cpr.2009.11.004
62. Roy-Byrne PP, Vitaliano PP, Cowley DS, Luciano G, Zheng Y, Dunner DL (1992) Coping in panic and major depressive disorder. Relative effects of symptom severity and diagnostic comorbidity. *J Nerv Ment Dis* 180 (3):179-183
63. Beck AT (1967) *Depression: Clinical, Experimental, and Theoretical Aspects*. Harper & Row, New York
64. Aldao A, Nolen-Hoeksema S (2012) The influence of context on the implementation of adaptive emotion regulation strategies. *Behav Res Ther* 50 (7-8):493-501. doi:10.1016/j.brat.2012.04.004
65. Roness A, Mykletun A, Dahl AA (2005) Help-seeking behaviour in patients with anxiety disorder and depression. *Acta Psychiatr Scand* 111 (1):51-58. doi:10.1111/j.1600-0447.2004.00433.x
66. Barker C, Pistrang N, Shapiro DA, Shaw I (1990) Coping and help-seeking in the UK adult population. *Br J Clin Psychol* 29 ( Pt 3):271-285
67. Brown J, Evans-Lacko S, Aschan L, Henderson MJ, Hatch SL, Hotopf M (2014) Seeking informal and formal help for mental health problems in the community: a secondary analysis from a psychiatric morbidity survey in South London. *BMC Psychiatry* 14 (1):275. doi:10.1186/PREACCEPT-2067133308130116
68. Nasser EH, Overholser JC (2005) Recovery from major depression: the role of support from family, friends, and spiritual beliefs. *Acta Psychiatr Scand* 111 (2):125-132. doi:10.1111/j.1600-0447.2004.00423.x
69. Clement S, Schauman O, Graham T, Maggioni F, Evans-Lacko S, Bezborodovs N, Morgan C, Rusch N, Brown JS, Thornicroft G (2015) What is the impact of mental health-related stigma on help-seeking? A systematic review of quantitative and qualitative studies. *Psychol Med* 45 (1):11-27. doi:10.1017/S0033291714000129
70. Barney LJ, Griffiths KM, Jorm AF, Christensen H (2006) Stigma about depression and its impact on help-seeking intentions. *Aust N Z J Psychiatry* 40 (1):51-54. doi:10.1111/j.1440-1614.2006.01741.x
71. Griffiths KM, Crisp DA, Barney L, Reid R (2011) Seeking help for depression from family and friends: a qualitative analysis of perceived advantages and disadvantages. *BMC Psychiatry* 11:196. doi:10.1186/1471-244X-11-196
72. van Wijngaarden B, Schene AH, Koeter MW (2004) Family caregiving in depression: impact on caregivers' daily life, distress, and help seeking. *J Affect Disord* 81 (3):211-222. doi:10.1016/S0165-0327(03)00168-X
73. Kirsch I (1990) *Changing expectations: A key to effective psychotherapy*. Brooks/Cole, Pacific Grove, CA
74. Kirsch IE (1999) *How expectancies shape experience*. American Psychological Association, Washington, DC
75. Mischel W, Shoda Y (1994) Personality psychology has two goals: Must it be two fields? . *Psychological Inquiry* (5):156–158
76. Catanzaro SJ, Wasch HH, Kirsch I, Mearns J (2000) Coping-related expectancies and dispositions as prospective predictors of coping responses and symptoms. *J Pers* 68 (4):757-788
77. Blake H (2012) Physical activity and exercise in the treatment of depression. *Front Psychiatry* 3:106. doi:10.3389/fpsy.2012.00106
78. Rethorst CD, Wipfli BM, Landers DM (2009) The antidepressive effects of exercise: a meta-analysis of randomized trials. *Sports Med* 39 (6):491-511. doi:10.2165/00007256-200939060-00004
79. Fox KR (1999) The influence of physical activity on mental well-being. *Public Health Nutr* 2 (3A):411-418

80. Mammen G, Faulkner G (2013) Physical activity and the prevention of depression: a systematic review of prospective studies. *Am J Prev Med* 45 (5):649-657. doi:10.1016/j.amepre.2013.08.001
81. Ortega E (2016) The "bioregulatory effect of exercise" on the innate/inflammatory responses. *J Physiol Biochem*. doi:10.1007/s13105-016-0478-4
82. Balchin R, Linde J, Blackhurst D, Rauch HL, Schonbachler G (2016) Sweating away depression? The impact of intensive exercise on depression. *J Affect Disord* 200:218-221. doi:10.1016/j.jad.2016.04.030
83. Krogh J, Nordentoft M, Sterne JA, Lawlor DA (2011) The effect of exercise in clinically depressed adults: systematic review and meta-analysis of randomized controlled trials. *J Clin Psychiatry* 72 (4):529-538. doi:10.4088/JCP.08r04913blu
84. Angst J, Gamma A, Benazzi F, Ajdacic V, Rossler W (2007) Melancholia and atypical depression in the Zurich study: epidemiology, clinical characteristics, course, comorbidity and personality. *Acta Psychiatr Scand Suppl* (433):72-84. doi:10.1111/j.1600-0447.2007.00965.x
85. Stewart JW, McGrath PJ, Fava M, Wisniewski SR, Zisook S, Cook I, Nierenberg AA, Trivedi MH, Balasubramani GK, Warden D, Lesser I, John Rush A (2010) Do atypical features affect outcome in depressed outpatients treated with citalopram? *Int J Neuropsychopharmacol* 13 (1):15-30. doi:10.1017/S1461145709000182
86. Frank E, Prien RF, Jarrett RB, Keller MB, Kupfer DJ, Lavori PW, Rush AJ, Weissman MM (1991) Conceptualization and rationale for consensus definitions of terms in major depressive disorder. Remission, recovery, relapse, and recurrence. *Arch Gen Psychiatry* 48 (9):851-855
87. Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE (2005) Lifetime prevalence and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 62 (6):593-602. doi:10.1001/archpsyc.62.6.593
88. Dawson MY, Michalak EE, Waraich P, Anderson JE, Lam RW (2004) Is remission of depressive symptoms in primary care a realistic goal? A meta-analysis. *BMC Fam Pract* 5:19. doi:10.1186/1471-2296-5-19
89. Santini ZI, Koyanagi A, Tyrovolas S, Mason C, Haro JM (2015) The association between social relationships and depression: a systematic review. *J Affect Disord* 175:53-65. doi:10.1016/j.jad.2014.12.049
90. Aldao A, Nolen-Hoeksema S (2010) Specificity of cognitive emotion regulation strategies: a transdiagnostic examination. *Behav Res Ther* 48 (10):974-983. doi:10.1016/j.brat.2010.06.002
91. Robinson MD, Clore GL (2002) Episodic and semantic knowledge in emotional self-report: evidence for two judgment processes. *J Pers Soc Psychol* 83 (1):198-215

**Table 1.** Associations between variables at baseline (T<sub>0</sub>) and current MDE (absence vs. presence) at follow-up (T<sub>1</sub>) according to marginal logistic regression models (*n* = 210).

Predictors at T <sub>0</sub>	<i>N</i>	Absence of current MDE at T <sub>1</sub>		
		<i>OR</i>	<i>95% CI</i>	<i>p</i>
<b>Demographics</b>				
<i>Sex</i>				
<i>Men</i>	72	1.32	0.68–2.55	0.41
<i>Women (ref.)</i>	138	--	--	--
<i>Age at baseline, y</i>	210	1.01	0.97–1.05	0.53
<i>Socio-economic status</i> <sup>1</sup>				
<i>Married</i>				
<i>Yes</i>	100	1.50	0.81–2.78	0.20
<i>No (ref.)</i>	110	--	--	--
<i>Living status</i>				
<i>Living with somebody</i>	152	1.16	0.59–2.27	0.66
<i>Living alone (ref.)</i>	58	--	--	--
<b>Childhood-related factors</b>				
<i>General childhood unhappiness</i> <sup>2</sup>	210	0.85	0.59–1.22	0.37
<i>Childhood adversities</i> <sup>3</sup>				
<i>No</i>	134	1.28	0.68–2.39	0.44
<i>Yes (ref.)</i>	76	--	--	--
<i>Childhood trauma</i> <sup>4</sup>				
<i>No trauma &lt;17y</i>	190	1.51	0.57–3.99	0.41
<i>Trauma &lt;17y (ref.)</i>	20	--	--	--
<b>Disorder-related factors</b>				
<i>Age of onset MDD, y</i>	210	1.01	0.99–1.04	0.22
<i>Depression subtype at baseline</i>				
<i>Atypical</i>	53	0.88	0.40–1.96	0.76
<i>Melancholic</i>	46	1.14	0.48–2.72	0.77
<i>Combined atypical-melancholic</i>	32	0.41	0.17–0.97	<b>0.04</b>
<i>Unspecified (ref.)</i>	79	--	--	--
<i>Number of depression symptoms during the current MDE</i> <sup>5</sup> (max. 9)	210	0.73	0.56–0.95	<b>0.02</b>



<i>Duration of all MDEs (including current episode</i> <sup>5</sup> )	210	1.00	1.00–1.00	0.60
<i>Duration of the current MDE</i> <sup>5</sup>	209	1.00	1.00–1.00	0.69
<i>Number of MDEs (including current episode</i> <sup>5</sup> )	210	1.01	0.95–1.08	0.67
<i>Current GAF score of all disorders at baseline</i>	208	1.05	1.02–1.08	<0.001
<i>Help from a doctor or other healthcare professional</i>				
Yes	160	0.53	0.24–1.17	0.11
No (ref.)	49	--	--	--
<i>Psychopharmacological treatment (antidepressants)</i>				
Yes	5	0.24	0.04–1.47	0.12
No (ref.)	205	--	--	--
<i>Anxiety disorder at baseline</i> <sup>6</sup>				
No	165	1.32	0.64–2.73	0.45
Yes (ref.)	44	--	--	--
<i>Drug abuse at baseline</i>				
No	203	1.84	0.30–11.31	0.51
Yes (ref.)	5	--	--	--
<i>Number of comorbid diagnoses</i> <sup>7</sup> (lifetime)	210	1.01	0.69–1.47	0.96

#### **Behavioural characteristics**

##### *Regular smoking*<sup>8</sup>

Never	91	1.46	0.66–3.24	0.35
Former	71	0.86	0.39–1.91	0.71
Current (ref.)	48	--	--	--

##### *Regular alcohol consumption*<sup>9</sup>

No	74	0.75	0.40–1.42	0.38
Yes (ref.)	130	--	--	--

##### *Regular physical activity*<sup>10</sup>

Yes	67	1.63	0.82–3.24	0.17
No (ref.)	143	--	--	--

#### **Personality characteristics**

<b><i>Coping</i></b>				
<i>Emotion-oriented coping</i>	144	0.82	0.72–0.92	<b>&lt;0.001</b>
<i>Informal help-seeking behaviour</i>	144	1.13	0.95–1.33	0.17
<i>Problem-oriented coping</i>	144	1.10	0.90–1.35	0.33
<b>Somatic characteristic</b>				
<b><i>Body Mass Index</i></b>	210	0.98	0.92–1.04	0.48

---

Bold:  $p \leq 0.05$

MDD: Major Depressive Disorder; MDE: major depressive episode; OR: odd ratio, CI; confidence interval, GAF: global assessment of functioning

<sup>1</sup> Socio-economic status following Hollingshead's index: A value of 3 represents an SES of middle class.

<sup>2</sup> Rated on a Likert Scale ranging from 1 (happy) to 4 (very unhappy)

<sup>3</sup> Parental quarrels, fear of parental punishment, foster home

<sup>4</sup> Any trauma

<sup>5</sup> Current episode at the baseline investigation

<sup>6</sup> Agoraphobia, generalized anxiety disorder, panic disorder, social phobia

<sup>7</sup> Alcohol abuse/dependence, drug abuse/dependence, agoraphobia, generalized anxiety disorder, panic disorder, social phobia

<sup>8</sup> At least 5 days of smoking per week

<sup>9</sup> At least once a week for at least 6 months

<sup>10</sup> At least 2-3 times per week

**Table 2.** Associations between variables at baseline (T<sub>0</sub>) and current MDE (absence vs. presence) at follow-up (T<sub>1</sub>) according to logistic regression models with simultaneous introduction of predictor variables (*n* = 143).

Predictors at T <sub>0</sub>	Absence of current MDE at T <sub>1</sub>		
	OR*	95% CI	<i>p</i>
<b><i>Coping</i></b>			
<i>Emotion-oriented coping</i>	0.77	0.66–0.91	<b>0.001</b>
<i>Informal help-seeking behaviour</i>	1.38	1.10–1.74	<b>0.01</b>
<b><i>Depression subtype</i></b>			
<i>Atypical</i>	1.19	0.37–3.84	0.77
<i>Melancholic</i>	1.60	0.45–5.62	0.46
<i>Combined atypical-melancholic</i>	1.32	0.30–5.85	0.72
<i>Unspecified (ref.)</i>	--	--	--
<i>Number of depression symptoms during the current MDE (max. 9)</i>	0.77	0.51–1.17	0.22
<i>Current GAF score of all disorders</i>	1.04	0.99–1.09	0.11
<i>Help from a doctor or other healthcare professional</i>	0.73	0.21–2.56	0.62
<i>Psychopharmacological treatment (antidepressants)</i>	0.10	0.01–1.69	0.11
<i>Regular physical activity</i>	2.48	0.94–6.57	0.07

\* Adjusted for sex, age at baseline and socio-economic status

Bold:  $p \leq 0.05$

MDE; major depressive episode; OR: odd ratio, CI; confidence interval; GAF: global assessment of functioning