



Strathprints Institutional Repository

Parkes, E.J. (2009) *A note on travelling-wave solutions to Lax's seventh-order KdV equation*. Applied Mathematics and Computation, 215 (2). pp. 864-865. ISSN 0096-3003

Strathprints is designed to allow users to access the research output of the University of Strathclyde. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. You may not engage in further distribution of the material for any profitmaking activities or any commercial gain. You may freely distribute both the url (<http://strathprints.strath.ac.uk/>) and the content of this paper for research or study, educational, or not-for-profit purposes without prior permission or charge.

Any correspondence concerning this service should be sent to Strathprints administrator: <mailto:strathprints@strath.ac.uk>

A note on travelling-wave solutions to Lax's seventh-order KdV equation

E.J. Parkes^a

^a*Department of Mathematics, University of Strathclyde, Glasgow G1 1XH, UK*

Abstract

Ganji and Abdollahzadeh [D.D. Ganji, M. Abdollahzadeh, Appl. Math. Comput. 206 (2008) 438–444] derived three supposedly new travelling-wave solutions to Lax's seventh-order KdV equation. Each solution was obtained by a different method. It is shown that any two of the solutions may be obtained trivially from the remaining solution. Furthermore it is noted that one of the solutions has been known for many years.

Key words: Travelling-wave solutions. Lax's seventh-order KdV equation.

PACS: 03.40.Kf.

Over the past two decades several expansion methods for finding travelling-wave solutions to nonlinear evolution equations have been proposed, developed and extended. The solutions to dozens of equations have been found by one or other of these methods. Reference [1] and references therein mention some of this activity. Unfortunately, many authors claim that their solutions are 'new', when the truth is that these solutions are merely 'old' solutions in a different guise. Recently, in a series of enlightening papers [2–4], Kudryashov has warned researchers and referees of the danger of not recognizing that apparently different solutions may simply be different forms of the same solution. He has provided numerous examples to illustrate this phenomenon. In the present note we point out one more example.

Recently, Ganji and Abdollahzadeh [1] presented three travelling-wave solutions to Lax's seventh-order KdV equation

$$u_t + (35u^4 + 70(u^2u_{xx} + uu_x^2) + 7(2uu_{xxx} + 3u_{xx}^2 + 4u_xu_{xxx}) + u_{xxxxx})_x = 0. \quad (1)$$

Each solution was derived by a different method. We wish to point out that the second and third of these solutions may be derived trivially from the first solution, and that the latter has been known for at least a dozen years.

Email address: `ejp@maths.strath.ac.uk` (E.J. Parkes).

The first solution to (1) is given by Eq. (4.5) in [1], namely

$$u = a_0 + 2\alpha^2 \operatorname{sech}^2 \xi, \quad \xi = \alpha(x - \beta t), \quad \beta = 64\alpha^6 + 224\alpha^4 a_0 + 280\alpha^2 a_0^2 + 140a_0^3, \quad (2)$$

where a_0 and α are arbitrary. It was derived by using the sech-function expansion method.

The authors formulated the sec-function expansion method, applied it to (1), and obtained another solution given by (4.8) in [1], namely

$$u = a_0 - 2\alpha^2 \sec^2 \xi, \quad \xi = \alpha(x - \beta t), \quad \beta = -64\alpha^6 + 224\alpha^4 a_0 - 280\alpha^2 a_0^2 + 140a_0^3, \quad (3)$$

where a_0 and α are arbitrary.

Finally, they used the ‘rational function in exp’ method to get (4.12) in [1], namely

$$u = a_0 + \frac{2\alpha^2}{(1 + e^\xi)} - \frac{2\alpha^2}{(1 + e^\xi)^2}, \quad \xi = \alpha(x - \beta t), \quad \beta = \alpha^6 + 14\alpha^4 a_0 + 70\alpha^2 a_0^2 + 140a_0^3. \quad (4)$$

Firstly, we note that (3) may be obtained from (2) by using the transformation $\alpha \rightarrow i\alpha$ and the identity $\operatorname{sech}(iX) = \sec(X)$. Thus the effort involved in applying the sec-function method, as described in [1], is superfluous.

Secondly, we observe that the identity

$$\tanh X = 1 - \frac{2}{(1 + e^{2X})} \quad (5)$$

implies that any solution expressed as a series in $\tanh X$ may be written as a series in inverse powers of $1 + e^{2X}$. In particular, (4) may be obtained from (2) by using the transformation $\alpha \rightarrow \alpha/2$ and the identity (5).

Written in terms of the tanh function, (2) becomes

$$u = b_0 - 2\alpha^2 \tanh^2 \xi, \quad \xi = \alpha(x - \beta t), \quad \beta = -4(96\alpha^6 - 196\alpha^4 b_0 + 140\alpha^2 b_0^2 - 35b_0^3), \quad (6)$$

where b_0 and α are arbitrary. This solution has been known for at least a dozen years. For example it is given by Eq. (6.4) in [5] where it was derived by using the Automated Tanh-Function Method (ATFM), a Mathematica package designed to take the drudgery out of applying the tanh-function expansion method by hand. The particular case given by putting $a_0 = 0$ in (2) or, equivalently, $b_0 = 2\alpha^2$ in (6), namely

$$u = 2\alpha^2 \operatorname{sech}^2[\alpha(x - 64\alpha^6 t)] \quad (7)$$

is mentioned in [1] and [5]. In the latter, it is noted that (7) agrees with the solution given over twenty years ago in Table 3 in [6] with appropriate choices of parameter values.

References

- [1] D.D. Ganji, M. Abdollahzadeh, Exact travelling solutions for the Lax's seventh-order KdV equation by sech method and rational exp-function method, *Appl. Math. Comput.* 206 (2008) 438–444.
- [2] N.A. Kudryashov, N.B. Loguinova, Be careful with the Exp-function method, *Commun. Nonlinear Sci. Numer. Simulat.* 14 (2009) 1881–1890.
- [3] N.A. Kudryashov, On “new travelling wave solutions” of the KdV and KdV-Burgers equations, *Commun. Nonlinear Sci. Numer. Simulat.* 14 (2009) 1891–1900.
- [4] N.A. Kudryashov, Seven common errors in finding exact solutions of nonlinear differential equations, *Commun. Nonlinear Sci. Numer. Simulat.* 14 (2009) 3507–3529.
- [5] E.J. Parkes, B.R. Duffy, An automated tanh-function method for finding solitary wave solutions to nonlinear evolution equations, *Comput. Phys. Commun.* 98 (1996) 288–300.
- [6] W. Hereman, P.P. Banerjee, A. Korpel, G. Assanto, A. Van Immerzeele, A. Meerpoel, Exact solitary wave solutions of non-linear evolution and wave equations using a direct algebraic method, *J. Phys. A: Math. Gen.* 19 (1986) 607–628.