

Mémoire de Maîtrise en médecine No 2482

Unscheduled consultations: a cross-sectional study of patients using walk-in emergency clinics

Etudiante

Marie Vetterli

Tuteur

Dr Philippe Staeger

Centre de médecine générale et urgences, PMU

Co-tuteur

Dr Olivier Hugli

Service des urgences, CHUV

Expert

Pr Olivier Rutschmann

Service des urgences, HUG

Lausanne, janvier 2016

Swiss Medical Weekly

Unscheduled consultations: a cross-sectional study of patients using walk-in emergency clinics --Manuscript Draft--

Manuscript Number:	
Article Type:	Original article
Full Title:	Unscheduled consultations: a cross-sectional study of patients using walk-in emergency clinics
Corresponding Author:	Philippe Staeger, MD, MPH Policlinique médicale universitaire Lausanne, SWITZERLAND
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Policlinique médicale universitaire
Corresponding Author E-Mail:	Philippe.Staeger@hospvd.ch
Manuscript Region of Origin:	SWITZERLAND
Order of Authors:	Marie Vetterli Olivier Hugli, MD, MPH Philippe Staeger, MD, MPH
Order of Authors Secondary Information:	
Author Comments:	
Abstract:	<p>Questions under study/principles Switzerland experiences a strong increase of the unscheduled medical consultations which participates to the congestion of the hospital emergency departments. In this context, many walk-in emergency clinics have been established but less is known about the characteristics of the patients who visit these structures.</p> <p>Methods First, retrospective data about frequentation between 2011 and 2014 of three walk-in emergency clinics in Lausanne were analysed. Secondly, a questionnaire about sociodemographic data, access to care, patient's usual health status, and their global resources to solve their health problem was submitted during one week in the waiting room of each clinic from 1-20 September 2014, to patients aged 16 or older.</p> <p>Results The overall number of consultations increased globally by 6.9%, whereas Lausanne's population only increased by 2.9%. 305 (87%) patients were included for the questionnaire. The mean age of participants was 40.6 years old, 50% were women and 65% were Swiss. 76% of patients had a primary care physician (PCP), 38.7% of them said they had try to contact him in the last 24h for their problem. Among them, 81% did not get an appointment on the same day.</p> <p>Conclusions Our study shows that many patients suffering from a non-life-threatening health problem use walk-in emergency clinics as their PCP. Walk-in emergency clinics seem to respond to patient's needs and to the change in the way that care is consumed.</p>
Keywords:	Walk-in; unscheduled care; ambulatory care; primary care physician; questionnaire; Switzerland
Additional Information:	
Question	Response
I confirm that neither this manuscript, nor	Yes

any other with substantially similar content by one or more of the same authors, has been published, accepted or is currently being assessed by another journal with a view to publication.	
I certify that all authors have participated sufficiently (1) in the conception and design, or acquisition of data, or analysis and interpretation of data; (2) drafting the article or revising it critically for important intellectual content; and (3) final approval of the version to be published.	Yes
Please specify the study design (randomized controlled trial / cohort study / case-control study / cross-sectional study / economic evaluation / diagnostic test study)	cross-sectional study
Trial registration number: We encourage the registration of clinical trials in a primary registry that participates in WHO's International Clinical Trial Registry Platform. The ICMJE defines a clinical trial as any research project that prospectively assigns human subjects to intervention or concurrent comparison or control groups to study the cause-and-effect relationship between a medical intervention and a health outcome. Medical interventions include drugs, surgical procedures, devices, behavioral treatments, process-of-care changes, and the like. Please include the trial registration number and the name of the trial registry at the end of the abstract too.	None
We require every article reporting results of prospective research using human subjects or samples or results of animal research to include a statement that the study obtained ethics approval, including the name of the ethics committee(s) or institutional review board(s) and the number/ID of the approval(s). Where ethical approval is not required, the manuscript should include a clear statement of this and the reason why.	Name of the ethics committee(s) or institutional review board(s): Human Research Ethics Committee of the Canton Vaud Number/ID of the approval(s): Our study was approved by the ethics committee even though it has ruled that our study did not fall under the definition of article 2 of the "Human Research Act" since it was designed to be conducted anonymously. (see attached file)
Have you obtained consent from the patient if there is an unavoidable risk of breach of privacy? (Please note that the manuscript will not be reviewed until this consent is received.)	No - - Informed consent is not necessary for this paper.
Have figures or tables from other publications been used? (Permission to reprint figures or tables from other publications must be obtained by the author prior to submission of the manuscript. A copy of the permission from the copyright holder has to be sent to the editorial office.)	No - No copyrighted material has been used.
Please disclose outside financial support or other financial relationships (both	None

<p>personal and institutional) that could be viewed as presenting a potential conflict of interest in connection with the submitted manuscript. A conflict of interest statement is published with each paper. (We do not need to receive signed copies of the authors' forms now, but if your manuscript is accepted for publication the editorial office will ask you to return the authors' forms signed by each author.)</p>	
<p>Do you apply for rapid publication? (If you feel that for reasons of general or public interest your manuscript requires more expeditious publication and quick processing through peer review, you may inform the Editorial Board indicating: the reasons the authors think the manuscript should be published immediately and the implications of the findings.)</p>	<p>No</p>
<p>Suggested Reviewers:</p>	<p>Aris Exadaktylos University Hospital Bern</p> <ul style="list-style-type: none"> - He has a strong interest for the theme of our article -> ha has participated to many articles on the same subject -> we cited some - He knows the Swiss health system

Unscheduled consultations: a cross-sectional study of patients using walk-in emergency clinics

M. Vetterli ¹, O. Hugli ², P. Staeger ³

¹ Faculty of Biology and Medicine, Lausanne University Hospital, Switzerland

² Emergency Department, Lausanne University Hospital, Switzerland

³ Department of Ambulatory Care and Community Medicine, Lausanne University Hospital, Switzerland

Date of draft: 19.01.2016

Hospital or institute: Lausanne University Hospital

Funding/potential competing interests: none

Correspondence:

Philippe Staeger, MD, MPH

Policlinique médicale universitaire, Bugnon 44, CH-1011 Lausanne, Switzerland.

Philippe.Staeger@hospvd.ch, 021/314'60'60

ABSTRACT

Questions under study/principles

Switzerland experiences a strong increase of the unscheduled medical consultations which participates to the congestion of the hospital emergency departments. In this context, many walk-in emergency clinics have been established but less is known about the characteristics of the patients who visit these structures.

Methods

First, retrospective data about frequentation between 2011 and 2014 of three walk-in emergency clinics in Lausanne were analysed. Secondly, a questionnaire about sociodemographic data, access to care, patient's usual health status, and their global resources to solve their health problem was submitted during one week in the waiting room of each clinic from 1-20 September 2014, to patients aged 16 or older.

Results

The overall number of consultations increased globally by 6.9%, whereas Lausanne's population only increased by 2.9%. 305 (87%) patients were included for the questionnaire. The mean age of participants was 40.6 years old, 50% were women and 65% were Swiss. 76% of patients had a primary care physician (PCP), 38.7% of them said they had try to contact him in the last 24h for their problem. Among them, 81% did not get an appointment on the same day.

Conclusions

Our study shows that many patients suffering from a non-life-threatening health problem use walk-in emergency clinics as their PCP. Walk-in emergency clinics seem to respond to patient's needs and to the change in the way that care is consumed.

Key words

Walk-in - unscheduled care - ambulatory care - primary care physician - questionnaire - Switzerland

INTRODUCTION

1 In Switzerland, as in numerous Western countries over the last few years, the healthcare
2 system has experienced a strong increase in the number of unscheduled medical
3 consultations [1,2]. So-called *walk-in patients* go directly to emergency units without a prior
4 medical opinion or referral [3], contributing to the constant increase in cases dealt with by
5 hospital emergency departments (EDs) and their almost constant state of congestion [4].
6 According to a recent study in Switzerland, the overall number of cases dealt with by EDs
7 increased by 26% between 2007 and 2011, with increases of 16% and 32% in the number
8 of consultations that resulted in hospitalisation or an outpatient consultation, respectively [5].

9
10
11
12
13
14
15
16
17
18
19
20
21 Other reasons contributing to this increase in hospital ED patients include a growing and
22 aging population [6,7], more numerous cases of chronic diseases [8], as well as a shortage
23 of primary care physicians (PCPs) and the difficulties they face when dealing with patients in
24 an emergency [9].

25
26
27
28
29
30
31
32 One means of reducing the reliance on hospital EDs is better access to primary care [10].
33 Walk-in primary care clinics thus represent an interesting alternative solution for healthcare
34 for walk-in patients. Indeed, these clinics are recognised for their accessibility, longer
35 opening hours and the possibility to consult a physician without an appointment [11,12]. At
36 the beginning of the 2000s, the United Kingdom's National Health Service attempted to
37 respond to overcrowded hospital EDs [11,13] by developing walk-in clinics based on models
38 used in the USA and in Canada for forty years [14,15]. In Switzerland, where personal
39 health insurance is obligatory and the patient is free to choose a healthcare provider, the
40 past few years have seen the establishment of many walk-in emergency clinics.

41
42
43
44
45
46
47
48
49
50
51
52
53
54 Numerous studies have investigated the characteristics of patients with non-life threatening
55 conditions who resort to hospital EDs and why they do so [3,16–19]. However, there are
56 currently few data in Switzerland about patients who choose to use urban walk-in
57
58
59
60
61
62
63
64
65

1 emergency clinics. The present study aims to identify why patients consult in three of
2 Lausanne's walk-in emergency clinics and to gather sociodemographic data and information
3 on their global resources.
4
5
6
7

8 **METHODOLOGY**

9

10 **Setting**

11 This study took place in 2014 in Lausanne, a city with more than 140,000 habitants, four
12 private hospital clinics and a university hospital centre. Three walk-in clinics situated in
13 different neighbourhoods in the city were examined for this study. Two are private and were
14 opened in 1992 and 1999; the third is public and was set up in 2010. Consultations in these
15 three clinics are delivered by physicians and they have similar healthcare cost
16 reimbursement systems with insurance companies.
17
18
19
20
21
22
23
24
25
26
27

28 **Study design**

29 This cross-sectional study was made up of two parts. The first part studied the changing
30 frequentation of the three walk-in emergency clinics between 2011 and 2014 by analysing
31 retrospective data on the number of consultations and how these were spread across the
32 seven days of the week.
33
34
35
36
37
38
39
40

41 The second part consisted of a survey carried out using a questionnaire filled in by patients
42 in the clinic waiting rooms and a review of their medical file after the consultation. The
43 survey took place over three consecutive weeks, from September 1st to 20th 2014, with one
44 week spent in each walk-in clinic, from Monday to Saturday. Survey periods lasted for six
45 hours, varying day-to-day, but always within the usual opening hours for medical practices
46 (08h00-18h30). Data by walk-in clinic were collected in matching periods according to the
47 day. The survey period included no public or school holidays.
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Participants

1 All patients aged 16 years old or above in the waiting rooms of walk-in emergency clinics
2
3 were asked to participate in the study and answer the questionnaire. They were all awaiting
4
5 a medical consultation for non-life-threatening conditions, as defined by levels 3 and 4 of the
6
7 Swiss Emergency Triage Scale [16]. The investigator did not participate in their treatment.
8
9
10 Potential participants were excluded from the study if they were fluent in neither French nor
11
12 English, if there was insufficient time in which to fill in the questionnaire, if they refused to
13
14 take part or if they were incapable of giving informed consent (Figure 1). Participants
15
16 completed the questionnaire themselves, and the investigator was on hand to assist them if
17
18 necessary.
19
20
21
22

Questionnaire

23
24
25 The questionnaire was based on one developed by a British research team and was used
26
27 with the authors' consent [17]. It was expanded with elements taken from other
28
29 questionnaires used specifically in research into Lausanne's ED and a walk-in emergency
30
31 clinic [16,20] (see Annex 1). It was divided into five sections: i) access to care; ii) patient's
32
33 usual health status; iii) reason for coming for a consultation; iv) the walk-in emergency clinic
34
35 and patients' expectations; and v) sociodemographic data.
36
37
38
39
40

41 The investigator extracted the reasons for the consultation, the diagnoses proposed and the
42
43 subsequent treatments prescribed from the patients' electronic medical records (see Annex
44
45 2).
46
47
48

Statistical analysis

49
50
51 The results from the three walk-in emergency clinics were pooled. Categorical data are
52
53 described as percentages and continuous data are presented as averages with their
54
55 standard deviation (SD). Comparisons between groups were made using the unpaired
56
57 Student t-test for quantitative variables and the chi-squared test or Fisher's exact test for
58
59
60
61
62
63
64
65

1 categorical data. The trend over time was evaluated using linear regression. A value of
2 P < 0.05 was considered to show a significant difference. All these statistical analyses were
3 made using Stata 13.1 (StataCorp LP, College Station, TX, USA).
4
5

6 **Ethical approval**

7 This study was approved by the Human Research Ethics Committee of the Canton Vaud.
8
9 Data were collected in an anonymous format and no data was treated individually.
10
11
12
13
14
15

16 **RESULTS**

17 **Trends over time**

18 The analysis of retrospective data shows that the overall number of consultations and the
19 number of consultations in walk-in clinics II and III did not increase significantly during the
20 period from 2011 to 2014 (P = 0.17). In walk-in clinic I, which opened in 2010 and was the
21 newest, consultations had increased by 44% (P = 0.04). The number of patients consulting
22 in walk-in clinic II actually decreased in 2014 (Figure 2).
23
24
25
26
27
28
29
30
31

32 The two busiest days in all three walk-in emergency clinics were Mondays (a mean of 65.1
33 patients between 2011–2014) and Thursdays (58.4 patients). The two weekend days were
34 the least busy, with a mean of 49.2 consultations on Saturdays and 44.9 on Sundays
35 (Figure 3).
36
37
38
39
40
41
42
43
44

45 **Questionnaire**

46 During the survey period, 374 potential participants were identified, 352 were eligible and
47 305 (87%) were included (Figure 1).
48
49
50
51
52

53 The mean age of participants was 40.6 years old (SD 17.9), with an equal split between
54 men and women (Table 1). On average, three quarters of patients had a PCP. These
55 patients were 11 years older than those who did not and were more likely to be women.
56
57
58
59
60
61
62
63
64
65

1 However, this correlation with age was not linear: 73% of patients aged 16–24 years old had
2 a PCP (N = 45), decreasing to 58% for those 25–34 years old (N = 42), and rising to 100%
3 for patients 55–64 years old (N = 23) and participants over 75 (N = 19) (P < 0.001).
4

5 With regards to patients' origins, 65% were Swiss, 27% were from European countries and
6 8.1% were from outside Europe. Among Swiss patients, 82% had a PCP, against 62% of
7 foreign patients (P < 0.001). The longer foreign patients had lived in Switzerland, the more
8 likely they were to have a PCP. Almost all of the foreign patients had a Swiss residence or
9 work permit, and only a few were students or tourists. The majority of participants were
10 employed and almost half of them were educated to secondary school leaving diploma level
11 or above. The proportion of patients with a university or technical college level of education
12 was greater among those who did not have a PCP.
13
14
15
16
17
18
19
20
21
22
23
24

25 The great majority of participants judged their usual health status to be good or very good
26 (83%) and only 1% judged it to be poor or very poor. Half of the patients with a PCP
27 indicated that they had only seen him once or not at all in the previous 12 months.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

66 More than one in three patients with a PCP stated that they had tried to contact him about
67 their health problem in the last 24 hours. In more than half of these cases, the physician's
68 practice was closed and only a quarter of the patients managed to arrange an appointment,
69 although not for the same day – the median delay for an appointment was three days. Only
70 one patient in twenty actually managed to speak to their PCP on the telephone, and 4.4%
71 had an appointment with him before going to the walk-in emergency clinic.

72 On average, 79% of participants had attempted a treatment before coming to the walk-in
73 emergency clinic. Nearly half tried to rest or took non-prescribed medicines as a first
74 solution, followed by one in five who used a home remedy. One in ten participants had taken

1 a medicine prescribed by a physician and the same proportion had tried a treatment based
2 on complementary medicine (Table 3).

3 The main sources of advice to participants before their consultation at a walk-in emergency
4 clinic were family members or partners, followed by friends and work colleagues. Nearly one
5 in five patients had visited a pharmacy before coming to the emergency clinic, and one in
6 ten joined a physician's practice. Twenty-two (7.7%) participants had checked the Internet
7 for information, and only seven (2.4%) patients had tried to telephone a medical call centre.

8 Had the walk-in emergency clinic not be available, the majority of the patients stated that
9 they would have gone to another emergency clinic or ED, and only a quarter of those with a
10 PCP would have consulted him.
11
12
13
14
15
16
17
18
19
20
21
22
23
24

25 **DISCUSSION**

26
27 An analysis of the statistics from the three walk-in emergency clinics showed an overall
28 increase in the number of consultations of 6.9% between 2011 and 2014, although
29 Lausanne's population only increased by 2.9% during that period [21]. The detailed results
30 by clinic showed that the number of consultations at the oldest and most peripheral clinic
31 decreased by 8%, but they increased by 44% at the newest, most central clinic. Certain
32 elements may help to explain this variation. Clinic II was under renovation between 2013
33 and 2014, and there was a lack of readily available parking; it is also more difficult to reach
34 by public transport. This indicates that proximity and ease of access probably play a
35 significant role when choosing where to go for an unscheduled emergency consultation,
36 whether at a walk-in emergency clinic or an ED [22]. The newest walk-in emergency clinic
37 probably benefitted from its newness and became better known each year.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53

54 The slight decrease in the overall growth curve for consultations in 2014 might be explained
55 in part by the increased number of walk-in emergency clinics in the Lausanne area since the
56 start of that year.
57
58
59
60
61
62
63
64
65

1 The increase in consultations between 2011 and 2014 followed the trend seen in most EDs,
2 although at a lower level. Lausanne university hospital's ED registered an increase in
3 consultations of 33% between 2005 and 2013, including a 34% increase in consultations for
4 non-life-threatening conditions [16].
5
6

7
8 In all three walk-in emergency clinics, from 2011 to 2014, the mean number of consultations
9 was greater on Mondays and Thursdays. Monday is traditionally the busiest day for EDs
10 [23], probably because patients find it impossible to contact their PCP over the weekend. In
11 Switzerland, many medical practices are traditionally closed on Thursdays, further limiting
12 access to care.
13
14
15
16
17
18
19
20
21
22

23 The present study provides new data about the profiles of patients attending walk-in
24 emergency clinics. The population that participated in the study was young, employed,
25 predominantly indigenous, with a high level of education and perceived its health status to
26 be good. This corresponds to other observational studies in Europe [13,20] and Canada
27 [15]. Men and women were equally represented in the sample, and the indigenous Swiss
28 population was over-represented in the sample with respect to Lausanne's general
29 population [21]. However, the present sample's sociodemographic data showed that patient
30 profiles compared quite well to those of a recent study of patients with non-life-threatening
31 conditions attending Lausanne's university hospital ED [16], except for a few differences; the
32 population attending the ED was slightly older (mean of 44.5 years old vs 40.6), had a lower
33 rate of employment (51.9% vs 64%) and had a higher proportion of foreigners (43.1% vs
34 34.9%). The greater proportion of non-Europeans (17.3% vs 8.1%) treated in the ED may
35 indicate that they resort to this kind healthcare structure because they lack knowledge about
36 existing alternative walk-in emergency clinics that can deal with non-life-threatening
37 conditions. They may also have different habits when it comes to consuming healthcare
38 [24].
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 The proportion of patients who indicated that they had a PCP was high, despite the fact that
2 *gatekeeping* is not obligatory in Switzerland [1]. This result was comparable to that of the
3 Lausanne university hospital ED study [16] and very close to the British research results that
4 were the basis for the present questionnaire [17]. A similar proportion has been found in
5 another Swiss study [3]. The 38.7% rate of contact with a PCP before the emergency
6 consultation was also close to the British research results [17]. Three quarters of the patient
7 sample would have gone to another walk-in emergency clinic rather than to their PCP, had
8 the initial clinic been unavailable. This shows their determination to consult a physician
9 rapidly. Nevertheless, PCPs in local practices remain essential actors in healthcare
10 networks in cases of non-life-threatening emergencies; they could offer an alternative to
11 EDs and walk-in emergency clinics. However, the literature reveals that unscheduled
12 consultations have moved away from PCP practices to walk-in emergency structures
13 [25,26]. The present results show that, according to patients, general practitioners are not
14 available enough for unscheduled appointments; 83% of patients' who called their PCP's
15 practice for an appointment on the same day did not get one (the practice was either closed
16 or an appointment could only be given for some days in the future). However, physicians
17 themselves do not agree with their patients' perceptions of difficult access – 62% of Swiss
18 physicians estimate that > 80% of their patients can indeed consult them on the same day
19 or the next day in an emergency. Some 78% of them have extended hours of consultation
20 [27]. However, it could be that the number of available emergency appointments is
21 insufficient simply due to the shortage of PCPs [9,28]. The question about whether patients
22 overuse walk-in emergency clinics or whether their needs are not being met by PCPs
23 remains unanswered in Switzerland. The issue of timely access to emergency care is
24 central to patients' expectations about their PCP [29]. A very recent report on the issue of
25 access to emergency care [30], prepared by the Institute of Medicine in the USA, suggests
26 in particular the use of approaches such as same-day scheduling [31] or more futuristic
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 ideas, such as virtual "visits" [32]. These approaches should help to reduce the chronic
2 overcrowding in hospital EDs and ensure continuity of care [14,31]. The present study also
3 showed that medical call centres are currently a little-used solution, rarely considered by the
4 patients who attend walk-in clinics, whether for advice or as an alternative when a clinic is
5 not available. These results are consistent with those of a Canadian study [33].
6
7
8
9

10
11
12 The present study has some limitations. Although the three walk-in emergency clinics
13 chosen were among the largest in Lausanne and were spread across different
14 neighbourhoods, their small number limited the conclusions that could be drawn. Although
15 the questionnaire used was based on an existing one in English, it has not yet been
16 validated. For reasons of study feasibility, the investigator could only spend one week in
17 each clinic and only for a limited number of hours per day. The study sample is relatively
18 small, therefore, and a little restricted in terms of internal validity. A certain number of
19 patients were unable to answer the questionnaire because of language difficulties, and this
20 may have caused a selection bias. However, 11 out of 352 eligible patients was only a small
21 part of the sample. Despite an excellent rate of participation, there were some missing data
22 and incomplete questionnaires. Finally, the data from Saturdays may have biased the
23 results as only a small number of PCPs work on weekends – responses may be a little
24 different from other days.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46 **CONCLUSION**

47
48 This study shows that many patients suffering from a non-life threatening health problem
49 use walk-in emergency clinics as their PCP. These clinics seem to respond to patients'
50 needs and to the change in the way care is consumed. Many patients contacted their PCP
51 before going to the walk-in clinic, but without being able to get an appointment on the same
52
53
54
55
56
57
58
59
60
61
62
63
64
65

day. This finding should motivate PCP to consolidate their prominent roles as indispensable parts of the primary emergency care system.

Acknowledgments

We would like to thank Dr M Eddé and the "Centres d'urgences médico-chirurgicales de Vidy et de La Source", Dr J Perdrix and the "Permanence PMU-Flon", Pr. O Rutschmann, Service des urgences, Hôpitaux Universitaires de Genève, for their precious contribution to this study, and Dr S Gnani et al., Imperial College London, for the permission to use their questionnaire.

REFERENCES

1. Chmiel C, Huber CA, Rosemann T, Zoller M, Eichler K, Sidler P, et al. Walk-ins seeking treatment at an emergency department or general practitioner out-of-hours service: a cross-sectional comparison. *BMC Health Serv Res*. 2011;11:94.
2. Dohn MN, Dohn AL. What is our plan for acute unscheduled care? *Ann Intern Med*. 2013 Oct 15;159(8):576.
3. Müller U, Winterhalder R, Businger A, Zimmermann H, Exadaktylos AK. Why do walk-in patients prefer a busy urban emergency department during office hours? A pilot survey of 200 consecutive patients from Switzerland. *Swiss Med Wkly*. 2012;142:w13565.
4. Sanchez B, Hirzel AH, Bingisser R, Ciurea A, Exadaktylos A, Lehmann B, et al. State of Emergency Medicine in Switzerland: a national profile of emergency departments in 2006. *Int J Emerg Med*. 2013;6(1):23.
5. Vilpert S. Consultations dans un service d'urgence en Suisse. Neuchâtel: Observatoire suisse de la santé. 2013 Report No. 3/2013. French/German.
6. Carron PN, Hugli OW, Schreyer N, Yersin B. Access of elderly patients in the emergency department: demographic evolution and ethical perspectives. *Rev Médicale Suisse*. 2006 Aug 9;2(75):1840–3. French.
7. Roberts DC, McKay MP, Shaffer A. Increasing rates of emergency department visits for elderly patients in the United States, 1993 to 2003. *Ann Emerg Med*. 2008 Jun;51(6):769–74.
8. Langer S, Chew-Graham C, Hunter C, Guthrie EA, Salmon P. Why do patients with long-term conditions use unscheduled care? A qualitative literature review. *Health Soc Care Community*. 2013 Jul;21(4):339–51.
9. Seematter-Bagnoud L, Junod J, Jaccard Ruedin H, Roth M, Foletti C, Santos-Eggimann B. Offre et recours aux soins médicaux ambulatoires en Suisse - Projections à l'horizon 2030. Neuchâtel: Observatoire suisse de la santé. 2008. Document de travail 33.
10. Weinick RM, Burns RM, Mehrotra A. Many emergency department visits could be managed at urgent care centers and retail clinics. *Health Aff Proj Hope*. 2010 Sep;29(9):1630–6.
11. Salisbury C, Chalder M, Scott TM, Pope C, Moore L. What is the role of walk-in centres in the NHS? *BMJ*. 2002 Feb 16;324(7334):399–402.
12. Rubin G. Unscheduled care following attendance at Minor Illness and Injury Units (MIU): cross-sectional survey. *J Eval Clin Pract*. 2012 Feb;18(1):100–3.
13. Salisbury C, Munro J. Walk-in centres in primary care: a review of the international literature. *Br J Gen Pract J R Coll Gen Pract*. 2003 Jan;53(486):53–9.
14. Chang JE, Brundage SC, Chokshi DA. Convenient Ambulatory Care--Promise, Pitfalls, and Policy. *N Engl J Med*. 2015 Jul 23;373(4):382–8.
15. Howard M, Goertzen J, Kaczorowski J, Hutchison B, Morris K, Thabane L, et al. Emergency Department and Walk-in Clinic Use in Models of Primary Care Practice with Different After-Hours Accessibility in Ontario. *Healthc Policy Polit Santé*. 2008 Aug;4(1):73–88.
16. Diserens L, Egli L, Fustinoni S, Santos-Eggimann B, Staeger P, Hugli O. Emergency department visits for non-life-threatening conditions: evolution over 13 years in a Swiss urban teaching hospital. *Swiss Med Wkly*. 2015;145:w14123.
17. Amiel C, Williams B, Ramzan F, Islam S, Ladbrooke T, Majeed A, et al. Reasons for attending an urban urgent care centre with minor illness: a questionnaire study. *Emerg Med J EMJ*. 2014 Jan 13.

18. Bardelli P, Kaplan V. Non-urgent encounters in a Swiss medical emergency unit. *Swiss Med Wkly.* 2013;143:w13760.
19. FitzGerald G, Toloo GS, Aitken P, Keijzers G, Scuffham P. Public use and perceptions of emergency departments: A population survey. *Emerg Med Australas EMA.* 2015 Aug;27(4):336–42.
20. Labгаа I, Locatelli I, Bischoff T, Gilgien W, Staeger P, Cornuz J, et al. Patients satisfaction in an academic walk-in centre: a new model of residents training achieved by family doctors. *BMC Res Notes.* 2014;7:874.
21. Statistique Vaud. Population de Lausanne. [cited 2015 Oct 21]. Available from: <http://www.scris-lausanne.vd.ch/Default.aspx?DomID=2022>
22. Henneman PL, Garb JL, Capraro GA, Li H, Smithline HA, Wait RB. Geography and travel distance impact emergency department visits. *J Emerg Med.* 2011 Mar;40(3):333–9.
23. Asaro PV, Lewis LM, Boxerman SB. The impact of input and output factors on emergency department throughput. *Acad Emerg Med Off J Soc Acad Emerg Med.* 2007 Mar;14(3):235–42.
24. Clément N, Businger A, Martinolli L, Zimmermann H, Exadaktylos AK. Referral practice among Swiss and non-Swiss walk-in patients in an urban surgical emergency department. *Swiss Med Wkly.* 2010;140:w13089.
25. Pitts SR, Carrier ER, Rich EC, Kellermann AL. Where Americans get acute care: increasingly, it's not at their doctor's office. *Health Aff Proj Hope.* 2010 Sep;29(9):1620–9.
26. Lang T, Davido A, Diakité B, Agay E, Viel JF, Flicoteaux B. Using the hospital emergency department as a regular source of care. *Eur J Epidemiol.* 1997 Feb;13(2):223–8.
27. Schoen C, Osborn R, Squires D, Doty M, Rasmussen P, Pierson R, et al. A survey of primary care doctors in ten countries shows progress in use of health information technology, less in other areas. *Health Aff Proj Hope.* 2012 Dec;31(12):2805–16.
28. Güntensperger U, Pinzello-Hürlimann R, Martina B, Ciurea A, Muff B, Gutzwiller J-P. Primary care emergency services utilization in German-speaking Switzerland: a population-based cross-sectional study. *Swiss Med Wkly.* 2010;140:w13111.
29. Bodenheimer T, Ghorob A, Willard-Grace R, Grumbach K. The 10 building blocks of high-performing primary care. *Ann Fam Med.* 2014 Apr;12(2):166–71.
30. Committee on Optimizing Scheduling in Health Care, Institute of Medicine. *Transforming Health Care Scheduling and Access: Getting to Now* [Internet]. Kaplan G, Lopez MH, McGinnis JM, editors. Washington (DC): National Academies Press (US); 2015.
31. Murray M, Berwick DM. Advanced access: reducing waiting and delays in primary care. *JAMA.* 2003 Feb 26;289(8):1035–40.
32. Pearl R. Kaiser Permanente Northern California: current experiences with internet, mobile, and video technologies. *Health Aff Proj Hope.* 2014 Feb;33(2):251–7.
33. Lafrance M, Leduc N. Prior use of a telephone-nursing triage service by patients of emergency services. *Rev Épidémiologie Santé Publique.* 2002 Dec;50(6):561–70.

Table 1

Table 1: Characteristics of patients with or without a primary care physician				
	PCP	No PCP	Total	p-value*
Patients: n (%)	232 (76.1)	73 (23.9)	305	
- Mean age, years (SD)	43.1 (18.9)	32.4 (11.1)	40.6 (17.9)	< 0.001
- Women, n (%)	127 (54.7)	25 (34.2)	152 (49.8)	0.002
Nationality: (n)	214	70	284	
- Swiss, n (%)	152 (71)	33 (47.1)	185 (65.1)	0.001
- Citizen of European country, n (%)	48 (22.4)	28 (40)	76 (26.8)	
- Citizen of non-European country, n (%)	14 (6.5)	9 (12.9)	23 (8.1)	
Non-Swiss resident in Switzerland for: (n)	63	37	100	
- < 1 year, n (%)	6 (9.5)	8 (21.6)	14 (14)	0.019
- 1-5 years, n (%)	14 (22.2)	14 (37.8)	28 (28)	
- > 5 years, n (%)	41 (65.1)	15 (40.5)	56 (56)	
- Unknown duration, n (%)	2 (3.2)	0	2 (2)	
Residency status in Switzerland: (n)	66	37	103	
- Residence, settlement or working permit, n (%)	63 (95.5)	36 (97.3)	99 (96)	1.000
- Temporary status (tourist, student), n (%)	3 (4.5)	1 (2.7)	4 (3.9)	
Highest level of education: (n)	218	70	288	
- None, n (%)	1 (0.5)	1 (1.4)	2 (0.7)	0.057
- Compulsory schooling, n (%)	52 (23.9)	11 (15.7)	63 (21.9)	
- Apprenticeship or upper-secondary vocational school, n (%)	71 (32.6)	18 (25.7)	89 (30.9)	
- Baccalaureate (secondary school), n (%)	29 (13.3)	8 (11.4)	37 (12.8)	
- Tertiary level education, university, n (%)	58 (26.6)	32 (45.7)	90 (31.3)	
- Other, n (%)	1 (0.5)	0	1 (0.3)	
- Missing, n (%)	6 (2.7)	0	6 (2.1)	
Professional occupation: (n)	216	70	286	
- Working, n (%)	130 (60.2)	53 (75.7)	183 (64)	0.008
- Stay-at-home mother or father, n (%)	6 (2.8)	2 (2.9)	8 (2.8)	
- Retired, n (%)	33 (15.3)	2 (2.9)	35 (12.2)	
- Unemployed, n (%)	7 (3.2)	5 (7.1)	12 (4.2)	
- Beneficiary of social allowance, n (%)	9 (4.2)	0	9 (3.1)	
- In training, n (%)	31 (14.4)	8 (11.4)	39 (13.6)	
PCP = primary care physician; SD = standard deviation				
* p-value for PCP vs no PCP				


Table 2

Table 2: Patients and their primary care physician	
	Total
Satisfaction with the overall service received by the patients at their PCP's practice: (n)	230
- Very satisfied, n (%)	86 (37.4)
- Quite satisfied, n (%)	90 (39.1)
- Neither satisfied nor dissatisfied, n (%)	32 (13.9)
- Quite dissatisfied, n (%)	10 (4.3)
- Very dissatisfied, n (%)	4 (1.7)
- Missing, n (%)	8 (3.5)
Number of visits to the PCP during the last year: (n)	232
- At least every week, n (%)	2 (0.9)
- At least every month, n (%)	15 (6.5)
- 3-4 times, n (%)	92 (39.7)
- Once, n (%)	77 (33.2)
- Never, n (%)	40 (17.2)
- Other, n (%)	4 (1.7)
- Missing, n (%)	2 (0.9)
Contact with the PCP's practice within the last 24 hours, n (%)	89 (38.7)*
Results of the contact with the PCP's practice: (n)	88
- The practice was closed, n (%)	51 (58)
- I was offered an appointment in x days, n (%)	22 (25)
- I spoke to my doctor by phone, n (%)	5 (5.7)
- I was given an appointment and saw my doctor, n (%)	4 (4.5)
- Other, n (%)	6 (6.8)
PCP = primary care physician	
* Among patients with a PCP, n = 230	

Table 3

Table 3: Patients' resources and solutions to solve their health problem	
	Total
Self-treatment attempted before going to the walk-in emergency clinic: (n)	288
Over-the-counter medicine from a pharmacy, n (%)	131 (45.5)
Bed rest, n (%)	119 (41.3)
Home remedies (teas/herbal teas, poultice...), n (%)	58 (20.1)
Prescription medicine, n (%)	30 (10.4)
Complementary medicine (homeopathy, herbal medicine, aromatherapy, acupuncture...), n (%)	29 (10.1)
Other, n (%)	21 (7.3)
Before going to the walk-in emergency clinic, advice was obtained from: n (%)	287
Family member or partner, n (%)	97 (33.8)
Friend, n (%)	57 (19.9)
Pharmacist, n (%)	51 (17.8)
Work colleague, n (%)	40 (13.9)
PCP's practice, n (%)	29 (10.1)
Internet, n (%)	22 (7.7)
Healthcare call centre, n (%)	7 (2.4)
Other, n (%)	15 (5.2)
Alternatives: if the walk-in emergency clinic had been unavailable, patients would have: n (%)	282
Gone to another walk-in emergency clinic or emergency department, n (%)	213 (75.5)
Gone to their family doctor, n (%)	56 (26.4)*
Gone to see the pharmacist, n (%)	24 (8.5)
Looked after the problem themselves, n (%)	24 (8.5)
Phoned a healthcare call centre, n (%)	4 (1.4)
Called their family doctor to organise a house call, n (%)	1 (0.5)*
Autre, n (%)	23 (8.2)
PCP = primary care physician	
* Among patients with a PCP, n = 212	

Figure 1



* Levels 1 and 2 of the Swiss Emergency Triage Scale

Figure 2




Figure 3




FIGURE LEGENDS

Fig 1: Study flow chart

Fig 2: Change in the number of unscheduled consultations between 2011 and 2014

Fig 3: Proportions of the total number of consultations by day of the week from 2011 to 2014

Accès et utilisation des permanences d'urgences ambulatoires

QUESTIONNAIRE PATIENT

N° :

A. ACCES AUX SOINS

1. Où vous trouviez-vous avant de vous rendre aux urgences ?
 - | A la maison
 - | Au travail
 - | Autre (décrivez s'il vous plait) :
 -

2. Actuellement, avez-vous un médecin traitant ?
 - | Oui
 - | Non → allez à la question 8

3. Avez-vous contacté le cabinet de votre médecin traitant ces dernières 24 heures pour le voir pour votre problème ?
 - | Oui
 - | Non → allez à la question 5

4. **Si oui**, cochez la case qui décrit le mieux ce qui s'est passé.
 - | On m'a donné un rendez-vous et j'ai vu mon médecin
 - | J'ai parlé à mon médecin par téléphone
 - | On m'a proposé un rendez-vous dans jours
 - | Le cabinet était fermé
 - | Autre (décrivez s'il vous plait) :
 -

5. Sur une échelle de 1 à 10, quelle est l'importance pour vous de voir le **même** médecin chaque fois que vous avez un problème de santé (entourez le chiffre correspondant à votre évaluation) ?

Pas du tout important					Très important				
1	2	3	4	5	6	7	8	9	10

6. Concernant mon suivi au cabinet médical je suis...
 - | Très satisfait
 - | Plutôt satisfait
 - | Ni satisfait, ni insatisfait
 - | Plutôt insatisfait
 - | Très insatisfait

7. Est-ce que votre degré de satisfaction quant à la prise en charge de votre médecin traitant a joué un rôle dans votre décision de venir ici aujourd'hui ?
 - | Oui
 - | Non

8. Si vous êtes venu(e) à la permanence d'urgences aujourd'hui, c'est parce que (cochez **toutes** les cases qui vous correspondent) :

Accès

- | Cette permanence d'urgences est pour moi d'un accès pratique
- | C'est l'endroit le plus proche de ma maison ou de mon travail
- | J'ai l'habitude de consulter dans cette permanence où j'ai un dossier
- | Je n'ai pas pensé aller ailleurs
- | Je m'attendais à peu d'attente
- | Je ne connais pas d'autre endroit où me rendre en urgence

Compétences

- | La permanence m'a été recommandée par un ami, ma famille ou un collègue
- | C'est le meilleur endroit pour mon type de problème
- | Je voulais un deuxième avis
- | J'ai **plus** confiance en les conseils et traitements donnés ici plutôt que ceux donnés par mon médecin traitant
- | En consultant ici, je pense accéder directement aux spécialistes

Rendez-vous

- | Il n'est pas nécessaire de prendre rendez-vous
- | Les horaires d'ouverture sont pratiques pour moi

Mon médecin traitant

- | Je ne voulais pas déranger mon médecin traitant
- | C'est plus rapide que d'obtenir un rendez-vous chez mon médecin
- | Je n'ai pas pu avoir de RDV avec mon médecin
- | Le traitement donné par mon médecin ne me convient pas
- | Mon médecin traitant ne prend pas en charge ce genre de cas
- | Autre (décrivez s'il vous plait) :
- |

B. VOTRE SANTE

1. Dans les 5 dernières années, combien de fois vous êtes-vous rendus aux urgences ?
 - | Au moins chaque semaine
 - | Au moins chaque mois
 - | 3-4 fois
 - | Une fois
 - | Jamais (avant de venir ici aujourd'hui)
 - | Autre (*décrivez s'il vous plait*) :
 -

2. **Si vous avez un médecin traitant**, combien de fois avez-vous vu votre médecin dans l'année écoulée ?
 - | Au moins chaque semaine
 - | Au moins chaque mois
 - | 3-4 fois
 - | Une seule fois
 - | Jamais
 - | Autre (*décrivez s'il vous plait*) :
 -

3. Comment jugez-vous votre état de santé en général ?
 - | Très bon
 - | Bon
 - | Moyen
 - | Mauvais
 - | Très mauvais

4. Avez-vous des maladies chroniques, problèmes de santé ou handicaps qui vous limitent dans vos activités ou votre travail au quotidien ?
 - | Oui
 - | Non

5. Combien prenez-vous de types de médicaments différents chaque jour ?
 - | Aucun
 - | Entre 1 et 2
 - | Entre 3 et 5
 - | Plus de 5

C. VOTRE PROBLEME

1. Avant de venir ici aujourd'hui, avez-vous essayé quelque chose pour résoudre votre problème ?
Si oui, cochez **toutes** les cases qui vous concernent :

- Repos
- Médicament sans ordonnance
- Médicament prescrit par un médecin
- Médecine complémentaire (homéopathie, phytothérapie, aromathérapie, acupuncture...)
- "Remède maison ou de grand-mère" (cataplasme, thés/tisanes...)
- Autre (*décrivez s'il vous plait*) :
-

2. Avant de venir ici, avez-vous obtenu des conseils auprès des sources suivantes pour votre problème ?

Si oui, cochez **toutes** les cases qui vous correspondent :

- Membre de la famille ou partenaire
- Ami
- Collègue de travail
- Pharmacie
- Centrale téléphonique sanitaire (*merci de préciser*)
 - centrale téléphonique des médecins de garde (CTMG)
 - centrale Medgate
 - centrale Medi24
 - 144
 - autre
- Cabinet médical
- Internet
- Autre (*décrivez s'il vous plait*) :
-

3. De quand date le problème de santé qui vous amène à la permanence d'urgences ?

- Aujourd'hui
- 1 à 2 jours
- 3 à 7 jours
- Plus de 7 jours

4. Sur une échelle de 1 à 10, comment évaluez-vous la gravité du problème de santé pour lequel vous êtes venu aujourd'hui (*entourez le chiffre correspondant à votre évaluation*) ?

Pas du tout grave Très grave

1 2 3 4 5 6 7 8 9 10

5. Sur une échelle de 1 à 10, quelle est votre degré d'inquiétude concernant le problème de santé pour lequel vous êtes venu aujourd'hui (*entourez le chiffre correspondant à votre évaluation*) ?

Pas du tout inquiet Très inquiet

1 2 3 4 5 6 7 8 9 10

D. LA PERMANENCE D'URGENCES

1. Etes-vous déjà venu à cette permanence d'urgences auparavant ?

- | Oui
- | Non → *allez à la question 4*

2. **Si oui**, comment évaluez-vous votre précédente expérience dans cette permanence ?

- | Excellente
- | Bonne
- | Moyenne
- | Mauvaise

3. Sur une échelle de 1 à 10, à quel point votre précédente expérience dans cette permanence a-t-elle joué un rôle dans votre décision de revenir aujourd'hui (*entourez le chiffre correspondant à votre évaluation*) ?

N'a pas du tout joué de rôle

A joué un rôle très important

1 2 3 4 5 6 7 8 9 10

4. En venant ici aujourd'hui, à quoi vous attendiez-vous ?

Cochez **toutes** les cases qui vous correspondent :

- | Uniquement recevoir des conseils
- | Une prescription de médicaments (ordonnance)
- | Une prise de sang
- | Un examen radiologique
- | Voir un spécialiste
- | Un certificat médical
- | Autre (*décrivez s'il vous plait*) :
-

5. Si le service ici n'avait pas été disponible aujourd'hui, qu'auriez-vous fait ?

Cochez **toutes** les cases qui vous correspondent :

- | Je serais allé chez mon médecin traitant
- | J'aurais appelé mon médecin traitant pour une visite à domicile
- | Je serais allé à la pharmacie
- | Je serais allé à un autre centre d'urgences
- | J'aurais appelé une centrale téléphonique sanitaire (CTMG, Medgate, Medi42...)
- | Je me serais occupé du problème moi-même
- | Autre (*décrivez s'il vous plait*) :
-

E. DONNES SOCIODEMOGRAPHIQUES

1. Sexe :
 - | Masculin
 - | Féminin

2. Quelle est votre **nationalité** ?
 - | Suisse
 - | Etrangère, en Suisse depuis moins d'un an
 - | Etrangère, en Suisse depuis un à cinq ans
 - | Etrangère, en Suisse depuis six à dix ans
 - | Etrangère, en Suisse depuis plus de dix ans
 - | Je ne souhaite pas répondre à cette question

3. Si vous êtes de nationalité étrangère, merci de préciser laquelle :
.....

4. Si vous êtes de nationalité étrangère, quel est votre **statut de séjour** en Suisse ?
 - | Permis d'établissement (Permis B ou C)
 - | Statut requérant d'asile / "Cas Dublin" (Permis N)
 - | Statut débouté de l'asile / Non entrée en matière
 - | Statut d'admission provisoire (Permis F)
 - | Statut "de passage" (étudiants, touristes, etc.)
 - | Sans Papier
 - | Autre :

5. Quelle est la **formation** la plus élevée que vous avez terminée ? *Une seule réponse SVP*
 - | Aucune
 - | Ecole obligatoire
 - | Apprentissage ou école professionnelle (brevet, CFC)
 - | Maturité ou baccalauréat
 - | Université/HES
 - | Autre :

6. Quelle est votre **situation professionnelle** actuelle ? *Une seule réponse SVP*
 - | En activité à temps plein
 - | En activité à temps partiel
 - | En activité mais en arrêt maladie actuellement
 - | Femme/homme au foyer
 - | AVS/Retraité(e)
 - | A l'assurance chômage
 - | Bénéficiaire d'une rente AI
 - | Bénéficiaire d'autres prestations sociales
 - | En formation

F. Avez-vous des remarques ou commentaires ?

Merci pour votre participation !

Accès et utilisation des permanences d'urgences ambulatoires

QUESTIONNAIRE INVESTIGATEUR

N° :

A compléter par l'investigateur, une seule réponse par question

1. Degré d'urgence au tri
 - | 3
 - | 4

2. Le questionnaire destiné au patient a-t-il été rempli ?
 - | Oui, par le patient
 - | Oui, par les proches du patient
 - | Oui, avec l'aide de l'investigateur
 - | Non, dans ce cas précisez le motif :
 - |

3. Motif de consultation :
 - |

4. Diagnostic final retenu :
 - |

5. Type d'affection ayant motivé la consultation par catégorie :
 - | 1. Maladies infectieuses et parasitaires (sauf grippe et infections de l'appareil respiratoire qui sont classées dans 10)
 - | 2. Tumeurs
 - | 3. Maladies du sang, des organes hématopoïétiques et désordres immunitaires
 - | 4. Maladies endocriniennes, de la nutrition et du métabolisme
 - | 5. Troubles mentaux et du comportement
 - | 6. Maladies du système nerveux
 - | 7. Maladies de l'œil et ses annexes
 - | 8. Maladies de l'oreille
 - | 9. Maladies de l'appareil circulatoire
 - | 10. Maladies de l'appareil respiratoire
 - | 11. Maladies de l'appareil digestif
 - | 12. Maladies de la peau et du tissu cellulaire sous-cutané
 - | 13. Maladies du système ostéoarticulaire, des muscles et du tissu conjonctif
 - | 14. Maladies des organes génito-urinaires
 - | 15. Complications de la grossesse, de l'accouchement et des suites de couches
 - | 16. Certaines affections dont l'origine se situe dans la période périnatale
 - | 17. Anomalies congénitales
 - | 18. Symptômes, signes et états morbides
 - | 19. Lésions traumatiques
 - | 20. Empoisonnement (= exposition accidentelle) à des substances médicinales ou non médicinales
 - | 21. Intoxication intentionnelle par des substances médicinales ou non médicinales
 - | 22. Effets secondaires de médicaments (administrés à des fins thérapeutiques)
 - | 23. Incidents d'une procédure ou d'un matériel
 - | 24. Autre diagnostic ne rentrant dans aucune des catégories ci-dessus :

6. Attitude à la fin de la consultation :

Destination :

- | Domicile
- | Hospitalisation (Hôpital / Clinique de)
- | Autre :

Traitement :

- | Prescription de médicaments
- | Physiothérapie
- | Autre :

Référé à un spécialiste :

- | Oui
- | Non

Arrêt de travail :

- | Oui
- | Non



**Commission cantonale d'éthique
de la recherche sur l'être humain**

Av. de Chailly 23, 1012 Lausanne

Prof. P. Francioli, Président
Prof. R. Darioli, Past-President

Secrétariat central
Tél. 021 316 18 30/31/32/33
Fax 021 316 18 37
E-mail: secretariat.cer@vd.ch

Dr Philippe Staeger
Médecin adjoint
PMU
CHUV – BU44 / 06 / 2108
1011 Lausanne

Lausanne, le 25 août 2014
PF/ns

Votre étude : Accès et utilisation des permanences d'urgences ambulatoires

Monsieur et cher Collègue,

Nous avons bien reçu votre protocole susmentionné ainsi que ses annexes et vous en remercions.

Après examen de votre dossier, la Commission considère que cette étude n'entre pas dans le champ d'application de la Loi relative à la recherche sur l'être humain tel que défini à son article 2, car il s'agit d'une enquête anonyme.

Je vous confirme cependant que votre approche respecte les principes éthiques généraux et que la CER-VD ne s'oppose pas au déroulement de cette recherche.

Veillez recevoir, Monsieur et cher Collègue, mes salutations les meilleures.

Prof. Patrick Francioli
Président

Copie : Mme Marie Vetterli, Etudiante en Master, marie.vetterli@unil.ch