



Strathprints Institutional Repository

Furman, Brian L. and Ong, W.K. and Pyne, Nigel J. (2010) *Cyclic AMP signalling in pancreatic islets*. *Advances in Experimental Medicine and Biology*, 654. pp. 281-304. ISSN 0065-2598

Strathprints is designed to allow users to access the research output of the University of Strathclyde. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. You may not engage in further distribution of the material for any profitmaking activities or any commercial gain. You may freely distribute both the url (<http://strathprints.strath.ac.uk/>) and the content of this paper for research or study, educational, or not-for-profit purposes without prior permission or charge.

Any correspondence concerning this service should be sent to Strathprints administrator: <mailto:strathprints@strath.ac.uk>

Chapter 13

Cyclic AMP Signalling in Pancreatic Islets

Brian Furman, Wee Kiat Ong, and Nigel Pyne

Abstract Cyclic 3'5'AMP (cAMP) is an important physiological amplifier of glucose-induced insulin secretion by the pancreatic islet β -cell, where it is formed by the activity of adenylyl cyclases, which are stimulated by glucose, through elevation in intracellular calcium concentrations, and by the incretin hormones (GLP-1 and GIP). cAMP is rapidly degraded in the pancreatic islet β -cell by various cyclic nucleotide phosphodiesterase (PDE) enzymes. Many steps involved in glucose-induced insulin secretion are modulated by cAMP, which is also important in regulating pancreatic islet β -cell differentiation, growth and survival. This chapter discusses the formation, destruction and actions of cAMP in the islets with particular emphasis on the β -cell.

Keywords Cyclic AMP · Adenylyl cyclase · Phosphodiesterase · Insulin secretion · Protein kinase A · Epac · GLP-1

13.1 Introduction

Interest in the role of cyclic 3'5' AMP (cAMP) in regulating insulin secretion dates back more than 40 years, since Turtle and Kipnis [1] showed increases in cAMP in isolated islets in response to glucagon. Increases in islet β -cell cyclic AMP occur in response to nutrients, especially glucose. Glucose has been widely shown to increase intracellular levels of cAMP in islets and various insulin-secreting cell lines [2–6]. Although cyclic AMP does not appear to be essential for glucose-induced insulin secretion [3, 7–9], it is established as an important intracellular amplifier of this process [10–12]. Several hormones exert their effects on insulin secretion through increased β -cell cAMP levels. These include glucose-dependent

B. Furman (✉)

Strathclyde Institute of Pharmacy and Biomedical Sciences, University of Strathclyde, Glasgow G4ONR, Scotland, UK

insulinotropic polypeptide (GIP) and glucagon-like peptide 1 (GLP-1) which are collectively referred to as the incretins, and which are also secreted in response to nutrients [13–16]. GLP-1 and GIP serve to augment meal-related insulin secretion [17]. Their physiological importance is evident from observations that mice lacking receptors for both incretin hormones show marked glucose intolerance and impairment of insulin secretion [18]. This chapter focuses largely on cAMP in the β -cell. Much less is known about the role of cAMP in other islet cells, although there is some information on this in relation to glucagon and somatostatin secretion/synthesis and these aspects will be addressed briefly at the end of the chapter.

13.2 Control of cAMP Levels in the β -Cell

The level of cyclic AMP in the β -cell depends on the balance between its formation through the activity of adenylyl cyclases (ACs) and its destruction by cyclic nucleotide phosphodiesterases (CN-PDEs). This is summarized in Fig. 13.1 and discussed below.

13.2.1 Formation of Cyclic AMP in the β -Cell

Glucose-induced elevations in intracellular cAMP are probably secondary to changes in the concentration of calcium, which is itself elevated as a result of a number of mechanisms but primarily by Ca^{2+} influx through voltage-sensitive Ca^{2+} channels in response to membrane depolarization, following closure of ATP-sensitive potassium channels. Hormone-induced formation of cAMP results from stimulation of seven transmembrane G-protein-coupled receptors (GPCRs), leading to activation of the G_s protein and dissociation of the $G\alpha\beta\gamma$ heterotrimeric complex and sequential activation of adenylyl cyclases [19]. The β -cell expresses several GPCRs coupled to G_s , stimulation of which leads to elevation in the β -cell level of cAMP. These include receptors for GLP-1, GIP, PACAP as well as the receptor GPR119 (see below). On the other hand, reductions in cAMP occur in response to several agents that activate GPCRs coupled to G_i , for example adrenaline [20], PGE_2 [21] and NPY (Y_1) [22]. There is also evidence for the role of the pertussis toxin-insensitive G-protein G_z in the reduction of cAMP and inhibition of insulin secretion in response to prostaglandin E^1 [23].

GLP-1, through stimulation of its Class II GPCR, activates AC with consequent production of intracellular cAMP [24, 25]. Oxyntomodulin, which like GLP-1, is derived from the proglucagon gene, also binds to the GLP-1 receptor, increases cAMP levels and stimulates insulin secretion [26]. There is also evidence for coupling to G_i/G_o , and, in various, non- β -cell systems to other G-proteins ($G_{q/11\alpha}$), although the physiological significance of this remains to be established. Sonoda et al. [27] identified an unusual role for β -arrestin-1 in coupling the GLP-1 receptor to

91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135




Fig. 13.1 Summary of the mechanisms for the formation and destruction of cAMP in the pancreatic islet β -cell. Glucose is transported into the β -cell using GLUT2 and is then metabolized generating ATP. This results in closure of the K_{ATP} channel, membrane depolarization and calcium influx through voltage-sensitive calcium channels. Calcium is also mobilized from intracellular stores by Ca^{2+} (calcium-induced calcium release – not shown). The increased cytosolic-free Ca^{2+} triggers exocytosis. These processes are amplified through increases in cAMP effected both through activation of adenylyl cyclases by glucose itself (through calcium-activated adenylyl cyclase – type VIII- AC VIII) and by the incretin hormones GLP-1 and GIP, acting through G-protein-coupled receptors in the β -cell membrane. Endogenous agonists for the G-protein-coupled receptor GPR119 include oleoylethanolamide (OEA). Activation of GLP-1 receptors acts synergistically with glucose in activating AC VIII and also activates other adenylyl cyclases, including soluble adenylyl cyclase (not shown). Activation of adenylyl cyclases increases the formation of cAMP which activates PKA and Epac which mediate the actions of cAMP in the cell. PKA/Epac facilitates calcium-induced calcium release which in turn may also activate AC VIII. The destruction of cAMP is effected through various phosphodiesterases (PDEs). Ca^{2+} activates PDE1 whereas PKA activates PDE3B, which is also activated by other signals generated through the IGF-1 and leptin receptors, as well as, possibly, the insulin receptor. On the other hand, PDE3B may be inhibited by increases in cGMP, allowing cross-talk between cGMP and cAMP signalling. Roles for other PDEs (PDE4, 8B and 10A) have been proposed (modified from [54])

adenylyl cyclase in INS-1 cells, thereby increasing cAMP and stimulating insulin secretion.

GIP produces its biological effects by interacting with its Class II G-protein-coupled receptor coupled to the production of cyclic AMP [28–30]. The pancreatic

136 islet β -cell GIP receptor is down-regulated by exposure to high concentrations of
137 glucose, which prevents the GIP-induced elevation in intracellular cAMP [31]. This
138 is hypothesized to explain the lack of response of diabetic patients to the peptide.

139 PACAP is expressed in nerve fibres and the pancreatic islets and is a potent
140 stimulator of insulin secretion [32, 33] through activation of adenylyl cyclase [34].
141 There are several receptors for PACAP, with the PAC1 receptor (PAC1-R) and
142 VPAC2 receptor (VPAC2-R) thought to be the most important in relation to insulin
143 secretion [35].

144 GPR119 is a Class I GPCR, the expression of which is restricted largely to
145 pancreatic islets, although lesser amounts of message are detected in the human
146 gastrointestinal tract and in the rodent brain [36–38]. The potential endogenous
147 ligands for this receptor so far identified are oleoyl lysophosphatidylcholine and
148 oleoylethanolamide, although there is as yet no evidence that they are available in
149 sufficient concentrations in the blood to stimulate the β -cell GRP119 receptor in
150 vivo. The receptor is coupled through G_s to adenylyl cyclase, and its activation
151 produces an increase in cAMP and stimulation of insulin secretion.

153 13.2.1.1 Adenylyl Cyclases in the Pancreatic Islet β -Cell

154 There are at least nine different membrane-bound isoforms of AC, described as
155 AC I–AC IX and expressed in mammalian cells [39, 40]. An additional, soluble
156 form is also expressed in certain mammalian cells [41]. RT-PCR studies, as well as
157 immunohistochemical staining, using rat and human islets, rat β -cells, and clonal
158 β -cell lines have shown expression of AC II [42] and III, IV, V, VI, VII and VIII
159 [5, 43–45]. All isoforms of adenylyl cyclase, apart from ACIX, are activated by the
160 diterpene forskolin, which produces marked increases in cAMP in numerous cell
161 types [46, 47]. There are three calcium-activated ACs (AC1, ACIII and ACVIII),
162 and the presence of calcium–calmodulin-activated ACVIII probably explains activa-
163 tion of cyclic AMP formation in response to glucose, which rapidly elevates $[Ca^{2+}]_i$.
164 This AC isoform is synergistically activated by both $G_s\alpha$ and calcium/calmodulin
165 [48]. Thus, the combination of glucose and GLP-1 increases cAMP accumulation in
166 rat isolated primary β -cells or clonal β -cell lines more markedly than either alone,
167 the effect being reduced if calcium entry through voltage-sensitive L-type channels
168 is prevented using verapamil [45]. The expression of type VI (but not types II, III or
169 V) adenylyl cyclase was increased along with the expression of the GLP-1 receptor
170 rat pups fed a high-carbohydrate diet for 12 days [42]. These findings provide some
171 circumstantial evidence that the type VI adenylyl cyclase may be associated with
172 GLP-1 signalling. More recently, a role for soluble AC was proposed to explain
173 the different kinetics of cAMP formation in response to glucose and GLP-1 in
174 INS-1E cells. GLP-1 produced a rapid increase as a result of activation of transmem-
175 brane AC, whereas the increase in cAMP in response to glucose was delayed and
176 was attributed to activation of the calcium, bicarbonate and ATP-sensitive soluble
177 AC [6].

179 Paradoxically, acetylcholine, which increases insulin secretion through stimula-
180 tion of muscarinic receptors coupled to phospholipase C/protein kinase C pathways,

181 also activated adenylyl cyclases and elevated cAMP content in islets from GK-
182 diabetic rats [49]. The insulin secretory response to acetylcholine in these islets was
183 blocked by inhibitors of adenylyl cyclase or PKA inhibitors. The abnormal nature of
184 the islet in these rats may somehow has facilitated cross-talk resulting in activation
185 of a calcium-sensitive adenylyl cyclase, or a PKC-sensitive adenylyl cyclase, e.g.
186 ACII [40], in response to acetylcholine.

190 **13.2.2 Destruction of cAMP in the Pancreatic Islet β -Cell -Cyclic** 191 **Nucleotide Phosphodiesterases**

194 Cyclic nucleotide phosphodiesterases (CN-PDEs) provide the only known means
195 for the rapid inactivation of the cyclic nucleotides cAMP and cGMP in most cells.
196 There are now known to be at least 100 PDE enzymes derived from 11 known
197 gene families (PDE1-11). The enzymes show differences in their tissue distribution,
198 substrate selectivities (cGMP vs cAMP), kinetics, regulation, and susceptibility to
199 pharmacological inhibition. There are several excellent reviews [50–53], and the
200 properties of those PDE enzymes present in pancreatic islets have been reviewed
201 elsewhere [54, 55]. The key observations are summarized in this chapter, together
202 with more recent findings.


203 Several PDE isoforms, including PDE1 [56–61], PDE3B [59–67], PDE4 [59, 60,
204 64] and PDE8B [68], contribute to the total β -cell PDE activity, and several of these
205 isoforms regulate glucose-induced insulin secretion and other cAMP-mediated β -
206 cell functions in islets and in cell lines [see 54, 55 for references]. There is much
207 evidence from RT-PCR, immunostaining, siRNA and biochemical and functional
208 studies using selective inhibitors that PDE3B plays a key role in both islets and
209 insulin-secreting cell lines in terms of regulating insulin secretion [54, 55, 61, 63–
210 66]. Additional evidence for the role of PDE3B in regulating β -cell cAMP and
211 insulin secretion was obtained by over-expressing PDE3B in the INS-1 β -cell line
212 and in islets and by using transgenic animals over-expressing PDE3B in the β -
213 cell. These in vitro and in vivo studies clearly showed that glucose-induced, as
214 well as GLP-1-induced, insulin secretion was impaired by PDE3B over-expression.
215 Interestingly, both endogenous and over-expressed PDE3B was found to be located
216 in insulin granules and the plasma membrane [67]. In vitro, the over-expression of
217 PDE3B markedly reduced cAMP-induced exocytosis and animals over-expressing
218 PDE3B in islets showed markedly impaired glucose tolerance [65–67]. In addition,
219 activation of PDE3B appears to mediate the effect of IGF-1 [63] and leptin [69] in
220 inhibiting insulin secretion.

221 The role of cGMP in regulating insulin secretion is not established, but several
222 studies have shown that nitric oxide, acting through a soluble guanylyl cyclase and
223 GMP formation, augments insulin secretion through several mechanisms shared
224 with cAMP (see Section 13.3.1) [70–73]. These observations might be explained
225 by cGMP-dependent inhibition of PDE3B and concomitant increases in [cAMP]_i.

226 Although evidence for the importance of PDE3B is widely supported there is also
227 evidence, but no consensus, for roles for other PDEs. Roles for PDE1C and PDE4
228 have been suggested on the basis of the use of either selective inhibitors [59, 64] or
229 siRNA [64]. Depletion of PDE8B using siRNA produced a marked enhancement of
230 glucose-induced insulin secretion from INS-1E cells [64, 68] and rat islets [68]. A
231 role for PDE10A has been proposed and selective inhibitors have been patented for
232 the treatment of diabetes [74], but there is no consensus on the expression of this
233 PDE in the β -cell, and in one study [64] selective knockdown of PDE10A failed to
234 modify glucose-induced insulin secretion in INS-1 cells.

235 236 237 238 **13.2.3 Dynamics of cAMP Formation and Destruction**

239
240 Real-time measurements of changes in cAMP in β -cells or islets have been hugely
241 facilitated by the development of new technologies, particularly the development of
242 genetically encoded fluorescence resonance energy transfer (FRET)-based biosen-
243 sors and the associated imaging techniques. These have either been transiently
244 transfected into β -cell lines or primary β -cells [5, 75–78] or been incorporated
245 in vivo by generating a transgenic mouse expressing a pancreatic β -cell-targeted
246 cAMP reporter which was inducible in response to tetracycline [4]. In MIN6 β -
247 cells, the use of the biosynthetic FRET-based cAMP sensor Epac1-camps, together
248 with FURA-2 to detect $[Ca^{2+}]_i$, showed a close coupling of changes in cAMP and
249 $[Ca^{2+}]_i$ [5]. Exendin-4 and forskolin induced pronounced FRET signals. Formation
250 of cAMP in response to these agents was preceded by increases in $[Ca^{2+}]_i$ and
251 was dependent upon extracellular calcium. Moreover, increases in $[Ca^{2+}]_i$ evoked
252 by other agents (carbachol, K^+ , and tolbutamide) also stimulated cAMP formation.
253 Simultaneous imaging of $[Ca^{2+}]_i$ and cAMP during glucose stimulation (in the pres-
254 ence of TEA) revealed a tight coupling between oscillations in $[Ca^{2+}]_i$ and cAMP
255 with peak cAMP concentrations being seen at the nadir of $[Ca^{2+}]_i$. The data are
256 consistent with the possibility that Ca^{2+} -activated adenylyl cyclases (AC VIII or
257 AC III) and PDEs (PDE1C?) contribute to the oscillatory changes in cAMP seen
258 in these studies. How this concept fits with the widely accepted role of PDE3B in
259 regulating the cAMP pool relevant to insulin secretion (Section 13.2.2) remains to
260 be determined. Other experimental studies (Fig. 13.2) and mathematical modelling
261 have supported these ideas [75]. Imaging of the islets from transgenic mice express-
262 ing a β -cell-targeted reporter showed a rapid, biphasic and concentration-dependent
263 (5.5–35 mM) increase in cAMP in response to glucose. This preceded increases in
264 $[Ca^{2+}]_i$ and was independent of extracellular $[Ca^{2+}]$ [4]. In INS-1 cells, GLP-1 pro-
265 duced marked oscillations in cAMP at low concentrations (0.3–1 nM) with higher
266 concentrations (10 nM) producing more sustained elevations [77]. GLP-1 also pro-
267 duced marked Ca^{2+} spiking, which rapidly followed the increases in cAMP. This
268 pattern of changes in cAMP and Ca^{2+} was mimicked by application of short pulses
269 of the non-selective PDE inhibitor, IBMX. The rapidity of the cAMP-induced Ca^{2+}
270 signal suggests a close proximity of the cAMP to the sites of calcium entry/release



AJP - Cell Physiology

292 **Fig. 13.2** Ca^{2+} and cAMP oscillations in glucose-stimulated MIN6 cells. Simultaneous imaging
293 of cytosolic cAMP concentration ($[\text{cAMP}]_i$; *top trace*, R_{485/535}) and cytosolic Ca^{2+} concentration
294 ($[\text{Ca}^{2+}]_i$; *bottom trace*, R_{340/380}) in a single MIN6 cell stimulated with 20 mM glucose and 20 mM
295 tetraethylammonium chloride (TEA). Note that second messenger oscillations were out of phase,
296 with each $[\text{Ca}^{2+}]_i$ spike coupled to a rapid and transient reduction in $[\text{cAMP}]_i$. (Reproduced from
297 Fridlyand LE, Harbeck MC, Roe MW, Philipson LH. Regulation of cAMP dynamics by Ca^{2+} and
298 G protein-coupled receptors in the pancreatic beta-cell: a computational approach. Am J Physiol
299 Cell Physiol 293: C1924–33, 2007 [75] with permission)

300
301 (see next section). On the other hand, translocation of the catalytic subunit of PKA to
302 the nucleus occurred relatively slowly and only in response to sustained increases in
303 cAMP. Glucose also induced oscillations of intracellular cAMP levels in MIN6 and
304 mouse primary β -cells. These oscillations correlated with pulsatile insulin secretion
305 and both cAMP oscillations and pulsatile insulin release were reduced by inhibiting
306 adenylyl cyclases [78]. Forskolin, glucagon and IBMX all augmented the frequency
307 of glucose-induced oscillations in $[\text{Ca}^{2+}]_i$ in mouse pancreatic islets [79]

13.2.4 Intracellular Compartmentalization of cAMP Formation, Action and Degradation

310
311
312
313
314 It is now established that intracellular cAMP is not uniformly distributed in the cell
315 and exists in different cellular locations to fulfil different functions. Local generation,

316 hydrolysis and activity of cAMP are ensured by spatial distribution into compart-
317 ments, or signalling complexes, of adenylyl cyclases, PDEs and effector proteins,
318 as well as phosphatases that terminate the activity of various kinases (e.g. 80, 81).
319 This spatial anchoring of signalling complexes is effected by a family of A-kinase
320 anchoring proteins (AKAPs). Recent work has suggested the importance of AKAPs
321 in the insulin-secreting β -cell. Peptides that competitively inhibit the interaction
322 between the regulatory subunit of PKA and the AKAP inhibited GLP-1-induced
323 insulin secretion from rat islets without modifying its ability to elevate intracellu-
324 lar cAMP [9]. Expression of this inhibitory peptide in the clonal rat β -cell line,
325 RINm5F, resulted in a redistribution of the PKA regulatory subunit and inhibited
326 elevations in $[Ca^{2+}]_i$ and insulin secretion in response to a cAMP analogue.
327 Expression of an AKAP (AKAP18) in clonal insulin-secreting cells (RINm5f) aug-
328 mented GLP-1-induced insulin release, whereas expression of a mutant form in
329 these cells was inhibitory [82]. These findings were supported by others [83] who
330 used a cell-permeable peptide (TAT-AKAPis) to competitively inhibit PKA–AKAP
331 interactions in INS-1 cells. This peptide disrupted PKA–AKAP interactions and
332 inhibited both glucagon-induced augmentation of insulin secretion and phosphory-
333 lation of p44/p42 MAPKs and cAMP response element binding protein. While rela-
334 tively little is known about the role of phosphatases in terminating phosphorylation-
335 mediated actions of cAMP in the pancreatic islet β -cell [84], there is evidence that
336 the AKAP AKAP79 (the human homologue of AKAP150) is important in targeting
337 the serine–threonine phosphatase PP2B to PKA-sensitive target proteins [85].
338
339
340

341 13.3 Functions of Cyclic AMP in the Pancreatic Islet β -Cell

342
343 cAMP modulates a number of β -cell functions including insulin secretion, insulin
344 synthesis, β -cell replication, and β -cell apoptosis. Actions of cAMP in general
345 are mediated by at least two distinct mechanisms. The first of these is through
346 protein kinase A (PKA)-mediated phosphorylation [86]. However, a second, and
347 PKA-independent, effect of cAMP on insulin secretion [87–88] is mediated by the
348 cyclic AMP-binding proteins known either as cAMP-regulated guanine nucleotide
349 exchange factors (GEFs) or as exchange proteins activated by cAMP (Epacs)
350 which target the small G-protein Rap1 [86]. Interestingly, most of the β -cell
351 Rap1, at least in MIN6 cells, appears to be co-localized with insulin secretory
352 granules [89]. When activated by cAMP, Epac, which exists as two isoforms
353 (Epac1 and Epac2) exchanges GDP for GTP and activates downstream sign-
354 nalling. The pancreatic islet β -cell expresses both Epac1 and Epac2 [90]. Antisense
355 oligodeoxynucleotides against Epac reduced the effect of a permeant cAMP ana-
356 logue in augmenting glucose-induced insulin secretion in pancreatic islets [91].
357 Studies using selective inhibitors/activators of PKA, selective activators of Epac
358 or the use of dominant-negative forms of Epac are revealing the roles of Epacs
359 vs PKA in the β -cell. Novel cAMP analogues, such as 8-(4-chlorophenylthio)-2'-
360 *O*-methyladenosine-3'-5'-cyclic monophosphate (8-pCPT-2'-*O*-Me-cAMP), and its

361 much more cell-permeant acetoxy methyl ester [92] activate Epac but not PKA,
362 having a 100-fold lower affinity for PKA relative to Epac [86]. Similarly, cAMP
363 analogues such as N6-Bnz-cAMP selectively activate PKA relative to Epac. Both
364 Epac and PKA mediate the effects of cAMP on insulin secretion. However, at least
365 in INS-1 cells, PKA-mediated effects account for the greater proportion of cAMP
366 effects [92]. There is evidence for interaction between PKA-mediated and Epac-
367 mediated effects in augmenting insulin secretion in native β -cells [93]. Some of
368 the reported discrepancies may be explained by the poor cell permeability of some
369 Epac-selective cAMP analogues [92].

370 The cyclic AMP-mediated effects of GIP and GLP-1 on insulin secretion involve
371 both PKA [24] and PKA-independent actions. The latter are probably mediated
372 through Epac, as evidenced by the comparative effects of the PKA inhibitor H89
373 and antisense oligodeoxynucleotides (ODNs) against Epac in reducing incretin-
374 augmented insulin secretion [91, 94]. Interestingly, Epac-dependent effects of
375 cAMP on insulin release are impaired in islets from mice lacking the SUR subunit
376 of the K_{ATP} channel [94, 95].

377

378

379 **13.3.1 Insulin Secretion**

380

381 Malaisse's group was the first to systematically examine the actions of cAMP
382 on insulin secretion [96, 97]. Elevations in cAMP in the β -cell augment glucose-
383 induced insulin secretion at several sites in the secretory pathway.

384

385

386 **13.3.1.1 Effects on the β -Cell ATP-Sensitive Potassium Channel**

387 The β -cell ATP-sensitive potassium channel (K_{ATP} channel) plays a fundamental
388 role in glucose-induced insulin secretion. Elevation of cAMP in the β -cell using
389 GLP-1, forskolin, or the non-selective PDE inhibitor IBMX inhibits the β -cell K_{ATP}
390 channel promoting depolarization of the cell [98–103]. This effect was reported to
391 be mediated via PKA in INS-1 cells [101] through phosphorylation of the SUR1
392 subunit. On the other hand, Epac was found to inhibit this channel in both human
393 β -cells and INS-1 cells, producing a leftward shift in the ATP-concentration–effect
394 curve [102, 103]. The same study [103] suggested a PKA-mediated *activation* of
395 the ATP-sensitive K channel.

396

397

398 **13.3.1.2 Voltage-Sensitive Potassium Channels**

399 Activation of voltage-sensitive potassium channels contribute to a restoration of
400 the β -cell membrane potential and a termination of insulin secretion. GIP, acting
401 through a PKA-dependent mechanism, reduced K currents through voltage-sensitive
402 potassium channels in HEK cells transfected with the GIP receptor and Kv1.4 chan-
403 nels, as well as in human islets and INS-1 cells [104]. GLP-1 and the GLP-1 mimetic
404 exendin-4 also inhibited voltage-dependent K currents effects again being PKA
405 dependent as evidenced by the preventative effects of PKA inhibition [105, 106]

13.3.1.3 Elevations in Intracellular Calcium $[Ca^{2+}]_i$

Increases in $[Ca^{2+}]_i$ can be effected through two main mechanisms, namely influx through voltage-sensitive Ca^{2+} channels and mobilization of Ca^{2+} from intracellular stores and cAMP influences both these mechanisms in the β -cell.

Voltage-Sensitive Ca^{2+} Channels

Entry of Ca^{2+} through L-type voltage-sensitive calcium channels in response to membrane depolarization is an important trigger for exocytosis. Agents elevating cAMP as well as cAMP itself augment the opening of channel and increase calcium influx [99, 107–109] through PKA-dependent mechanisms. This is consistent with observations that forskolin and IBMX were shown to produce phosphorylation of the cardiac-type alpha 1 subunit of the voltage-sensitive calcium channel in a mouse β -cell line β TC3 [110].

Mobilization of Ca^{2+} from Intracellular Stores

Calcium-Induced Calcium Release

In addition to facilitating calcium entry, agents that elevate β -cell cAMP also promote calcium-induced Ca^{2+} release [111–116]. For example, the uncaging of calcium from a membrane-permeable caged calcium (NP EGTA) produced a large, transient increase in $[Ca^{2+}]_i$ but only in the presence of the GLP-1 mimetic exendin 4 or the adenylyl cyclase activator forskolin. This could be replicated by non-selective cAMP analogues or those that selectively activated either PKA or Epac. The effects of exendin-4 were relatively insensitive to the PKA inhibitor H89 but were inhibited by expression of a dominant-negative Epac2 [116], suggesting an important role of Epac2 in the sensitizing effect of cAMP on calcium-induced Ca^{2+} release. The importance of non-PKA-dependent effects of GLP-1 in elevating $[Ca^{2+}]_i$ was also reported previously [117].

The mechanism whereby cAMP promotes calcium-induced Ca^{2+} release may be through activation of the ryanodine channel in the ER [93, 112, 113] and/or through phosphorylation of the IP_3 receptor [118]. The interaction of cAMP, via PKA, with IP_3 receptors is supported by the finding that 2-aminoethoxydiphenyl borate, a cell-permeable IP_3 -receptor antagonist, blocked the PKA-mediated cAMP amplification of calcium-induced Ca^{2+} release [119].

Generation of Ca^{2+} -Mobilizing Second Messengers

GLP-1 was shown to increase intracellular production of nicotinic acid adenine dinucleotide phosphate (NAADP) and cyclic ADP-ribose (ADPR) through cAMP mechanisms mediated by both PKA and Epac [120]. The production of the second messengers, cyclic ADPR and NAADP, is catalyzed by ADPR cyclases. Both mobilize Ca^{2+} from intracellular stores and NAADP stimulates insulin secretion. The

451 relative role of cyclic ADPR and NAADP in producing cAMP-mediated increases
452 in $[Ca^{2+}]_i$; remain to be determined.

453

454

13.3.1.4 Direct Effect on Exocytosis

455

456 Ammala et al. [107] and Gillis and Misler [121] were the first to demonstrate that
457 cAMP produced direct effects on exocytosis. This effect was suggested to repre-
458 sent the most important effect of cAMP on insulin release [107]. Both GIP and
459 GLP-1 promote PKA-dependent and PKA-independent exocytosis, independently
460 of changes in calcium entry [87, 99, 122]. Moreover, photo release of caged cAMP
461 produces a marked increase in granule exocytosis that is independent of changes
462 in $[Ca^{2+}]_i$ [87, 99, 123, 124]. GLP-1 and cAMP augmented depolarization-induced
463 exocytosis, and the effects of cAMP were mediated through both PKA-dependent
464 and PKA-independent, Epac-mediated effects [95]. cAMP also enhanced exocyto-
465 sis in single INS-1 cells, the effect being augmented by inhibition of PDE3 [65].
466 In permeabilized rat islets cAMP enhanced calcium-induced insulin secretion, inde-
467 pendently of changes in $[Ca^{2+}]_i$; this effect was largely dependent on Epac as it
468 was mimicked by an Epac-selective, but not by a PKA selective, cAMP analogue
469 and was unaffected by a PKA inhibitor [125]. Use of two-photon extracellular polar
470 tracer (TEP) imaging and electron microscopy showed different roles of PKA or
471 Epac in the enhancement by cAMP of calcium-evoked exocytosis of small compared
472 with large, secretory vesicles [124]. Effects of cAMP on large vesicle exocytosis
473 appeared to be PKA dependent, whereas effects on small vesicles were mediated
474 via Epac.

475

476 There are different pools of insulin secretory granules in the β -cell. The first
477 phase of glucose-induced insulin secretion is due to the release of granules docked at
478 the membrane in a readily releasable pool and the second phase is dependent on the
479 mobilization of granules to refill this readily releasable pool. The effects of cAMP,
480 which augments both first and second phases of insulin secretion, are at least partly
481 attributable to an expansion and refilling of the readily releasable pool [126–128].
482 Knockout of Epac2 specifically blocks the first phase of glucose-induced granule-
483 plasma membrane fusions, suggesting the importance of cAMP signalling through
484 Epac2 in this phase [89]. This supports earlier findings that the augmentation by
485 cAMP of short depolarizations was Epac dependent, whereas the effect on longer
486 depolarizations was largely PKA dependent and was more sensitive to cAMP [95].
487 The second phase of exocytosis appears to be mediated via both PKA and Epac
488 [95, 127, 128], although a PKA dependency of the first phase of glucose-induced
489 exocytosis has also been reported [123].

489

490

13.3.1.5 Activation of Protein Kinase C

491

492 Protein kinase C (PKC) is another second messenger contributing to the regula-
493 tion of insulin secretion, and one study suggests that PKC may mediate some of
494 the insulin secretory effects of agents that elevate cAMP. Thus, GLP-1 was shown
495 to activate the translocation of PKC α and PKC ϵ in INS-1 cells and its effects are

496 mimicked by forskolin. This activation was Ca^{2+} dependent, and it was hypothe-
497 sized that it was effected through mobilization of Ca^{2+} as a result, for example, of
498 PKA sensitization of the IP_3 channel and consequent Ca^{2+} -mediated activation of
499 phospholipase C [129].

500
501

502 **13.4 Role of cAMP in Insulin Synthesis and in β -Cell** 503 **Differentiation, Proliferation, and Survival**

504

505 The incretin GLP-1, acting to an important extent through cAMP effector mecha-
506 nisms, increases insulin synthesis, promotes β -cell proliferation and inhibits β -cell
507 apoptosis [25], although there is evidence for cAMP-independent effects [130].
508 Indeed much of the evidence for the importance of cAMP in these processes is
509 derived from studies using GLP-1 and exendin-4. The finding that mice with a β -
510 cell-specific deficiency in the α subunit of G_s showed reduced β -cell mass, reduced
511 islet content of insulin, reduced β -cell proliferation, and increased β -cell apoptosis,
512 and marked hyperglycaemia suggests the fundamental importance of responsiveness
513 to incretin hormones [131] in β -cell homeostasis.

514 Glucose-mediated increases in insulin synthesis involve the phosphorylation of
515 the transcription factor pancreatic duodenal homeobox-1 (PDX-1) and its transloca-
516 tion to the nucleus [132]. There is strong evidence for the importance of cAMP,
517 acting through PKA-dependent mechanisms, in mediating the ability of GLP-1
518 to increase β -cell levels of PDX-1, stimulate its translocation to the nucleus and
519 consequently activate the insulin gene promoter [133]. PDX-1 expression is itself
520 required for the generation of cAMP in response to exendin-4 through controlling
521 the expression of the GLP-1 receptor and the G_s protein α subunit [134].

522 CREB (cAMP response element binding protein) is the key transcriptional acti-
523 vator that mediates the effects of cAMP on gene regulation and its effects in
524 regulating islet β -cell proliferation and survival. cAMP, through a PKA-dependent
525 mechanism, and glucose act synergistically to regulate CREB activation in MIN6
526 or INS-1 cells [135, 136]. This appears to involve cAMP/PKA and glucose-induced
527 modulation of the phosphorylation status of TORC2, a key co-activator of CREB,
528 and the stimulation of its translocation to the nucleus [135, 136].

529
530

531 **13.4.1 Immediate Early Response Genes**

532

533 Cyclic AMP appears to mediate the effects of glucose in stimulating the β -cell
534 expression of immediate early response genes such as *c-myc* [137] and *c-fos* [138],
535 which probably play an important role in the effects of glucose in regulating the
536 gene expression of metabolic enzymes, cell growth, and apoptosis. In Min6 insulin-
537 secreting cells Glauser et al. [139] identified 592 targets and 1278 immediate early
538 genes responding to co-stimulation with glucose and cAMP (chlorophenylthio-
539 cAMP, a cell-permeant cAMP analogue) and suggested an important role for
540 the transcription factor AP-1. Indeed, the AP-1-regulated gene sulfiredoxin was

541 identified among the targets that were sequentially induced in primary cells from
542 rat islets. In the same context, cAMP also amplifies the effect of glucose in
543 stimulating the MAPK/ERK pathway [6, 140–142]. The augmentation of glucose-
544 induced activation of ERK in response to GLP-1 required both influx of Ca^{2+}
545 through voltage-dependent calcium channels and was PKA dependent [143] and
546 GIP activates this kinase pathway through cyclic AMP and PKA [144].

549 **13.4.2 Protection Against β -Cell Apoptosis and Stimulation** 550 **of β -Cell Proliferation**

552 There is abundant evidence for suppression of β -cell apoptosis by agents that elevate
553 cAMP, including GLP-1, GIP, exendin-4, ghrelin and obestatin [135, 145–151]. This
554 appears to be PKA mediated [148, 149]. Paradoxically, some β -cell lines were made
555 more susceptible to apoptosis following exposure to dibutyryl cyclic AMP [152] or
556 the cyclic AMP-elevating agent forskolin [153]. The anti-apoptotic effects of cAMP
557 are mediated, in part, by increased expression of the anti-apoptotic proteins Bcl-2
558 and Bcl-xL [135, 146], and are PKA dependent [135, 146, 151]. The anti-apoptotic
559 effects also involve caspase inhibition [147]. Inhibition of cytokine-mediated nitric
560 oxide production by β -cells [154] may also be implicated.

561 In addition to preventing apoptosis of β -cells, the incretin hormones and other
562 agents elevating cAMP promote β -cell proliferation through PKA-dependent mech-
563 anisms [134, 155, 156]. This effect appears to involve expression of cyclin D1 [155,
564 157] and cyclin A2 [134]. In this context, there may be an interaction of cAMP with
565 Wnt signalling, which plays an important role in β -cell proliferation and survival
566 with upregulation of cyclins D1 and D2 [158]. Thus, GLP-1 and exendin-4 acti-
567 vated Wnt signalling in INS-1 cells and in isolated islets [159]. Exendin-induced
568 β -cell proliferation was inhibited by blocking β -catenin or the transcription factor
569 TCF7L2, critical mediators of Wnt signalling [159].

570 An additional mechanism whereby cAMP modulates β -cell proliferation may
571 be through regulation of the CREB antagonists cAMP response element modulator
572 CREM- α and ICER1 and the dual specificity phosphatase DUSP14, a negative reg-
573 ulator of the MAPK/ERK1/2 pathway. Thus, genes for these proteins were rapidly
574 and strongly upregulated by GLP-1 in a β -cell line and in rat primary β -cells, an
575 effect that was mimicked by forskolin and blocked by the PKA inhibitor H89 but
576 not by an Epac inhibitor. shRNA-mediated knockdown of CREM- α or DUSP14,
577 or expression of a dominant-negative DUSP14, augmented GLP-1-induced β -cell
578 proliferation [156].

581 **13.5 Possible Roles of cAMP in Other Islet Cell Types**

582 Relatively little is known about the role of cAMP in other islet cells, although there
583 is some information on its role in the glucagon-secreting and somatostatin-secreting
584
585

586 cells. Forskolin was shown to stimulate glucagon secretion from rat islets [160].
587 GLP-1 (and GIP) augmented depolarization-evoked exocytosis from rat α -cells;
588 this effect was accompanied by elevations in intracellular cAMP, increases in Ca^{2+}
589 currents and was mediated by PKA [161]. Exposure of an α -cell line (INR1-
590 G9) expressing recombinant GLP-1 receptors to GLP-1 increased the formation
591 of cAMP and elevated free cytosolic $[\text{Ca}^{2+}]$ [162]. In the same cell line, an
592 Epac-selective cAMP analogue stimulated the expression of the glucagon gene pro-
593 moter and stimulated glucagon production, although not glucagon secretion [163].
594 Moreover, a dominant-negative Epac-2 attenuated forskolin-stimulated expression
595 of the glucagon gene promoter in the InR1-G9 cells [163]. While these data indi-
596 cate a stimulatory effect of GLP-1 on glucagon synthesis and secretion, GLP-1 is
597 known to inhibit glucagon secretion, an action likely to contribute to its therapeutic
598 effect in the treatment of diabetes [164]. The inhibition of glucagon secretion by
599 GLP-1 is thus likely to be mediated by a paracrine action in the islets, for example,
600 through stimulation of somatostatin secretion, which markedly inhibits glucagon
601 release [165]. In this context, GLP-1, oxyntomodulin and glucagon were shown
602 to potently stimulate somatostatin secretion from somatostatin-secreting cell lines
603 (RIN T3; RIN 1048-38) and to stimulate the accumulation of cAMP [166, 167].
604 Increases in cAMP levels in response to forskolin, theophylline or dibutyryl cAMP
605 were shown to be associated with increased somatostatin release from isolated islets
606 [168].

607 Glucagon itself stimulates glucagon release by activating glucagon, rather than
608 GLP-1, receptors, through cAMP-dependent mechanisms involving both PKA and
609 Epac [169].

610 Adrenaline, or isoprenaline, acting through β -adrenoceptors, augmented
611 depolarization-evoked glucagon secretion from rat primary α -cells [170]. This effect
612 was mimicked by forskolin and was PKA dependent. As in the β -cell the PKA-
613 dependent effects appear to involve more than one mechanism, including increased
614 Ca^{2+} entry and augmentation of the effects of Ca^{2+} . Photo release of caged cAMP
615 increased exocytosis even when intracellular $[\text{Ca}^{2+}]$ was clamped [170]. These data
616 were supported by observations using mouse primary α -cells, in which adrenaline-
617 induced increases in α -cell $[\text{Ca}^{2+}]_i$ were mediated, in part, by elevations in cAMP
618 and activation of PKA [171].

619

620

621

622 13.6 Conclusion

623

624 cAMP is clearly an important mediator/modulator of many β -cell functions from
625 hormone secretion to proliferation, survival and synthetic functions and is also likely
626 to be important in other islet cell types. Further work will elucidate the precise
627 mechanisms whereby PKA and Epac, the known mediators of the effects of cAMP,
628 exert their effects on these cellular processes. Novel ways of targeting cAMP mech-
629 anisms through small molecules, rather than peptides, may open up new treatments
630 for diabetes mellitus. Small molecules targeting the GRP119 receptor are under

development [37]. A number of non-peptide agents that act both as direct agonists and allosteric modulators of the GLP-1 receptor are also being examined [172].

References

1. Turtle J, Kipnis D. An adrenergic receptor mechanism for the control of cyclic 3',5' adenosine monophosphate synthesis in tissues. *Biochem Biophys Res Commun* 1967;28:797–802.
2. Charles M, Fanska R, Schmid F, Forsham P, Grodsky G. Adenosine 3',5'-monophosphate in pancreatic islets: glucose-induced insulin release. *Science* 1973;179:569–571.
3. Grill V, Cerasi E. Activation by glucose of adenyl cyclase in pancreatic islets of the rat. *FEBS Lett* 1973;33:311–4.
4. Kim J, Roberts C, Berg S, Caicedo A, Roper S, Chaudhari N. Imaging cyclic AMP changes in pancreatic islets of transgenic reporter mice. *PLoS ONE* 2008;3:e2127.
5. Landa LJ, Harbeck M, Kaihara K, Chepurny O, Kitiphongspattana K, Graf O, Nikolaev V, Lohse M, Holz G, Roe M. Interplay of Ca^{2+} and cAMP signaling in the insulin-secreting MIN6 beta-cell line. *J Biol Chem* 2005;280:31294–302.
6. Ramos L, Zippin J, Kamenetsky M, Buck J, Levin L. Glucose and GLP-1 stimulate cAMP production via distinct adenylyl cyclases in INS-1E insulinoma cells. *J Gen Physiol* 2008;132:329–38.
7. Sharp G. The adenylate cyclase-cyclic AMP system in islets of Langerhans and its role in the control of insulin release. *Diabetologia* 1979;16:287–96.
8. Persaud S, Jones P, Howell S. Glucose-stimulated insulin secretion is not dependent on activation of protein kinase A. *Biochem Biophys Res Commun* 1990;173:833–9.
9. Lester L, Langeberg L, Scott J. Anchoring of protein kinase A facilitates hormone-mediated insulin secretion. *Proc Natl Acad Sci U S A* 1997;94:14942–7.
10. Holz G, Habener J. Signal transduction crosstalk in the endocrine system: pancreatic beta-cells and the glucose competence concept. *Trends Biochem Sci* 1992;17:388–93.
11. Howell S, Jones P, Persaud S. Regulation of insulin secretion: the role of second messengers. *Diabetologia* 1994;37 Suppl 2:S30–5.
12. Braun M, Ramracheya R, Johnson P, Rorsman P. Exocytotic properties of human pancreatic beta-cells. *Ann N Y Acad Sci* 2009;1152:187–93.
13. MacIntosh C, Horowitz M, Verhagen M, Smout A, Wishart J, Morris H, Goble E, Morley J, Chapman I. Effect of small intestinal nutrient infusion on appetite, gastrointestinal hormone release, and gastric myoelectrical activity in young and older men. *Am J Gastroenterol* 2001;96:997–1007.
14. Brubaker P, Anini Y. Direct and indirect mechanisms regulating secretion of glucagon-like peptide-1 and glucagon-like peptide-2. *Can J Physiol Pharmacol* 2003;81:1005–12.
15. Feinle C, Chapman I, Wishart J, Horowitz M. Plasma glucagon-like peptide-1 (GLP-1) responses to duodenal fat and glucose infusions in lean and obese men. *Peptides* 2002;23:1491–95.
16. Wolfe M, Zhao K, Glazier K, Jarboe L, Tseng C. Regulation of glucose-dependent insulinotropic polypeptide release by protein in the rat. *Am J Physiol Gastrointest Liver Physiol* 2000;279:G561–6.
17. Thorens B. Expression cloning of the pancreatic beta cell receptor for the gluco-incretin hormone glucagon-like peptide 1. *Proc Natl Acad Sci U S A* 1992;89:8641–5.
18. Preitner F, Ibberson M, Franklin I, Binnert C, Pende M, Gjinovci A, Hansotia T, Drucker D, Wollheim C, Burcelin R, Thorens B. Gluco-incretins control insulin secretion at multiple levels as revealed in mice lacking GLP-1 and GIP receptors. *J Clin Invest* 2004;113:635–45.
19. Selbie L, Hill S. G protein-coupled-receptor cross-talk: the fine-tuning of multiple receptor-signalling pathways. *Trends Pharmacol Sci* 1998;19:87–93.

- 676 20. Yamazaki S, Katada T, Ui M. Alpha 2-adrenergic inhibition of insulin secretion via interference
677 with cyclic AMP generation in rat pancreatic islets. *Mol Pharmacol* 1982;21:648–53.
- 678 21. Robertson R, Tsai P, Little S, Zhang H, Walseth T. Receptor-mediated adenylate cyclase-
679 coupled mechanism for PGE₂ inhibition of insulin secretion in HIT cells. *Diabetes*
1987;36:1047–53.
- 680 22. Morgan D, Kulkarni R, Hurley J, Wang Z, Wang R, Ghatei M, Karlsen A, Bloom S, Smith D.
681 Inhibition of glucose stimulated insulin secretion by neuropeptide Y is mediated via the Y1
682 receptor and inhibition of adenylyl cyclase in RIN 5AH rat insulinoma cells. *Diabetologia*
683 1998;41:1482–91.
- 684 23. Kimple M, Nixon A, Kelly P, Bailey C, Young K, Fields T, Casey P. A role for G_z in
685 pancreatic islet β -cell biology. *J Biol Chem* 2005;280:31708–13.
- 686 24. Drucker D, Philippe J, Mojsov S, Chick W, Habener J. Glucagon-like peptide I stimulates
687 insulin gene expression and increases cyclic AMP levels in a rat islet cell line. *Proc Natl*
Acad Sci U S A 1987;84:3434–8.
- 688 25. Doyle M, Egan J. Mechanisms of action of glucagon-like peptide 1 in the pancreas.
689 *Pharmacol Ther* 2007;113:546–93.
- 690 26. Maida A, Lovshin J, Baggio L, Drucker D. The glucagon-like peptide-1 receptor agonist
691 oxyntomodulin enhances beta-cell function but does not inhibit gastric emptying in mice.
Endocrinology 2008;149:5670–8.
- 692 27. Sonoda N, Imamura T, Yoshizaki T, Babendure J, Lu J, Olefsky J. Beta-Arrestin-1 mediates
693 glucagon-like peptide-1 signaling to insulin secretion in cultured pancreatic beta cells. *Proc*
Natl Acad Sci U S A 2008;105:6614–9.
- 694 28. Amiranoff B, Vauclin-Jacques N, Laburthe M. Functional GIP receptors in a hamster pan-
695 creatic beta cell line, In 111: specific binding and biological effects. *Biochem Biophys Res*
Commun 1984;123:671–6.
- 696 29. Siegel E, Creutzfeldt W. Stimulation of insulin release in isolated rat islets by GIP in
697 physiological concentrations and its relation to islet cyclic AMP content. *Diabetologia*
698 1985;28:857–61.
- 699 30. Wheeler M, Gelling R, McIntosh C, Georgiou J, Brown J, Pederson R. Functional expres-
700 sion of the rat pancreatic islet glucose-dependent insulinotropic polypeptide receptor: ligand
701 binding and intracellular signaling properties. *Endocrinology* 1995;136:4629–9.
- 702 31. Zhou J, Livak M, Bernier M, Muller D, Carlson O, Elahi D, Maudsley S, Egan J. Ubiquitination
703 is involved in glucose-mediated downregulation of GIP receptors in islets. *Am J Physiol*
Endocrinol Metab 2007;293:E538–47.
- 704 32. Yada T, Sakurada M, Ihida K, Nakata M, Murata F, Arimura A, Kikuchi M. Pituitary adeny-
705 late cyclase activating polypeptide is an extraordinarily potent intra-pancreatic regulator of
706 insulin secretion from islet beta-cells. *J Biol Chem* 1994;269:1290–3.
- 707 33. Ahrén B. Role of pituitary adenylate cyclase-activating polypeptide in the pancreatic
708 endocrine system. *Ann N Y Acad Sci*. 2008;1144:28–35.
- 709 34. Borboni P, Porzio O, Pierucci D, Cicconi S, Magnaterra R, Federici M, Sesti G, Lauro
710 D, D'Agata V, Cavallaro S, Marlier L. Molecular and functional characterization of pitu-
711 itary adenylate cyclase-activating polypeptide (PACAP-38)/vasoactive intestinal polypeptide
712 receptors in pancreatic beta-cells and effects of PACAP-38 on components of the insulin
secretory system. *Endocrinology* 1999;140:5530–7.
- 713 35. Yamada S, Komatsu M, Sato Y, Yamauchi K, Kojima I, Aizawa T, Hashizume K. Time-
714 dependent stimulation of insulin exocytosis by 3',5'-cyclic adenosine monophosphate in the
715 rat islet beta-cell. *Endocrinology* 2002;143:4203–9.
- 716 36. Soga T, Ohishi T, Matsui T, Saito T, Matsumoto M, Takasaki J, Matsumoto S, Kamohara
717 M, Hiyama H, Yoshida S, Momose K, Ueda Y, Matsushima H, Kobori M, Furuichi
718 K. Lysophosphatidylcholine enhances glucose-dependent insulin secretion via an orphan
G-protein-coupled receptor. *Biochem Biophys Res Commun* 2005;326:744–51.
- 719 37. Overton H, Babbs A, Doel S, Fyfe M, Gardner L, Griffin G, Jackson H, Procter M,
720 Rasamison C, Tang-Christensen M, Widdowson P, Williams G, Reynet C. Deorphanization

- of a G protein-coupled receptor for oleylethanolamide and its use in the discovery of small-molecule hypophagic agents. *Cell Metab* 2006;3:167–75.
38. Chu Z, Jones R, He H, Carroll C, Gutierrez V, Lucman A, Moloney M, Gao H, Mondala H, Bagnol D, Unett D, Liang Y, Demarest K, Semple G, Behan D, Leonard J. A role for beta-cell-expressed G protein-coupled receptor 119 in glycemic control by enhancing glucose-dependent insulin release. *Endocrinology* 2007;148:2601–9.
39. Hanoune J, Defer N. Regulation and role of adenylyl cyclase isoforms. *Annu Rev Pharmacol* 2001;Toxicol.;41:145–74.
40. Willoughby D, Cooper D. Organization and Ca²⁺ regulation of adenylyl cyclases in cAMP microdomains. *Physiol Rev* 2007;87:965–1010.
41. Kamenetsky M, Middelhaufe S, Bank E, Levin L, Buck J, Steegborn C. Molecular details of cAMP generation in mammalian cells: a tale of two systems. *J Mol Biol* 2006;362:623–39.
42. Srinivasan M, Aalinkeel R, Song F, Lee B, Laychock S, Patel M. Adaptive changes in insulin secretion by islets from neonatal rats raised on a high-carbohydrate formula. *Am J Physiol Endocrinol Metab* 2000;279:E1347–57.
43. Leech C, Castonguay M, Habener J. Expression of adenylyl cyclase subtypes in pancreatic beta-cells. *Biochem Biophys Res Commun* 1999;254:703–6.
44. Guenifi A, Portela-Gomes G, Grimelius L, Efendić S, Abdel-Halim S. Adenylyl cyclase isoform expression in non-diabetic and diabetic Goto-Kakizaki (GK) rat pancreas. Evidence for distinct overexpression of type-8 adenylyl cyclase in diabetic GK rat islets. *Histochem Cell Biol* 2000;113:81–9.
45. Delmeire D, Flamez D, Hinke S, Cali J, Pipeleers D, Schuit F. Type VIII adenylyl cyclase in rat beta cells: coincidence signal detector/generator for glucose and GLP-1. *Diabetologia* 2003;46:1383–93.
46. Seamon K, Daly J. Forskolin: its biological and chemical properties. *Adv Cyclic Nucleotide Protein Phosphorylation Res* 1986;20:1–150.
47. Insel P, Ostrom R. Forskolin as a tool for examining adenylyl cyclase expression, regulation, and G protein signaling. *Cell Mol Neurobiol* 2003;23:305–14.
48. Cali J, Zwaagstra J, Mons N, Cooper D, Krupinski J. Type VIII adenylyl cyclase. A Ca²⁺/calmodulin-stimulated enzyme expressed in discrete regions of rat brain. *J Biol Chem* 1994;269:12190–5.
49. Dolz M, Bailbé D, Giroix M, Calderari S, Gangnerau M, Serradas P, Rickenbach K, Irminger J, Portha B. Restitution of defective glucose-stimulated insulin secretion in diabetic GK rat by acetylcholine uncovers paradoxical stimulatory effect of beta-cell muscarinic receptor activation on cAMP production. *Diabetes* 2005;54:3229–37.
50. Perry M, Higgs G. Chemotherapeutic potential of phosphodiesterase inhibitors. *Curr Opin Chem Biol* 1998;2:472–81.
51. Soderling S, Beavo J. Regulation of cAMP and cGMP signaling. new phosphodiesterases and new functions. *Curr Opin Cell Biol* 2000;12:174–9.
52. Mehats C, Andersen C, Filopanti M, Jin S, Conti M. Cyclic nucleotide phosphodiesterases and their role in endocrine cell signaling. *Trends Endocrinol Metab* 2002;13:29–35.
53. Conti M, Beavo J. Biochemistry and physiology of cyclic nucleotide phosphodiesterases: essential components in cyclic nucleotide signaling. *Annu Rev Biochem* 2007;76:481–511.
54. Pyne N, Furman B. Cyclic nucleotide phosphodiesterases in pancreatic islets. *Diabetologia* 2003;46:1179–89.
55. Furman B, Pyne N, Flatt P, O'Harte F. Targeting beta-cell cyclic 3'5' adenosine monophosphate for the development of novel drugs for treating type 2 diabetes mellitus. A review. *J Pharm Pharmacol* 2004;56:1477–92.
56. Sugden M, Ashcroft S. Cyclic nucleotide phosphodiesterase of rat pancreatic islets. Effects of Ca²⁺, calmodulin and trifluoperazine. *Biochem J* 1981;197:459–64.
57. Capito K, Hedekov C, Thams P. Cyclic AMP phosphodiesterase activity in mouse pancreatic islets. Effects of calmodulin and phospholipids. *Acta Endocrinol (Copenh)* 1986;111:533–38.

- 766 58. Lipson L, Oldham S. The role of calmodulin in insulin secretion: the presence of a
767 calmodulin-stimulatable phosphodiesterase in pancreatic islets of normal and pregnant rats.
768 *Life Sci* 1983;32:775–80.
- 769 59. Han P, Werber J, Surana M, Fleischer N, Michaeli T. The calcium/calmodulin-dependent
770 phosphodiesterase PDE1C down-regulates glucose-induced insulin secretion. *J Biol Chem*
771 1999;274:22337–44.
- 772 60. Ahmad M, Flatt P, Furman B, Pyne N. The role of the cyclic GMP-inhibited cyclic AMP-
773 specific phosphodiesterase (PDE3) in regulating clonal BRIN-BD11 insulin secreting cell
774 survival. *Cell Signal* 2000;12:541–48.
- 775 61. Shafiee-Nick R, Pyne N, Furman B. Effects of type-selective phosphodiesterase inhibitors
776 on glucose-induced insulin secretion and islet phosphodiesterase activity. *Br J Pharmacol*
777 1995;115:1486–92.
- 778 62. Parker J, VanVolkenburg M, Ketchum R, Brayman K, Andrews K. Cyclic AMP phospho-
779 diesterases of human and rat islets of Langerhans: contributions of types III and IV to the
780 modulation of insulin secretion. *Biochem Biophys Res Commun* 1995;217:916–23.
- 781 63. Zhao A, Zhao H, Teague J, Fujimoto W, Beavo J. Attenuation of insulin secretion by insulin-
782 like growth factor 1 is mediated through activation of phosphodiesterase 3B. *Proc Natl Acad Sci U S A*
783 1997;94:3223–28.
- 784 64. Waddleton D, Wu W, Feng Y, Thompson C, Wu M, Zhou Y, Howard A, Thornberry N, Li
785 J, Mancini J. Phosphodiesterase 3 and 4 comprise the major cAMP metabolizing enzymes
786 responsible for insulin secretion in INS-1 (832/13) cells and rat islets. *Biochem Pharmacol*
787 2008;76:884–93.
- 788 65. Härndahl L, Jing X, Ivarsson R, Degerman E, Ahrén B, Manganiello V, Renström E, Holst
789 L. Important role of phosphodiesterase 3B for the stimulatory action of cAMP on pancreatic
790 beta-cell exocytosis and release of insulin. *J Biol Chem* 2002;277:37446–55.
- 791 66. Härndahl L, Wierup N, Enerbäck S, Mulder H, Manganiello V, Sundler F, Degerman E,
792 Ahrén B, Holst L. Beta-cell-targeted overexpression of phosphodiesterase 3B in mice causes
793 impaired insulin secretion, glucose intolerance, and deranged islet morphology. *J Biol Chem*
794 2004;279:15214–22.
- 795 67. Walz H, Härndahl L, Wierup N, Zmuda-Trzebiatowska E, Svennelid F, Manganiello V, Ploug
796 T, Sundler F, Degerman E, Ahrén B, Holst L. Early and rapid development of insulin resistance,
797 islet dysfunction and glucose intolerance after high-fat feeding in mice overexpressing
798 phosphodiesterase 3B. *J Endocrinol* 2006;189:629–41.
- 799 68. Dov A, Abramovitch E, Warwar N, Neshet R. Diminished phosphodiesterase-8B potentiates
800 biphasic insulin response to glucose. *Endocrinology* 2008;149:741–8.
- 801 69. Zhao A, Bornfeldt K, Beavo J. Leptin inhibits insulin secretion by activation of phosphodi-
802 esterase 3B. *J Clin Invest* 1998;102:869–73.
- 803 70. Grapenjiesser E, Gylfe E, Dansk H, Hellman B. Nitric oxide induces synchronous Ca^{2+}
804 transients in pancreatic beta cells lacking contact. *Pancreas* 2001;23:387–92.
- 805 71. Smukler S, Tang L, Wheeler M, Salapatek A. Exogenous nitric oxide and endoge-
806 nous glucose-stimulated beta-cell nitric oxide augment insulin release. *Diabetes* 2002;51:
807 3450–60.
- 808 72. Kaneko Y, Ishikawa T, Amano S, Nakayama K. Dual effect of nitric oxide on cytosolic Ca^{2+}
809 concentration and insulin secretion in rat pancreatic beta-cells. *Am J Physiol Cell Physiol*
810 2003;284:C1215–22.
73. Sunouchi T, Suzuki K, Nakayama K, Ishikawa T. Dual effect of nitric oxide on ATP-sensitive
 K^+ channels in rat pancreatic beta cells. *Pflugers Arch* 2008;456:573–9.
74. Cantin L, Magnuson S, Gunn D, Barucci N, Breuhaus M, Bullock W, Burke J, Claus T, Daly
M, Decarr L, Gore-Willse A, Hoover-Litty H, Kumarasinghe E, Li Y, Liang S, Livingston J,
Lowinger T, Macdougall M, Ogutu H, Olague A, Ott-Morgan R, Schoenleber R, Tersteegen
A, Wickens P, Zhang Z, Zhu J, Zhu L, Sweet L. PDE-10A inhibitors as insulin secretagogues.
Bioorg Med Chem Lett 2007;17:2869–73.
75. Fridlyand LE, Harbeck MC, Roe MW, Philipson LH. Regulation of cAMP dynamics by
 Ca^{2+} and G protein-coupled receptors in the pancreatic β -cell. a computational approach.
Am J Physiol Cell Physiol. 2007;293:C1924–33.

- 811 76. Dyachok O, Isakov Y, Sagetorp J, Tengholm A. Oscillations of cyclic AMP in hormone-
812 stimulated insulin-secreting beta-cells. *Nature* 2006;439:349–52.
- 813 77. Dyachok O, Sagetorp J, Isakov Y, Tengholm A. cAMP oscillations restrict protein kinase A
814 redistribution in insulin-secreting cells. *Biochem Soc Trans* 2006;34:498–501.
- 815 78. Dyachok O, Idevall-Hagren O, Sagetorp J, Tian G, Wuttke A, Arriemerlou C, Akusjarvi G,
816 Gylfe E, Tengholm A. Glucose-induced cyclic AMP oscillations regulate pulsatile insulin
secretion. *Cell Metab* 2008;8:26–37.
- 817 79. Baltrusch S, Lenzen S. Regulation of $[Ca^{2+}]_i$ oscillations in mouse pancreatic islets by
818 adrenergic agonists. *Biochem Biophys Res Commun* 2007;363:1038–43.
- 819 80. Jarnaess E, Tasken K. Spatiotemporal control of cAMP signalling processes by anchored
820 signalling complexes. *Biochem Soc Trans* 2007;35:931–37.
- 821 81. Dodge-Kafka K, Kapiloff M. The mAKAP signaling complex: integration of cAMP,
822 calcium, and MAP kinase signaling pathways. *Eur J Cell Biol* 2006;85:593–602.
- 823 82. Fraser I, Tavalin S, Lester L, Langeberg L, Westphal A, Dean R, Marrion N, Scott J. A novel
824 lipid-anchored A-kinase Anchoring Protein facilitates cAMP-responsive membrane events.
825 *EMBO J* 1998;17:2261–72.
- 826 83. Faruque O, Le-Nguyen D, Lajoix A, Vives E, Petit P, Bataille D, Hani e-H. Cell-permeable
827 peptide-based disruption of endogenous PKA-AKAP complexes: a tool for studying the
828 molecular roles of AKAP-mediated PKA subcellular anchoring. *Am J Physiol Cell Physiol*
829 2009;296:C306–16.
- 830 84. Jones PM, Persaud SJ. Protein kinases, protein phosphorylation, and the regulation of insulin
831 secretion from pancreatic β -cells. *Endocr Rev.* 1998;429–461.
- 832 85. Lester LB, Faux MC, Nauert JB, Scott JD. Targeted protein kinase A and PP-2B regulate
833 insulin secretion through reversible phosphorylation. *Endocrinology.* 2001;142(3):1218–27.
- 834 86. Kopperud R, Krakstad C, Selheim F, Doskeland S. cAMP effector mechanisms. Novel twists
835 for an 'old' signaling system. *FEBS Lett* 2003;546:121–6.
- 836 87. Renstrom E, Eliasson L, Rorsman P. Protein kinase A-dependent and independent stimula-
837 tion of exocytosis by cAMP in mouse pancreatic B-cells. *J Physiol* 1997;502:105–18.
- 838 88. Seino S, Shibasaki T. PKA-dependent and PKA-independent pathways for cAMP-regulated
839 exocytosis. *Physiol Rev* 2005;85:1303–42.
- 840 89. Shibasaki T, Takahashi H, Miki T, Sunaga Y, Matsumura K, Yamanaka M, Zhang C,
841 Tamamoto A, Satoh T, Miyazaki J, Seino S. Essential role of Epac2/Rap1 signaling in regula-
842 tion of insulin granule dynamics by cAMP. *Proc Natl Acad Sci U S A;* 2007;104:19333–193.
- 843 90. Holz G. Epac: A new cAMP-binding protein in support of glucagon-like peptide-1 receptor-
844 mediated signal transduction in the pancreatic beta-cell. *Diabetes* 2004;53:5–13.
- 845 91. Kashima Y, Miki T, Shibasaki T, Ozaki N, Miyazaki M, Yano H, Seino S. Critical role
846 of cAMP-GEFII—Rim2 complex in incretin-potentiated insulin secretion. *J Biol Chem*
847 2001;276:46046–53.
- 848 92. Chepurny O, Leech C, Kelley G, Dzhura I, Dzhura E, Li X, Rindler M, Schwede F,
849 Genieser H, Holz G. Enhanced Rap1 activation and insulin secretagogue properties of an
850 acetoxymethyl ester of an Epac-selective cyclic AMP analog in rat INS-1 cells: Studies with
851 8-pCPT-2'-O-Me-cAMP-AM. *J Biol Chem*, 2009.
- 852 93. Liu G, Jacobo S, Hilliard N, Hockerman G. Differential modulation of Cav1.2 and Cav1.3-
853 mediated glucose-stimulated insulin secretion by cAMP in INS-1 cells: distinct roles for
854 exchange protein directly activated by cAMP 2 (Epac2) and protein kinase A. *J Pharmacol*
855 *Exp Ther* 2006;318:152–60.
94. Nakazaki M, Crane A, Hu M, Seghers V, Ullrich S, Aguilar-Bryan L, Bryan J. cAMP-
activated protein kinase-independent potentiation of insulin secretion by cAMP is impaired
in SUR1 null islets. *Diabetes* 2002;51:3440–9.
95. Eliasson L, Ma X, Renstrom E, Barg S, Berggren P, Galvanovskis J, Gromada J, Jing X,
Lundquist I, Salehi A, Sewing S, Rorsman P. SUR1 regulates PKA-independent cAMP-
induced granule priming in mouse pancreatic B-cells. *J Gen Physiol* 2003;121:181–97.
96. Malaisse W, Malaisse-Lagae F, Mayhew D. A possible role for the adenyl cyclase system in
insulin secretion. *J Clin Invest* 1967;46:1724–34.

- 856 97. Brisson G, Malaisse-Lagae F, Malaisse W. The stimulus-secretion coupling of glucose-
857 induced insulin release. VII. A proposed site of action for adenosine-3',5'-cyclic monophos-
858 phate. *J Clin Invest* 1972;51:232-41.
- 859 98. Holz G, Kühtreiber W, Habener J. Pancreatic beta-cells are rendered glucose-competent by
860 the insulinotropic hormone glucagon-like peptide-1(7-37). *Nature* 1993;361:362-5.
- 861 99. Gromada J, Bokvist K, Ding W, Holst J, Nielsen J, Rorsman P. Glucagon-like peptide 1 (7-
862 36) amide stimulates exocytosis in human pancreatic beta-cells by both proximal and distal
863 regulatory steps in stimulus-secretion coupling. *Diabetes* 1998;47:57-65.
- 864 100. He L, Mears D, Atwater I, Kitasato H. Glucagon induces suppression of ATP-sensitive K⁺
865 channel activity through a Ca²⁺/calmodulin-dependent pathway in mouse pancreatic β-cells.
866 *J Membr Biol* 1998;166:237-44.
- 867 101. Light P, Manning Fox J, Riedel M, Wheeler M. Glucagon-like peptide-1 inhibits pancreatic
868 ATP-sensitive potassium channels via a protein kinase A- and ADP-dependent mechanism.
869 *Mol Endocrinol* 2002;16:2135-44.
- 870 102. Kang G, Chepurny O, Malester B, Rindler M, Rehmann H, Bos J, Schwede F, Coetzee W,
871 Holz G. cAMP sensor Epac as a determinant of ATP-sensitive potassium channel activity in
872 human pancreatic β cells and rat INS-1 cells. *J Physiol* 2006;573:595-609.
- 873 103. Kang G, Leech C, Chepurny O, Coetzee W, Holz G. Role of the cAMP sensor Epac as a
874 determinant of KATP channel ATP sensitivity in human pancreatic β-cells and rat INS-1
875 cells. *J Physiol* 2008;586:1307-19.
- 876 104. Kim S, Choi W, Han J, Warnock G, Fedida D, McIntosh C. A novel mechanism for
877 the suppression of a voltage-gated potassium channel by glucose-dependent insulinotropic
878 polypeptide: protein kinase A-dependent endocytosis. *J Biol Chem* 2005;280:28692-700.
- 879 105. MacDonald P, Salapatek A, Wheeler M. Glucagon-like peptide-1 receptor activation antag-
880 onizes voltage-dependent repolarizing K⁺ currents in β-cells: a possible glucose-dependent
881 insulinotropic mechanism. *Diabetes* 51 Suppl 2002;3:S443-47.
- 882 106. MacDonald P, Wang X, Xia F, El-Kholy W, Targonsky E, Tushima R, Wheeler M.
883 Antagonism of rat β-cell voltage-dependent K⁺ currents by exendin 4 requires dual activa-
884 tion of the cAMP/protein kinase A and phosphatidylinositol 3-kinase signaling pathways.
885 *J Biol Chem* 2003;278:52446-53.
- 886 107. Ammälä C, Ashcroft F, Rorsman P. Calcium-independent potentiation of insulin release by
887 cyclic AMP in single beta-cells. *Nature* 1993;363:356-58.
- 888 108. Kanno T, Suga S, Wu J, Kimura M, Wakui M. Intracellular cAMP potentiates voltage-
889 dependent activation of L-type Ca²⁺ channels in rat islet beta-cells. *Pflugers Arch*
890 1998;435:578-80.
- 891 109. Suga S, Kanno T, Nakano K, Takeo T, Dobashi Y, Wakui M. GLP-I (7-36) amide augments
892 Ba²⁺ current through L-type Ca²⁺ channel of rat pancreatic β-cell in a cAMP-dependent
893 manner. *Diabetes* 1997;46:1755-60.
- 894 110. Leiser M, Fleischer N. cAMP-dependent phosphorylation of the cardiac-type alpha 1 sub-
895 unit of the voltage-dependent Ca²⁺ channel in a murine pancreatic β-cell line. *Diabetes*
896 1996;45:1412-8.
- 897 111. Gromada J, Dissing S, Bokvist K, Renström E, Frøkjær-Jensen J, Wulff B, Rorsman P.
898 Glucagon-like peptide I increases cytoplasmic calcium in insulin-secreting beta TC3-cells
899 by enhancement of intracellular calcium mobilization. *Diabetes* 1995;44:767-74.
- 900 112. Islam M, Leibiger I, Leibiger B, Rossi D, Sorrentino V, Ekström T, Westerblad H, Andrade
F, Berggren P. In situ activation of the type 2 ryanodine receptor in pancreatic β cells requires
cAMP-dependent phosphorylation. *Proc Natl Acad Sci U S A* 1998;95:6145-50.
113. Holz G, Leech C, Heller R, Castonguay M, Habener J. cAMP-dependent mobilization of
intracellular Ca²⁺ stores by activation of ryanodine receptors in pancreatic beta-cells. A Ca²⁺
signaling system stimulated by the insulinotropic hormone glucagon-like peptide-1(7-37).
J Biol Chem 1999;274:14147-56.
114. Kang G, Chepurny O, Holz G. cAMP-regulated guanine nucleotide exchange factor
II (Epac2) mediates Ca²⁺-induced Ca²⁺ release in INS-1 pancreatic β-cells. *J Physiol*
2001;536:375-85.

- 901 115. Kang G, Joseph J, Chepurny O, Monaco M, Wheeler M, Bos J, Schwede F, Genieser H, Holz
902 G. Epac-selective cAMP analog 8-pCPT-2'-O-Me-cAMP as a stimulus for Ca^{2+} -induced
903 Ca^{2+} release and exocytosis in pancreatic β -cells. *J Biol Chem* 2003;278:8279–85.
- 904 116. Kang G, Chepurny O, Rindler M, Collis L, Chepurny Z, Li W, Harbeck M, Roe M, Holz G.
905 A cAMP and Ca^{2+} coincidence detector in support of Ca^{2+} -induced Ca^{2+} release in mouse
906 pancreatic β cells. *J Physiol* 2005;566:173–88.
- 907 117. Bode H, Moormann B, Dabew R, Göke B. Glucagon-like peptide 1 elevates cytosolic
908 calcium in pancreatic beta-cells independently of protein kinase A. *Endocrinology*
909 1999;140:3919–27.
- 910 118. Tsuboi T, da Silva Xavier G, Holz G, Jouaville L, Thomas A, Rutter G. Glucagon-like
911 peptide-1 mobilizes intracellular Ca^{2+} and stimulates mitochondrial ATP synthesis in
912 pancreatic MIN6 β -cells. *Biochem J* 2003;369:287–99.
- 913 119. Dyachok O, Gylfe E. Ca^{2+} -induced Ca^{2+} release via inositol 1,4,5-trisphosphate receptors
914 is amplified by protein kinase A and triggers exocytosis in pancreatic β -cells. *J Biol Chem*
915 2004;279:45455–61.
- 916 120. Kim BJ, Park KH, Yim CY, Takasawa S, Okamoto H, Im MJ, Kim UH Generation of nicotinic
917 acid adenine dinucleotide phosphate and cyclic ADP-ribose by glucagon-like peptide-1
918 evokes Ca^{2+} signal that is essential for insulin secretion in mouse pancreatic islets. *Diabetes*
919 2008;57:868–78.
- 920 121. Gillis K, Mislis S. Enhancers of cytosolic cAMP augment depolarization-induced exocytosis
921 from pancreatic B-cells: evidence for effects distal to Ca^{2+} entry. *Pflugers Arch*
922 1993;424:195–7.
- 923 122. Ding W, Gromada J. Protein kinase A-dependent stimulation of exocytosis in mouse pancreatic
924 beta-cells by glucose-dependent insulintropic polypeptide. *Diabetes* 1997;46:615–21.
- 925 123. Hatakeyama H, Kishimoto T, Nemoto T, Kasai H, Takahashi N. Rapid glucose sensing
926 by protein kinase A for insulin exocytosis in mouse pancreatic islets. *J Physiol* 2006;570:
927 271–82.
- 928 124. Hatakeyama H, Takahashi N, Kishimoto T, Nemoto T, Kasai H. Two cAMP-dependent
929 pathways differentially regulate exocytosis of large dense-core and small vesicles in mouse
930 β -cells. *J Physiol* 2007;582:1087–98.
- 931 125. Hashiguchi H, Nakazaki M, Koriyama N, Fukudome M, Aso K, Tei C. Cyclic AMP/cAMP-
932 GEF pathway amplifies insulin exocytosis induced by Ca^{2+} and ATP in rat islet beta-cells.
933 *Diabetes Metab Res Rev* 2006;22:64–71.
- 934 126. Kwan E, Gaisano H. Glucagon-like peptide 1 regulates sequential and compound exocytosis
935 in pancreatic islet β -cells. *Diabetes* 2005;54:2734–43.
- 936 127. Kwan E, Xie L, Sheu L, Ohtsuka T, Gaisano H. Interaction between Munc13-1 and RIM
937 is critical for glucagon-like peptide-1 mediated rescue of exocytotic defects in Munc13-1
938 deficient pancreatic beta-cells. *Diabetes* 2007;56:2579–88.
- 939 128. Kwan E, Gao X, Leung Y, Gaisano H. Activation of exchange protein directly activated by
940 cyclic adenosine monophosphate and protein kinase A regulate common and distinct steps in
941 promoting plasma membrane exocytic and granule-to-granule fusions in rat islet beta cells.
942 *Pancreas* 2007;35:e45–54.
- 943 129. Suzuki Y, Zhang H, Saito N, Kojima I, Urano T, Mogami H. Glucagon-like peptide
944 1 activates protein kinase C through Ca^{2+} -dependent activation of phospholipase C in
945 insulin-secreting cells. *J Biol Chem* 2006;281:28499–507.
- 946 130. Chepurny O, Hussain M, Holz G. Exendin-4 as a stimulator of rat insulin I gene promoter
947 activity via bZIP/CRE interactions sensitive to serine/threonine protein kinase inhibitor Ro
948 31-8220. *Endocrinology* 2002;143:2303–13.
- 949 131. Xie T, Chen M, Zhang Q, Ma Z, Weinstein L. β -cell-specific deficiency of the stimulatory G
950 protein α -subunit $\text{G}_s\alpha$ leads to reduced β -cell mass and insulin-deficient diabetes. *Proc Natl
951 Acad Sci U S A.* 2007;104:19601–6.
- 952 132. Elrick L, Docherty K. Phosphorylation-dependent nucleocytoplasmic shuttling of pancreatic
953 duodenal homeobox-1. *Diabetes* 2001;50:2244–52.

- 946 133. Wang X, Zhou J, Doyle M, Egan J. Glucagon-like peptide-1 causes pancreatic duodenal
947 homeobox-1 protein translocation from the cytoplasm to the nucleus of pancreatic β -cells by
948 a cyclic adenosine monophosphate/protein kinase A-dependent mechanism. *Endocrinology*
2001;142:1820–27.
- 949 134. Song W, Schreiber W, Zhong E, Liu F, Kornfeld B, Wondisford F, Hussain M. Exendin-4
950 stimulation of cyclin A2 in β -cell proliferation. *Diabetes* 2008;57:2371–81.
- 951 135. Kim S, Nian C, Widenmaier S, McIntosh C. Glucose-dependent insulinotropic polypeptide-
952 mediated up-regulation of beta-cell antiapoptotic Bcl-2 gene expression is coordinated
953 by cyclic AMP (cAMP) response element binding protein (CREB) and cAMP-responsive
954 CREB coactivator 2. *Mol Cell Biol* 2008;28:1644–56.
- 955 136. Jansson D, Ng A, Fu A, Depatie C, Al Azzabi M, Screaton R. Glucose controls CREB
956 activity in islet cells via regulated phosphorylation of TORC2. *Proc Natl Acad Sci U S A*
2008;105:10161–66.
- 957 137. Jonas J, Laybutt D, Steil G, Trivedi N, Pertusa J, Van de Castele M, Weir G, Henquin J.
958 High glucose stimulates early response gene c-Myc expression in rat pancreatic beta cells. *J*
959 *Biol Chem* 2001;276:35375–81.
- 960 138. Susini S, Roche E, Prentki M, Schlegel W. Glucose and glucocorticoid peptides synergize
961 to induce c-fos, c-jun, junB, zif-268, and nur-77 gene expression in pancreatic beta(INS-1)
962 cells. *FASEB J* 1998;12:1173–82.
- 963 139. Glauser D, Brun T, Gauthier B, Schlegel W. Transcriptional response of pancreatic beta
964 cells to metabolic stimulation: large scale identification of immediate-early and secondary
965 response genes. *BMC Mol Biol* 2007;8:54.
- 966 140. Frödin M, Sekine N, Roche E, Filloux C, Prentki M, Wollheim C, Van Obberghen E.
967 Glucose, other secretagogues, and nerve growth factor stimulate mitogen-activated protein
968 kinase in the insulin-secreting beta-cell line, INS-1. *J Biol Chem* 1995;270:7882–89.
- 969 141. Benes C, Roisin M, Van Tan H, Creuzet C, Miyazaki J, Fagard R. Rapid activation
970 and nuclear translocation of mitogen-activated protein kinases in response to physio-
971 logical concentration of glucose in the MIN6 pancreatic beta cell line. *J Biol Chem*
972 1998;273:15507–13.
- 973 142. Benes C, Poitout V, Marie J, Martin-Perez J, Roisin M, Fagard R. Mode of regulation of
974 the extracellular signal-regulated kinases in the pancreatic beta-cell line MIN6 and their
975 implication in the regulation of insulin gene transcription. *Biochem J* 1999;340 (Pt 1):
976 219–25.
- 977 143. Gomez E, Pritchard C, Herbert T. cAMP-dependent protein kinase and Ca^{2+} influx through
978 L-type voltage-gated calcium channels mediate Raf-independent activation of extracellular
979 regulated kinase in response to glucagon-like peptide-1 in pancreatic β -cells. *J Biol Chem*
2002;277:48146–51.
- 980 144. Ehses J, Pelech S, Pederson R, McIntosh C. Glucose-dependent insulinotropic polypep-
981 tide activates the Raf-Mek1/2-ERK1/2 module via a cyclic AMP/cAMP-dependent protein
982 kinase/Rap1-mediated pathway. *J Biol Chem* 2002;277:37088–97.
- 983 145. Drucker D. Glucagon-like peptides: regulators of cell proliferation, differentiation, and
984 apoptosis. *Mol Endocrinol* 2003;17:161–71.
- 985 146. Hui H, Nourparvar A, Zhao X, Perfetti R. Glucagon-like peptide-1 inhibits apoptosis of
986 insulin-secreting cells via a cyclic 5'-adenosine monophosphate-dependent protein kinase A-
987 and a phosphatidylinositol 3-kinase-dependent pathway. *Endocrinology* 2003;144:1444–55.
- 988 147. Ehses J, Casilla V, Doty T, Pospisilik J, Winter K, Demuth H, Pederson R, McIntosh C.
989 Glucose-dependent insulinotropic polypeptide promotes beta-(INS-1) cell survival via cyclic
990 adenosine monophosphate-mediated caspase-3 inhibition and regulation of p38 mitogen-
activated protein kinase. *Endocrinology* 2003;144:4433–45.
148. Ranta F, Avram D, Berchtold S, Düfer M, Drews G, Lang F, Ullrich S. Dexamethasone
induces cell death in insulin-secreting cells, an effect reversed by exendin-4. *Diabetes*
2006;55:1380–90.
149. Granata R, Settanni F, Biancone L, Trovato L, Nano R, Bertuzzi F, Destefanis S, Annunziata
M, Martinetti M, Catapano F, Ghè C, Isgaard J, Papotti M, Ghigo E, Muccioli G. Acylated

- 991 and unacylated ghrelin promote proliferation and inhibit apoptosis of pancreatic beta-cells
992 and human islets: involvement of 3',5'-cyclic adenosine monophosphate/protein kinase A,
993 extracellular signal-regulated kinase 1/2, and phosphatidylinositol 3-Kinase/Akt signaling.
994 *Endocrinology* 2007;148:512–29.
- 995 150. Granata R, Settanni F, Gallo D, Trovato L, Biancone L, Cantaluppi V, Nano R, Annunziata
996 M, Campiglia P, Arnoletti E, Ghè C, Volante M, Papotti M, Muccioli G, Ghigo E. Obestatin
997 promotes survival of pancreatic β -cells and human islets and induces expression of genes
998 involved in the regulation of β -cell mass and function. *Diabetes* 2008;57:967–79.
- 999 151. Ferdaoussi M, Abdelli S, Yang J, Cornu M, Niederhauser G, Favre D, Widmann C, Regazzi
1000 R, Thorens B, Waeber G, Abderrahmani A. Exendin-4 protects β -cells from interleukin-1
1001 beta-induced apoptosis by interfering with the c-Jun NH2-terminal kinase pathway. *Diabetes*
1002 2008;57:1205–15.
- 1003 152. Loweth A, Williams G, Scarpello J, Morgan N. Heterotrimeric G-proteins are implicated in
1004 the regulation of apoptosis in pancreatic β -cells. *Exp Cell Res* 1996;229:69–76.
- 1005 153. Ahmad M, Abdel-Wahab YH, Tate R, Flatt PR, Pyne NJ, Furman BL. Effect of type-
1006 selective inhibitors on cyclic nucleotide phosphodiesterase activity and insulin secretion in
1007 the clonal insulin secreting cell line BRIN-BD11. *Br J Pharmacol*. 2000;129:1228–34.
- 1008 154. Andersen H, Mauricio D, Karlsen A, Mandrup-Poulsen T, Nielsen J, Nerup J. Interleukin-1
1009 β -induced nitric oxide production from isolated rat islets is modulated by D-glucose and
1010 3-isobutyl-1-methyl xanthine. *Eur J Endocrinol* 1996;134:251–9.
- 1011 155. Friedrichsen B, Neubauer N, Lee Y, Gram V, Blume N, Petersen J, Nielsen J, Møldrup A.
1012 Stimulation of pancreatic β -cell replication by incretins involves transcriptional induction of
1013 cyclin D1 via multiple signalling pathways. *J Endocrinol* 2006;188:481–92.
- 1014 156. Klinger S, Poussin C, Debril M, Dolci W, Halban P, Thorens B. Increasing GLP-1-induced
1015 β -cell proliferation by silencing the negative regulators of signaling cAMP response element
1016 modulator-alpha and DUSP14. *Diabetes* 2008;57:584–93.
- 1017 157. Kim M, Kang J, Park Y, Ryu G, Ko S, Jeong I, Koh K, Rhie D, Yoon S, Hahn S, Kim
1018 M, Jo Y. Exendin-4 induction of cyclin D1 expression in INS-1 β -cells: involvement of
1019 cAMP-responsive element. *J Endocrinol* 2006;188:623–33.
- 1020 158. Welters H, Kulkarni R. Wnt signaling: relevance to β -cell biology and diabetes. *Trends*
1021 *Endocrinol Metab*. 2008;349–55.
- 1022 159. Liu Z, Habener J. Glucagon-like peptide-1 activation of TCF7L2-dependent Wnt signaling
1023 enhances pancreatic β cell proliferation. *J Biol Chem* 2008;283:8723–35.
- 1024 160. Hii C, Howell S. Role of second messengers in the regulation of glucagon secretion from
1025 isolated rat islets of Langerhans. *Mol Cell Endocrinol* 1987;50:37–44.
- 1026 161. Ding W, Renström E, Rorsman P, Buschard K, Gromada J. Glucagon-like peptide I and
1027 glucose-dependent insulinotropic polypeptide stimulate Ca^{2+} -induced secretion in rat α -cells
1028 by a protein kinase A-mediated mechanism. *Diabetes* 1997;46:792–800.
- 1029 162. Dillon J, Lu M, Bowen S, Homan L. The recombinant rat glucagon-like peptide-1 receptor,
1030 expressed in an α -cell line, is coupled to adenylyl cyclase activation and intracellular calcium
1031 release. *Exp Clin Endocrinol Diabetes* 2005;113:182–9.
- 1032 163. Islam D, Zhang N, Wang P, Li H, Brubaker P, Gaisano H, Wang Q, Jin T. Epac is involved
1033 in cAMP-stimulated proglucagon expression and hormone production but not hormone
1034 secretion in pancreatic α - and intestinal L-cell lines. *Am J Physiol Endocrinol Metab*
1035 2009;296:E174–81.
- 1036 164. Dunning B, Foley J, Ahren B. Alpha cell function in health and disease: influence of
1037 glucagon-like peptide-1. *Diabetologia* 2005;48:1700–13.
- 1038 165. Gromada J, Høy M, Buschard K, Salehi A, Rorsman P. Somatostatin inhibits exocytosis
1039 in rat pancreatic α -cells by G(i2)-dependent activation of calcineurin and depriming of
1040 secretory granules. *J Physiol* 2001;535:519–32.
- 1041 166. Gros L, Thorens B, Bataille D, Kervran A. Glucagon-like peptide-1-(7-36) amide, oxynto-
1042 modulin, and glucagon interact with a common receptor in a somatostatin-secreting cell line.
1043 *Endocrinology*. 1993;133:631–8.

- 1036 167. Fehmann H, Janssen M, Göke B. Interaction of glucagon-like peptide-I (GLP-I) and
1037 galanin in insulin (beta TC-1)- and somatostatin (RIN T3)-secreting cells and evidence
1038 that both peptides have no receptors on glucagon (INR1G9)-secreting cells. *Acta Diabetol*
1039 1995;32:176–81.
- 1040 168. Patel Y, Papachristou D, Zingg H, Farkas E. Regulation of islet somatostatin secretion and
1041 gene expression: selective effects of adenosine 3',5'-monophosphate and phorbol esters in
1042 normal islets of Langerhans and in a somatostatin-producing rat islet clonal cell line 1027
1043 B2. *Endocrinology* 1991;128:1754–62.
- 1044 169. Ma X, Zhang Y, Gromada J, Sewing S, Berggren P, Buschard K, Salehi A, Vikman J,
1045 Rorsman P, Eliasson L. Glucagon stimulates exocytosis in mouse and rat pancreatic α -cells
1046 by binding to glucagon receptors. *Mol Endocrinol* 2005;19:198–12.
- 1047 170. Gromada J, Bokvist K, Ding WG, Barg S, Buschard K, Renström E, Rorsman P. Adrenaline
1048 stimulates glucagon secretion in pancreatic A-cells by increasing the Ca^{2+} current and the
1049 number of granules close to the L-type Ca^{2+} channels. *J Gen Physiol* 1997;110:217–28.
- 1050 171. Vieira E, Liu Y, Gylfe E. Involvement of alpha1 and beta-adrenoceptors in adrenaline stimulation
1051 of the glucagon-secreting mouse alpha-cell. *Naunyn Schmiedebergs Arch Pharmacol*
1052 2004;369:179–83.
- 1053 172. Knudsen L, Kiel D, Teng M, Behrens C, Bhumralkar D, Kodra J, Holst J, Jeppesen C,
1054 Johnson M, de Jong J, Jorgensen A, Kercher T, Kostrowicki J, Madsen P, Olesen P, Petersen
1055 J, Poulsen F, Sidelmann U, Sturis J, Truesdale L, May J, Lau J. Small-molecule agonists for
1056 the glucagon-like peptide 1 receptor. *Proc Natl Acad Sci U S A* 104: 2007;937–42.
- 1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080