

Strathprints Institutional Repository

McGinty, Sean and Mckee, Sean (2011) *A comment on pulsatile pipe flow*. Applied Mathematical and Computational Sciences, 3 (2). pp. 217-225. ISSN 0976-1586

Strathprints is designed to allow users to access the research output of the University of Strathclyde. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. You may not engage in further distribution of the material for any profitmaking activities or any commercial gain. You may freely distribute both the url (<http://strathprints.strath.ac.uk/>) and the content of this paper for research or study, educational, or not-for-profit purposes without prior permission or charge.

Any correspondence concerning this service should be sent to Strathprints administrator: <mailto:strathprints@strath.ac.uk>

A comment on pulsatile pipe flow

S. M c G I N T Y and S. M c K E E

Department of Mathematics,
University of Strathclyde, Glasgow, G1 1XH, Scotland.
smck@maths.strath.ac.uk

This article is concerned with analytic solutions of flows in cylindrical and annular pipes subject to an arbitrary time dependent pressure gradient and arbitrary initial flow.

Consider unsteady flow within a circular pipe driven by a time-dependent pressure gradient. The flow is unidirectional with $u(r, t)$ in the axial direction satisfying

$$\frac{\partial u}{\partial t} = g(t) + \nu \left(\frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} \right), \quad (1)$$

with

$$u(a, t) = 0 \quad ; \quad u(0, t) \text{ bounded, all } t. \quad (2)$$

Here ν , ρ and a are respectively the kinematic viscosity, the density and the pipe radius and $\rho g(t) = -\partial p / \partial z$, with $-\partial p / \partial z$ the pressure gradient (z is the axial direction). This problem has a long history dating back to Sexl (1928), Lambossy (1952), Womersley (1955 a,b) and Uchida (1956). Other authors who have worked on closely related problems including pressure gradients which are impulses, exponentially decreasing functions etc. include Syzmanski (1932), Ito (1953), Lance (1956), Sanyal (1956), Verma (1960) and Smith (1997). Drazin and Riley (2006) provide the solution

$$u(r, t) = \frac{P}{4\mu} (a^2 - r^2) - \operatorname{Re} \left\{ \frac{ic}{\omega\rho} \left(1 - \frac{J_0(i^{3/2}kr)}{J_0(i^{3/2}ka)} \right) e^{i\omega t} \right\}, \quad k^2 = \frac{\omega}{\nu}, \quad (3)$$

where, here,

$$\frac{\partial p}{\partial z} = -P - \operatorname{Re} \{ c e^{i\omega t} \}, \quad c = c_1 - ic_2 \quad (4)$$

which is essentially the form given by Sexl (1928), Uchida (1956) and Womersley (1955a). This form is ubiquitous in the literature and yet it is unsatisfactory as it assumes a very specific initial condition. The problem one wishes to solve is (1) with (2) together with an arbitrary initial condition, let us say

$$u(r, 0) = f(r). \quad (5)$$

In fact it is relatively straightforward (see Appendix) to write down a solution to (1) subject to (2) and (5):

$$u(r, t) = 2 \sum_{n=1}^{\infty} \frac{J_0(\alpha_n r) e^{-\nu \alpha_n^2 t}}{a J_1(\alpha_n a)} \left\{ \frac{\int_0^a r f(r) J_0(\alpha_n r) dr}{a J_1(\alpha_n a)} + \frac{\int_0^t e^{\nu \alpha_n^2 t'} g(t') dt'}{\alpha_n} \right\}, \quad (6)$$

where $\alpha_n, n = 1, 2, \dots$, are the countably infinite roots of $J_0(\alpha_n a) = 0$.

The volume flow rate is given by

$$Q = 4\pi \sum_{n=1}^{\infty} \frac{e^{-\nu\alpha_n^2 t}}{\alpha_n} \left\{ \frac{\int_0^a r f(r) J_0(\alpha_n r) dr}{a J_1(\alpha_n a)} + \frac{\int_0^t e^{\nu\alpha_n^2 t'} g(t') dt'}{\alpha_n} \right\}. \quad (7)$$

We observe immediately that we have a solution for any initial flow and time-dependent pressure gradient, with no possible singularities: $J_1(\alpha_n a)$ is never zero since the zeros of $J_0(z)$ and $J_1(z)$ interlace. We note that when the flow is initially quiescent, we obtain the same expression for the flow as Szymanski (1932) and, additionally, if $\partial p/\partial z = \text{constant}$, then as $t \rightarrow \infty$ we retrieve Poiseuille flow, since

$$a^2 - r^2 = \frac{8}{a} \sum_{n=1}^{\infty} \frac{J_0(\alpha_n r)}{\alpha_n^3 J_1(\alpha_n a)}. \quad (8)$$

Of course, because of the generality of the above expressions, flows subject to impulsive pressure gradients and gradients subject to exponential decay, for example, are easily derived from (6): indeed, the solution obtained by Ito (1953) for a linear pressure gradient can be shown to be a special case of (6).

We may also consider flow in an annular region subject to an arbitrary time-dependent pressure gradient where the outer radius is a while the inner radius is b ($b < a$). We obtain

$$u(r, t) = \pi \sum_{n=1}^{\infty} \frac{V_0(\alpha_n r) e^{-\nu\alpha_n^2 t}}{1 + X_0(\alpha_n)} \left\{ \frac{\pi\alpha_n^2 \int_b^a r f(r) V_0(\alpha_n r) dr}{2(1 - X_0(\alpha_n))} + \int_0^t e^{\nu\alpha_n^2 t'} g(t') dt' \right\} \quad (9)$$

where

$$V_0(\alpha_n r) = J_0(\alpha_n r) Y_0(\alpha_n b) - J_0(\alpha_n b) Y_0(\alpha_n r),$$

$$X_0(\alpha_n) = J_0(\alpha_n b) / J_0(\alpha_n a), \quad n = 1, 2, \dots,$$

where here α_n are the countably infinite zeros of $V_0(\alpha_n a) = 0$. The corresponding volumetric flow rate is

$$Q = 4\pi \sum_{n=1}^{\infty} \frac{e^{-\nu\alpha_n^2 t} (1 - X_0(\alpha_n))}{\alpha_n^2 (1 + X_0(\alpha_n))} \left\{ \frac{\pi\alpha_n^2 \int_b^a r f(r) V_0(\alpha_n r) dr}{2(1 - X_0(\alpha_n))} + \int_0^t e^{\nu\alpha_n^2 t'} g(t') dt' \right\}. \quad (10)$$

It is straightforward to show that as $b \rightarrow 0$ expressions (9) and (10) reduce to (6) and (7) respectively.

Appendix

The eigenfunctions of the related homogeneous problem satisfy

$$\frac{d}{dr} \left(r \frac{d\phi(r)}{dr} \right) + \alpha^2 r \phi(r) = 0, \quad (A 1)$$

$$\phi(a) = 0 \quad (A 2)$$

and admit the general solution

$$\phi(r) = A J_0(\alpha r) + B Y_0(\alpha r). \quad (A 3)$$

Boundedness and the no slip condition then imply, respectively, that $B = 0$ and

$$J_0(\alpha a) = 0, \quad (A 4)$$

yielding, through superposition, the general solution

$$u(r, t) = \sum_{n=1}^{\infty} a_n(t) J_0(\alpha_n r). \quad (\text{A } 5)$$

With $f(r) = \sum_{n=1}^{\infty} a_n(0) J_0(\alpha_n r)$ we deduce, using orthogonality, that

$$a_n(0) = \frac{2}{(J_1(\alpha_n a))^2} \int_0^a r f(r) J_0(\alpha_n r) dr. \quad (\text{A } 6)$$

Substitution of (A 5) into the inhomogeneous problem results in

$$\sum_{n=1}^{\infty} \left\{ \frac{da_n(t)}{dt} + \nu \alpha_n^2 a_n(t) \right\} J_0(\alpha_n r) = g(t). \quad (\text{A } 7)$$

From orthogonality we deduce that

$$a_n(t) = a_n(0) e^{-\nu \alpha_n^2 t} + \frac{2}{a \alpha_n J_1(\alpha_n a)} \int_0^t g(t') e^{-\nu \alpha_n^2 (t-t')} dt', \quad (\text{A } 8)$$

giving rise to the solution (6) provided in the main text. The derivation of (9) is similar except that the orthogonal functions employed are $V_0(\alpha_n r)$.

REFERENCES

- DRAZIN, P. & RILEY, N. 2006 *The Navier–Stokes Equations: A Classification of Flows and Exact Solutions*. Cambridge University Press.
- ITO, H. 1953 Theory of laminar flow through a pipe with non-steady pressure gradients. *Rep. Inst. High-speed Mech., Tohoku University*, No.3, 163–180.
- LAMBOSSY, P. 1952 Oscillations forcées d'un liquide incompressible et visqueux dans un tube rigide et horizontal. Calcul de la force de frottement. *Helv. Phys. Acta* **25**, 371–386.
- LANCE, G.N. 1956 Motion of a viscous fluid in a tube which is subjected to a series of pulses. *Quart. Appl. Math.* **14**, 312–315.
- SANYAL, L. 1956 The flow of viscous liquid in a circular tube under pressure gradients varying exponentially with time. *Indian J. Phys.* **30**, No.2, 57–61.
- SEXL, T. 1928 Über einige Integrale der für die achsensymmetrischen Strömungen in Röhren charakteristischen Differentialgleichung. *Ann. d. Physik.* **87** 570–580.
- SMITH, S. H. 1997 Time-dependent Poiseuille flow. *SIAM Rev.* **39**, 511–513.
- SZYMANSKI, P. 1932 Some exact solutions of the hydrodynamic equations of a viscous fluid in the case of a cylindrical tube. *J. Math. Pure Appl.* **11**, 67–101.
- UCHIDA, S. 1956 The pulsating viscous flow superimposed on the steady laminar motion of incompressible fluid in a circular pipe. *Z. Angew. Math. Phys.* **7**, 403–422.
- VERMA, P.D. 1960 The pulsating viscous flow superimposed on the steady laminar motion of incompressible fluid between two co-axial cylinders. *Proc. Natn. Inst. Sci. India* **26**, 447–458.
- WOMERSLEY, J.R. 1955a Method for the calculation of velocity, rate of flow and viscous drag in arteries when the pressure gradient is known. *J. Physiol.* **127**, 553–563.
- WOMERSLEY, J.R. 1955b Oscillating motion of a viscous liquid in a thin-walled elastic tube—I: The linear approximation for long waves. *Phil Mag.* **46**, No.373, 199–221.