



Strathprints Institutional Repository

Fawcett, J. and Moralee, A.C. and Percy, Jonathan and Salafia, Vittoria and Vincent, M.A. and Hillier, I.H. (2004) *A surprising steric effect on a tandem cycloaddition/ring-opening reaction : rapid syntheses of difluorinated analogues of (hydroxymethyl)conduritols*. Chemical Communications, 2004 (9). pp. 1062-1063. ISSN 1359-7345

Strathprints is designed to allow users to access the research output of the University of Strathclyde. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. You may not engage in further distribution of the material for any profitmaking activities or any commercial gain. You may freely distribute both the url (<http://strathprints.strath.ac.uk/>) and the content of this paper for research or study, educational, or not-for-profit purposes without prior permission or charge.

Any correspondence concerning this service should be sent to Strathprints administrator: <mailto:strathprints@strath.ac.uk>

A surprising steric effect on a tandem cycloaddition/ring-opening reaction: rapid syntheses of difluorinated analogues of (hydroxymethyl)conduritols†

John Fawcett,^a Andrew C. Moralee,^a Jonathan M. Percy,^{*a} Vittoria Salafia,^a Mark A. Vincent^b and Ian H. Hillier^b

^a Department of Chemistry, University of Leicester, University Road, Leicester, UK LE1 7RH.
E-mail: jmp29@le.ac.uk

^b Department of Chemistry, University of Manchester, Manchester, UK M13 9PL.
E-mail: Ian.Hillier@man.ac.uk

Received (in Cambridge, UK) 27th January 2004, Accepted 13th March 2004
First published as an Advance Article on the web 8th April 2004


Difluorinated analogues of (hydroxymethyl)conduritols can be synthesised from selected furans and a difluorinated dienophile in two reaction steps.

The literature contains many examples of fluorinated building blocks.¹ These are either commercially available compounds or readily prepared intermediates that can be manipulated using the important C–C bond-forming reactions of modern synthetic chemistry. Few, if any, of these building blocks allow the rapid synthesis of complex fluorine-containing molecules.

We identified cyclic carbonate **4** as a side product during the formation of *endo* and *exo* cycloadducts **2** and **3** via the furan Diels–Alder reaction of rare difluorinated dienophile **1** in the presence of sub-stoichiometric amounts of stannic chloride (Scheme 1).³

We were interested in this product, as it complements species we could obtain by either hydrostannylation/stannate ring opening according to Lautens⁴ or *via* procedures in which sulfur electrophiles and reductive desulfonation/ring-opening chemistry are used.⁵ Furthermore, though the yield is modest, the product arises in a single reaction step from cycloaddition, followed by highly controlled ring-opening, so the sequence is particularly concise. We therefore decided to investigate further and found that **4** (26%) replaced *exo*-**3** when the cycloaddition was carried out with stoichiometric Lewis acid.† However, non-aqueous work-up of a reaction solution led to the exclusive presence of **2** and **3**, as revealed by ¹⁹F NMR: we were only able to observe **4** after *aqueous* work-up. Substituted 2-methyl-, 2,3-dimethyl- and 2,5-dimethylfurans behaved quite differently, failing to afford cyclic carbonates. Instead, only *exo* cycloadducts were obtained and all attempts to force the reactions led to decomposition.


There is little in the literature that might help to predict how the Lewis acid would bind to **2/3**. A search of the literature revealed only 5- and 6-membered chelates involving stannic chloride. Denmark and Fu have obtained structures for bis-phosphoramidate ligand–stannic chloride complexes,⁶ though these include much larger rings. We therefore attempted to grow crystals of a complex between the reactive *exo*-**3** and stannic chloride. Diffusion of hexane into a DCM solution of **3** and stannic chloride at room temperature produced air-sensitive crystals of **7a** (Fig. 1), which were transferred rapidly to the cold diffractometer stage.§ Clearly,



Scheme 1 Furan Diels–Alder reaction of difluorinated dienophile **1**. (i) 25 mol% SnCl₄, furan (2 equiv.), DCM, rt.

† Electronic supplementary information (ESI) available: calculations and Cartesian coordinates for **3–7** (Me substituted for Et throughout); data for **8–10**. See <http://www.rsc.org/suppdata/cc/b4/b401245c/>

this cannot be an intermediate on a direct pathway to **4**. Scheme 2 presents our proposed mechanism for the formation of **4**.



Scheme 2 Proposed mechanism for the formation of cyclic carbonate **4**.

To trigger ring opening, stannic chloride must first bind to the bridging ether oxygen, which is a poor donor. To compensate for this, 6-membered chelate formation (in **5a**) involving carbonyl and ether oxygen donors, which is impossible from the *endo* cycloadduct, is proposed. Bridge cleavage, which is strain relieving, may then occur reversibly. The proposed intermediate **6a** is transformed into **4** when hydrolysis removes it from the equilibrium.⁷

We have used electronic structure calculations to investigate a number of aspects of this proposed mechanism.⁸ Minima and transition structures were calculated at the B3LYP/6-31G** level, including the effect of the solvent, DCM, using the polarisable continuum model (PCM). These calculations first showed that **5a** and **7a** differed in energy by only 21 kJ mol^{−1} in favour of **7a**, thus confirming the proposed equilibrium between these structures. The proposed intermediate **6a** was confirmed as an energy minimum and the transition structure linking it to reactant **5a** was located, the barrier being 80 kJ mol^{−1}. This transition structure showed a high degree of bond cleavage (C–O = 2.06 Å), suggesting that the effect of methylation at the bridgehead position is steric rather than electronic.† To investigate this further, the calculations were repeated for the methylated molecules. We now find **7b** to be more stable than **5b** by 42 kJ mol^{−1}, with the barrier to the formation of





Fig. 1 Crystal structure of air-sensitive complex **7a** formed between *exo*-**3** and stannic chloride.



Scheme 3 Diels-Alder reaction of **1** with 3-bromofuran and adduct processing: (i) 3-bromofuran (2 equiv.), SnCl_4 (1.0 equiv.), CH_2Cl_2 , rt, 48 h; (ii) $\text{PhB}(\text{OH})_2$, Bu_4NBr , $\text{Pd}(\text{OAc})_2$, H_2O , 60 W, 20 min; (iii) DIBAL-H (6 equiv.), THF, reflux; (iv) *E*- $\text{PhHC}=\text{CHB}(\text{OH})_2$, Bu_4NBr , $\text{Pd}(\text{OAc})_2$, H_2O , μW 60 W, 100 °C, 20 min.

6b (from **5b**) being increased to 95 kJ mol^{-1} . Both these effects will contribute to the observed failure to yield the corresponding carbonate from the 2-methyl (or 2,3-dimethylfuran) cycloadducts.

Cyclic carbonate **9§** and *endo*-cycloadduct **8§** were obtained in moderate yield from the reaction between **1** and 3-bromofuran, a seldom-used diene⁹ (Scheme 3), as verified by X-ray diffraction analysis of single crystals. This result is in line with our previous conclusion that the steric hindrance at the bridgehead carbon where C–O scission occurs is more important for ring opening than electronic effects in the transition state.

The use of the vinyl bromide as a locus for the generation of diversity is an attractive possibility that we propose to explore more fully, since preliminary experiments have yielded encouraging results. Cycloadduct **8** reacted with phenylboronic acid under Leadbetter's aqueous microwave-mediated conditions¹⁰ to afford **10**. Exposure of **9** to excess DIBAL-H provided tetrol **11** in moderate yield; a styryl group was then introduced *via* the coupling reaction.

Potentially, these reactions afford extremely rapid access to a very wide range of functionally rich difluorinated cyclohexene derivatives, though there is considerable scope for optimisation.

We thank the EPSRC (GR/M94922) for grants (fellowship to A. C. M., studentship to V. S.).

Notes and references

‡ Cycloaddition with stoichiometric tin(IV) chloride: preparation of **3**. Tin(IV) chloride (2 mmol, 1.9 mL of a 1.03 M solution in DCM) was added slowly to a mixture of alkenoate **1** (2 mmol, 0.5 g) and furan (4 mmol, 0.3 mL) in DCM (3 mL); the mixture was stirred at room temperature for 36 h. The mixture was diluted with water (5 mL) and extracted with ethyl acetate (3×15 mL). The combined organic extracts were dried (MgSO_4) and concentrated *in vacuo* to leave an orange oil (0.61 g), which was purified by column chromatography (20% diethyl ether in light petroleum) to afford *endo* adduct **2a** (0.22 g, 34%) and cyclic carbonate **3** (0.14 g, 26%) as cubic

crystals; mp 61–62 °C; R_f (20% diethyl ether in light petroleum) 0.05; (Found: C, 45.66; H, 3.70; $\text{C}_{10}\text{H}_{10}\text{F}_2\text{O}_6$ requires: C, 45.46; H, 3.82%); ν_{max} (film)/ cm^{-1} : 3538m br (OH), 2996m (C–H), 1846m (C=O), 1750m (C=O), 1473w (C–C), 1315m (C–O), 1229m (C–O), 1118m (C–O); δ_{H} (300 MHz, CDCl_3): 6.15–6.09 (1H, m), 5.96–5.91 (1H, m), 5.48 (1H, d, J 4.0), 4.68–4.61 (1H, m), 4.42–4.32 (2H, m), 3.66 (1H, br s), 1.34 (3H, t, J 7.2 Hz); δ_{C} (75 MHz, CDCl_3): 164.2 (d, $^4J_{\text{C-F}}$ 1.7), 151.3, 133.5 (dd, $^3J_{\text{C-F}}$ 5.9, 1.4), 122.0 (d, $^4J_{\text{C-F}}$ 1.1), 116.6 (dd, $^1J_{\text{C-F}}$ 256.6, 250.9), 80.6 (dd, $^2J_{\text{C-F}}$ 33.1, 23.5), 76.8 (d, $^3J_{\text{C-F}}$ 2.3), 65.2 (dd, $^2J_{\text{C-F}}$ 27.1, 20.9 Hz), 64.2, 13.7; δ_{F} (282 MHz, CDCl_3): –115.3 (1F, d, $^2J_{\text{F-F}}$ 261.5), –125.9 (1F, dd, $^2J_{\text{F-F}}$ 261.5, $^3J_{\text{F-H}}$ 13.4 Hz); m/z (CI): 283 (100%, $[\text{M} + \text{Na}]^+$).

§ *Crystal data* for **6** (mp not measured): $\text{C}_{14}\text{H}_{19}\text{Cl}_4\text{F}_2\text{NO}_5\text{Sn}$, $M = 579.8$; monoclinic, $a = 10.4879(11)$, $b = 18.8071(19)$, $c = 10.7203(11)$ Å, $\beta = 102.993(2)^\circ$, $U = 2060.4(4)$ Å³; $T = 150(2)$ K; space group $P2_1/n$, $Z = 4$, $\mu(\text{Mo-K}\alpha) 1.802$ mm^{–1}; 10 971 reflections measured, 4033 unique ($R_{\text{int}} = 0.0261$), which were used in all calculations; $R_1 = 0.0278$, $wR_2 = 0.0651$; the final $wR(F^2)$ was 0.0639 (all data). For **8** (mp 71–72 °C): $\text{C}_{14}\text{H}_{18}\text{BrF}_2\text{NO}_5$, $M = 398.2$; monoclinic, $a = 11.6660(9)$, $b = 8.0535(6)$, $c = 18.3629(14)$ Å, $\beta = 102.7150(10)^\circ$, $U = 1682.9(2)$ Å³; $T = 150(2)$ K; space group $P2_1/c$, $Z = 4$, $\mu(\text{Mo-K}\alpha) 2.486$ mm^{–1}; 12 743 reflections measured, 3310 unique ($R_{\text{int}} = 0.0269$), which were used in all calculations; $R_1 = 0.0417$, $wR_2 = 0.0914$; the final $wR(F^2)$ was 0.0888 (all data). For **9** (mp 123–125 °C): $\text{C}_{10}\text{H}_9\text{BrF}_2\text{NO}_6$, $M = 343.1$; monoclinic, $a = 7.0091(8)$, $b = 11.5134(13)$, $c = 14.9095(17)$ Å, $\beta = 94.404(2)^\circ$, $U = 1199.6(2)$ Å³; $T = 150(2)$ K; space group $P2_1$, $Z = 4$, $\mu(\text{Mo-K}\alpha) 3.474$ mm^{–1}; 8057 reflections measured, 4055 unique ($R_{\text{int}} = 0.0236$), which were used in all calculations; $R_1 = 0.0328$, $wR_2 = 0.0768$; the final $wR(F^2)$ was 0.0768 (all data). CCDC 229969–229971. See <http://www.rsc.org/suppdata/cc/b4/b401245c/> for crystallographic data in CIF or other electronic format.

- J. M. Percy, *Top. Curr. Chem.*, 1997, **193**, 131.
- For the only previous example of a difluorinated alkenoate dienophile, see: J. Leroy, H. Molines and C. Wakselman, *J. Org. Chem.*, 1987, **52**, 290; the application of DFT methods to a monofluorinated dienophile is described in: M. Essers, C. Muck-Lichtenfeld and G. Haufe, *J. Org. Chem.*, 2002, **67**, 4715.
- P. J. Crowley, A. C. Moralee, J. M. Percy and N. S. Spencer, *Synlett*, 2000, 1737.
- M. Lautens and W. Klute, *Angew. Chem., Int. Ed. Engl.*, 1996, **35**, 442; A. Arany, P. J. Crowley, J. Fawcett, M. B. Hursthouse, B. M. Kariuki, M. E. Light, A. C. Moralee, J. M. Percy and V. Salafia, *Org. Biomol. Chem.*, 2004, 455.
- P. J. Crowley, J. Fawcett, B. M. Kariuki, A. C. Moralee, J. M. Percy and V. Salafia, *Org. Lett.*, 2002, **4**, 4125.
- S. E. Denmark and J. P. Fu, *J. Am. Chem. Soc.*, 2003, **125**, 2208; for a 7-membered chelate involving ether oxygen ligands, see: K. Ishihara, H. Ishibashi and H. Yamamoto, *J. Am. Chem. Soc.*, 2002, **124**, 3647.
- A referee suggested an alternative explanation in which water reacts with **5a**, which then progresses irreversibly. We cannot disprove this suggestion, but must point out that our hypothesis does appear to be strongly supported by the computational findings.
- M. J. Frisch, G. W. Trucks, H. B. Schlegel, G. E. Scuseria, M. A. Robb, J. R. Cheeseman, V. G. Zakrzewski, J. A. Montgomery, Jr., R. E. Stratmann, J. C. Burant, S. Dapprich, J. M. Millam, A. D. Daniels, K. N. Kudin, M. C. Strain, O. Farkas, J. Tomasi, V. Barone, M. Cossi, R. Cammi, B. Mennucci, C. Pomelli, C. Adamo, S. Clifford, J. Ochterski, G. A. Petersson, P. Y. Ayala, Q. Cui, K. Morokuma, P. Salvador, J. J. Dannenberg, D. K. Malick, A. D. Rabuck, K. Raghavachari, J. B. Foresman, J. Cioslowski, J. V. Ortiz, A. G. Baboul, B. B. Stefanov, G. Liu, A. Liashenko, P. Piskorz, I. Komaromi, R. Gomperts, R. L. Martin, D. J. Fox, T. Keith, M. A. Al-Laham, C. Y. Peng, A. Nanayakkara, M. Challacombe, P. M. W. Gill, B. G. Johnson, W. Chen, M. W. Wong, J. L. Andres, C. Gonzalez, M. Head-Gordon, E. S. Replogle and J. A. Pople, Gaussian 98, Gaussian, Inc., Pittsburgh, PA, USA, 2001.
- M. G. Van Campen and J. R. Johnson, *J. Am. Chem. Soc.*, 1933, **55**, 430; D. G. Batt, D. G. Jones and S. La Greca, *J. Org. Chem.*, 1991, **56**, 6704.
- N. E. Leadbeater and M. Marco, *J. Org. Chem.*, 2003, **68**, 888–892.