

*Corynebacterium marinum* sp. nov. isolated from coastal sediment  
Zong-Jun Du, Elizabeth M. Jordan, Alejandro P. Rooney, Guan-Jun Chen  
and Brian Austin<sup>4</sup>

International Journal of Systematic and Evolutionary Microbiology,  
Volume 60, Issue 8, pp. 1944 - 1947  
<http://hdl.handle.net/1893/2715>

This is an author manuscript that has been accepted for publication in International Journal of Systematic and Evolutionary Microbiology, copyright International Union of Microbiological Societies, but has not been copy-edited, formatted or proofed.

Cite this article as appearing in International Journal of Systematic and Evolutionary Microbiology.

This version of the manuscript may not be duplicated or reproduced, other than for personal use or within the rule of 'Fair Use of Copyrighted Materials' (section 17, Title 17, US Code), without permission from the copyright owner, International Union of Microbiological Societies. The Society for General Microbiology and the International Union of Microbiological Societies disclaim any responsibility or liability for errors or omissions in this version of the manuscript or in any version derived from it by any other parties.

The final copy-edited, published article, which is the version of record, can be found at <http://ijs.sgmjournals.org>, and is freely available without a subscription 24 months after publication. [http://ijs.sgmjournals.org/misc/self\\_archiving.dtl](http://ijs.sgmjournals.org/misc/self_archiving.dtl)

1 ***Corynebacterium marinum* sp. nov. isolated from coastal sediment in Qingdao,**  
2 **China**

3 Zong-Jun Du,<sup>1</sup> Elizabeth M. Jordan<sup>2</sup>, Alejandro P Rooney,<sup>3</sup> Guan-Jun Chen,<sup>1</sup> and Brian Austin<sup>2,4</sup>

4 <sup>1</sup> College of Marine Science, Shandong University at Weihai, Weihai 264209, P. R. China

5 <sup>2</sup> School of Life Sciences, Heriot-Watt University, Riccarton, Edinburgh EH14 4AS, UK

6 <sup>3</sup> National Center for Agricultural Utilization Research, Agricultural Research Service, U.S.

7 Department of Agriculture, 1815 North University Street, Peoria, IL 61604, USA

8 <sup>4</sup> Institute of Aquaculture, University of Stirling, Stirling, FK9 4LA, Scotland, UK

9

10 **Correspondence** Brian Austin [brian.austin@stir.ac.uk](mailto:brian.austin@stir.ac.uk)

11

12

13

14

15 **Running title:** *Corynebacterium marinum* sp. nov.

16

---

17 The GenBank accession number for the 16S rRNA gene sequence of *Corynebacterium marinum*

18 D7015<sup>T</sup> is DQ219354. A representative phase-contrast micrograph of strain D7015<sup>T</sup> and a table

19 showing the cellular fatty acid profiles of strain D7015<sup>T</sup> and related species are available as

20 supplementary material in IJSEM Online.

21 A taxonomic study was performed on strain D7015<sup>T</sup>, which was isolated from coastal sediment  
22 close to a coal-fired power station in Qingdao, China. Strain D7015<sup>T</sup> comprised Gram-positive,  
23 non-motile diphtheroid rods, which grew in the presence of 0-8% (w/v) NaCl and at 4-37°C, with  
24 optimum growth at 1% (w/v) NaCl and 30-32°C. The G+C content was 65.0 mol%. The major fatty  
25 acids were C<sub>18:1ω9c</sub> (56.18%), C<sub>16:0</sub> (38.02%), C<sub>16:1ω7c</sub> (4.45%), C<sub>18:0</sub> (1.0%) and C<sub>14:0</sub> (0.35%).  
26 On the basis of the morphological, physiological and phylogenetic characteristics, strain D7015<sup>T</sup>  
27 was classified in the genus *Corynebacterium*. It exhibited a 16S rRNA gene sequence similarity of  
28 95.9% and a DNA:DNA relatedness value of 20.4% with *Corynebacterium halotolerans* DSM  
29 44683<sup>T</sup>. Strain D7015<sup>T</sup> was sufficiently different from hitherto described *Corynebacterium* species  
30 to be considered as a novel species. The name *Corynebacterium marinum* sp. nov. is proposed, with  
31 strain D7015<sup>T</sup> (=CGMCC 1.6998<sup>T</sup> =NRRL B-24779<sup>T</sup>) as the type strain.

---

32  
33 The genus *Corynebacterium* was proposed by Lehmann & Neumann (1896) and represents a large  
34 group of Gram-positive, asporogenous, rod-shaped bacteria with high DNA G+C content.  
35 Corynebacteria have been isolated from a wide range of environments, namely dairy products, soil,  
36 sewage, sediments, plant materials and aquatic sources, but the majority of novel species described  
37 in recent years have originated from human or animal clinical samples (e.g. Renaud *et al.*, 2007;  
38 Yassin & Siering, 2008; Yassin, 2009; Funke *et al.*, 2009). However, some corynebacterial species  
39 have been isolated from the marine environment, and they may occur as part of the indigenous flora  
40 of marine animals. For example, *Corynebacterium phocae* and *Corynebacterium caspium* were  
41 isolated from seals (Pascual *et al.*, 1998; Collins *et al.*, 2004), *Corynebacterium spheniscorum* and  
42 *Corynebacterium sphenisci* were recovered from wild penguins (Goyache *et al.*, 2003a, 2003b),  
43 and *Corynebacterium maris* Coryn-1<sup>T</sup> was found in the mucus of the coral, *Fungia granulose*  
44 (Ben-Dov *et al.*, 2009). Here, we report the taxonomic characteristics of a novel *Corynebacterium*  
45 species that originated from coastal sediment close to a coal-fired power station in Qingdao, China.

46

47 Strain D7015<sup>T</sup> was isolated from coastal sediment in 2000 (Du *et al.*, 2002). The isolate was grown  
48 aerobically at 28°C on marine 2216E agar (MA; Difco) for 48 h. Cultures were maintained on MA  
49 slants at room temperature, and stock cultures were kept in tryptone soya broth (Oxoid)  
50 supplemented with 1% (w/v) NaCl (= TNB) and 20% (v/v) glycerol at -70°C. Identification of  
51 strain D7015<sup>T</sup> was performed as described by Jordan *et al.* (2007). For phenotypic tests, the strain  
52 was grown on MA for 48 h at 28°C, and cells were resuspended in saline for use as an inoculum.  
53 Tolerance of 1, 3, 5, 7, 8 and 10% (w/v) NaCl was assessed on appropriately modified tryptone  
54 soya agar (TSA; Oxoid). Growth in the absence of NaCl was assessed on plate count agar (PCA;  
55 Oxoid). Inoculated plates were incubated at 28°C for up to 5 days. The effects of different  
56 temperatures on growth were assessed on TSA plates supplemented with 1.0% (w/v) NaCl (TNA)  
57 with incubation at 4, 10, 15, 28, 30, 32, 37, 42 and 45°C. The reduction of nitrate was assessed in  
58 nitrate broth, prepared according to the method of Cowan & Steel (1974), and incubated at room  
59 temperature for 10 days. Oxidase and catalase activities were determined by using standard  
60 methods. The culture was characterized biochemically using the API Coryne, API 50CH and API  
61 ZYM systems according to the manufacturer's instructions (bioMérieux). Measuring turbidity was  
62 used to evaluate the growth of strain D7015<sup>T</sup> in the API 50CH and the API 20NE systems. The API  
63 50 CH strips were read after 7 days incubation at 28°C.

64

65 The almost-complete 16S rRNA gene sequence (1444 nt) of strain D7015<sup>T</sup> was obtained using the  
66 universal primers 27f and 1492r (MWG Biotech; Lane, 1991). The 16S rRNA gene sequence of  
67 strain D7015<sup>T</sup> was submitted to GenBank and EMBL to search for similar sequences using the  
68 BLAST algorithm. A phylogenetic dendrogram of strain D7015<sup>T</sup> and some closely related  
69 members of the genus *Corynebacterium* based on 16S rRNA gene homology was constructed using  
70 the neighbor-joining method of the MEGA software version 4.1 (Tamura *et al.*, 2007). The

71 resultant tree topologies were evaluated by bootstrap analysis based on 1000 replicates.

72

73 Cellular fatty acids were determined on a 3 day old culture grown on marine 2216E agar plates after  
74 incubation at 28°C. The fatty acids were extracted, methylated and analyzed using the standard  
75 MIDI (Microbial Identification) system (Sasser, 1990). The G+C content of the DNA was  
76 determined directly by high pressure liquid chromatography according to a method described  
77 previously (Tamaoka & Komagata, 1984; Mesbah *et al.*, 1989). DNA:DNA hybridization between  
78 strain D7015<sup>T</sup> and *Corynebacterium halotolerans* DSM 44683<sup>T</sup> was carried out by applying the  
79 optical renaturation method (De Ley *et al.*, 1970; Huss *et al.*, 1983; Jahnke, 1992) under optimal  
80 hybridization conditions.

81

82 Microscopic examination of strain D7015<sup>T</sup> suggested that it belonged to the genus  
83 *Corynebacterium*, as cells stained Gram-positive and produced short, diphtheroid rods with some  
84 of the cells arranged in a V formation due to their snapping division (Collins & Cummins, 1986). A  
85 representative phase-contrast micrograph of strain D7015<sup>T</sup> is available as supplementary material  
86 in IJSEM Online. Strain D7015<sup>T</sup> was non-motile and non-endospore-forming. The isolate was  
87 catalase-positive and oxidase-negative. The following results of carbon source assimilation were  
88 positive in API 50CH: aesculin ferric citrate, salicin, D-maltose and glycogen. Strain D7015<sup>T</sup>  
89 displayed a numerical profile of 3200127 with the commercial API Coryne system, which  
90 corresponded to a “doubtful” identification as *Corynebacterium glucuronolyticum* (with a  
91 confidence level of 99.2%). However, further biochemical analyses, using API ZYM and  
92 additional phenotypic tests revealed that strain D7015<sup>T</sup> could be distinguished from *C.*  
93 *glucuronolyticum* on the basis of its ability to produce  $\alpha$ -chymotrypsin and its inability to produce  
94 esterase lipase (C4). Strain D7015<sup>T</sup> produced acid from glucose, maltose, sucrose and glycogen,  
95 but not from ribose, xylose, mannitol and lactose. Strain D7015<sup>T</sup> was also different from *C.*

96 *spheniscorum* notably in the inability of the latter to reduce nitrate, or produce  $\beta$ -glucuronidase,  $\alpha$ -  
97 chymotrypsin or naphthol-AS-BI-phosphohydrolase. The complete morphological and  
98 biochemical characteristics for strain D7015<sup>T</sup> are given in the species description.

99

100 Phylogenetic analyses performed with nearly complete sequences of members of closely related  
101 species (Fig. 1) showed that no sequence available in the GenBank database exhibited more than  
102 96% similarity. The sequence similarity observed between the new isolate and its closest relative,  
103 *C. halotolerans*, was 95.9%, which is a lower value than the borderline used for defining bacterial  
104 species (i.e. 97%) as proposed by Stackebrandt & Goebel (1994). *C. halotolerans* was isolated  
105 from a saline soil sample in China (Chen *et al.*, 2004). For strain D7015<sup>T</sup>, growth was not observed  
106 at 10% (w/v) NaCl. However, optimum growth of *C. halotolerans* was in 10% (w/v) KCl, NaCl or  
107 MgCl<sub>2</sub>·6H<sub>2</sub>O. These two strains were also different biochemically, with strain D7015<sup>T</sup> able to  
108 hydrolyse starch and Tween 20-80, but unable to produce esterase lipase (C4). The DNA:DNA  
109 relatedness value of  $20.4 \pm 0.1\%$  (experiment repeated twice) was significantly lower than 70%,  
110 which is considered to be the threshold value for the delineation of genomic species (Wayne *et al.*,  
111 1987). Clearly, strain D7015<sup>T</sup> differed biochemically from all of its closest relatives, both in its  
112 ability to hydrolyse casein and in its inability to produce esterase lipase C4. The phenotypic  
113 features that differentiate strain D7015<sup>T</sup> from its closest phylogenetic relatives are provided in  
114 Table 1.

115

116 The cellular fatty acids in strain D7015<sup>T</sup> were C<sub>18:1 $\omega$ 9c</sub> (56.18%), C<sub>16:0</sub> (38.02%), C<sub>16:1 $\omega$ 7c</sub> (4.45%),  
117 C<sub>18:0</sub> (1.0%) and C<sub>14:0</sub> (0.35%). The genomic DNA G+C content was 65.0 mol%.

118

119 Based on these molecular, chemotaxonomic and phenotypic results, it is proposed that strain  
120 D7015<sup>T</sup> should be classified as a new species of the genus *Corynebacterium*, for which the name

121 *Corynebacterium marinum* sp. nov. is proposed.

122

123 **Description of *Corynebacterium marinum* sp. nov.**

124 *Corynebacterium marinum* (ma.ri'num. L. neut. adj. *marinum* of the sea, marine)

125

126 Cells are short Gram-positive, non-motile, diphtheroid rods; some of the cells are arranged in a V  
127 formation. Colonies on marine 2216E agar medium are circular, erose, convex, yellow and of a  
128 creamy consistency and are 0.5-1.5 mm in diameter after 48 h at 28°C. Facultatively anaerobic,  
129 catalase-positive and oxidase-negative. Bacteria are methyl-red negative, Voges-Proskauer  
130 positive, reduce nitrate, and lyse horse blood cells. Aesculin and urea are not hydrolysed, but casein  
131 is digested and starch and pullulan are hydrolysed. Gelatin is not liquefied. Tween 20-80 are  
132 hydrolysed. Cells grow in the presence of 0-8% (w/v) NaCl and at 4-37°C. Prolific growth occurs  
133 at 30-32°C in media that contain 1% (w/v) NaCl. Using the API 50CH system, aesculin ferric  
134 citrate, salicin, D-maltose, and glycogen are utilized. Acid is produced from glucose, maltose,  
135 sucrose and glycogen, but not from ribose, xylose, mannitol or lactose. Activities for esterase lipase  
136 (C8), leucine arylamidase,  $\alpha$ -chymotrypsin, naphthol-as-bi-phosphohydrolase, pyrazinamidase and  
137  $\beta$ -glucuronidase are positive. Esterase lipase (C4), lipase (C14), valine arylamidase, cystine  
138 arylamidase, trypsin, alkaline phosphatase, pyrrolidonyl arylamidase, acid phosphatase,  
139  $\alpha$ -galactosidase,  $\beta$ -galactosidase,  $\alpha$ -glucosidase,  $\beta$ -glucosidase, N-acetyl- $\beta$ -glucosaminidase,  
140  $\alpha$ -mannosidase and  $\alpha$ -fucosidase are negative. The type strain is resistant to nalidixic acid (30  $\mu$ g),  
141 nitrofurantoin (50  $\mu$ g), sulphamethizole (200  $\mu$ g), tetracycline (100  $\mu$ g) and cotrimoxazole (25  $\mu$ g),  
142 but sensitive to ampicillin (25  $\mu$ g), chloramphenicol (50  $\mu$ g), gentamycin (10  $\mu$ g), kanamycin (30  
143  $\mu$ g), carbenicillin (100  $\mu$ g) and streptomycin (25  $\mu$ g) as determined by antibiotic discs. Major fatty  
144 acids produced are C<sub>18:1 $\omega$ 9c</sub> (56.18%), C<sub>16:0</sub> (38.02%), C<sub>16:1 $\omega$ 7c</sub> (4.45%), C<sub>18:0</sub> (1.0%) and C<sub>14:0</sub>  
145 (0.35%). The percentage whole-cell fatty acid compositions of strain D7015<sup>T</sup> and related species is

146 available as supplementary material in IJSEM Online. The genomic DNA G+C content is 65.0  
147 mol% for the type strain.

148

149 The type strain is D7015<sup>T</sup> (=CGMCC 1.6998<sup>T</sup> = NRRL B-24779<sup>T</sup>), which was isolated from coastal  
150 sediment close to a coal-fired power station in Qingdao, China.

151

## 152 **Acknowledgements**

153 This work was supported by a Darwin Initiative Project of the UK (No.162/8/065) , the National  
154 Science Foundation of China (Project no. 40730847), Special Non-profit Research Project from  
155 Ministry of Agriculture of China (nyhyzx 07-046), and a grant (MBSMAT-2009-07) from key lab  
156 of marine bioactive substance and modern analytical technique, State Oceanic Administration  
157 (SOA), China.

158

## 159 **References**

160 **Ben-Dov, E., Ben-Yosef, D. Z., Pavlov, V. & Kushmaro, A. (2009).** *Corynebacterium*  
161 *maris* sp. nov., a marine bacterium isolated from the mucus of the coral *Fungia granulose*. *Int J*  
162 *Syst Evol Microbiol*, published online ahead of print on 21 July 2009 as doi:ijs. 0.007468-0.

163 **Chen, H. H., Li, W. J., Tang, S. K., Kroppenstedt, R. M., Stackebrandt, E., Xu, L. H. & Jiang,**  
164 **C. L. (2004).** *Corynebacterium halotolerans* sp. nov., isolated from saline soil in the west of China.  
165 *Int J Syst Evol Microbiol* 54, 779–782.

166 **Collins, M. D. & Cummins, C. S. (1986).** Genus *Corynebacterium* Lehmann and Neumann  
167 1896, 350<sup>AL</sup>. In *Bergey's Manual of Systematic Bacteriology*, vol. 2, pp. 1266–1276. Edited by P.  
168 H. A. Sneath, N. S. Mair, M. E. Sharpe & J. G. Holt. Baltimore: Williams & Wilkins.

169 **Collins, M. D., Hoyles L., Foster G. & Falsen E. (2004).** *Corynebacterium caspium* sp. nov.,  
170 from a Caspian seal (*Phoca caspica*). *Int J Syst Evol Microbiol* 54, 925–928.



171 **Cowan, S. T. & Steel, K. J. (1974).** In *Cowan and Steel's Manual for the Identification of*  
172 *Medical Bacteria*, 2nd edn. Revised by S. T. Cowan. Cambridge, UK: Cambridge University Press.

173 **De Ley, J., Cattoir, H. & Reynaerts, A. (1970).** The quantitative measurement of DNA  
174 hybridization from renaturation rates. *Eur J Biochem* 12, 133–142.

175 **Du, Z., Li, Y., Yu, D., Wang, X., Chen, J., Robertson, P. A. W., Austin, B. & Xu, H.**  
176 **(2002).** Heterotrophic bacterial floras in industrial area and unpolluted marine environments  
177 around Qingdao. *Journal of Ocean University of Qingdao* 1, 59–62.

178 **Fudou, R., Jojima Y., Seto, A., Yamada, K., Kimura, E., Nakamatsu, T., Hiraishi, A. &**  
179 **Yamanaka, S. (2002).** *Corynebacterium efficiens* sp. nov., a glutamic acid-producing species  
180 from soil and vegetables. *Int J Syst Evol Microbiol* 52, 1127–1131.

181 **Funke, G., Frodl, R., Bernard, K. A. & Englert, R. (2009).** *Corynebacterium freiburgense*  
182 sp. nov., isolated from a wound obtained from a dog bite. *Int J Syst Evol Microbiol* 59, 2054–2057.

183 **Goyache, J., Vela, A. I., Collins, M. D., Ballesteros, C., Briones, V., Moreno, J., Yorio,**  
184 **P., Domínguez, L., Hutson, R. & Fernández-Garayzábal, J. F. (2003a).** *Corynebacterium*  
185 *spheniscorum* sp. nov., isolated from the cloacae of wild penguins. *Int J Syst Evol Microbiol* 53,  
186 43–46.

187 **Goyache, J., Ballesteros, C., Vela, A. I., Collins, M. D., Briones, V., Hutson, R. A.,**  
188 **Potti, J., García-Borboroglu, P., Domínguez, L. & Fernández-Garayzábal, J. F.**  
189 **(2003b).** *Corynebacterium sphenisci* sp. nov., isolated from wild penguins. *Int J Syst Evol*  
190 *Microbiol* 53, 1009–1012.

191 **Huss, V. A. R., Festl, H. & Schleifer, K.-H. (1983).** Studies on the spectrophotometric  
192 determination of DNA hybridization from renaturation rates. *Syst Appl Microbiol* 4, 184–192.

193 **Jahnke, K.-D. (1992).** BASIC computer program for evaluation of spectroscopic DNA  
194 renaturation data from Gilford System 2600 spectrophotometer on a PC/XT/AT type personal  
195 computer. *J Microbiol Methods* 15, 61–73.

196 **Jordan, E. M., Thompson, F. L., Zhang, X.-H., Li, Y., Vancanneyt, M., Kroppenstedt**  
197 **R. M., Priest, F. G. & Austin, B. (2007).** *Sneathiella chinensis* gen. nov., sp. nov., a novel  
198 marine alphaproteobacterium isolated from coastal sediment in Qingdao, China. *Int J Syst Evol*  
199 *Microbiol* 57, 114–121.

200 **Lane, D. J. (1991).** 16S/23S rRNA sequencing. In *Nucleic Acid Techniques in Bacterial*  
201 *Systematics*, pp. 115–176. Edited by E. Stackebrandt & M. Goodfellow. Chichester: Wiley.

202 **Lehmann, K. B. & Neumann, R. (1896).** *Atlas und Grundriss der Bakteriologie und Lehrbuch*  
203 *der Speziellen Bakteriologischen Diagnostik*. Munich: J. F. Lehmann (in German).

204 **Mesbah, M., Premachandran, U. & Whitman, W. B. (1989).** Precise measurement of the  
205 G+C content of deoxyribonucleic acid by high-performance liquid chromatography. *Int J Syst*  
206 *Bacteriol* 39, 159–167.

207 **Pascual, C., Foster, G., Alvarez, N. & Collins, M. D. (1998).** *Corynebacterium phocae* sp.  
208 nov., isolated from the common seal (*Phoca vitulina*). *Int J Syst Evol Microbiol* 48, 601–604.

209 **Renaud, F. N. R., Coustumier, A. L., Wilhem, N., Aubel, D., Riegel, P., Bollet, C. &**  
210 **Freney, J. (2007).** *Corynebacterium hansenii* sp. nov., an  $\alpha$ -glucosidase-negative bacterium  
211 related to *Corynebacterium xerosis*. *Int J Syst Evol Microbiol* 57, 1113–1116.

212 **Sasser, M. (1990).** Identification of bacteria by gas chromatography of cellular fatty acids.  
213 *Technical Note 101*. Newark, DE: MIDI.

214 **Stackebrandt, E. & Goebel, B. M. (1994).** Taxonomic note: a place for DNA-DNA  
215 reassociation and 16S rRNA sequence analysis in the present species definition in bacteriology. *Int*  
216 *J Syst Bacteriol* 44, 846–849.

217 **Tamaoka, J. & Komagata, K. (1984).** Determination of DNA-base composition by  
218 reversed-phase high-performance liquid-chromatography. *FEMS Microbiol Lett* 25, 125–128.

219 **Tamura, K., Dudley J., Nei, M. & Kumar, S. (2007).** MEGA4: Molecular Evolutionary  
220 Genetics Analysis (MEGA) software version 4.0. *Molecular Biology and Evolution* 24, 1596-1599.

221 **Wayne, L. G., Brenner, D. J., Colwell, R. R. & 9 other authors (1987).** International  
222 Committee on Systematic Bacteriology. Report of the ad hoc committee on reconciliation of  
223 approaches to bacterial systematics. *Int J Syst Bacteriol* 37, 463–464.

224 **Yassin, A. F. & Siering, C. (2008).** *Corynebacterium sputi* sp. nov., isolated from the sputum  
225 of a patient with pneumonia. *Int J Syst Evol Microbiol* 58, 2876–2879.

226 **Yassin, A. F. (2009).** *Corynebacterium ulceribovis* sp. nov., isolated from the skin of the udder  
227 of a cow with a profound ulceration. *Int J Syst Evol Microbiol* 59, 34–37.

228

229 **Table 1.** Comparison of strain D7015<sup>T</sup> with the phylogenetically related species of the genus  
 230 *Corynebacterium*  
 231 Species: 1, Strain D7015<sup>T</sup>, 2, *Corynebacterium efficiens* DSM 44549<sup>T</sup> (Fudou *et al.*, 2002); 3,  
 232 *Corynebacterium halotolerans* DSM 44683<sup>T</sup> (Chen *et al.*, 2004). +, positive; -, negative; ND, no  
 233 data available.

<b>Characteristic</b>	<b>1</b>	<b>2</b>	<b>3</b>
<b>Starch hydrolysis</b>	+	ND	-
<b>Esterase lipase (C4)</b>	-	ND	+
<b>Growth at 45 °C</b>	-	+	ND
<b>Growth in 10% (w/v) NaCl</b>	-	+	+
<b>Acid produced from:</b>			
<b>ribose</b>	-	+	-
<b>maltose</b>	+	+	-
<b>glycogen</b>	+	-	-
<b>DNA G+C content (mol%)</b>	65.0	59.0-60.2	63.0

234  
 235  
 236  
 237  
 238  
 239  
 240  
 241  
 242  
 243  
 244

245 **Figure legend:**


246 **Fig. 1.** Neighbor-joining phylogeny of *Corynebacterium marinum* and closely related species.

247 Numbers along branches represent bootstrap values; only values greater than 50% are shown.

248 GenBank accession numbers for the 16S rRNA sequences used in the tree reconstruction are given

249 in the parentheses next to each taxon. Bar, 0.01 substitutions per nucleotide.

250



251