

2

3 Lipid and fatty acid composition of parasitic caligid copepods belonging to  
4 the genus *Lepeophtheirus*

5

6 J. A. Tocher, J. R. Dick, J. E. Bron, A. P. Shinn and D. R. Tocher\*

7

8 Institute of Aquaculture, University of Stirling, Stirling FK9 4LA, Scotland, United Kingdom

9

10 Running title: Lipid composition of *Lepeophtheirus salmonis*

11

12 Key words: *Lepeophtheirus salmonis*, salmon lice, lipid, fatty acid, copepod

13

14 \* To whom correspondence should be addressed

15 Douglas Tocher

16 Institute of Aquaculture, University of Stirling, Stirling FK9 4LA, Scotland, UK. Phone: +44 1786  
17 467996; Fax: +44 1786 472133; E-mail: d.r.tocher@stir.ac.uk

18

19 **Abstract**

20 Sea lice are copepod ectoparasites that constitute a major barrier to the sustainability and economic  
21 viability of marine finfish aquaculture operations worldwide. In particular, the salmon louse,  
22 *Lepeophtheirus salmonis*, poses a considerable problem for salmoniculture in the northern  
23 hemisphere. The free-swimming nauplii and infective copepodids of *L. salmonis* are lecithotrophic,  
24 subsisting principally on maternally-derived lipid reserves. However, the lipids and fatty acids of  
25 sea lice have been sparsely studied and therefore the present project aimed to investigate the lipid  
26 and fatty acid composition of sea lice of the genus *Lepeophtheirus* obtained from a variety of fish  
27 hosts. Total lipid was extracted from eggs and adult female *Lepeophtheirus salmonis* obtained from  
28 both wild and farmed Atlantic salmon (*Salmo salar* L.) sampled at two time points, in the mid  
29 1990's and in 2009. In addition, *L. salmonis* from wild sea trout (*Salmo trutta* L.) and *L.*  
30 *hippoglossi* from wild Atlantic halibut (*Hippoglossus hippoglossus* L.) were sampled and analyzed.  
31 The lipids of both females and egg strings of *Lepeophtheirus* were characterized by triacylglycerol  
32 (TAG) as the major neutral (storage) lipid with phosphatidylcholine and phosphatidylethanolamine  
33 as the major polar (membrane) lipids. The major fatty acids were 22:6n-3 (DHA), 18:1n-9 and 16:0,  
34 with lesser amounts of 20:5n-3, 22:5n-3 and 18:0. *L. salmonis* sourced from farmed salmon were

35 characterized by higher levels of 18:2n-6 and 18:3n-3 than lice from wild salmon. Egg strings had  
36 higher levels of TAG and lower DHA compared to females, whereas *L. hippoglossi* had lower  
37 levels of TAG and higher DHA than *L. salmonis*. The results demonstrate that the fatty acid  
38 compositions of lice obtained from wild and farmed salmon differ and that changes to the lipid and  
39 fatty acid composition of feeds for farmed salmon influence the louse compositions.

40

## 41 **1. Introduction**

42

43 Copepods are a group of small crustaceans found in most marine and freshwater habitats of the  
44 world. More than 14,000 species have thus far been described, with sizes typically being in the  
45 range of < 1 to 6 mm. Many copepods are herbivorous, often feeding on phytoplankton, whilst  
46 others may be detritivores, predators or commensals, some of which are fully parasitic. Copepods  
47 are considered to be the most numerous metazoans on the planet, exceeding the numbers of the  
48 other two hyperabundant groups: insects and nematodes. As such, they are generally assumed to  
49 constitute the predominant biomass of zooplankton and are the major food for many fish, larger  
50 crustaceans, sea mammals and seabirds (Skjoldal et al., 2004). Many copepods, particularly those in  
51 cold or deep waters, build up large lipid energy reserves through feeding on phytoplankton. These  
52 lipids are stored in oils sacs and / or as oil droplets. In some species lipid may accumulate to  
53 between 50-70 % of body dry weight (Kattner and Krause, 1989; Lee et al., 2006; Falk-Petersen et  
54 al., 2009), making copepods the principal source of dietary lipid for many plankton-feeding fish  
55 species.

56 Parasitic copepods of the genus *Lepeophtheirus* constitute one of the most serious pathogens  
57 of marine farmed salmonids the world (Johnson et al., 2004). It is estimated that sea lice infection  
58 by species belonging to the genera *Lepeophtheirus* and *Caligus*, cost the world's eight major  
59 salmon-producing countries, a combined total of over €300 million (Costello, 2009). Sea lice can  
60 pose a considerable risk to fish health and can inhibit growth, cause external damage and, in  
61 extreme cases, lead to mortality (Pike and Wadsworth, 1999). Sea lice are therefore a major  
62 constraint to farm production in coldwater salmoniculture and have also been suggested to be a  
63 threat to wild salmonid populations such as sea trout (Ford and Myers, 2008).

64 The species of copepod parasite of prime concern to mariculture and wild fisheries in  
65 Scotland is the salmon louse, *Lepeophtheirus salmonis*, which is the most pathogenic marine  
66 ectoparasite of Atlantic salmon (*Salmo salar* L.). The life cycle of this species is well characterized,  
67 comprising 10 stages separated by moults and five developmental phases (Kabata, 1979). After  
68 hatching from paired egg strings carried by host-attached adult females, the lice progress through  
69 two free-swimming planktonic nauplius stages before developing into copepodids, which infect a

70 new fish host (Schram, 1993). After attachment, development proceeds through four chalimus and  
71 two sexually differentiated preadult stages before sexual maturity is reached at the adult stage.  
72 During development on the fish host, lice survive by feeding exclusively on host material including  
73 mucus, skin and blood (Brandal et al., 1976; Jonsdottir et al., 1992). Off the host, free-living stages  
74 of *L. salmonis*, nauplii and copepodids, are sustained by body reserves until the infective copepodid  
75 larva attaches to a new host (Boxaspen, 2006).

76 Despite considerable research into the biology, genetics and control of sea lice (Pike and  
77 Wadsworth, 1999; Boxaspen, 2006), very little is known about lipids and lipid metabolism in *L.*  
78 *salmonis* and in parasitic copepods in general. The period of survival of the free-swimming  
79 copepodid stage is constrained by its endogenous energy supplies (Boxaspen, 2006). The available  
80 energy reserves of *L. salmonis* copepodids were estimated at 7800 cal g<sup>-1</sup> dry weight by bomb  
81 calorimetry, this being similar to reserves reported for copepodid stages of other parasitic and free-  
82 living copepods during winter (Tucker et al., 2000). This figure declined sharply between 1-2-day-  
83 old and 7-day-old copepodids with those at 7 days having substantially depleted reserves. In  
84 addition to their role as energy reserves, lipids are also key to the parasite's ability to immuno-  
85 modulate the host. In this respect, *L. salmonis* has been demonstrated to secrete prostaglandin E<sub>2</sub>, an  
86 arachidonic acid (ARA; 20:4n-6) metabolite (Fast et al., 2004).

87 In contrast to the parasitic copepods, there has been considerable research into lipid storage  
88 and lipid metabolism in free-living copepods (see Lee et al., 2006; Kattner et al., 2007). An initial  
89 comparison of lipids in free-living and parasitic species (Lee, 1975) reported that, unlike the lipids  
90 of many free-living marine copepods that use wax esters (WE) as their primary energy store, *L.*  
91 *salmonis* from Pacific coho (*Oncorhynchus kisutch*) and pink (*O. gorbuscha*) salmon and other  
92 parasitic caligid copepods used triacylglycerol (TAG) as their main energy store. This was  
93 confirmed for *L. salmonis* from Atlantic salmon by Tucker et al. (2000). Given the paucity of  
94 information concerning lipids in caligid copepods, the present study therefore investigated lipid  
95 class and fatty acid compositions of total lipids of sea lice of the genus *Lepeophtheirus* obtained  
96 from a variety of fish hosts.

97

## 98 **2. Materials and Methods**

### 99 *2.1. Samples and sampling*

100 Individual lice of the genus *Lepeophtheirus* were collected from infected fish obtained from various  
101 sites in Scottish waters (Table 1). Collected sea lice were maintained in fresh seawater (minimum  
102 33 ppt) at 10 °C with aeration for approximately 24 h prior to processing. This study used samples  
103 of *L. salmonis*, collected from wild and farmed Atlantic salmon (*Salmo salar* L.) at two time points  
104 (early summer 1995 and summer 2009), and also those collected from wild sea trout (*Salmo trutta*

105 L.) in summer 1996. Samples obtained in 2009 were used fresh whilst older samples were stored at  
106 -80°C prior to use. The Atlantic salmon were all sampled in sea water mostly from various sea lochs  
107 whereas the sea trout was sampled in freshwater. Samples of *L. hippoglossi* collected from wild  
108 Atlantic halibut (*Hippoglossus hippoglossus* L.) in summer 1998 were also examined. All lice  
109 samples were adult females without egg strings. For three samples from farmed salmon, egg strings  
110 were carefully removed and also used for analysis.

111 Individual fresh or frozen lice were processed by macerating them in 30 µl of  
112 homogenization buffer (1 Mm Tris-HCL, pH 7.0, 0.1 mM EDTA, 0.1mM 2-mercaptoethanol) using  
113 a pellet pestle (Anachem, Luton UK). Samples were then flash frozen in liquid nitrogen and stored  
114 at -70 °C until required. Salmon muscle samples were skinned and boned white muscle fillets that  
115 were flash frozen in liquid nitrogen and stored at -70 °C until required. The muscle samples were  
116 thawed and homogenized into a paté prior to lipid extraction.

117

## 118 2.2. Lipid extraction

119 Total lipid was prepared according to the method of Folch et al. (1957). Sea lice samples or 0.5 g  
120 samples of salmon muscle paté were added to 5 ml ice cold chloroform/methanol (2:1, by volume)  
121 containing 0.01 % butylated hydroxytoluene as an antioxidant, and were homogenized using an  
122 IKA Ultra-Turrax T8. Tubes of homogenate were left on ice for one hour. A further 1 ml of  
123 chloroform/methanol (2:1, v/v) was then added along with 1.5 ml aqueous KCl (0.88 %). Samples  
124 were left on ice for a further 5 min and were then centrifuged at 600 g<sub>ave</sub> for 5 min to separate the  
125 mixture. The lower organic layer was filtered through Whatman No. 1 filter paper into clean test  
126 tubes, the solvent evaporated under a stream of oxygen-free nitrogen (OFN) and the dry lipid  
127 extract re-suspended in chloroform/methanol (2:1, v/v)

128

## 129 2.3. Lipid class composition analysis

130 Lipid class separation was performed by high-performance thin-layer chromatography (HPTLC).  
131 The concentration of the lipid extracts was adjusted to 10 mg/ml in chloroform/methanol (2:1, v/v)  
132 and two µL of each sample loaded as 2 mm streaks, 1 cm up on HPTLC plates (10 cm x 10 cm x  
133 0.15 mm), precoated with silica gel 60 (Merck, Darmstadt, Germany). The plate was developed to  
134 approximately 5 cm with methyl acetate/isopropanol/chloroform/methanol/0.25 % aqueous KCl  
135 (25:25:25:10:9, by vol.) then, after drying in air for 30 min, developed fully with isohexane/diethyl  
136 ether/acetic acid (85:15:1, by vol.). The lipid classes were visualized and quantified by charring at  
137 160 °C for 15 min after spraying with 3 % (w/v) aqueous cupric acetate containing 8 % (v/v)  
138 phosphoric acid and quantified by densitometry using a Camag 3 TLC Scanner (Camag, Muttenz,  
139 Switzerland) and winCATS software (Henderson and Tocher, 1992). The identities of individual

140 lipid classes were confirmed by comparison with reference to the R<sub>f</sub> values of authentic standards  
141 run alongside samples on HPTLC plates and developed in the above solvent systems.

142

#### 143 *2.4. Fatty acid composition analysis*

144 Fatty acid composition was determined by gas chromatography analysis of fatty acid methyl esters  
145 (FAME) prepared by acid-catalyzed transmethylation (Christie, 1993). Approximately 0.5 mg of  
146 total lipid was taken into a clean test tube and the solvent evaporated under a stream of OFN before  
147 1 ml toluene and 2.5 ml of 1 % H<sub>2</sub>SO<sub>4</sub> in methanol were added and the sample incubated overnight  
148 at 50°C (Christie, 1993). After cooling, FAME were extracted by addition of 2 ml of 2 % KHCO<sub>3</sub>  
149 and 5 ml of isohexane/diethyl ether (1:1, v/v) containing 0.01% BHT, followed by thorough mixing  
150 and centrifugation at 600 g<sub>ave</sub> for 2 min. The upper layer was removed into a clean test tube and the  
151 lower layer further extracted with a further portion of 5 ml of isohexane/diethyl ether (1:1, v/v)  
152 without BHT. After centrifugation at 600 g<sub>ave</sub> for 2 min, the upper layer was removed and added to  
153 the previous upper layer, and the solvent evaporated under a stream of OFN, before the crude  
154 FAME extract was resuspended in 100 µl isohexane. The FAME were purified by thin-layer  
155 chromatography (TLC) prior to GC analysis. Samples were applied as 2 cm streaks to TLC plates  
156 (20 cm x 20 cm x 0.25 mm) and FAME separated from non-derivatized lipid classes using  
157 hexane/diethyl ether/acetic acid (90:10:1, by vol.) as developing solvent. The FAME were located  
158 on the plate by staining standards run at the side of the plate using an Iodine spray. The areas of  
159 silica containing FAME were scraped into clean test tubes and FAME eluted using  
160 isohexane/diethyl ether (1:1). After centrifugation the supernatant solvent was removed into a clean  
161 test tube, the solvent evaporated under OFN and the FAME resuspended in 200 µL of isohexane.  
162 The FAME were separated and quantified by gas-liquid capillary chromatography using a Fisons  
163 GC8600 gas chromatograph (Fisons Ltd., Crawley, U.K) equipped with a 30 m x 0.32 mm i.d.  
164 capillary column (CP Wax 52CB, Chrompak Ltd., London, U.K.) and on-column injection.  
165 Hydrogen was used as carrier gas and temperature programming was from 50 °C to 150 °C at 40 °C  
166 min<sup>-1</sup> and then to 230 °C at 2.0 °C min<sup>-1</sup>. Individual methyl esters were identified by comparison  
167 with known standards and by reference to published data (Ackman, 1980; Tocher and Harvie,  
168 1988). Data were collected and processed using the Chromcard for Windows (Version 1.19)  
169 computer package (Thermoquest Italia S.p.A., Milan, Italy).

170

#### 171 *2. 5. Materials*

172 BHT was obtained from Sigma Chemical Co. (Poole, U.K.). HPTLC (10 cm x 10 cm x 0.15 mm)  
173 and TLC (20 cm x 20 cm x 0.25 mm) plates, precoated with silica gel 60 (without fluorescent

174 indicator) were obtained from Merck (Darmstadt, Germany). All solvents were HPLC grade and  
175 were obtained from Fisher Scientific UK (Loughborough, England).

176

## 177 2.6. Statistical analysis

178 All data are presented as means  $\pm$  SD (n = 3). The significance of differences between samples  
179 were determined by one-way analysis of variance (ANOVA) followed, where appropriate, by  
180 Tukey's comparison test (Zar, 1999). Percentage data and data that were identified as non-  
181 homogeneous (Bartlett's test) were subjected to arcsine transformation before analysis. Differences  
182 were regarded as significant when  $P < 0.05$ .

183

## 184 3. Results

### 185 3.1. Effect of location on lipid class and fatty acid compositions of sea lice from farmed salmon

186 Triacylglycerol (TAG) was the major lipid class in salmon lice, constituting between 35 – 45 % of  
187 total lipid with 9-13 % cholesterol. Similar amounts of the major polar lipids, phosphatidylcholine  
188 (PC) and phosphatidylethanolamine (PE) were seen, with lesser amounts of other  
189 phosphoglycerides such as phosphatidylserine (PS) and phosphatidylinositol (PI), and the major  
190 sphingolipid, sphingomyelin (Table 2). There were few major differences in lipid class composition  
191 between samples and, although TAG levels varied, no significant differences in TAG or the  
192 proportions of total polar and neutral lipids were apparent. Free fatty acids (FFA) varied between 8  
193 and 11 %, but these values were possibly partly artifactual resulting from *post-mortem* lipolytic  
194 action during homogenization and storage.

195 The fatty acid composition of the lice was characterized by about 25 % saturated fatty acids,  
196 predominantly 16:0, around 35 % monounsaturated fatty acids, predominantly 18:1n-9, 2-3 % n-6  
197 polyunsaturated fatty acids (PUFA) including 20:4n-6 (ARA, arachidonic acid), and 27-32 % n-3  
198 PUFA (Table 3). No major differences between the fatty acid compositions of lice obtained from  
199 farmed salmon at different sites were observed. Saturated fatty acids (mainly 16:0) showed little  
200 variation (23-25 %), whereas monounsaturated fatty acids (largely 18:1n-9) showed slightly more  
201 (30-40 %), although none of the differences were significant (Table 3). There was more variability  
202 in the levels of certain PUFA, particularly 18:2n-6 and the main n-3 long-chain polyunsaturated  
203 fatty acid (LC-PUFA), 22:6n-3 (DHA, docosahexaenoic acid) (Table 3). There were no significant  
204 differences in the levels of the other important LC-PUFA, ARA and 20:5n-3 (EPA,  
205 eicosapentaenoic acid) between lice from the different populations of farmed salmon (Table 3).

206

### 207 3.2. Lipid class and fatty acid compositions of female sea lice and their egg strings

208 Very clear differences between the relative lipid class compositions of females and the egg strings  
209 were observed, with eggs containing significantly higher proportions of TAG and neutral lipids than  
210 females (Table 4). Thus, the lipid in egg strings comprised 80 % neutral lipid, predominantly TAG  
211 at around 70 % and around 7 % cholesterol. Consequently the proportions of all polar lipid classes  
212 were significantly lower in egg strings compared to females although the relative proportions were  
213 similar with PC and PE predominating (Table 4).

214 Compared to the female lice, the fatty acid compositions of the egg strings were  
215 characterized by higher proportions of monounsaturated fatty acids, particularly 18:1n-9 but also  
216 20:1, and lower levels of total PUFA, specifically n-3 PUFA, particularly DHA (Table 5). Few  
217 differences in fatty acid composition of the egg strings obtained from the different locations could  
218 be discerned.

219

### 220 *3.3. Effect of year of sampling on lipid class and fatty acid compositions of sea lice from wild and* 221 *farmed salmon*

222 Irrespective of sampling time point, the lipid class compositions of lice obtained from farmed or  
223 wild salmon showed few differences (Table 6). Clear trends were apparent for higher cholesterol  
224 levels in lice obtained from farmed fish and also for lower levels of FFA in the 2009 samples (Table  
225 6). Relatively few major effects of louse source (wild / farmed) or year were apparent in the fatty  
226 acid compositions of the sea lice. However, there was a clear trend for higher 18:2n-6 and 18:3n-3  
227 in the lice from the farmed fish compared to lice from wild fish (Table 7). The farmed fish also  
228 tended to have lower 16:0, 16:1n-7 and EPA, and higher 22:1 and 20:4n-3 compared to the lice  
229 from wild salmon. The levels of 18:2n-6, 18:3n-3, 20:4n-3 and DHA were higher in lice from  
230 farmed fish in 2009 compared to 1995 (Table 7). In contrast, there was essentially no difference in  
231 the fatty acid composition of lice obtained from wild salmon in 1995 and those samples obtained in  
232 2009.

233 The host-parasite transfer of fatty acids can be observed by comparing the compositions of  
234 female lice, their egg strings and the muscle of host salmon (Table 8). The higher level of 18:2n-6  
235 in the lice from farmed fish in 2009 is a reflection of the relatively high level of this fatty acid (> 5  
236 %) in the salmon, and this fatty acid is also present in eggs. However, there are differences in the  
237 fatty acid compositions of the three samples. For instance the lice and eggs have generally higher  
238 saturated fatty acids and LC-PUFA (especially ARA and DHA), and lower monounsaturated fatty  
239 acids and C18 PUFA (18:2n-6 and 18:3n-3) compared to the salmon muscle (Table 8). Total  
240 phospholipids were characterized by higher percentages of n-3 PUFA, total PUFA and saturated  
241 fatty acids compared to TAG, whereas TAG showed higher proportions of monounsaturated fatty

242 acids and n-6 PUFA (Table 9). This was the pattern in female lice, egg strings and salmon muscle,  
243 although it was most pronounced in the fish muscle.

244

### 245 3.4. Lipid class and fatty acid compositions of sea lice from salmon, sea trout and halibut

246 Essentially no differences in lipid class composition between *L. salmonis* obtained from wild  
247 salmon and wild sea trout were significant with the difference in lyso-PC (a lipolytic product of PC)  
248 probably being due to differences in sampling. In contrast, *L. hippoglossi* was characterized by  
249 significantly lower levels of TAG and total neutral lipids compared to *L. salmonis* samples (Table  
250 10). This was also reflected in higher proportions of PC, PE and sphingomyelin in halibut lice  
251 (Table 10).

252 The halibut lice had generally lower levels of saturated fatty acids than lice obtained from  
253 salmon (Table 11). However, there was a very clear variation in the levels of 18:1n-9 and total  
254 monoenes between the louse species, with levels decreasing significantly from salmon to trout to  
255 halibut (Table 11). Other than 24:1n-9, all the monoenes were significantly lower in halibut lice  
256 compared to the salmonid lice. In contrast, the levels of n-6, n-3 and total PUFA were significantly  
257 higher in halibut lice compared to the salmonid lice (Table 11). Interestingly, the higher n-3 PUFA  
258 in halibut was entirely due to higher DHA as levels of EPA and 22:5n-3 were significantly lower in  
259 halibut lice compared to the salmonid lice.

260

## 261 4. Discussion

262 In many free-living copepods, periods of dietary surplus and extensive feeding, such as occurs  
263 during plankton blooms, may foster phases of rapid growth. In some habitats, however, such as  
264 those of polar regions, phytoplankton blooms have only short durations and so many animals build  
265 up significant stored lipid reserves to carry them through periods of low food availability. These  
266 stored reserves function to allow survival through dark winters with low primary production, and  
267 are also used for reproductive purposes. Such lipid storage has been particularly noted for many  
268 zooplankton species (Lee et al., 2006). TAG is by far the most common form of energy store in krill  
269 species (Lee et al., 2006) and, in addition, is used as the main lipid store in most fish and marine  
270 mammals such as seals. However, a peculiarity in many marine organisms, such as calanoid  
271 copepods, is the occurrence of wax esters (WEs) as the main lipid store (Lee et al., 2006). Wax  
272 esters comprise long-chain fatty acids esterified to long-chain alcohols and there has been  
273 considerable discussion as to their advantages as storage products. It is suggested that WEs are  
274 suited to use as long-term energy reserves while TAG can be used for short-term energy supplies, or  
275 that WEs may be used for buoyancy regulation (Lee et al., 2006). However, few animals have only  
276 WEs as the only storage lipid. In many cases, there will also be various levels of TAG present

277 depending on developmental stage. The present study has confirmed sparse earlier data (Lee, 1975)  
278 suggesting that parasitic caligid copepods of the genus *Lepeophtheirus* sp. store their lipid  
279 essentially as TAG with only small, trace amounts of WE. This perhaps supports the hypothesis that  
280 WEs are only required for long-term storage as the utilization of stored lipid in the parasitic  
281 copepods will only be required over a short time period between fish hosts.

282 The major phospholipid classes (membrane lipids) in the sea lice were PC and PE as in fish  
283 and most animals in general, and there were no major differences in the relative percentages of  
284 these membrane lipids. The major difference in lipid class composition observed between different  
285 samples was in the proportion of neutral (storage) lipid, TAG (Tocher, 2003). The major fatty acids  
286 observed in sea lice and their egg strings were 16:0, 18:1n-9, EPA and DHA, with lesser amounts of  
287 18:0, 16:1n-7, 20:1, ARA and 22:5n-3, which is a pattern characteristic of fish (Tocher, 2003).  
288 Therefore, the general fatty acid composition of lice is suggested to be a reflection of the fatty acid  
289 composition of the fish host.

290 Some of the differences in fatty acid composition observed between the samples were also  
291 reflective of the differences in lipid class composition discussed above. Neutral lipids like TAG  
292 generally have higher levels of monounsaturated fatty acids (monoenes) and lower PUFA as they  
293 are used primarily as an energy store, whereas phospholipids have higher PUFA as they are  
294 membrane lipids and PUFA are essential for membrane function, possibly in terms of membrane  
295 fluidity but more so for enzyme, receptor and carrier protein activities (Tocher, 2003). Lee (1975)  
296 reported the fatty acid composition of TAG and phospholipids from *L. salmonis* and showed that  
297 TAG had higher monoenes and lower PUFA than phospholipids. In the present study, this pattern  
298 was also clearly observed in the salmon muscle, female lice and egg string samples. Indeed all the  
299 samples in the present study that contained higher proportions of TAG were characterized by higher  
300 proportions of monoenes and lower proportions of PUFA and, especially, LC-PUFA. This was  
301 observed in the salmon louse versus egg string comparisons and also in the halibut versus salmonid  
302 lice comparisons.

303 It is probable, however, that the sea lice can modify the fatty acids obtained from the host.  
304 The adult female lice and the salmon muscle had similar levels of TAG (Table 8) but the sea lice  
305 showed lower levels of C18 PUFA, 18:2n-6 and 18:3n-3 and higher levels of the LC-PUFA ARA  
306 and DHA. This may indicate that they were able to convert the shorter chain fatty acids to the LC-  
307 PUFA by fatty acid desaturation and elongation, or, alternatively, differences in PUFA composition  
308 could be generated by selective oxidation of C18 PUFA and selective retention of LC-PUFA in lice  
309 (Tocher, 2003). Previously, Lee (1975) reported that the DHA:EPA ratio in *L. salmonis* lipids were  
310 2:1 in TAG and 6:1 in phospholipid. In the present study, the DHA:EPA ratios in *L. salmonis* and

311 their egg strings were around 8:1 and 3:1 in phospholipids and TAG, respectively, and only 4:1 and  
312 2:1 in salmon muscle phospholipids and TAG.

313 The fatty acid composition of fish, and animals in general, is most strongly influenced by  
314 diet (Tocher, 2003). As salmon lice are considered to obtain all their nutrition from the fish host,  
315 differences in the fatty acid composition of the diet of the fish host could similarly be reflected in  
316 the composition of the lice, as observed in the present study. Thus, the fatty acid composition of lice  
317 from farmed salmon in 2009 showed a difference compared to lice from farmed fish in 1995,  
318 specifically in terms of increased levels of 18:2n-6 and 18:3n-3. This is because feeds for salmon  
319 farming are being changed to more sustainable formulations, in particular replacing the marine  
320 resources, fishmeal and fish oil, global supplies of which are limiting, with more sustainable plant  
321 meals and vegetable oils (Tacon and Metian, 2008). Currently, around 25 % of fish meal and 50 %  
322 of fish oil is now generally substituted with plant alternatives in salmon feeds. Lipids in fishmeal  
323 and fish oil have high EPA and DHA and low C18 PUFA whereas plant meals and oils contain no  
324 LC-PUFA whatsoever, and high levels of C18 fatty acids, especially 18:1n-9 and 18:2n-6 but also  
325 18:3n-3 (Gunstone and Harwood, 2007). Therefore, the increased levels of C18 PUFA in the  
326 farmed samples from 2009 were expected but it was surprising that levels of EPA and DHA were  
327 maintained. This may be due to the specific fish oil included in the diets, as fish oils with higher  
328 LC-PUFA sourced from the southern hemisphere, may act to compensate for reduced levels of fish  
329 oil in the feed (Pratoomyot et al., 2008). In contrast, no differences in fatty acid compositions of lice  
330 from farmed and wild salmon were observed in 1995 samples due to fact that salmon feeds at that  
331 time comprised almost exclusively fishmeal and fish oil and, therefore, farmed fish would be  
332 getting essentially the same diet (fish) as the wild fish. Similarly, no difference was observed in the  
333 fatty acid composition of lice from wild fish sampled in 1995 and 2009. Differences in sea louse  
334 composition resulting from changes in salmon feed components may not be simply of academic  
335 interest. It is recognised, for instance, that *L. salmonis* secretes immuno-modulatory products  
336 including the prostaglandin E<sub>2</sub> (Fast et al., 2004), a derivative of the omega-6 fatty acid ARA,  
337 which may function to protect the parasite from the host's immune response. Changes in the fatty  
338 acid profile of the parasite could therefore potentially affect, in a positive or negative fashion, the  
339 interaction between parasite and host, leading to changes in infection intensity and / or host injury.

340 In conclusion, the present study has confirmed that parasitic caligid copepods of the genus  
341 *Lepeophtheirus* store their lipid essentially as TAG. Egg strings had higher lipid contents than adult  
342 female lice and this was reflected in higher TAG levels. Differences in fatty acid composition  
343 observed between the samples partly reflected the differences in lipid content and TAG levels but it  
344 is also likely that the endogenous metabolism of the lice modifies their fatty acid composition.  
345 Changes in the fatty acid composition of salmon feeds is reflected in the fatty acid composition of

346 the lice and their egg strings and this may have consequences for the ability of lice to suppress the  
347 host's immune response. The fact that changes to the lipid and fatty acid composition of feeds for  
348 farmed salmon influence the composition of sea lice indicates that further investigation of lipid and  
349 fatty acid metabolism of *L. salmonis* is warranted.

350

### 351 **Acknowledgement**

352 JAT was supported by a Nuffield Science Bursary from The Nuffield Foundation.

353

### 354 **References**

355

- 356 Ackman R.G., 1980. Fish lipids. In: *Advances in Fish Science and Technology* (ed. by J.J. Connell),  
357 pp. 87-103. Fishing News Books, Farnham.
- 358 Boxaspen, K., 2006. A review of the biology and genetics of sea lice. *ICES J. Mar. Sci.* 63, 1304-  
359 1316.
- 360 Brandal, P.O., Egidius, E., Romslo, I., 1976. Host blood: a major food component for the parasitic  
361 copepod *Lepeophtheirus salmonis*, Krøyer, 1838 (Crustacea: Caligidae). *Nor. J. Zool.* 24, 341-  
362 343.
- 363 Christie W.W., 1993. Preparation of derivatives of fatty acids for chromatographic analysis. In:  
364 Christie, W.W. (Ed.), *Advances in Lipid Methodology – Two*. The Oily Press, Dundee. pp.69-  
365 111.
- 366 Costello, M.J., 2009. The global economic cost of sea lice to the salmonid farming industry. *J. Fish*  
367 *Dis.* 32, 115-118.
- 368 Falk-Petersen, S., Mayzaud, P., Kattner, G., Sargent, J. 2009. Lipids and life strategy of Arctic  
369 *Calanus*. *Marine Biology Research*, 5, 18-39.
- 370 Fast, M.D., Ross, N.W., Craft, C.A., Locke, S.J., MacKinnon, S.L., Johnson, S.C., 2004.  
371 *Lepeophtheirus salmonis*: characterization of prostaglandin E-2 in secretory products of the  
372 salmon louse by Rp-HPLC and mass spectrometry. *Exp. Parasitol.* 107, 5-13.
- 373 Folch, J., Lees, M., Sloane-Stanley, G.H., 1957. A simple method for the isolation and purification  
374 of total lipids from animal tissues. *J. Biol. Chem.* 226, 497-509.
- 375 Ford, J.S., Myers, R.A., 2008. A global assessment of salmon aquaculture impacts on wild  
376 salmonids. *PLoS Biology* 6, e33.
- 377 Gunstone, F.D., Harwood, J.L., 2007. Occurrence and characteristics of oils and fats. In: *The Lipid*  
378 *Handbook* (Gunstone, F.D., Harwood, J.L. and Dijkstra, A.J., Eds.) pp.37-142. CRC Press,  
379 Taylor & Francis, London.

- 380 Henderson, R.J., Tocher, D.R., 1992. Thin-layer chromatography. In: *Lipid Analysis: A Practical*  
381 *Approach* (ed. by R.J. Hamilton and S. Hamilton), pp. 65-111. IRL Press, Oxford, UK.
- 382 Jonsdottir, H., Bron, J.E., Wootten, R., Turnbull, J.T., 1992. The histopathology associated with the  
383 preadult and adult stages of *Lepeophtheirus salmonis* on the Atlantic salmon *Salmo salar* L. J.  
384 Fish Dis. 15, 521-527.
- 385 Johnson, S.C., Treasurer, J.W., Bravo, S., Nagasawa, K., Kabata, Z., 2004. A review of the impact  
386 of parasitic copepods on marine aquaculture. *Zoological Studies* 43, 229-243.
- 387 Kabata, Z., 1979. Parasitic Copepoda of British Fishes. Ray Society, London. 468 pp.
- 388 Kattner, G. and Krause, M. 1989. Seasonal variations of lipids (wax esters, fatty acids and alcohols)  
389 in Calanoid copepods from the North Sea. *Mar. Chem.*, 26, 261-275.
- 390 Kattner, G., Hagen, W., Lee, R.F., Campbell, R., Deibel, D., Falk-Petersen, S., Graeve, M., Hansen,  
391 B.W., Hirche, H.J., Jonsdottir, S.H., Madsen, M.L., Mayzaud, P., Muller-Navarra, D., Nichols,  
392 P.D., Paffenhofer, G.-A., Pond, D., Saito, H., Stubing, D., Virtue, P., 2007. Perspectives on  
393 marine zooplankton lipids. *Can. J. Fish. Aquat. Sci.* 64, 1628-1639.
- 394 Lee, R.F., 1975. Lipids of parasitic copepods associated with marine fish. *Comp. Biochem. Physiol.*  
395 52B, 363-364.
- 396 Lee, R.F., Hagen, W., Kattner, G., 2006. Lipid storage in marine zooplankton. *Mar. Ecol. Prog. Ser.*  
397 307, 273-306.
- 398 Pike, A.W., Wadsworth, S.L., 1999. Sealice on salmonids: Their biology and control. *Adv.*  
399 *Parasitol.* 44, 233-337.
- 400 Pratoomyot, J., Bendiksen, E.Å., Bell, J.G., Tocher, D.R., 2008. Comparison of effects of blended  
401 vegetable oils and contaminant-stripped fish oil on growth performance, composition, and  
402 gene expression in Atlantic salmon (*Salmo salar* L.). *Aquaculture* 280, 170-178.
- 403 Ritchie, G., 1997. The host transfer ability of *Lepeophtheirus salmonis* (Copepoda: Caligidae)  
404 from farmed Atlantic salmon, *Salmo salar* L. *J. Fish Dis.* 20, 153-157.
- 405 Rohde, K. (ed.), 2005. *Marine Parasitology*. CSIRO Publishing-CABI, Collingwood, Australia,  
406 565 p.
- 407 Schram, T.A., Supplemental descriptions of the developmental stages of *Lepeophtheirus*  
408 *salmonis* (Krøyer, 1837) (Copepoda: Caligidae). In Boxshall, G.A., Defaye, D. (Eds.),  
409 *Pathogens of Wild and Farmed Fish: Sea Lice*. Ellis Horwood: New York, pp. 30-47.
- 410 Skjoldal, H.R., Dalpadado, P., Dommasnes, A. 2004. Food webs and trophic interactions. In:  
411 Skjoldal, H. R., Sætre, R., Færnø, A., Misund, O. A., Røttingen, I. (Eds.), *The Norwegian*  
412 *Sea Ecosystem*. Trondheim, Norway: Tapir Academic Press. 447-506.
- 413 Tacon, A.G.J., Meitan, M., 2008. Global overview on the use of fish meal and fish oil in  
414 industrially compounded aquafeeds: Trends and future prospects. *Aquaculture* 285, 146-158.

- 415 Tocher D.R., 2003. Metabolism and functions of lipids and fatty acids in teleost fish. Rev. Fisheries  
416 Sci. 11, 107-184.
- 417 Tocher, D.R., Harvie, D.G., 1988. Fatty acid compositions of the major phosphoglycerides from  
418 fish neural tissues : (n-3) and (n-6) polyunsaturated fatty acids in rainbow trout (*Salmo*  
419 *gairdneri*, L.) and cod (*Gadus morhua*) brains and retinas. Fish Physiol. Biochem. 5, 229-  
420 239.
- 421 Tucker, C., Sommerville, C., Wootten, R., 2000. An investigation onto the larval energetics and  
422 settlement of sea louse, *Lepeophtheirus salmonis*, an ectoparasitic copepod of Atlantic salmon,  
423 *Salmo salar*. Fish Path. 35, 137-143.
- 424 Zar J.H., 1999. Biostatistical Analysis 4<sup>th</sup> Edition; Prentice-Hall; Upper Saddle River, New Jersey.  
425

Table 1. Details of the *Lepeophtheirus* von Nordmann, 1832 samples collected in Scottish waters and used for the current study.

Species	Host	Locality (date)	Latitude / longitude	Samples analysed
<i>Lepeophtheirus hippoglossi</i> (Krøyer, 1837)	<i>Hippoglossus hippoglossus</i> L. (W)	North Atlantic (06/98)	58° 52' 56.87"N / 7° 27' 04.63"W	Adult female lice (S)
<i>Lepeophtheirus salmonis</i> (Krøyer, 1837)	<i>Salmo trutta</i> L. (W)	River Ewe (07/96)	57° 50' 23.43"N / 5° 34' 56.21"W	Adult female lice (S)
	<i>Salmo salar</i> L. (W)	Loch Duich (06/95)	57° 13' 48.41"N / 5° 28' 02.04"W	Adult female lice (S)
		Armadale, Skye (06/09)	57° 03' 30.90"N / 5° 53' 09.19"W	Adult female lice (NS)
	<i>Salmo salar</i> L. (F)	Loch Duich (06/95)	57° 14' 49.70"N / 5° 29' 00.19"W	Adult female lice (S)
		Loch Fyne (site 2; 04/95)	56° 13' 40.35"N / 5° 02' 30.38"W	Adult female lice (S)
		Loch Fyne (site 1; 05/95)	56° 04' 02.78"N / 5° 17' 25.78"W	Eggstrings
		Loch na Keal, Mull (06/95)	56° 26' 02.05"N / 6° 12' 13.94"W	Adult female lice (S)
		Loch Linnhe (04/95)	56° 31' 38.61"N / 5° 32' 42.41"W	Adult female lice (S)
		Lumlash Bay, Arran (05/95)	55° 31' 48.38"N / 5° 06' 15.2"W	Adult female lice (S)
		Lumlash Bay, Arran (05/95)	55° 31' 48.38"N / 5° 06' 15.2"W	Eggstrings
		Shuna (04/95)	56° 13' 49.68"N / 5° 35' 20.15"W	Eggstrings
	Machrihanish (07/09)	55° 25' 24.45"N / 5° 44' 54.40"W	Adult female lice (S)	

F, Farmed; NS, not starved; S, starved for 24 hrs prior to processing; W, wild.

425

426

Table 2. Lipid class composition (percentage of total lipid) of *L. salmonis* obtained from Atlantic salmon farmed at various locations.

Lipid class	Duich	Linnhe	Mull	Arran	Shuna	Fyne
PC	13.5 ± 2.9	14.3 ± 1.5	14.6 ± 2.6	12.7 ± 0.3	12.6 ± 1.3	15.9 ± 0.9
PE	14.2 ± 1.8	12.2 ± 3.8	13.3 ± 0.9	10.8 ± 2.0	12.6 ± 1.9	14.6 ± 0.9
PS/PI/PA/CL	9.9 ± 1.9	7.8 ± 2.8	10.1 ± 3.8	7.7 ± 1.6	7.4 ± 2.7	8.0 ± 1.6
Sphingomyelin	3.5 ± 1.2	4.4 ± 0.9	3.5 ± 1.9	3.2 ± 0.3	2.9 ± 0.8	4.2 ± 0.2
LPC	0.5 ± 0.3 <sup>b</sup>	0.2 ± 0.2 <sup>b</sup>	0.8 ± 0.3 <sup>ab</sup>	1.2 ± 0.1 <sup>a</sup>	0.7 ± 0.2 <sup>ab</sup>	0.7 ± 0.2 <sup>ab</sup>
Total polar	41.5 ± 7.4	38.9 ± 5.0	42.2 ± 8.5	35.6 ± 3.5	36.2 ± 6.7	43.4 ± 2.5
Total neutral	58.5 ± 7.4	61.1 ± 5.0	57.8 ± 8.5	64.4 ± 3.5	63.8 ± 6.7	56.6 ± 2.5
Cholesterol	13.3 ± 0.8 <sup>a</sup>	10.8 ± 1.1 <sup>ab</sup>	12.7 ± 1.6 <sup>a</sup>	8.7 ± 0.5 <sup>b</sup>	11.7 ± 2.0 <sup>ab</sup>	10.9 ± 0.6 <sup>ab</sup>
Triacylglycerol	34.7 ± 8.4	40.8 ± 5.7	36.0 ± 10.6	45.4 ± 5.3	44.4 ± 6.5	35.0 ± 3.2
Free fatty acid	10.4 ± 1.8	9.4 ± 0.6	9.1 ± 3.4	10.2 ± 1.4	7.7 ± 1.4	10.7 ± 2.7
Steryl/wax ester	Trace	Trace	Trace	Trace	Trace	Trace

Results are means ± SD (n = 3). Values within a row with different superscript letters are significantly different (P < 0.05).

CL, cardiolipin; LPC, lyso-PC; PA, phosphatidic acid; PC, phosphatidylcholine; PE, phosphatidylethanolamine; PI, phosphatidylinositol; PS, phosphatidylserine.

426

427

Table 3. Fatty acid composition (percentage of total fatty acids) of *L. salmonis* obtained from Atlantic salmon farmed at various locations.

Fatty acid	Duich	Linnhe	Mull	Arran	Shuna	Fyne
14:0	2.1 ± 0.0 <sup>b</sup>	2.5 ± 0.2 <sup>b</sup>	2.2 ± 0.3 <sup>b</sup>	2.4 ± 0.2 <sup>b</sup>	2.4 ± 0.2 <sup>b</sup>	3.1 ± 0.1 <sup>a</sup>
16:0	17.9 ± 0.5	17.1 ± 2.8	16.8 ± 0.7	17.1 ± 0.9	16.6 ± 0.9	16.9 ± 1.9
18:0	3.9 ± 0.5	3.7 ± 0.8	3.7 ± 0.2	3.4 ± 0.3	2.9 ± 0.3	3.9 ± 0.9
Total saturated <sup>1</sup>	25.4 ± 0.9	24.1 ± 3.6	23.5 ± 1.0	23.5 ± 0.5	22.6 ± 0.6	25.0 ± 3.5
16:1n-9	1.7 ± 0.4	1.8 ± 1.2	2.1 ± 0.3	2.0 ± 0.2	2.2 ± 0.5	3.0 ± 0.1
16:1n-7	2.3 ± 0.2 <sup>b</sup>	3.1 ± 1.1 <sup>ab</sup>	2.7 ± 0.3 <sup>ab</sup>	3.1 ± 0.2 <sup>ab</sup>	3.7 ± 0.4 <sup>ab</sup>	3.6 ± 0.1 <sup>a</sup>
18:1n-9	22.5 ± 2.5 <sup>ab</sup>	24.8 ± 3.2 <sup>ab</sup>	23.2 ± 3.7 <sup>ab</sup>	28.2 ± 1.9 <sup>a</sup>	25.4 ± 2.7 <sup>ab</sup>	19.0 ± 3.7 <sup>b</sup>
18:1n-7	1.4 ± 0.2	1.4 ± 0.0	1.3 ± 0.3	1.6 ± 0.1	1.4 ± 0.2	1.5 ± 0.3
20:1 <sup>2</sup>	2.8 ± 0.5	2.9 ± 0.3	2.2 ± 0.2	2.7 ± 0.2	2.4 ± 0.1	2.6 ± 0.1
22:1 <sup>3</sup>	2.1 ± 0.4 <sup>a</sup>	1.0 ± 0.1 <sup>b</sup>	1.1 ± 0.3 <sup>b</sup>	1.0 ± 0.1 <sup>b</sup>	1.3 ± 0.1 <sup>b</sup>	1.4 ± 0.5 <sup>ab</sup>
24:1n-9	2.5 ± 1.2 <sup>a</sup>	1.7 ± 0.1 <sup>ab</sup>	1.2 ± 0.0 <sup>b</sup>	0.9 ± 0.2 <sup>b</sup>	0.9 ± 0.1 <sup>b</sup>	0.7 ± 0.3 <sup>b</sup>
Total monoenes	35.2 ± 3.2	36.6 ± 4.8	33.8 ± 4.0	39.6 ± 1.6	37.3 ± 2.2	30.0 ± 5.3
18:2n-6	1.0 ± 0.1 <sup>ab</sup>	0.4 ± 0.0 <sup>d</sup>	1.2 ± 0.1 <sup>a</sup>	0.8 ± 0.0 <sup>bc</sup>	0.6 ± 0.0 <sup>cd</sup>	1.4 ± 0.4 <sup>a</sup>
20:4n-6	1.2 ± 0.4	1.0 ± 0.1	1.0 ± 0.4	1.3 ± 0.2	0.9 ± 0.1	0.9 ± 0.1
Total n-6PUFA <sup>4</sup>	2.9 ± 0.5 <sup>a</sup>	2.0 ± 0.1 <sup>b</sup>	2.6 ± 0.5 <sup>ab</sup>	2.6 ± 0.3 <sup>ab</sup>	1.8 ± 0.1 <sup>b</sup>	2.8 ± 0.3 <sup>a</sup>
18:3n-3	0.3 ± 0.0 <sup>bc</sup>	0.3 ± 0.0 <sup>bc</sup>	0.5 ± 0.0 <sup>c</sup>	0.3 ± 0.0 <sup>bc</sup>	0.4 ± 0.0 <sup>b</sup>	0.7 ± 0.2 <sup>a</sup>
20:4n-3	0.6 ± 0.1 <sup>ab</sup>	0.4 ± 0.2 <sup>b</sup>	0.7 ± 0.1 <sup>b</sup>	0.7 ± 0.1 <sup>ab</sup>	0.8 ± 0.1 <sup>a</sup>	0.6 ± 0.1 <sup>ab</sup>
20:5n-3	4.3 ± 0.8	4.6 ± 1.1	4.8 ± 0.6	4.7 ± 0.1	5.2 ± 0.2	4.0 ± 0.4
22:5n-3	3.0 ± 0.5 <sup>ab</sup>	2.5 ± 0.9 <sup>b</sup>	3.5 ± 0.4 <sup>a</sup>	3.3 ± 0.6 <sup>ab</sup>	3.0 ± 0.3 <sup>ab</sup>	2.3 ± 0.7 <sup>b</sup>
22:6n-3	18.4 ± 3.1 <sup>b</sup>	20.8 ± 4.6 <sup>ab</sup>	25.3 ± 3.4 <sup>ab</sup>	20.6 ± 0.4 <sup>ab</sup>	23.0 ± 0.8 <sup>ab</sup>	25.8 ± 2.0 <sup>a</sup>
Total n-3PUFA <sup>5</sup>	26.9 ± 4.5	28.8 ± 6.0	35.1 ± 3.7	29.9 ± 0.8	32.7 ± 0.2	33.6 ± 3.0
Total PUFA	29.7 ± 4.5	30.8 ± 6.0	37.7 ± 3.7	32.5 ± 1.1	34.5 ± 0.1	36.4 ± 2.8

Results are means ± SD (n = 3). Values within a row with different superscript letters are significantly different (P < 0.05).

<sup>1</sup>, Totals include 15:0, 20:0 and 22:0 present in some samples at up to 0.7 %; <sup>2</sup>, Predominantly n-9 isomer;

<sup>3</sup>, Predominantly n-11 isomer; <sup>4</sup>, Totals include 18:3n-6, 20:2n-6, 20:3n-6, 22:4n-6 and 22:5n-6 present in some samples at up to 0.3 %; <sup>5</sup>, Totals include 18:3n-3 and 20:3n-3 present in some samples at up to 0.2 %; PUFA, polyunsaturated fatty acids.

427

428

Table 4. Lipid class compositions (percentage of total lipid) of *L. salmonis* females and their egg strings

Lipid class	Females			Egg strings		
	Arran	Shuna	Fyne	Arran	Shuna	Fyne
PC	12.7 ± 0.3 <sup>a</sup>	12.6 ± 1.3 <sup>a</sup>	15.9 ± 0.9 <sup>a</sup>	8.7 ± 0.8 <sup>b</sup>	9.3 ± 0.9 <sup>b</sup>	9.6 ± 0.8 <sup>b</sup>
PE	10.8 ± 2.0 <sup>a</sup>	12.6 ± 1.9 <sup>a</sup>	14.6 ± 0.9 <sup>a</sup>	6.4 ± 0.1 <sup>b</sup>	5.9 ± 0.6 <sup>b</sup>	6.4 ± 1.6 <sup>b</sup>
PS/PI/PA/CL	7.7 ± 1.6 <sup>a</sup>	7.4 ± 2.7 <sup>a</sup>	8.0 ± 1.6 <sup>a</sup>	4.5 ± 0.7 <sup>ab</sup>	2.8 ± 1.7 <sup>b</sup>	2.5 ± 0.7 <sup>b</sup>
Sphingomyelin	3.2 ± 0.3 <sup>a</sup>	2.9 ± 0.8 <sup>a</sup>	4.2 ± 0.2 <sup>a</sup>	1.4 ± 0.3 <sup>b</sup>	1.6 ± 0.2 <sup>b</sup>	1.5 ± 0.2 <sup>b</sup>
LPC	1.2 ± 0.1	0.7 ± 0.2	0.7 ± 0.2	0.3 ± 0.2	0.2 ± 0.3	0.5 ± 0.8
Total polar	35.6 ± 3.5 <sup>a</sup>	36.2 ± 6.7 <sup>a</sup>	43.4 ± 2.5 <sup>a</sup>	21.2 ± 0.2 <sup>b</sup>	19.7 ± 2.1 <sup>b</sup>	20.5 ± 2.5 <sup>b</sup>
Total neutral	64.4 ± 3.5 <sup>b</sup>	63.8 ± 6.7 <sup>b</sup>	56.6 ± 2.5 <sup>b</sup>	78.8 ± 0.2 <sup>a</sup>	80.3 ± 2.1 <sup>a</sup>	79.5 ± 2.5 <sup>a</sup>
Cholesterol	8.7 ± 0.5 <sup>b</sup>	11.7 ± 2.0 <sup>a</sup>	10.9 ± 0.6 <sup>a</sup>	7.0 ± 0.4 <sup>b</sup>	7.1 ± 0.2 <sup>b</sup>	6.6 ± 0.1 <sup>b</sup>
Triacylglycerol	45.4 ± 5.3 <sup>b</sup>	44.4 ± 6.5 <sup>b</sup>	35.0 ± 3.2 <sup>b</sup>	67.1 ± 2.3 <sup>a</sup>	70.2 ± 3.8 <sup>a</sup>	69.3 ± 3.0 <sup>a</sup>
Free fatty acid	10.2 ± 1.4 <sup>a</sup>	7.7 ± 1.4 <sup>ab</sup>	10.7 ± 2.7 <sup>a</sup>	4.7 ± 2.0 <sup>bc</sup>	3.1 ± 1.5 <sup>c</sup>	3.5 ± 0.7 <sup>c</sup>
Steryl/wax ester	Trace	Trace	Trace	Trace	Trace	Trace

Results are means ± SD (n = 3). Values within a row with different superscript letters are significantly different (P < 0.05).

CL, cardiolipin; LPC, lyso-PC; PA, phosphatidic acid; PC, phosphatidylcholine; PE, phosphatidylethanolamine;

PI, phosphatidylinositol; PS, phosphatidylserine.

428

429

Table 5. Fatty acid compositions (percentage of total fatty acids) of *L. salmonis* females and their egg strings

Fatty acid	Females			Egg strings		
	Arran	Shuna	Fyne	Arran	Shuna	Ardcastle
14:0	2.4 ± 0.2 <sup>b</sup>	2.4 ± 0.2 <sup>b</sup>	3.1 ± 0.1 <sup>a</sup>	2.4 ± 0.2 <sup>b</sup>	2.5 ± 0.2 <sup>b</sup>	1.5 ± 0.1 <sup>c</sup>
16:0	17.1 ± 0.9 <sup>ab</sup>	16.6 ± 0.9 <sup>ab</sup>	16.9 ± 1.9 <sup>ab</sup>	17.4 ± 1.4 <sup>a</sup>	16.9 ± 0.7 <sup>ab</sup>	14.5 ± 0.8 <sup>b</sup>
18:0	3.4 ± 0.3	2.9 ± 0.3	3.9 ± 0.9	3.2 ± 0.3	2.9 ± 0.1	3.4 ± 0.3
Total saturated <sup>1</sup>	23.5 ± 0.5 <sup>ab</sup>	22.6 ± 0.6 <sup>ab</sup>	25.0 ± 3.5 <sup>a</sup>	24.3 ± 2.6 <sup>a</sup>	23.2 ± 1.0 <sup>ab</sup>	19.9 ± 1.3 <sup>b</sup>
16:1n-9	2.0 ± 0.2 <sup>b</sup>	2.2 ± 0.5 <sup>b</sup>	3.0 ± 0.1 <sup>a</sup>	1.7 ± 0.1 <sup>b</sup>	2.0 ± 0.4 <sup>b</sup>	1.8 ± 0.1 <sup>b</sup>
16:1n-7	3.1 ± 0.2 <sup>b</sup>	3.7 ± 0.4 <sup>ab</sup>	3.6 ± 0.1 <sup>ab</sup>	3.5 ± 0.1 <sup>ab</sup>	3.9 ± 0.2 <sup>a</sup>	3.6 ± 0.3 <sup>ab</sup>
18:1n-9	28.2 ± 1.9 <sup>ab</sup>	25.4 ± 2.7 <sup>b</sup>	19.0 ± 3.7 <sup>b</sup>	30.5 ± 0.7 <sup>a</sup>	30.9 ± 1.9 <sup>a</sup>	30.7 ± 0.4 <sup>a</sup>
18:1n-7	1.6 ± 0.1 <sup>b</sup>	1.4 ± 0.2 <sup>b</sup>	1.5 ± 0.3 <sup>b</sup>	1.7 ± 0.1 <sup>b</sup>	1.5 ± 0.2 <sup>b</sup>	2.1 ± 0.0 <sup>a</sup>
20:1 <sup>2</sup>	2.7 ± 0.2 <sup>bc</sup>	2.4 ± 0.1 <sup>c</sup>	2.6 ± 0.1 <sup>bc</sup>	3.2 ± 0.1 <sup>ab</sup>	3.2 ± 0.2 <sup>ab</sup>	3.5 ± 0.3 <sup>a</sup>
22:1 <sup>3</sup>	1.0 ± 0.1	1.3 ± 0.1	1.4 ± 0.5	1.1 ± 0.5	1.4 ± 0.2	1.1 ± 0.1
24:1n-9	0.9 ± 0.2	0.9 ± 0.1	0.7 ± 0.3	0.9 ± 0.1	1.0 ± 0.2	1.1 ± 0.3
Total monoenes	39.6 ± 1.6 <sup>bc</sup>	37.3 ± 2.2 <sup>c</sup>	30.0 ± 5.3 <sup>c</sup>	42.6 ± 0.4 <sup>ab</sup>	43.9 ± 2.3 <sup>ab</sup>	44.0 ± 0.6 <sup>a</sup>
18:2n-6	0.8 ± 0.0 <sup>b</sup>	0.6 ± 0.0 <sup>b</sup>	1.4 ± 0.4 <sup>a</sup>	0.7 ± 0.0 <sup>b</sup>	0.5 ± 0.0 <sup>b</sup>	0.7 ± 0.0 <sup>b</sup>
20:4n-6	1.3 ± 0.2 <sup>ab</sup>	0.9 ± 0.1 <sup>c</sup>	0.9 ± 0.1 <sup>c</sup>	1.4 ± 0.1 <sup>a</sup>	1.0 ± 0.1 <sup>bc</sup>	1.5 ± 0.2 <sup>a</sup>
Total n-6PUFA <sup>4</sup>	2.6 ± 0.3 <sup>b</sup>	1.8 ± 0.1 <sup>c</sup>	2.8 ± 0.3 <sup>ab</sup>	2.6 ± 0.2 <sup>b</sup>	1.8 ± 0.1 <sup>c</sup>	3.1 ± 0.1 <sup>a</sup>
18:3n-3	0.3 ± 0.0 <sup>b</sup>	0.4 ± 0.0 <sup>b</sup>	0.7 ± 0.2 <sup>a</sup>	0.4 ± 0.0 <sup>b</sup>	0.3 ± 0.0 <sup>b</sup>	0.3 ± 0.0 <sup>b</sup>
20:4n-3	0.7 ± 0.1	0.8 ± 0.1	0.6 ± 0.1	0.7 ± 0.1	0.7 ± 0.1	0.6 ± 0.1
20:5n-3	4.7 ± 0.1	5.2 ± 0.2	4.0 ± 0.4	4.8 ± 0.3	5.1 ± 0.5	5.2 ± 0.8
22:5n-3	3.3 ± 0.6	3.0 ± 0.3	2.3 ± 0.7	2.5 ± 0.6	2.4 ± 0.1	2.7 ± 0.5
22:6n-3	20.6 ± 0.4 <sup>ab</sup>	23.0 ± 0.8 <sup>a</sup>	25.8 ± 2.0 <sup>a</sup>	17.3 ± 1.9 <sup>b</sup>	17.6 ± 0.4 <sup>b</sup>	18.8 ± 3.1 <sup>ab</sup>
Total n-3PUFA <sup>5</sup>	29.9 ± 0.8 <sup>ab</sup>	32.7 ± 0.2 <sup>a</sup>	33.6 ± 3.0 <sup>a</sup>	25.9 ± 2.9 <sup>b</sup>	26.4 ± 1.0 <sup>ab</sup>	27.8 ± 4.4 <sup>ab</sup>
Total PUFA	32.5 ± 1.1	34.5 ± 0.1	36.4 ± 2.8	28.5 ± 3.1	28.2 ± 1.1	30.9 ± 4.4

Results are means ± SD (n = 3). Values within a row with different superscript letters are significantly different (P < 0.05).

<sup>1</sup>, Totals include 15:0, 20:0 and 22:0 present in some samples at up to 0.7 %; <sup>2</sup>, Predominantly n-9 isomer; <sup>3</sup>, Predominantly n-11 isomer; <sup>4</sup>, Totals include 18:3n-6, 20:2n-6, 20:3n-6, 22:4n-6 and 22:5n-6 present in some samples at up to 0.3 %;

<sup>5</sup>, Totals include 18:3n-3 and 20:3n-3 present in some samples at up to 0.2 %; PUFA, polyunsaturated fatty acids.

429

430

Table 6. Lipid class composition (percentage of total lipid) of *L. salmonis* obtained from wild and farmed Atlantic salmon in 1995 and 2009

Lipid class	Wild		Farmed	
	1995	2009	1995	2009
PC	13.9 ± 1.5	12.5 ± 0.1	13.5 ± 2.9	13.2 ± 0.5
PE	14.1 ± 2.6	12.6 ± 1.7	14.2 ± 1.8	13.3 ± 1.6
PS/PI/PA/CL	5.2 ± 3.0	6.3 ± 4.5	9.9 ± 1.9	6.8 ± 0.6
Sphingomyelin	3.7 ± 0.8	3.3 ± 0.3	3.5 ± 1.2	3.1 ± 0.2
LPC	0.3 ± 0.2 <sup>ab</sup>	0.5 ± 0.1 <sup>a</sup>	0.5 ± 0.3 <sup>a</sup>	0.0 ± 0.0 <sup>b</sup>
Total polar	37.2 ± 5.3	35.2 ± 3.1	41.5 ± 7.4	36.5 ± 2.0
Total neutral	62.8 ± 5.3	64.8 ± 3.1	58.5 ± 7.4	63.5 ± 2.0
Cholesterol	10.5 ± 0.7 <sup>b</sup>	8.3 ± 0.5 <sup>c</sup>	13.3 ± 0.8 <sup>a</sup>	10.6 ± 0.3 <sup>b</sup>
Triacylglycerol	42.2 ± 7.5	49.1 ± 4.5	34.7 ± 8.4	47.1 ± 2.8
Free fatty acid	10.1 ± 1.5 <sup>a</sup>	7.4 ± 1.3 <sup>ab</sup>	10.4 ± 1.8 <sup>a</sup>	5.8 ± 0.5 <sup>b</sup>
Steryl/wax ester	Trace	Trace	Trace	Trace

Both wild and farmed samples were obtained from Loch Duich in 1995, and from Armadale (wild) and Machrihanish (farmed) in 2009. Results are means ± SD (n = 3).

Values within a row with different superscript letters are significantly different ( $P < 0.05$ ).

CL, cardiolipin; LPC, lyso-PC; PA, phosphatidic acid; PC, phosphatidylcholine; PE, phosphatidylethanolamine; PI, phosphatidylinositol; PS, phosphatidylserine.

Table 7. Fatty acid composition (percentage of total fatty acids) of *L. salmonis* obtained from wild and farmed Atlantic salmon in 1995 and 2009.

Fatty acid	Wild		Farmed	
	1995	2009	1995	2009
14:0	2.0 ± 0.1 <sup>ab</sup>	2.0 ± 0.1 <sup>ab</sup>	2.1 ± 0.0 <sup>a</sup>	1.8 ± 0.1 <sup>b</sup>
16:0	18.7 ± 0.7 <sup>a</sup>	18.3 ± 1.4 <sup>ab</sup>	17.9 ± 0.5 <sup>ab</sup>	16.2 ± 0.3 <sup>b</sup>
18:0	3.8 ± 0.1	3.7 ± 0.4	3.9 ± 0.5	3.6 ± 0.1
Total saturated <sup>1</sup>	25.1 ± 1.1	24.7 ± 2.1	25.4 ± 0.9	22.6 ± 0.3
16:1n-9	1.7 ± 0.3	1.8 ± 0.1	1.7 ± 0.4	1.5 ± 0.1
16:1n-7	2.7 ± 0.1 <sup>b</sup>	3.2 ± 0.1 <sup>a</sup>	2.3 ± 0.2 <sup>c</sup>	2.1 ± 0.0 <sup>c</sup>
18:1n-9	25.0 ± 1.2	27.0 ± 2.7	22.5 ± 2.5	21.8 ± 1.0
18:1n-7	1.5 ± 0.1	1.7 ± 0.1	1.4 ± 0.2	1.4 ± 0.3
20:1 <sup>2</sup>	2.2 ± 0.2	2.4 ± 0.1	2.8 ± 0.5	2.2 ± 0.1
22:1 <sup>3</sup>	1.0 ± 0.0 <sup>b</sup>	0.6 ± 0.1 <sup>c</sup>	2.1 ± 0.4 <sup>a</sup>	1.0 ± 0.0 <sup>b</sup>
24:1n-9	1.1 ± 0.1	1.1 ± 0.1	2.5 ± 1.2	1.2 ± 0.1
Total monoenes	35.3 ± 1.2	37.6 ± 2.9	35.2 ± 3.2	31.2 ± 1.0
18:2n-6	0.4 ± 0.1 <sup>c</sup>	0.3 ± 0.0 <sup>c</sup>	1.0 ± 0.1 <sup>b</sup>	2.2 ± 0.1 <sup>a</sup>
20:4n-6	1.2 ± 0.3	1.4 ± 0.1	1.2 ± 0.4	1.5 ± 0.1
Total n-6PUFA <sup>4</sup>	2.3 ± 0.1	2.0 ± 0.2	2.9 ± 0.5	4.4 ± 0.2
18:3n-3	0.2 ± 0.0 <sup>c</sup>	0.2 ± 0.0 <sup>c</sup>	0.3 ± 0.0 <sup>b</sup>	0.5 ± 0.0 <sup>a</sup>
20:4n-3	0.4 ± 0.1 <sup>b</sup>	0.4 ± 0.1 <sup>b</sup>	0.6 ± 0.1 <sup>b</sup>	1.0 ± 0.0 <sup>a</sup>
20:5n-3	5.1 ± 0.2 <sup>ab</sup>	6.0 ± 0.5 <sup>a</sup>	4.3 ± 0.8 <sup>b</sup>	5.5 ± 0.1 <sup>ab</sup>
22:5n-3	4.3 ± 0.3 <sup>a</sup>	3.5 ± 0.3 <sup>ab</sup>	3.0 ± 0.5 <sup>b</sup>	3.5 ± 0.2 <sup>ab</sup>
22:6n-3	22.1 ± 1.2 <sup>ab</sup>	20.9 ± 2.7 <sup>ab</sup>	18.4 ± 3.1 <sup>b</sup>	25.7 ± 1.2 <sup>a</sup>
Total n-3PUFA <sup>5</sup>	32.2 ± 1.6 <sup>ab</sup>	31.1 ± 3.5 <sup>ab</sup>	26.9 ± 4.5 <sup>b</sup>	36.8 ± 1.3 <sup>a</sup>
Total PUFA	34.5 ± 1.6 <sup>ab</sup>	33.1 ± 3.8 <sup>ab</sup>	29.7 ± 4.5 <sup>b</sup>	41.1 ± 1.4 <sup>a</sup>

Both wild and farmed samples were obtained from Loch Duich in 1995, and from Armadale (wild) and Machrihanish (farmed) in 2009. Results are means ± SD (n = 3).

Values within a row with different superscript letters are significantly different (P < 0.05).

<sup>1</sup>, Totals include 15:0, 20:0 and 22:0 present in some samples at up to 0.7 %;

<sup>2</sup>, Predominantly n-9 isomer; <sup>3</sup>, Predominantly n-11 isomer; <sup>4</sup>, Totals include 18:3n-6, 20:2n-6, 20:3n-6, 22:4n-6 and 22:5n-6 present in some samples at up to 0.3 %; <sup>5</sup>, Totals include 18:3n-3 and 20:3n-3 present in some samples at up to 0.3 %; PUFA, polyunsaturated fatty acids.

Table 8. Lipid content (percentage of wet weight), triacylglycerol (TAG) content (percentage of total lipid), and fatty acid composition (percentage of total fatty acids) of *L. salmonis* and their egg strings along with the muscle of the Atlantic salmon from which they were collected.

Fatty acid	Salmon muscle	Female lice	Egg strings
Lipid content	3.5 ± 0.1 <sup>b</sup>	1.6 ± 0.2 <sup>c</sup>	6.1 ± 1.2 <sup>a</sup>
TAG	54.4 ± 4.4 <sup>b</sup>	47.1 ± 2.8 <sup>b</sup>	72.0 ± 1.4 <sup>a</sup>
14:0	3.2 ± 0.5 <sup>a</sup>	1.8 ± 0.1 <sup>b</sup>	1.9 ± 0.1 <sup>b</sup>
16:0	11.6 ± 0.2 <sup>b</sup>	16.2 ± 0.3 <sup>a</sup>	16.4 ± 0.4 <sup>a</sup>
18:0	3.0 ± 0.0 <sup>c</sup>	3.6 ± 0.1 <sup>b</sup>	3.9 ± 0.1 <sup>a</sup>
Total saturated <sup>1</sup>	18.4 ± 0.8 <sup>b</sup>	22.6 ± 0.3 <sup>a</sup>	22.7 ± 0.4 <sup>a</sup>
16:1n-9	0.0 ± 0.0 <sup>c</sup>	1.5 ± 0.1 <sup>a</sup>	1.3 ± 0.0 <sup>b</sup>
16:1n-7	3.6 ± 0.0 <sup>a</sup>	2.1 ± 0.0 <sup>b</sup>	2.1 ± 0.1 <sup>b</sup>
18:1n-9	16.1 ± 0.4 <sup>c</sup>	21.8 ± 1.0 <sup>b</sup>	23.6 ± 0.4 <sup>a</sup>
18:1n-7	2.1 ± 0.1 <sup>a</sup>	1.4 ± 0.3 <sup>b</sup>	1.6 ± 0.1 <sup>b</sup>
20:1 <sup>2</sup>	8.7 ± 0.3 <sup>a</sup>	2.2 ± 0.1 <sup>c</sup>	2.9 ± 0.0 <sup>b</sup>
22:1 <sup>3</sup>	9.7 ± 0.5 <sup>a</sup>	1.0 ± 0.0 <sup>b</sup>	1.2 ± 0.0 <sup>b</sup>
24:1n-9	2.0 ± 0.9	1.2 ± 0.1	0.9 ± 0.1
Total monoenes	42.3 ± 0.4 <sup>a</sup>	31.2 ± 1.0 <sup>c</sup>	33.6 ± 0.7 <sup>b</sup>
18:2n-6	5.3 ± 0.1 <sup>a</sup>	2.2 ± 0.1 <sup>b</sup>	2.2 ± 0.1 <sup>b</sup>
20:4n-6	0.7 ± 0.0 <sup>b</sup>	1.5 ± 0.1 <sup>a</sup>	1.7 ± 0.1 <sup>a</sup>
Total n-6PUFA <sup>4</sup>	7.0 ± 0.2 <sup>a</sup>	4.4 ± 0.2 <sup>b</sup>	4.9 ± 0.2 <sup>b</sup>
18:3n-3	1.2 ± 0.0 <sup>a</sup>	0.5 ± 0.0 <sup>b</sup>	0.6 ± 0.0 <sup>b</sup>
20:4n-3	1.5 ± 0.1 <sup>a</sup>	1.0 ± 0.0 <sup>b</sup>	1.0 ± 0.1 <sup>b</sup>
20:5n-3	5.6 ± 0.3 <sup>ab</sup>	5.5 ± 0.1 <sup>b</sup>	6.3 ± 0.4 <sup>a</sup>
22:5n-3	1.3 ± 1.5 <sup>b</sup>	3.5 ± 0.2 <sup>a</sup>	3.0 ± 0.1 <sup>ab</sup>
22:6n-3	14.4 ± 0.7 <sup>c</sup>	25.7 ± 1.2 <sup>a</sup>	21.6 ± 0.5 <sup>b</sup>
Total n-3PUFA <sup>5</sup>	25.9 ± 2.7 <sup>b</sup>	36.8 ± 1.3 <sup>a</sup>	32.8 ± 0.8 <sup>a</sup>
Total PUFA	32.9 ± 2.9 <sup>b</sup>	41.1 ± 1.4 <sup>a</sup>	37.8 ± 0.9 <sup>a</sup>

Samples were obtained from Machrihanish in 2009. Results are means ± SD (n = 3). Values within a row with different superscript letters are significantly different (P < 0.05). <sup>1</sup>, Totals include 15:0, 20:0 and 22:0 present in some samples at up to 0.7 %; <sup>2</sup>, Predominantly n-9 isomer; <sup>3</sup>, Predominantly n-11 isomer; <sup>4</sup>, Totals include 18:3n-6, 20:2n-6, 20:3n-6, 22:4n-6 and 22:5n-6 present in some samples at up to 0.3 %; <sup>5</sup>, Totals include 18:4n-3 and 20:3n-3 present in some samples at up to 0.3 %; PUFA, polyunsaturated fatty acids.

Table 9. Fatty acid compositions (percentage of total fatty acids) of phospholipids and triacylglycerols of *L. salmonis* and their egg strings along with the muscle of the Atlantic salmon from which they were collected

	Phospholipids			Triacylglycerols		
	Salmon muscle	Female lice	Egg strings	Salmon muscle	Female lice	Egg strings
14:0	1.3 ± 0.1	2.9 ± 0.2	4.6 ± 0.0	4.4 ± 0.0	1.5 ± 0.0	1.7 ± 0.0
16:0	19.6 ± 0.1	18.5 ± 0.5	18.2 ± 0.4	11.6 ± 0.0	17.0 ± 1.1	18.5 ± 0.3
18:0	5.9 ± 0.1	3.2 ± 0.1	2.8 ± 0.1	2.8 ± 0.0	4.3 ± 0.2	4.6 ± 0.1
Total saturated <sup>1</sup>	27.3 ± 0.1	25.3 ± 0.6	26.2 ± 0.5	19.4 ± 0.1	23.3 ± 1.3	25.3 ± 0.2
16:1n-9	0.0 ± 0.0	1.7 ± 0.1	1.7 ± 0.0	0.0 ± 0.0	1.5 ± 0.0	1.3 ± 0.0
16:1n-7	1.2 ± 0.0	1.9 ± 0.0	1.5 ± 0.1	4.5 ± 0.0	2.4 ± 0.0	2.3 ± 0.1
18:1n-9	5.7 ± 0.0	16.1 ± 0.7	16.6 ± 1.4	19.9 ± 0.0	28.3 ± 0.9	27.7 ± 0.0
18:1n-7	1.7 ± 0.0	1.3 ± 0.4	0.9 ± 0.0	2.3 ± 0.1	1.9 ± 0.2	1.8 ± 0.2
20:1 <sup>2</sup>	1.4 ± 0.0	1.2 ± 0.0	1.3 ± 0.1	10.9 ± 0.0	3.2 ± 0.0	3.6 ± 0.0
22:1 <sup>3</sup>	0.6 ± 0.0	0.6 ± 0.0	0.9 ± 0.9	12.3 ± 0.2	1.4 ± 0.0	1.4 ± 0.0
24:1n-9	1.1 ± 0.0	0.9 ± 0.1	2.4 ± 2.1	1.0 ± 0.0	0.9 ± 0.1	0.7 ± 0.0
Total monoenes	11.7 ± 0.2	23.7 ± 0.5	25.5 ± 1.8	51.0 ± 0.1	39.6 ± 0.6	39.0 ± 0.2
18:2n-6	1.6 ± 0.0	1.6 ± 0.0	1.1 ± 0.0	6.1 ± 0.0	2.8 ± 0.0	2.4 ± 0.1
20:3n-6	0.9 ± 0.3	0.2 ± 0.0	0.3 ± 0.1	0.2 ± 0.0	0.3 ± 0.0	0.3 ± 0.0
20:4n-6	1.7 ± 0.0	1.0 ± 0.1	0.6 ± 0.0	0.5 ± 0.0	1.8 ± 0.2	2.0 ± 0.0
Total n-6 PUFA	5.1 ± 0.4	3.4 ± 0.2	2.8 ± 0.6	7.9 ± 0.0	5.6 ± 0.3	5.4 ± 0.1
18:3n-3	0.6 ± 0.0	0.5 ± 0.0	0.5 ± 0.0	1.3 ± 0.0	0.7 ± 0.0	0.6 ± 0.0
18:4n-3	0.3 ± 0.0	0.2 ± 0.0	0.1 ± 0.0	1.9 ± 0.0	0.4 ± 0.0	0.3 ± 0.0
20:4n-3	1.0 ± 0.0	0.7 ± 0.0	0.6 ± 0.0	1.5 ± 0.0	1.2 ± 0.0	1.1 ± 0.0
20:5n-3	11.1 ± 0.0	4.8 ± 0.0	4.1 ± 0.1	4.4 ± 0.0	6.1 ± 0.3	6.5 ± 0.2
22:5n-3	2.0 ± 0.0	4.1 ± 0.0	3.6 ± 0.2	2.3 ± 0.0	3.3 ± 0.3	2.8 ± 0.0
22:6n-3	40.1 ± 0.3	36.8 ± 0.9	36.1 ± 2.0	9.6 ± 0.0	18.7 ± 1.1	18.2 ± 0.5
Total n-3 PUFA	55.3 ± 0.3	47.1 ± 0.9	45.0 ± 1.9	21.2 ± 0.0	30.6 ± 1.7	29.7 ± 0.2
Total PUFA <sup>6</sup>	61.0 ± 0.1	51.0 ± 1.1	48.3 ± 1.2	29.7 ± 0.0	37.1 ± 1.9	35.7 ± 0.1

Samples were obtained from Machrihanish in 2009. Results are means ± SD (n = 3). Values within a row with different superscript letters are significantly different (P < 0.05). <sup>1</sup>, Totals include 15:0, 20:0 and 22:0 present in some samples at up to 0.4 %; <sup>2</sup>, Predominantly n-9 isomer; <sup>3</sup>, Predominantly n-11 isomer; <sup>4</sup>, Totals include 18:3n-6, 20:2n-6, 22:4n-6 and 22:5n-6 present in some samples at up to 0.5 %; <sup>5</sup>, Totals include 20:3n-3 present in some samples at up to 0.2 %; <sup>6</sup>, Includes C16 PUFA; PUFA, polyunsaturated fatty acids.

Table 10. Lipid class composition (percentage of total lipid) of *Lepeophtheirus spp.* from Atlantic salmon, sea trout and halibut

Lipid class	Atlantic salmon	sea trout	halibut
PC	13.9 ± 1.5 <sup>b</sup>	14.2 ± 1.7 <sup>b</sup>	18.6 ± 0.7 <sup>a</sup>
PE	14.1 ± 2.6 <sup>b</sup>	13.7 ± 1.8 <sup>b</sup>	20.0 ± 1.4 <sup>a</sup>
PS/PI/PA/CL	5.2 ± 3.0	7.6 ± 3.7	12.4 ± 5.1
Sphingomyelin	3.7 ± 0.8 <sup>b</sup>	5.3 ± 1.6 <sup>ab</sup>	8.1 ± 1.3 <sup>a</sup>
LPC	0.3 ± 0.2 <sup>b</sup>	1.3 ± 0.2 <sup>a</sup>	0.7 ± 0.2 <sup>b</sup>
Total polar	37.2 ± 5.3 <sup>b</sup>	42.0 ± 7.4 <sup>b</sup>	59.7 ± 5.7 <sup>a</sup>
Total neutral	62.8 ± 5.3 <sup>a</sup>	58.0 ± 7.4 <sup>a</sup>	40.3 ± 5.7 <sup>b</sup>
Cholesterol	10.5 ± 0.7	12.1 ± 3.1	11.6 ± 1.4
Triacylglycerol	42.2 ± 7.5 <sup>a</sup>	39.9 ± 12.1 <sup>a</sup>	18.2 ± 3.1 <sup>b</sup>
Free fatty acid	10.1 ± 1.5	6.1 ± 1.6	10.5 ± 1.4
Steryl/wax ester	Trace	Trace	Trace

Results are means ± SD (n = 3). Values within a row with different superscript letter are significantly different (P < 0.05). CL, cardiolipin; LPC, lyso-PC; PA, phosphatidic acid; PC, phosphatidylcholine; PE, phosphatidylethanolamine; PI, phosphatidylinositol; PS, phosphatidylserine.

Table 11. Fatty acid composition (percentage total fatty acids) of *Lepeophtheirus spp.* from Atlantic salmon, sea trout and halibut.

Fatty acid	Atlantic salmon	sea trout	halibut
14:0	2.0 ± 0.1 <sup>a</sup>	1.0 ± 0.3 <sup>b</sup>	0.5 ± 0.2 <sup>b</sup>
16:0	18.7 ± 0.7 <sup>a</sup>	17.4 ± 1.0 <sup>ab</sup>	15.0 ± 2.0 <sup>b</sup>
18:0	3.8 ± 0.1 <sup>b</sup>	4.9 ± 0.7 <sup>ab</sup>	5.1 ± 0.4 <sup>a</sup>
Total saturated <sup>1</sup>	25.1 ± 1.1	24.9 ± 1.0	21.6 ± 3.2
16:1n-9	1.7 ± 0.3	1.6 ± 0.1	1.4 ± 0.1
16:1n-7	2.7 ± 0.1 <sup>a</sup>	2.4 ± 0.2 <sup>a</sup>	0.9 ± 0.2 <sup>b</sup>
18:1n-9	25.0 ± 1.2 <sup>a</sup>	18.8 ± 4.3 <sup>b</sup>	8.8 ± 0.4 <sup>c</sup>
18:1n-7	1.5 ± 0.1 <sup>b</sup>	1.8 ± 0.1 <sup>a</sup>	0.8 ± 0.1 <sup>c</sup>
20:1 <sup>2</sup>	2.2 ± 0.2 <sup>a</sup>	1.9 ± 0.1 <sup>a</sup>	1.4 ± 0.2 <sup>b</sup>
22:1 <sup>3</sup>	1.0 ± 0.0	0.8 ± 0.3	0.5 ± 0.2
24:1n-9	1.1 ± 0.1	2.3 ± 1.1	3.6 ± 1.7
Total monoenes	35.3 ± 1.2 <sup>a</sup>	29.7 ± 2.8 <sup>b</sup>	17.6 ± 2.1 <sup>c</sup>
18:2n-6	0.4 ± 0.1	0.8 ± 0.4	0.9 ± 0.2
20:4n-6	1.2 ± 0.3	1.1 ± 0.1	0.9 ± 0.1
Total n-6PUFA <sup>4</sup>	2.3 ± 0.1 <sup>b</sup>	3.3 ± 0.4 <sup>a</sup>	3.8 ± 0.4 <sup>a</sup>
18:3n-3	0.2 ± 0.0	0.3 ± 0.0	0.2 ± 0.1
20:4n-3	0.4 ± 0.1 <sup>a</sup>	0.5 ± 0.0 <sup>a</sup>	0.3 ± 0.0 <sup>b</sup>
20:5n-3	5.1 ± 0.2 <sup>a</sup>	5.7 ± 0.1 <sup>a</sup>	4.2 ± 0.4 <sup>b</sup>
22:5n-3	4.3 ± 0.3 <sup>a</sup>	2.5 ± 0.2 <sup>b</sup>	1.4 ± 0.4 <sup>c</sup>
22:6n-3	22.1 ± 1.2 <sup>b</sup>	22.2 ± 2.1 <sup>b</sup>	35.8 ± 4.9 <sup>a</sup>
Total n-3PUFA <sup>5</sup>	32.2 ± 1.6 <sup>b</sup>	31.5 ± 2.1 <sup>b</sup>	42.1 ± 5.6 <sup>a</sup>
Total PUFA	34.5 ± 1.6 <sup>b</sup>	34.8 ± 2.4 <sup>b</sup>	45.9 ± 5.5 <sup>a</sup>

Results are means ± SD (n = 3). Values within a row with a different superscript letter are significantly different (P < 0.05). <sup>1</sup>, Totals include 15:0, 20:0 and 22:0 present in some samples at up to 0.7 %; <sup>2</sup>, Predominantly n-9 isomer; <sup>3</sup>, Predominantly n-11 isomer; <sup>4</sup>, Totals include 18:3n-6, 20:2n-6, 20:3n-6, 22:4n-6 and 22:5n-6 present in some samples at up to 0.3 %; <sup>5</sup>, Totals include 18:3n-3 and 20:3n-3 present in some samples at up to 0.2 %; PUFA, polyunsaturated fatty acids.