

THE UNIVERSITY OF WARWICK

Original citation:

Ferraz-Leite, S., Ortner, Christoph and Praetorius, D.. (2010) Convergence of simple adaptive Galerkin schemes based on $h - h/2$ error estimators. Numerische Mathematik, Vol.116 (No.2). pp. 291-316.

Permanent WRAP url:

<http://wrap.warwick.ac.uk/48591>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes this work of researchers of the University of Warwick available open access under the following conditions.

This article is made available under the Creative Commons Attribution- 3.0 Unported (CC BY 3.0) license and may be reused according to the conditions of the license. For more details see <http://creativecommons.org/licenses/by/3.0/>

A note on versions:

The version presented in WRAP is the published version, or, version of record, and may be cited as it appears here.

For more information, please contact the WRAP Team at: publications@warwick.ac.uk

warwick**publications**wrap

highlight your research

<http://wrap.warwick.ac.uk/>

Convergence of simple adaptive Galerkin schemes based on $h - h/2$ error estimators

S. Ferraz-Leite · C. Ortner · D. Praetorius

Received: 9 April 2009 / Revised: 26 December 2009 / Published online: 30 March 2010
© The Author(s) 2010. This article is published with open access at Springerlink.com

Abstract We discuss several adaptive mesh-refinement strategies based on $(h - h/2)$ -error estimation. This class of adaptive methods is particularly popular in practise since it is problem independent and requires virtually no implementational overhead. We prove that, under the saturation assumption, these adaptive algorithms are convergent. Our framework applies not only to finite element methods, but also yields a first convergence proof for adaptive boundary element schemes. For a finite element model problem, we extend the proposed adaptive scheme and prove convergence even if the saturation assumption fails to hold in general.

Mathematics Subject Classification (2000) 65N30 · 65N38 · 65N50 · 65N12

1 Introduction

Introductory remarks

Let \mathcal{H} be a real Hilbert space with scalar product $\langle\langle \cdot, \cdot \rangle\rangle$ and associated energy norm $\|\| \cdot \|\|$. Given a right-hand side $F \in \mathcal{H}^*$, we aim at a numerical approximation of the unique solution $u \in \mathcal{H}$ of

$$\langle\langle u, v \rangle\rangle = F(v) \quad \text{for all } v \in \mathcal{H}. \quad (1)$$

S. Ferraz-Leite · D. Praetorius (✉)
Institute for Analysis and Scientific Computing, Vienna University of Technology,
Wiedner Hauptstrasse 8-10, 1040 Vienna, Austria
e-mail: Dirk.Praetorius@tuwien.ac.at

S. Ferraz-Leite
e-mail: Samuel.Ferraz-Leite@tuwien.ac.at

C. Ortner
Mathematical Institute, University of Oxford, 24-29 St Giles', Oxford OX1 3LB, UK
e-mail: ortner@maths.ox.ac.uk

To that end, let X_ℓ be a finite-dimensional subspace of \mathcal{H} , which in many applications is based on a triangulation \mathcal{T}_ℓ of some physical domain. Let $u_\ell \in X_\ell$ be the corresponding Galerkin solution, obtained by solving the finite-dimensional linear system

$$\langle\langle u_\ell, v_\ell \rangle\rangle = F(v_\ell) \quad \text{for all } v_\ell \in X_\ell. \quad (2)$$

Throughout, the index $\ell \in \mathbb{N}_0 := \{0, 1, 2, \dots\}$ denotes the step of an adaptive algorithm, and associated quantities are discrete and numerically computable.

Whereas uniform refinement of the meshes \mathcal{T}_ℓ usually leads to convergence $u_\ell \rightarrow u$ as $\ell \rightarrow \infty$, this cannot be expected, in general, if the meshes are only refined locally by means of an adaptive algorithm. We say that an adaptive mesh-refining algorithm is *convergent* if, and only if, $u_\ell \rightarrow u$ as $\ell \rightarrow \infty$. In this sense, convergence of an adaptive algorithm has first been proven by Dörfler [12], where also the marking criterion (7) used in Algorithm 1 below is introduced. The latter work considered the residual error estimator for a P1-finite element discretization of the Poisson problem, and it is assumed that data oscillations on the initial mesh are sufficiently small. In [19], the resolution of the data oscillations is included into the adaptive algorithm. The convergence analysis is based on reliability and the so-called *discrete local efficiency* of the residual error estimator, which relies on an *interior node property* for the local mesh-refinement. The main idea of the convergence proof then is to show that the error is contractive up to the data oscillations. Cascon et al. [10] recently presented a new convergence proof under weaker conditions. They showed that a weighted sum of error and error estimator satisfies a contraction property *without* requiring (discrete local) efficiency of the estimator. In all of the cited works [10, 12, 19], however, the focus was only on residual-based error estimators for conforming finite element schemes.

$(h - h/2)$ -Error estimation and adaptive algorithm

In contrast to the preceding works, we study the $(h - h/2)$ -error estimation strategy which is a well-known technique for the a posteriori estimation of the error in the energy norm $\|u - u_\ell\|$; see [16] in the context of ordinary differential equations, and the works of Bank [3–5] or the monograph [1, Chapter 5] in the context of the finite element method. Let $\widehat{\mathcal{T}}_\ell$ be a uniform refinement of \mathcal{T}_ℓ and let \widehat{X}_ℓ be the associated finite-dimensional subspace of \mathcal{H} with corresponding Galerkin solution $\widehat{u}_\ell \in \widehat{X}_\ell$. Then, the $(h - h/2)$ -error estimator

$$\eta_\ell := \|\widehat{u}_\ell - u_\ell\| \quad (3)$$

is a computable quantity [11] which can be used to estimate the error $\|u - u_\ell\|$. By use of the Galerkin orthogonality property it is easy to see that

$$\eta_\ell \leq \|u - u_\ell\| \leq C_{\text{rel}} \eta_\ell, \quad (4)$$

where the upper bound is equivalent to the so-called *saturation assumption*

$$\|u - \widehat{u}_\ell\| \leq C_{\text{sat}} \|u - u_\ell\| \quad \text{with some uniform constant } C_{\text{sat}} \in (0, 1), \quad (5)$$

see e.g. [1, Theorem 5.1]. We remark that the saturation assumption dates back to the early work [5], but may fail to hold in general [6, 13].

To steer an adaptive mesh-refinement, we additionally assume that $\varrho_\ell(T)$ is a computable quantity which (at least heuristically) measures the local error between the exact solution u and a discrete solution $u_\ell \in X_\ell$ on an element $T \in \mathcal{T}_\ell$. To link these so-called *refinement indicators* with η_ℓ , we assume that

$$C_1^{-1} \eta_\ell \leq \varrho_\ell := \left(\sum_{T \in \mathcal{T}_\ell} \varrho_\ell(T)^2 \right)^{1/2} \leq C_2 \eta_\ell \quad \text{for all } \ell \in \mathbb{N}, \quad (6)$$

with ℓ -independent constants $C_1, C_2 > 0$. The adaptive mesh-refining algorithm then reads as follows:

Algorithm 1 Let an initial triangulation \mathcal{T}_0 with associated discrete space X_0 and an adaptivity parameter $\theta \in (0, 1)$ be given. For $\ell = 0, 1, 2, \dots$ do the following:

- (a) Compute the discrete solution $u_\ell \in X_\ell$.
- (b) Compute the refinement indicators $\varrho_\ell(T)$ for all $T \in \mathcal{T}_\ell$.
- (c) Stop if u_ℓ is sufficiently accurate (e.g., ϱ_ℓ resp. η_ℓ is sufficiently small).
- (d) Otherwise, choose a set $\mathcal{M}_\ell \subseteq \mathcal{T}_\ell$ of marked elements such that

$$\theta \sum_{T \in \mathcal{T}_\ell} \varrho_\ell(T)^2 \leq \sum_{T \in \mathcal{M}_\ell} \varrho_\ell(T)^2. \quad (7)$$

- (e) Refine at least the marked elements $T \in \mathcal{M}_\ell$, generate a new mesh $\mathcal{T}_{\ell+1}$ and an associated discrete space $X_{\ell+1}$, increase the counter $\ell \mapsto \ell + 1$, and go to (a).

Outline

In the present paper, we adapt the arguments from [10] to $(h - h/2)$ -based error estimation techniques and investigate the convergence of Algorithm 1 steered by $(h - h/2)$ -based refinement indicators. In Sect. 2, linear finite elements for the numerical solution of the 2D Poisson problem are studied. We consider the indicators

$$\begin{aligned} \tilde{\mu}_\ell(T) &:= \|\nabla \widehat{u}_\ell - \Pi_\ell(\nabla \widehat{u}_\ell)\|_{L^2(T)} \\ \text{and } \mu_\ell(T) &:= \|\nabla(\widehat{u}_\ell - I_\ell \widehat{u}_\ell)\|_{L^2(T)} \quad \text{for } T \in \mathcal{T}_\ell, \end{aligned} \quad (8)$$

where $\Pi_\ell : L^2(\Omega) \rightarrow \mathcal{P}^0(\mathcal{T}_\ell)$ is the L^2 -orthogonal projection operator onto piecewise constants and $I_\ell : \mathcal{C}(\overline{\Omega}) \rightarrow \mathcal{S}^1(\mathcal{T}_\ell)$ the nodal interpolation operator onto piecewise linear and globally continuous functions. Under the saturation assumption (5), we prove

that Algorithm 1, steered by $\varrho_\ell = \tilde{\mu}_\ell$, and for any choice of $\theta \in (0, 1)$, guarantees the contraction property

$$\Delta_{\ell+1} \leq \kappa \Delta_\ell \quad \text{with} \quad \Delta_\ell := \| \|u - u_\ell\|^2 + \| \|u - \widehat{u}_\ell\|^2 + \gamma \varrho_\ell^2 \tag{9}$$

and with appropriate constants $\kappa, \gamma \in (0, 1)$. In particular, the adaptive algorithm is convergent (Theorem 4). We also consider Algorithm 1 steered by $\varrho_\ell = \mu_\ell$, where convergence $\lim_\ell u_\ell = u$ follows for arbitrary $\theta \in (0, 1)$, but a contraction property (9) is only proven for θ sufficiently large (Theorem 5).

In [13], it has been observed that small data oscillations imply the saturation assumption. In Sect. 2.6, we incorporate this observation in the convergence analysis in the sense that we enhance the refinement indicators $\mu_\ell(T)$ and $\tilde{\mu}_\ell(T)$ by the local data oscillations $\text{osc}_\ell(T)$ defined in (40) below. Using these combined refinement indicators, e.g. $\varrho_\ell^2 = \tilde{\mu}_\ell^2 + \text{osc}_\ell^2$, the data is adaptively resolved by Algorithm 1, and we obtain convergence $u_\ell \rightarrow u$ as $\ell \rightarrow \infty$, independently of the saturation assumption (Theorem 6).

In addition to the adaptive finite element method, our arguments also apply to boundary integral formulations discussed in Sect. 3. As model example serves the elliptic integral formulation of the Laplace problem with Dirichlet boundary conditions in 2D and 3D. We use the $(h - h/2)$ -based error indicators

$$\mu_\ell(T) := \left\| h^{1/2}(\widehat{u}_\ell - u_\ell) \right\|_{L^2(T)} \quad \text{and} \quad \tilde{\mu}_\ell(T) := \left\| h^{1/2}(\widehat{u}_\ell - \Pi_\ell \widehat{u}_\ell) \right\|_{L^2(T)} \tag{10}$$

from [14] to steer the adaptive mesh-refinement. Under the saturation assumption (5), we prove that the associated error quantity Δ_ℓ satisfies the contraction property (9) for both, the $\tilde{\mu}_\ell$ -steered and the μ_ℓ -steered adaptive algorithm (Theorem 7–8). This yields a first convergence proof for the adaptive boundary element method.

Finally, in Sect. 4, numerical experiments for both, finite element and boundary element methods, are performed. The results verify our analysis empirically and suggest the validity of the saturation assumption even in the context of boundary element methods.

2 Finite element method

2.1 Model problem

We consider the elliptic model problem

$$\begin{aligned} -\Delta u &= f && \text{in } \Omega, \\ u &= 0 && \text{on } \Gamma_D, \\ \partial_n u &= g && \text{on } \Gamma_N. \end{aligned} \tag{11}$$

Here, Ω is a bounded Lipschitz domain in \mathbb{R}^2 with polygonal boundary. The boundary Γ is split into a Dirichlet boundary Γ_D and a Neumann boundary Γ_N which satisfy

$\Gamma = \bar{\Gamma}_D \cup \bar{\Gamma}_N$ as well as $\Gamma_D \cap \Gamma_N = \emptyset$. Moreover, we assume that Γ_D has positive surface measure $|\Gamma_D| > 0$ so that (11) admits a unique weak solution. The energy scalar product of the weak formulation of (11) reads

$$\langle\langle u, v \rangle\rangle := \int_{\Omega} \nabla u \cdot \nabla v \, dx \quad \text{for all } u, v \in \mathcal{H} = \{u \in H^1(\Omega) : u|_{\Gamma_D} = 0\}, \quad (12)$$

and the right-hand side is given by

$$F(v) = \int_{\Omega} f v \, dx + \int_{\Gamma_N} g v \, ds, \quad (13)$$

where ds denotes integration along the boundary. In particular, the energy norm reads $\|v\| = \|\nabla v\|_{L^2(\Omega)}$. We consider the lowest-order Galerkin scheme, where \mathcal{T}_ℓ is a regular triangulation of Ω in the sense of Ciarlet and where $X_\ell = \{v_\ell \in \mathcal{S}^1(\mathcal{T}_\ell) : v_\ell|_{\Gamma_D} = 0\}$ with $\mathcal{S}^1(\mathcal{T}_\ell) = \{v_\ell \in C(\Omega) : \forall T \in \mathcal{T}_\ell \, v_\ell|_T \text{ is affine}\}$.

To fix notation, \mathcal{N}_ℓ denotes the set of nodes of \mathcal{T}_ℓ , and \mathcal{E}_ℓ denotes the set of edges of \mathcal{T}_ℓ . Moreover, $\mathcal{E}_{N,\ell}$ denotes the set of all edges, which belong to the Neumann boundary Γ_N . The local mesh-width $h_\ell \in L^\infty(\Omega)$ is defined by $h_\ell(x) = \text{diam}(T)$ for $T \in \mathcal{T}_\ell$ and $x \in \text{interior}(T)$. Moreover, $h_\ell(x) = \text{diam}(E)$ if x belongs to the relative interior of an edge $E \in \mathcal{E}_\ell$. Finally, $\mathcal{P}^p(\mathcal{T}_\ell)$ denotes the space of all \mathcal{T}_ℓ -piecewise polynomials of total degree $\leq p$, and $\mathcal{S}^1(\mathcal{T}_\ell) = \mathcal{P}^1(\mathcal{T}_\ell) \cap C(\Omega)$.

2.2 Mesh-refinement

The essential point in the analysis below is that all sons $T' \in \mathcal{T}_{\ell+1}$ of a marked element $T \in \mathcal{M}_\ell$ also belong to the uniform refinement $\widehat{\mathcal{T}}_\ell$ of \mathcal{T}_ℓ . We therefore generate both meshes $\mathcal{T}_{\ell+1}$ and $\widehat{\mathcal{T}}_\ell$ by the same refinement strategy with respect to the marked elements \mathcal{M}_ℓ and $\widehat{\mathcal{M}}_\ell := \mathcal{T}_\ell$, respectively.

Our mesh-refinement is a variant of the newest vertex bisection algorithm (cf. Fig. 1), using an edge-based formulation: For a marked element T , we mark all of its edges for refinement. To avoid hanging nodes and to ensure regularity of the refined triangulation, we then proceed recursively: If any edge of an element T' is marked for refinement, we also mark its reference edge. This recursion necessarily terminates after finitely many steps with the result that, for each element in which at least one edge is marked for refinement, the reference edge is marked as well. We then proceed with the refinement rules for newest vertex bisection shown in Fig. 2.

For further details on newest vertex bisection, the reader is referred to [23, Chapter 4] or the original works [2, 17, 21]. We stress that all generated meshes are uniformly shape regular, see Fig. 3.

However, to guarantee the saturation assumption, we use $\text{bise}_5(T)$ instead of $\text{bise}_3(T)$ if an element is marked for refinement, cf. Fig. 1. We refer to [13] and the following section for the fact that uniform bise_5 -refinement asymptotically yields the saturation assumption, whereas bise_3 does not.




Fig. 1 For each triangle $T \in \mathcal{T}$, there is one *reference edge*, indicated by the *doubleline* (*left*). Bisection of T is achieved by halving the reference edge, where its midpoint becomes a new node. The reference edges of the son triangles are opposite to this newest vertex (*left-middle*). Further steps of newest vertex bisection lead to steps, which are denoted by $\text{bisec}_3(T)$ and $\text{bisec}_5(T)$, respectively (*right-middle*, *right*)




Fig. 2 Refinement by newest vertex bisection: We consider one triangle T and assume that certain edges, but at least the reference edge, are marked for refinement (*top*). After refinement, the element is split into 2, 3, or 4 son triangles, respectively (*bottom*). Throughout, the reference edges are indicated by a *doubleline*



Fig. 3 Refinement by newest vertex bisection only leads to finitely many interior angles for the family of all possible triangulations obtained by arbitrary newest vertex bisections. To see this, we start from a macro element (*left*), where the bottom edge is the reference edge. Using iterated newest vertex bisection, one observes that only four similarity classes of triangles occur, which are indicated by the *colouring*. After three steps of bisections (*right*), no additional similarity class appears (color figure online)

We stress that refinement rules based on newest vertex bisection ensure the inclusions $X_\ell \subset X_{\ell+1} \subseteq \widehat{X}_\ell \subset \widehat{X}_{\ell+1}$, which are crucial in our analysis. We remark that almost all mesh-refinement strategies will yield $X_\ell \subset X_{\ell+1}$ and $X_\ell \subset \widehat{X}_\ell$. However, the inclusions $X_{\ell+1} \subset \widehat{X}_\ell \subset \widehat{X}_{\ell+1}$ may be violated. For instance, they fail in general for the popular red-green-blue refinement strategies [23, Chapter 04].

2.3 Saturation assumption

In [13], it is stated (without a proof) that the saturation assumption (5) holds for the homogeneous Dirichlet problem (11) with $\Gamma_D = \partial\Omega$, if $\widehat{\mathcal{T}}_\ell$ is obtained by uniform red-refinement of \mathcal{T}_ℓ and if data oscillations are sufficiently small. Here, red-refinement means that each triangle $T \in \mathcal{T}_\ell$ is divided into four similar son triangles $T_1, \dots, T_4 \in \widehat{\mathcal{T}}_\ell$. The notion *data oscillations* is understood as

$$\widetilde{\text{osc}}_\ell^2 = \sum_{z \in \mathcal{N}_\ell} \|h_\ell(f - f_{\ell,z})\|_{L^2(\omega_{\ell,z})}^2, \tag{14}$$

where $\omega_{\ell,z} := \bigcup \{T \in \mathcal{T}_\ell : z \in T\}$ is the patch of a node z with respect to \mathcal{T}_ℓ and where $f_{\ell,z} := |\omega_{\ell,z}|^{-1} \int_{\omega_{\ell,z}} f \, dx$ denotes the patch-wise integral mean of f .

For $\Gamma_N = \emptyset$ and with bise_5 replacing red-refinement, an analogous result is already contained in [19], although not stated explicitly; compare Eq. (15) below and [19, Lemma 4.2]. For the mixed boundary value problem (11), we sketch the proof of the saturation assumption for the convenience of the reader.

Proposition 2 *Let $\widehat{\mathcal{T}}_\ell$ be obtained by uniform bise_5 -refinement of \mathcal{T}_ℓ , and let $u_\ell \in X_\ell$ and $\widehat{u}_\ell \in \widehat{X}_\ell$ be the corresponding Galerkin solutions to (11). Then, there exists a constant $\widetilde{C}_{\text{sat}} \in (0, 1)$ such that*

$$\|u - \widehat{u}_\ell\|^2 \leq \widetilde{C}_{\text{sat}}^2 \|u - u_\ell\|^2 + \text{osc}_\ell^2, \tag{15}$$

where the data oscillations are defined by

$$\text{osc}_\ell^2 = \|h_\ell(f - \Pi_\ell f)\|_{L^2(\Omega)}^2 + \|h_\ell^{1/2}(g - \Pi_\ell g)\|_{L^2(\Gamma_N)}^2. \tag{16}$$

Here, Π_ℓ denotes the L^2 -orthogonal projection onto $\mathcal{P}^0(\mathcal{T}_\ell)$ and $\mathcal{P}^0(\mathcal{E}_{N,\ell})$, respectively. Consequently, if

$$\text{osc}_\ell^2 \leq \lambda \|u - u_\ell\|^2, \tag{17}$$

for some $\lambda > 0$ with $C_{\text{sat}}^2 = \widetilde{C}_{\text{sat}}^2 + \lambda < 1$, then the saturation assumption (5) is satisfied.

Sketch of Proof To prove (15), we consider the residual-based error estimator [23]

$$\varrho_\ell^2 = \sum_{T \in \mathcal{T}_\ell} \varrho_\ell(T)^2$$

with local contributions

$$\varrho_\ell(T)^2 = \|h_\ell f\|_{L^2(T)}^2 + \|h_\ell^{1/2}[\partial_n u_\ell]\|_{L^2(\partial T \cap \Omega)}^2 + \|h_\ell^{1/2}(g - \partial_n u_\ell)\|_{L^2(\partial T \cap \Gamma_N)}^2,$$

where $[\cdot]$ denotes the jump over an interior edge $E \in \mathcal{E}_\ell$. It is well-known that ϱ_ℓ is reliable,

$$\|u - u_\ell\| \leq \widetilde{C}_{\text{rel}} \varrho_\ell. \tag{18}$$

and locally efficient up to oscillation terms,

$$\begin{aligned} \varrho_\ell(T)^2 \leq \widetilde{C}_{\text{eff}}^2 & \left(\|\nabla(u - u_\ell)\|_{L^2(\omega_T)}^2 + \|h_\ell(f - \Pi_\ell f)\|_{L^2(\omega_T)}^2 \right. \\ & \left. + \|h_\ell^{1/2}(g - \Pi_\ell g)\|_{L^2(\partial T \cap \Gamma_N)}^2 \right), \end{aligned} \tag{19}$$

where $\omega_T = \{T\} \cup \bigcup \{T' \in \mathcal{T}_\ell : T \cap T' \in \mathcal{E}_\ell\}$ denotes the element patch of $T \in \mathcal{T}_\ell$, cf. [23]. The proof of (19) goes back to Verfürth [22] and is obtained by use of appropriate bubble functions. First,

$$\|h_\ell \Pi_\ell f\|_{L^2(T)} \lesssim \|\nabla u - \nabla u_\ell\|_{L^2(T)} + \|h_\ell(f - \Pi_\ell f)\|_{L^2(T)} \tag{20}$$

follows by use of the element bubble $b_T \in \mathcal{P}^3(T) \cap H_0^1(T)$. Second, for an interior edge $E \in \mathcal{E}_\ell$, let $E = T \cap T'$ and denote by $\omega_E = T \cup T'$ the corresponding edge patch. Then, by use of the edge bubble $b_E \in \mathcal{P}^2(\{T, T'\}) \cap H_0^1(\omega_E)$, one can prove that

$$\|h_\ell^{1/2}[\partial_n u_\ell]\|_{L^2(E)} \lesssim \|\nabla u - \nabla u_\ell\|_{L^2(\omega_E)} + \|h_\ell f\|_{L^2(\omega_E)}. \tag{21}$$

Third, for a Neumann edge $E \in \mathcal{E}_{N,\ell}$ with corresponding element $T \in \mathcal{T}_\ell$, using again the edge bubble $b_E \in \mathcal{P}^2(T)$ with $b_E|_{\partial T \setminus E} = 0$, one shows that

$$\begin{aligned} \|h_\ell^{1/2}(\Pi_\ell g - \partial_n u_\ell)\|_{L^2(E)} &\lesssim \|\nabla u - \nabla u_\ell\|_{L^2(T)} + \|h_\ell f\|_{L^2(T)} \\ &\quad + \|h_\ell^{1/2}(g - \Pi_\ell g)\|_{L^2(E)}. \end{aligned} \tag{22}$$

Using the triangle inequality and (20)–(22), one obtains (19).

In all of the three estimates (20)–(22), the term ∇u on the right-hand side stems from the weak form (1) applied with the properly weighted bubble functions $v \in \{b_T, b_E\}$. It has already been observed by Dörfler [12] that one may use the hat function $v = b_E \in \widehat{X}_\ell$ for the midpoint of an edge $E \in \mathcal{E}_\ell$ instead of the edge bubble function. This allows to use the Galerkin formulation (2) for \widehat{X}_ℓ instead of the continuous variational form (1) and leads to $\nabla \widehat{u}_\ell$ instead of ∇u in (21)–(22). Finally, it was noticed in [19] that one may use the hat function $v = b_T \in \widehat{X}_\ell$ which corresponds to the interior node of $\text{bisec}_5(T)$ instead of the element bubble, cf. Fig. 1. This leads to $\nabla \widehat{u}_\ell$ instead of ∇u in (20). With these two observations, one can therefore use Verfürth’s proof to verify the local discrete efficiency

$$\begin{aligned} \varrho_\ell(T)^2 &\leq \widetilde{C}_{\text{eff}}^2 \left(\|\nabla(\widehat{u}_\ell - u_\ell)\|_{L^2(\omega_T)}^2 + \|h_\ell(f - \Pi_\ell f)\|_{L^2(\omega_T)}^2 \right. \\ &\quad \left. + \|h_\ell^{1/2}(g - \Pi_\ell g)\|_{L^2(\partial T \cap \Gamma_N)}^2 \right), \end{aligned} \tag{23}$$

for all $T \in \mathcal{T}_\ell$, even with the same constant as in (19). Summation over all elements $T \in \mathcal{T}_\ell$ results in

$$\varrho_\ell^2 \leq 3\widetilde{C}_{\text{eff}}^2 \left(\|\widehat{u}_\ell - u_\ell\|^2 + \text{osc}_\ell^2 \right) \tag{24}$$

since each element $T \in \mathcal{T}_\ell$ is at most counted three times.

Now, we are in a position to verify (15): The combination of (18) and (24) yields

$$\|u - u_\ell\|^2 \leq \tilde{C}_{\text{rel}}^2 \varrho_\ell^2 \leq 3\tilde{C}_{\text{eff}}^2 \tilde{C}_{\text{rel}}^2 \left(\|\widehat{u}_\ell - u_\ell\|^2 + \text{osc}_\ell^2 \right).$$

With $C := 3\tilde{C}_{\text{eff}}^2 \tilde{C}_{\text{rel}}^2$, the Galerkin orthogonalities for \widehat{X}_ℓ and $X_\ell \subset \widehat{X}_\ell$ prove

$$\|u - \widehat{u}_\ell\|^2 = \|u - u_\ell\|^2 - \|\widehat{u}_\ell - u_\ell\|^2 \leq (C - 1)\|\widehat{u}_\ell - u_\ell\|^2 + C \text{osc}_\ell^2.$$

Thus, a further application of the Galerkin orthogonality yields

$$C \|u - \widehat{u}_\ell\|^2 \leq (C - 1)\|u - u_\ell\|^2 + C \text{osc}_\ell^2$$

and concludes the proof with $\tilde{C}_{\text{sat}}^2 = (C - 1)/C$.

2.4 Refinement indicators

In this section, we discuss the refinement indicators

$$\tilde{\mu}_\ell(T) = \|\nabla \widehat{u}_\ell - \Pi_\ell(\nabla \widehat{u}_\ell)\|_{L^2(T)} \quad \text{and} \quad \mu_\ell(T) = \|\nabla(\widehat{u}_\ell - I_\ell \widehat{u}_\ell)\|_{L^2(T)}, \quad (25)$$

as well as

$$\eta_\ell(T) = \|\nabla \widehat{u}_\ell - \nabla u_\ell\|_{L^2(T)}. \quad (26)$$

Here, $\Pi_\ell : L^2(\Omega)^2 \rightarrow \mathcal{P}^0(\mathcal{T}_\ell)^2$ denotes the L^2 -orthogonal projection onto the \mathcal{T}_ℓ -piecewise constant fields, and $I_\ell : C(\overline{\Omega}) \rightarrow X_\ell$ denotes the nodal interpolation operator. Since Π_ℓ is even the \mathcal{T}_ℓ -piecewise orthogonal projection and $\nabla u_\ell \in \mathcal{P}^0(\mathcal{T}_\ell)^2$, it holds that

$$\tilde{\mu}_\ell(T) \leq \mu_\ell(T) \quad \text{and} \quad \tilde{\mu}_\ell(T) \leq \eta_\ell(T) \quad \text{for all } T \in \mathcal{T}_\ell. \quad (27)$$

Together with the best approximation property of the Galerkin solution $u_\ell \in X_\ell$, this provides the global estimates

$$\tilde{\mu}_\ell = \|\nabla \widehat{u}_\ell - \Pi_\ell(\nabla \widehat{u}_\ell)\|_{L^2(\Omega)} \leq \eta_\ell \leq \mu_\ell = \|\nabla(\widehat{u}_\ell - I_\ell \widehat{u}_\ell)\|_{L^2(\Omega)}. \quad (28)$$

The following proposition states the converse inequalities.

Proposition 3 *There is a constant $C_3 \geq 1$ which only depends on the smallest interior angle in \mathcal{T}_ℓ such that*

$$\mu_\ell(T) \leq C_3 \tilde{\mu}_\ell(T) \quad \text{for all } T \in \mathcal{T}_\ell, \quad (29)$$

so that

$$C_3^{-1} \mu_\ell \leq \tilde{\mu}_\ell \leq \eta_\ell \leq \mu_\ell \quad (30)$$

In particular, the error estimators η_ℓ , $\tilde{\mu}_\ell$, and μ_ℓ are equivalent in the sense of (6).

Proof Let C_∞ denote an upper bound on the norm of the interpolation operator I_ℓ in the $W^{1,\infty}$ -semi-norm, valid in each element $T \in \mathcal{T}_\ell$, more precisely, we require

$$\|\nabla(I_\ell \widehat{v}_\ell)\|_{L^\infty(T)} \leq C_\infty \|\nabla \widehat{v}_\ell\|_{L^\infty(T)} \quad \text{for all } \widehat{v}_\ell \in \widehat{X}_\ell \quad \text{and } T \in \mathcal{T}_\ell.$$

We note that, for rectangular elements, $C_\infty = 1$. It is clear that, in general, the constant is bounded only in terms of the mesh quality.

Furthermore, we note that our refinement strategy yields a uniform constant $\nu > 0$ so that $|T'| \geq \nu|T|$ whenever T' is a son of T . Let v be an affine function defined on T with gradient $\nabla v = \Pi_\ell \widehat{u}_\ell$, then

$$\begin{aligned} \|\nabla(1 - I_\ell)\widehat{u}_\ell\|_{L^2(T)}^2 &= \|(1 - \Pi_\ell)\nabla(1 - I_\ell)\widehat{u}_\ell\|_{L^2(T)}^2 + \|\Pi_\ell \nabla(1 - I_\ell)\widehat{u}_\ell\|_{L^2(T)}^2 \\ &= \tilde{\mu}_\ell(T)^2 + \|\nabla v - \nabla I_\ell \widehat{u}_\ell\|_{L^2(T)}^2 \\ &\leq \tilde{\mu}_\ell(T)^2 + |T|C_\infty^2 \|\nabla v - \nabla \widehat{u}_\ell\|_{L^\infty(T)}^2 \\ &\leq \tilde{\mu}_\ell(T)^2 + \nu^{-1}C_\infty^2 \sum_{T' \subset T} |T'| \|\Pi_\ell - 1\| \|\nabla \widehat{u}_\ell\|_{L^\infty(T')}^2 \\ &= (1 + \nu^{-1}C_\infty^2)\tilde{\mu}_\ell(T)^2. \end{aligned}$$

This establishes (29) with $C_3 = (1 + \nu^{-1}C_\infty^2)^{1/2}$. □

2.5 Convergence of AFEM under the saturation assumption

We first prove convergence of the adaptive finite element scheme, based on the refinement indicators $\tilde{\mu}_\ell(T)$ from (25).

Theorem 4 *Suppose that we use the indicators $\tilde{\mu}_\ell(T)$, defined in (25), in Algorithm 1. Under the saturation assumption (5) and for any choice of $\theta \in (0, 1)$, there are constants $\kappa, \gamma \in (0, 1)$ such that*

$$\tilde{\Delta}_\ell := \| \|u - u_\ell\|^2 + \| \|u - \widehat{u}_\ell\|^2 + \gamma \tilde{\mu}_\ell^2 \tag{31}$$

satisfies the contraction property

$$\tilde{\Delta}_{\ell+1} \leq \kappa \tilde{\Delta}_\ell. \tag{32}$$

In particular, it holds that $\lim_{\ell \rightarrow \infty} \| \|u - \widehat{u}_\ell\| = \lim_{\ell \rightarrow \infty} \| \|u - u_\ell\| = 0 = \lim_{\ell \rightarrow \infty} \tilde{\mu}_\ell$.

Proof Application of the triangle inequality yields

$$\begin{aligned} \tilde{\mu}_{\ell+1} &= \|(1 - \Pi_{\ell+1})(\nabla \widehat{u}_{\ell+1})\|_{L^2(\Omega)} \\ &\leq \|(1 - \Pi_{\ell+1})(\nabla \widehat{u}_\ell)\|_{L^2(\Omega)} + \|(1 - \Pi_{\ell+1})(\nabla \widehat{u}_{\ell+1} - \nabla \widehat{u}_\ell)\|_{L^2(\Omega)} \\ &\leq \|\nabla \widehat{u}_\ell - \Pi_{\ell+1}(\nabla \widehat{u}_\ell)\|_{L^2(\Omega)} + \|\nabla \widehat{u}_{\ell+1} - \nabla \widehat{u}_\ell\|_{L^2(\Omega)}, \end{aligned}$$

where we have used orthogonality of $\Pi_{\ell+1}$ in the final estimate. According to the mesh-refinement rule from Sect. 2.2, and since ∇u_ℓ is constant on the six sons $T' \in \mathcal{T}_{\ell+1} \cap \widehat{\mathcal{T}}_\ell$ of $T \in \mathcal{M}_\ell$, it holds that

$$\|\nabla \widehat{u}_\ell - \Pi_{\ell+1}(\nabla \widehat{u}_\ell)\|_{L^2(T)} = 0 \quad \text{for all } T \in \mathcal{M}_\ell.$$

Moreover, $\mathcal{T}_{\ell+1}$ -piecewise orthogonality of $\Pi_{\ell+1}$ yields

$$\|\nabla \widehat{u}_\ell - \Pi_{\ell+1}(\nabla \widehat{u}_\ell)\|_{L^2(T)} \leq \|\nabla \widehat{u}_\ell - \Pi_\ell(\nabla \widehat{u}_\ell)\|_{L^2(T)} = \widetilde{\mu}_\ell(T) \quad \text{for all } T \in \mathcal{T}_\ell \setminus \mathcal{M}_\ell.$$

Finally, Dörfler’s marking strategy (7) provides

$$\sum_{T \in \mathcal{T}_\ell \setminus \mathcal{M}_\ell} \widetilde{\mu}_\ell(T)^2 = \widetilde{\mu}_\ell^2 - \sum_{T \in \mathcal{M}_\ell} \widetilde{\mu}_\ell(T)^2 \leq (1 - \theta) \widetilde{\mu}_\ell^2. \tag{33}$$

The combination of the three foregoing observations reads

$$\begin{aligned} \|\nabla \widehat{u}_\ell - \Pi_{\ell+1}(\nabla \widehat{u}_\ell)\|_{L^2(\Omega)}^2 &= \sum_{T \in \mathcal{T}_\ell \setminus \mathcal{M}_\ell} \|\nabla \widehat{u}_\ell - \Pi_{\ell+1}(\nabla \widehat{u}_\ell)\|_{L^2(T)}^2 \\ &\leq \sum_{T \in \mathcal{T}_\ell \setminus \mathcal{M}_\ell} \widetilde{\mu}_\ell(T)^2 \leq \rho \widetilde{\mu}_\ell^2 \end{aligned}$$

with $\rho := 1 - \theta < 1$. For any $\delta > 0$, Young’s inequality thus proves

$$\widetilde{\mu}_{\ell+1}^2 \leq (1 + \delta)\rho \widetilde{\mu}_\ell^2 + (1 + \delta^{-1})\|\widehat{u}_{\ell+1} - \widehat{u}_\ell\|^2.$$

To abbreviate the notation, we define $e_\ell := \|u - u_\ell\|^2$ and $E_\ell := \|u_{\ell+1} - u_\ell\|^2$ as well as $\widehat{e}_\ell := \|u - \widehat{u}_\ell\|^2$ and $\widehat{E}_\ell := \|\widehat{u}_{\ell+1} - \widehat{u}_\ell\|^2$.

Since $\mathcal{T}_{\ell+1}$ is obtained by refinement of \mathcal{T}_ℓ it holds that $X_\ell \subsetneq X_{\ell+1}$, and therefore, Galerkin orthogonality implies that $e_{\ell+1} + E_\ell = e_\ell$. Similarly, since $\widehat{X}_\ell \subsetneq \widehat{X}_{\ell+1}$, we also have $\widehat{e}_{\ell+1} + \widehat{E}_\ell = \widehat{e}_\ell$.

With this notation, the last inequality can be rewritten as

$$\widetilde{\mu}_{\ell+1}^2 \leq (1 + \delta)\rho \widetilde{\mu}_\ell^2 + C_\delta(E_\ell + \widehat{E}_\ell), \tag{34}$$

where $C_\delta = (1 + \delta^{-1})$. The additional term E_ℓ will appear for the adaptive boundary element schemes discussed below, hence we already include it in our present analysis.

We now choose $\delta > 0$ such that $(1 + \delta)\rho < 1$. Next, we choose $\gamma \in (0, 1)$ such that $\gamma C_\delta \leq 1$. Then, Galerkin orthogonality and choice of γ prove

$$\begin{aligned} \widetilde{\Delta}_{\ell+1} &= e_{\ell+1} + \widehat{e}_{\ell+1} + \gamma \widetilde{\mu}_{\ell+1}^2 = e_\ell + \widehat{e}_\ell - (E_\ell + \widehat{E}_\ell) + \gamma \widetilde{\mu}_{\ell+1}^2 \\ &\leq e_\ell + \widehat{e}_\ell + \gamma(1 + \delta)\rho \widetilde{\mu}_\ell^2. \end{aligned} \tag{35}$$

Note that so far only the last term on the right-hand side is contractive. Therefore, we choose an additional parameter $\beta > 0$ with $\beta + (1 + \delta)\rho < 1$ and write

$$e_\ell + \gamma(1 + \delta)\rho \tilde{\mu}_\ell^2 = (e_\ell - \gamma\beta\tilde{\mu}_\ell^2) + \gamma[\beta + (1 + \delta)\rho]\tilde{\mu}_\ell^2.$$

Recall that the saturation assumption (5) is equivalent to the reliability (4) of the $(h - h/2)$ error estimator η_ℓ . Thus, the equivalence (6) of η_ℓ and $\tilde{\mu}_\ell$ proves reliability of $\tilde{\mu}_\ell$, i.e. $e_\ell = \|u - u_\ell\|^2 \leq C_4 \tilde{\mu}_\ell^2$, and hence

$$\tilde{\Delta}_{\ell+1} \leq (1 - \gamma\beta C_4^{-1})e_\ell + \widehat{e}_\ell + \gamma[\beta + (1 + \delta)\rho]\tilde{\mu}_\ell^2.$$

Finally, we choose $\varepsilon > 0$ so that $\varepsilon < \gamma\beta C_4^{-1}$. Since $\widehat{e}_\ell \leq e_\ell$, we obtain

$$\tilde{\Delta}_{\ell+1} \leq (1 - \gamma\beta C_4^{-1} + \varepsilon)e_\ell + (1 - \varepsilon)\widehat{e}_\ell + \gamma[\beta + (1 + \delta)\rho]\tilde{\mu}_\ell^2 \leq \kappa \tilde{\Delta}_\ell,$$

where $\kappa = \max \{ 1 - \gamma\beta C_4^{-1} + \varepsilon, 1 - \varepsilon, \beta + (1 + \delta)\rho \} < 1$. □

Remark Note that, in view of (35) and $(1 + \delta)\rho < 1$, the error quantity $\tilde{\Delta}_\ell$ is monotonically decreasing, i.e. $\tilde{\Delta}_{\ell+1} \leq \tilde{\Delta}_\ell$, even if the saturation assumption fails to hold.

Remark We stress that only the estimator reduction (34) depends on the precise mathematical setting as well as on the refinement indicators chosen. The remainder of the proof works in a general framework so that we shall only verify (34) in the following, whenever we aim to prove convergence by verification of a contraction property for some error quantity.

Next, we observe that the previous theorem also implies convergence of the μ_ℓ -based AFEM. We stress that the following result is weaker than Theorem 4 in the sense that the latter provides a contraction property (32) for some error quantity $\tilde{\Delta}_\ell = \|u - u_\ell\|^2 + \|u - \widehat{u}_\ell\|^2 + \gamma\tilde{\mu}_\ell^2$ which then necessarily satisfies $\lim_\ell \tilde{\Delta}_\ell = 0$. In contrast, the following result only states $\lim_\ell \Delta_\ell = 0$, where $\Delta_\ell = \|u - u_\ell\|^2 + \|u - \widehat{u}_\ell\|^2 + \gamma\mu_\ell^2$. The contraction property of Δ_ℓ is only proven for sufficiently large $\theta \in (0, 1)$. This, however, seems to be suboptimal: We stress that $\theta = 1$ generically leads to uniform mesh-refinement, whereas small $\theta \ll 1$ yields highly adapted meshes.

Theorem 5 *Suppose that we use the indicators $\mu_\ell(T)$ defined in (25) in Algorithm 1, and assume that the saturation assumption (5) holds. Then, for any choice of $\theta \in (0, 1)$,*

$$\lim_{\ell \rightarrow \infty} \|u - \widehat{u}_\ell\| = \lim_{\ell \rightarrow \infty} \|u - u_\ell\| = 0 = \lim_{\ell \rightarrow \infty} \mu_\ell. \tag{36}$$

Provided that $\theta \in (0, 1)$ is sufficiently large, there are constants $\kappa, \gamma \in (0, 1)$ such that the weighted error quantity

$$\Delta_\ell := \|u - u_\ell\|^2 + \|u - \widehat{u}_\ell\|^2 + \gamma\mu_\ell^2 \tag{37}$$

satisfies the contraction property

$$\Delta_{\ell+1} \leq \kappa \Delta_\ell. \tag{38}$$

Proof of Theorem 5 (Convergence) From (29)–(30), we infer

$$\sum_{T \in \mathcal{M}_\ell} \mu_\ell(T)^2 \leq C_3 \sum_{T \in \mathcal{M}_\ell} \tilde{\mu}_\ell(T)^2 \quad \text{as well as} \quad \tilde{\mu}_\ell^2 \leq \mu_\ell^2$$

with some constant $C_3 \geq 1$. Therefore, μ_ℓ -based Dörfler marking (7) with constant $\theta \in (0, 1)$ implies $\tilde{\mu}_\ell$ -based Dörfler marking with constant $\tilde{\theta} = \theta/C_3 \in (0, 1)$, namely

$$\theta \tilde{\mu}_\ell^2 \leq \theta \mu_\ell^2 \leq \sum_{T \in \mathcal{M}_\ell} \mu_\ell(T)^2 \leq C_3 \sum_{T \in \mathcal{M}_\ell} \tilde{\mu}_\ell(T)^2.$$

Therefore, Theorem 4 implies that $\lim_\ell \|u - \hat{u}_\ell\| = \lim_\ell \|u - u_\ell\| = 0 = \lim_\ell \tilde{\mu}_\ell$. From the equivalence of $\tilde{\mu}_\ell$ and μ_ℓ , we obtain $\lim_\ell \mu_\ell = 0$.

Proof of Theorem 5 (Contraction property) A scaling argument with respect to \mathcal{T}_ℓ and continuity of $I_{\ell+1}$ onto the finite dimensional space $\hat{X}_{\ell+1}$ provide some constant $C_5 > 0$ with

$$\|\nabla I_{\ell+1}(\hat{u}_{\ell+1} - \hat{u}_\ell)\|_{L^2(T)} \leq C_5 \|\nabla(\hat{u}_{\ell+1} - \hat{u}_\ell)\|_{L^2(T)} \quad \text{for all } T \in \mathcal{T}_\ell.$$

The triangle inequality thus yields

$$\begin{aligned} \mu_{\ell+1} = \|\hat{u}_{\ell+1} - I_{\ell+1}\hat{u}_{\ell+1}\| &\leq \|\hat{u}_{\ell+1} - \hat{u}_\ell\| + \|\hat{u}_\ell - I_{\ell+1}\hat{u}_\ell\| + \|I_{\ell+1}(\hat{u}_{\ell+1} - \hat{u}_\ell)\| \\ &\leq \|\hat{u}_\ell - I_{\ell+1}\hat{u}_\ell\| + (C_5 + 1) \|\hat{u}_{\ell+1} - \hat{u}_\ell\|. \end{aligned}$$

Copying the proof of Proposition 3 verbatim, we obtain a constant C_6 such that

$$\|\nabla(\hat{u}_\ell - I_{\ell+1}\hat{u}_\ell)\|_{L^2(T)} \leq C_6 \|\nabla\hat{u}_\ell - \Pi_{\ell+1}(\nabla\hat{u}_\ell)\|_{L^2(T)},$$

and in particular,

$$C_6^{-1} \|\nabla(\hat{u}_\ell - I_{\ell+1}\hat{u}_\ell)\|_{L^2(T)} \leq \|\nabla\hat{u}_\ell - \Pi_\ell(\nabla\hat{u}_\ell)\|_{L^2(T)} = \tilde{\mu}_\ell(T) \quad \text{for all } T \in \mathcal{T}_\ell. \tag{39}$$

From $\tilde{\mu}_\ell(T) \leq \mu_\ell(T)$, we thus infer

$$\|\nabla(\hat{u}_\ell - I_{\ell+1}\hat{u}_\ell)\|_{L^2(T)} \leq C_6 \mu_\ell(T) \quad \text{for all } T \in \mathcal{T}_\ell.$$

The refinement strategy gives

$$\|\nabla(\hat{u}_\ell - I_{\ell+1}\hat{u}_\ell)\|_{L^2(T)} = 0 \quad \text{for all } T \in \mathcal{M}_\ell.$$

The combination of the latter two estimates proves

$$\begin{aligned} \|\widehat{u}_\ell - I_{\ell+1}\widehat{u}_\ell\|^2 &= \sum_{T \in \mathcal{T}_\ell \setminus \mathcal{M}_\ell} \|\nabla(\widehat{u}_\ell - I_{\ell+1}\widehat{u}_\ell)\|_{L^2(T)}^2 \\ &\leq C_6^2 \sum_{T \in \mathcal{T}_\ell \setminus \mathcal{M}_\ell} \mu_\ell(T)^2 \leq C_6^2(1 - \theta) \mu_\ell^2, \end{aligned}$$

where the final estimate follows from the Dörfler marking, cf. (33). We assume that $\theta \in (0, 1)$ is sufficiently large to guarantee $\rho = C_6^2(1 - \theta) < 1$. Using the same notation as in the proof of Theorem 4, we obtain the estimator reduction (34) with $C_\delta = (1 + \delta^{-1})(C_5 + 1)^2$.

From a conceptual point of view, it seems more natural to consider the local contributions $\eta_\ell(T)$ of $\eta_\ell = \|\widehat{u}_\ell - u_\ell\|$ — instead of $\widetilde{\mu}_\ell(T)$ — for marking in step (d) of Algorithm 1.

We stress, however, that an adaptive algorithm will usually return \widehat{u}_ℓ instead of u_ℓ since $\|\widehat{u}_\ell - u_\ell\| \leq \|u - u_\ell\|$. One advantage of $\widetilde{\mu}_\ell$ and μ_ℓ from (25) over $\eta_\ell(T)$ now is that we do not need to compute the coarse-mesh solution $u_\ell \in X_\ell$. Instead, $\Pi_\ell(\nabla\widehat{u}_\ell)$, for instance, can simply be computed in linear complexity by $\Pi_\ell(\nabla\widehat{u}_\ell)|_T = (1/|T|) \int_T \nabla\widehat{u}_\ell \, dx$, for all $T \in \mathcal{T}_\ell$.

Remark Although numerical experiments give evidence that the steering of Algorithm 1 by $\eta_\ell(T)$ leads to a convergent adaptive scheme, we did not succeed to prove convergence for arbitrary but only for *sufficiently large* $\theta \in (0, 1)$.

For example, closely following the proof of the contraction property in Theorem 5 one can show that, if $(1 - \theta)C_6^2 < 1$, then the combined error quantity

$$\overline{\Delta}_\ell := \|u - u_\ell\|^2 + \|u - \widehat{u}_\ell\|^2 + \gamma \eta_\ell^2$$

satisfies a contraction property.

2.6 Convergence of AFEM without the saturation assumption

In this section, we include the resolution of the data oscillations into the adaptive algorithm. With

$$\text{osc}_\ell(T)^2 = \|h_\ell(f - \Pi_\ell f)\|_{L^2(T)}^2 + \|h_\ell^{1/2}(g - \Pi_\ell g)\|_{L^2(\partial T \cap \Gamma_N)}^2, \tag{40}$$

we consider

$$\widetilde{\zeta}_\ell^2 = \sum_{T \in \mathcal{T}_\ell} \widetilde{\zeta}_\ell(T)^2 = \widetilde{\mu}_\ell^2 + \text{osc}_\ell^2, \quad \text{where } \widetilde{\zeta}_\ell(T)^2 = \widetilde{\mu}_\ell(T)^2 + \text{osc}_\ell(T)^2. \tag{41}$$

By use of Proposition 2, we will show that this estimator enforces the saturation assumption to hold. The following theorem states convergence of the $\widetilde{\zeta}_\ell$ -based version of AFEM.

Theorem 6 *Suppose that we use the indicators $\tilde{\zeta}_\ell(T)$ defined in (41) in Algorithm 1. Then, for any $\theta \in (0, 1)$, there is a constant $\gamma \in (0, 1)$ such that the combined error quantity*

$$\tilde{Z}_\ell := \|u - u_\ell\|^2 + \|u - \hat{u}_\ell\|^2 + \gamma \tilde{\zeta}_\ell^2$$

satisfies $\lim_{\ell \rightarrow \infty} \tilde{Z}_\ell = 0$. In particular, it holds that $\lim_{\ell \rightarrow \infty} \|u - \hat{u}_\ell\| = \lim_{\ell \rightarrow \infty} \|u - u_\ell\| = 0 = \lim_{\ell \rightarrow \infty} \tilde{\zeta}_\ell$.

Proof Note that the data oscillations satisfy

$$\begin{aligned} \text{osc}_{\ell+1}(T)^2 &\leq \text{osc}_\ell(T)^2 \quad \text{for } T \in \mathcal{T}_\ell, \\ \text{as well as } \text{osc}_{\ell+1}(T)^2 &\leq \frac{1}{2} \text{osc}_\ell(T)^2 \quad \text{for } T \in \mathcal{M}_\ell. \end{aligned}$$

In particular, the data oscillations are monotonically decreasing. With the same arguments as, e.g., in the proof of Theorem 4 above, we thus obtain the estimator reduction

$$\tilde{\zeta}_{\ell+1}^2 \leq (1 + \delta)\rho \tilde{\zeta}_\ell^2 + 4(1 + \delta^{-1})\|\hat{u}_{\ell+1} - \hat{u}_\ell\|^2 \tag{42}$$

with $\rho = 1 - \theta/2$, for all $\delta > 0$. Arguing as above, we find $\gamma \in (0, 1)$ such that \tilde{Z}_ℓ is monotonically decreasing, i.e. $\tilde{Z}_{\ell+1} \leq \tilde{Z}_\ell$. Moreover, the saturation assumption

$$C_\ell := \frac{\|u - \hat{u}_\ell\|}{\|u - u_\ell\|} < 1$$

in the ℓ -step implies contraction $\tilde{Z}_{\ell+1} \leq \kappa_\ell \tilde{Z}_\ell$ for some $\kappa_\ell \in (0, 1)$, which depends on C_ℓ and which satisfies $\kappa_\ell \rightarrow 1$ if and only if $C_\ell \rightarrow 1$ as $\ell \rightarrow \infty$.

Provided that $\liminf_\ell C_\ell < 1$, the according subsequence thus satisfies $1 > \kappa \geq \kappa_{\ell_k}$. Therefore, monotonicity of \tilde{Z}_ℓ yields $\tilde{Z}_{\ell_{k+1}} \leq \tilde{Z}_{\ell_k} \leq \kappa \tilde{Z}_{\ell_k}$, whence $\lim_k \tilde{Z}_{\ell_k} = 0$. Again by monotonicity of \tilde{Z}_ℓ , the whole sequence satisfies $\lim_\ell \tilde{Z}_\ell = 0$.

We may thus restrict to consider the case $\lim_\ell C_\ell = \liminf_\ell C_\ell = 1$. Note that the Galerkin orthogonality $\|u - \hat{u}_\ell\|^2 + \eta_\ell^2 = \|u - u_\ell\|^2$ then implies $\lim_\ell \eta_\ell^2 = 0$, hence $\lim_\ell \tilde{\mu}_\ell^2 = 0$. For $\varepsilon > 0$, we choose a sufficiently large ℓ_0 such that $\tilde{\mu}_\ell^2 \leq \varepsilon$ for all $\ell \geq \ell_0$. Dörfler’s marking strategy provides

$$\begin{aligned} \text{osc}_{\ell+1}^2 &\leq \sum_{T \in \mathcal{T}_\ell \setminus \mathcal{M}_\ell} \text{osc}_\ell(T)^2 + \frac{1}{2} \sum_{T \in \mathcal{M}_\ell} \text{osc}_\ell(T)^2 \\ &= \text{osc}_\ell^2 - \frac{1}{2} \sum_{T \in \mathcal{M}_\ell} \tilde{\zeta}_\ell(T)^2 + \frac{1}{2} \sum_{T \in \mathcal{M}_\ell} \tilde{\mu}_\ell(T)^2 \\ &\leq \text{osc}_\ell^2 - \frac{\theta}{2} \tilde{\zeta}_\ell^2 + \frac{1}{2} \sum_{T \in \mathcal{M}_\ell} \tilde{\mu}_\ell(T)^2 \\ &\leq (1 - \theta/2) \text{osc}_\ell^2 + \varepsilon/2. \end{aligned}$$

By induction, the geometric series provides

$$\text{osc}_{\ell+k}^2 \leq (1 - \theta/2)^k \text{osc}_\ell^2 + (\varepsilon/2) \sum_{j=1}^k (1 - \theta/2)^j \leq (1 - \theta/2)^k \text{osc}_0^2 + \varepsilon/\theta,$$

where we have used the monotonicity of data oscillations $\text{osc}_{\ell+1}^2 \leq \text{osc}_\ell^2$. From this, we infer $\limsup_\ell \text{osc}_\ell^2 \leq \varepsilon/\theta$ for all $\varepsilon > 0$ and conclude $\lim_\ell \text{osc}_\ell^2 = 0$. Due to $\|u - \widehat{u}_\ell\| \leq \|u - u_\ell\|$ and $\lim_\ell \widetilde{\zeta}_\ell^2 = 0$, the estimate $\widetilde{Z}_\ell > C > 0$ then implies $\|u - u_\ell\|^2 \geq C/2$. In particular, Proposition 2 implies that the saturation assumption holds for all $\ell \geq \ell_0$, as soon as $\text{osc}_{\ell_0}^2 \leq \lambda C/4$. This, however, contradicts $\lim_\ell C_\ell = 1$ and thus proves $\lim_\ell \widetilde{Z}_\ell = 0$. □

Remark Using the arguments of Theorem 5 above, one may prove that, for arbitrary $\theta \in (0, 1)$, the indicators $\zeta_\ell(T)^2 := \mu_\ell(T)^2 + \text{osc}_\ell(T)^2$ also lead to a convergent version of AFEM.

3 Boundary element method

3.1 Symm’s integral equation

As model problem, we consider the first-kind integral equation

$$(Vu)(x) = \int_\Gamma G(x, y)u(y) \, ds_y = F(x) \quad \text{for } x \in \Gamma \tag{43}$$

with the weakly-singular integral kernel

$$G(x, y) = \begin{cases} -\frac{1}{2\pi} \log |x - y| & \text{for } d = 2, \\ +\frac{1}{4\pi} \frac{1}{|x-y|} & \text{for } d = 3. \end{cases} \tag{44}$$

Here, Γ is an open piece of the boundary $\partial\Omega$ of a Lipschitz domain Ω in \mathbb{R}^d , and ds denotes the integration along the arclength or on the manifold for $d = 2, 3$, respectively. For $d = 2$, we additionally assume $\text{diam}(\Omega) < 1$. We define the fractional-order Sobolev space

$$H^{1/2}(\Gamma) := \{\widetilde{u}|_\Gamma : \widetilde{u} \in H^1(\Omega)\}$$

with norm

$$\|u\|_{H^{1/2}(\Gamma)} := \inf \left\{ \|\widetilde{u}\|_{H^1(\Omega)} : \widetilde{u} \in H^1(\Omega) \text{ with } \widetilde{u}|_\Gamma = u \right\}.$$

Then, $H^{1/2}(\Gamma)$ is a Hilbert space, and we define $\widetilde{H}^{-1/2}(\Gamma)$ as the algebraic-topological dual of $H^{1/2}(\Gamma)$ with respect to the extended L^2 -scalar product $\langle \cdot, \cdot \rangle$. The single-layer

potential V is an elliptic isomorphism between $\mathcal{H} = \tilde{H}^{-1/2}(\Gamma)$ and $H^{1/2}(\Gamma)$, cf. [18]. The energy scalar product is thus given by

$$\langle\langle u, v \rangle\rangle := \langle Vu, v \rangle \quad \text{for all } u, v \in \mathcal{H} = \tilde{H}^{-1/2}(\Gamma). \tag{45}$$

We consider the lowest-order Galerkin scheme, where \mathcal{T}_ℓ is a triangulation of Γ and where $X_\ell := \mathcal{P}^0(\mathcal{T}_\ell)$ denotes the space of all \mathcal{T}_ℓ -piecewise constant functions on Γ .

As above, we define the local mesh-width function $h_\ell \in \mathcal{P}^0(\mathcal{T}_\ell)$ by $h_\ell|_T := \text{diam}(T)$ for $T \in \mathcal{T}_\ell$.

3.2 Mesh-refinement

We restrict to the case that the boundary elements $T \in \mathcal{T}_\ell$ are affine line segments for $d = 2$ and planar triangles for $d = 3$, respectively.

In 2D, a marked element $T \in \mathcal{M}_\ell$ is halved into two elements $T_1, T_2 \in \mathcal{T}_{\ell+1}$. To control the K -mesh constant

$$\kappa(\mathcal{T}_\ell) = \max \{h_\ell|_T / h_\ell|_{T'} : T, T' \in \mathcal{T}_\ell \text{ neighbouring elements}\}, \tag{46}$$

we recursively do some additional marking: If the neighbour $T' \in \mathcal{T}_\ell$ of a marked element $T \in \mathcal{M}_\ell$ satisfies $h_\ell|_{T'} / h_\ell|_T > \kappa(\mathcal{T}_0)$, we also mark the element T' for refinement. This ensures boundedness $\kappa(\mathcal{T}_\ell) \leq 2\kappa(\mathcal{T}_0)$ as $\ell \rightarrow \infty$.

In 3D, we use the mesh-refinement introduced in Sect. 2.2.

3.3 Refinement indicators

Note that the energy norm $\|\cdot\|$ is nonlocal in the sense that $\|\cdot\|^2 = \|\cdot\|_\Gamma^2$ is not equivalent to $\sum_{T \in \mathcal{T}_\ell} \|\cdot\|_T^2$, where $\|v\|_T^2 = \int_T \int_T u(x)G(x, y)u(y) ds_y ds_x$, cf. [7]. Therefore, the error estimator η_ℓ does not provide information for a local mesh-refinement directly. Instead, we consider the refinement indicators

$$\mu_\ell(T) = \left\| h_\ell^{1/2}(\widehat{u}_\ell - u_\ell) \right\|_{L^2(T)} \quad \text{as well as} \quad \tilde{\mu}_\ell(T) = \left\| h_\ell^{1/2}(\widehat{u}_\ell - \Pi_\ell \widehat{u}_\ell) \right\|_{L^2(T)} \tag{47}$$

for all $T \in \mathcal{T}_\ell$, where Π_ℓ denotes the L^2 -orthogonal projection onto $X_\ell = \mathcal{P}^0(\mathcal{T}_\ell)$. Using a local inverse estimate from [15] and a local approximation result from [9], one can prove estimator equivalence [14]

$$C_1^{-1} \eta_\ell \leq \tilde{\mu}_\ell = \left\| h_\ell^{1/2}(\widehat{u}_\ell - \Pi_\ell \widehat{u}_\ell) \right\|_{L^2(\Gamma)} \leq \mu_\ell = \left\| h_\ell^{1/2}(\widehat{u}_\ell - u_\ell) \right\|_{L^2(\Gamma)} \leq C_2 \eta_\ell. \tag{48}$$

The constant C_1 only depends on Γ , whereas C_2 stems from an inverse estimate and additionally depends on the polynomial degree and on the shape regularity constant of

the triangulation \mathcal{T}_ℓ . The proposed mesh-refinement from Sect. 3.2 thus ensures that C_2 remains uniformly bounded.

Remark Note that the definition of h_ℓ has to be consistent with the mesh-refining strategy in the sense that

$$h_{\ell+1}|_T \leq q h_\ell|_T \quad \text{for all } T \in \mathcal{M}_\ell \tag{49}$$

where $q \in (0, 1)$ is a uniform contraction rate. In [10], newest vertex bisection, with the slightly non-standard definition $h_\ell|_T = |T|^{1/d}$ instead of $h_\ell|_T = \text{diam}(T)$ is used to ensure (49). In our refinement strategy, marked elements are bisecc_5 -refined so that (49) holds with the usual definition of h_ℓ and $q = 1/2$.

3.4 Convergence of ABEM under the saturation assumption

In this section, we prove convergence of ABEM steered by either μ_ℓ or $\tilde{\mu}_\ell$ from (47). For both error estimators, we prove the estimator reduction (34).

Theorem 7 *Suppose that we use the indicators $\tilde{\mu}_\ell(T)$ defined in (47) in Algorithm 1. Under the saturation assumption (5), and for any choice of $\theta \in (0, 1)$, there are constants $\kappa, \gamma \in (0, 1)$ such that the combined error quantity*

$$\tilde{\Delta}_\ell := \| \|u - u_\ell\|^2 + \| \|u - \hat{u}_\ell\|^2 + \gamma \tilde{\mu}_\ell^2 \tag{50}$$

satisfies the contraction property

$$\tilde{\Delta}_{\ell+1} \leq \kappa \tilde{\Delta}_\ell. \tag{51}$$

In particular, it holds that $\lim_{\ell \rightarrow \infty} \| \|u - \hat{u}_\ell\| = \lim_{\ell \rightarrow \infty} \| \|u - u_\ell\| = 0 = \lim_{\ell \rightarrow \infty} \tilde{\mu}_\ell$.

Proof Recall that $X_\ell \subset X_{\ell+1} \subset \hat{X}_\ell \subset \hat{X}_{\ell+1}$. Moreover, we note the pointwise estimates $h_\ell/4 \leq \hat{h}_{\ell+1} \leq h_\ell/2$ and $h_\ell/16 \leq \hat{h}_{\ell+1} \leq h_\ell/4$ for $d = 2, 3$, respectively. Using the fact that the mesh-sizes h_ℓ and $\hat{h}_{\ell+1}$ are equivalent up to a multiplicative factor, and applying [15, Theorem 4.1], we obtain the inverse estimate

$$\left\| h_\ell^{1/2} \hat{v}_{\ell+1} \right\|_{L^2(\Gamma)} \leq C_{\text{inv}} \| \hat{v}_{\ell+1} \| \quad \text{for all } \hat{v}_{\ell+1} \in \hat{X}_{\ell+1}. \tag{52}$$

In 2D, the constant $C_{\text{inv}} > 0$ only depends on the mesh-ratio $\kappa(\hat{\mathcal{T}}_{\ell+1}) \leq 2\kappa(\mathcal{T}_0)$. In 3D, the constant $C_{\text{inv}} > 0$ only depends on the smallest interior angle in $\hat{\mathcal{T}}_{\ell+1}$, and in particular, since $\hat{\mathcal{T}}_{\ell+1}$ is obtained from \mathcal{T}_0 by certain newest vertex bisections, it depends only on the smallest interior angle in \mathcal{T}_0 .

We use the triangle inequality, the $\mathcal{T}_{\ell+1}$ -piecewise orthogonality of $\Pi_{\ell+1}$, and the inverse estimate (52) to obtain

$$\begin{aligned} \tilde{\mu}_{\ell+1} &= \left\| h_{\ell+1}^{1/2} (1 - \Pi_{\ell+1}) \widehat{u}_{\ell+1} \right\|_{L^2(\Gamma)} \\ &\leq \left\| h_{\ell+1}^{1/2} (1 - \Pi_{\ell+1}) \widehat{u}_{\ell} \right\|_{L^2(\Gamma)} + \left\| h_{\ell+1}^{1/2} (1 - \Pi_{\ell+1}) (\widehat{u}_{\ell+1} - \widehat{u}_{\ell}) \right\|_{L^2(\Gamma)} \\ &\leq \left\| h_{\ell+1}^{1/2} (1 - \Pi_{\ell+1}) \widehat{u}_{\ell} \right\|_{L^2(\Gamma)} + \left\| h_{\ell+1}^{1/2} (\widehat{u}_{\ell+1} - \widehat{u}_{\ell}) \right\|_{L^2(\Gamma)} \\ &\leq \left\| h_{\ell+1}^{1/2} (1 - \Pi_{\ell+1}) \widehat{u}_{\ell} \right\|_{L^2(\Gamma)} + C_{\text{inv}} \|\widehat{u}_{\ell+1} - \widehat{u}_{\ell}\|. \end{aligned}$$

To estimate the second term, we now proceed precisely as in the proof of Theorem 4 (AFEM) to verify that

$$\left\| h_{\ell+1}^{1/2} (\widehat{u}_{\ell} - \Pi_{\ell+1} \widehat{u}_{\ell}) \right\|_{L^2(\Gamma)}^2 \leq \rho \tilde{\mu}_{\ell}^2,$$

where $\rho = 1 - \theta$. Using the same notation as in the proof of Theorem 4, we obtain the estimator reduction (34) with $C_{\delta} = (1 + \delta^{-1}) C_{\text{inv}}^2$. □

Theorem 8 *Suppose that we use the indicators $\mu_{\ell}(T)$ defined in (47) in Algorithm 1. Under the saturation assumption (5), and for any choice of $\theta \in (0, 1)$, there are constants $\kappa, \gamma \in (0, 1)$ such that*

$$\Delta_{\ell} := \|u - u_{\ell}\|^2 + \|u - \widehat{u}_{\ell}\|^2 + \gamma \mu_{\ell}^2 \tag{53}$$

satisfies the contraction property

$$\Delta_{\ell+1} \leq \kappa \Delta_{\ell}. \tag{54}$$

In particular, Algorithm 1 is convergent, that is, $\lim_{\ell \rightarrow \infty} \|u - \widehat{u}_{\ell}\| = \lim_{\ell \rightarrow \infty} \|u - u_{\ell}\| = 0 = \lim_{\ell \rightarrow \infty} \mu_{\ell}$.

Proof The triangle inequality and the inverse estimate (52) prove

$$\begin{aligned} \mu_{\ell+1} &= \left\| h_{\ell+1}^{1/2} (\widehat{u}_{\ell+1} - u_{\ell+1}) \right\|_{L^2(\Gamma)} \\ &\leq \left\| h_{\ell+1}^{1/2} (\widehat{u}_{\ell} - u_{\ell}) \right\|_{L^2(\Gamma)} + \left\| h_{\ell+1}^{1/2} ((\widehat{u}_{\ell+1} - u_{\ell+1}) - (\widehat{u}_{\ell} - u_{\ell})) \right\|_{L^2(\Gamma)} \\ &\leq \left\| h_{\ell+1}^{1/2} (\widehat{u}_{\ell} - u_{\ell}) \right\|_{L^2(\Gamma)} + C_{\text{inv}} \|\widehat{u}_{\ell+1} - \widehat{u}_{\ell} - (u_{\ell+1} - u_{\ell})\|. \end{aligned}$$

The mesh-refinement rule from Sect. 3.2 yields the properties

$$\begin{aligned} h_{\ell+1}|_T &\leq \frac{1}{2} h_{\ell}|_T \quad \text{for all } T \in \mathcal{M}_{\ell}, \\ \text{and } h_{\ell+1}|_T &\leq h_{\ell}|_T \quad \text{for all } T \in \mathcal{T}_{\ell} \setminus \mathcal{M}_{\ell}. \end{aligned}$$

Dörfler’s marking strategy (7) gives

$$\begin{aligned}
 \|h_{\ell+1}^{1/2}(\widehat{u}_\ell - u_\ell)\|_{L^2(\Gamma)}^2 &= \sum_{T \in \mathcal{M}_\ell} \|h_{\ell+1}^{1/2}(\widehat{u}_\ell - u_\ell)\|_{L^2(T)}^2 \\
 &\quad + \sum_{T \in \mathcal{T}_\ell \setminus \mathcal{M}_\ell} \|h_{\ell+1}^{1/2}(\widehat{u}_\ell - u_\ell)\|_{L^2(T)}^2 \\
 &\leq \frac{1}{2} \sum_{T \in \mathcal{M}_\ell} \|h_\ell^{1/2}(\widehat{u}_\ell - u_\ell)\|_{L^2(T)}^2 \\
 &\quad + \sum_{T \in \mathcal{T}_\ell \setminus \mathcal{M}_\ell} \|h_\ell^{1/2}(\widehat{u}_\ell - u_\ell)\|_{L^2(T)}^2 \\
 &= -\frac{1}{2} \sum_{T \in \mathcal{M}_\ell} \mu_\ell(T)^2 + \sum_{T \in \mathcal{T}_\ell} \mu_\ell(T)^2 \\
 &\leq \left(1 - \frac{\theta}{2}\right) \sum_{T \in \mathcal{T}_\ell} \mu_\ell(T)^2 \\
 &= \rho \mu_\ell^2,
 \end{aligned}$$

with $\rho := (1 - \theta/2) < 1$. For any $\delta > 0$, Young’s inequality then proves

$$\begin{aligned}
 \mu_{\ell+1}^2 &\leq (1 + \delta)\rho \mu_\ell^2 + (1 + \delta^{-1})C_{\text{inv}}^2 \|\widehat{u}_{\ell+1} - \widehat{u}_\ell - (u_{\ell+1} - u_\ell)\|^2 \\
 &= (1 + \delta)\rho \mu_\ell^2 + (1 + \delta^{-1})C_{\text{inv}}^2 \left(\|\widehat{u}_{\ell+1} - \widehat{u}_\ell\|^2 + \|u_{\ell+1} - u_\ell\|^2\right),
 \end{aligned}$$

where we have used Galerkin orthogonality and $X_\ell \subset X_{\ell+1} \subseteq \widehat{X}_\ell \subset \widehat{X}_{\ell+1}$. Using the same notation as in the proof of Theorem 4, we now have proven the estimator reduction (34) with $C_\delta = (1 + \delta^{-1})C_{\text{inv}}^2$. □

4 Numerical experiments

In the following section, we present three numerical experiments to underline our theoretical findings. Throughout, the energy error is computed by the Galerkin orthogonality

$$\|u - u_\ell\|^2 = \|u\|^2 - \|u_\ell\|^2. \tag{55}$$

Note that discrete energies $\|u_\ell\|^2$ can be computed by use of the Galerkin matrix. The exact energy $\|u\|^2$ is extrapolated by Aitken’s Δ^2 -method applied to certain discrete energies $\|u_\ell\|^2$ corresponding to a sequence of uniformly refined meshes. Here, uniform refinement means that, for 2D FEM or 3D BEM, all triangles are split into 4 similar son triangles by halving its edges, whereas boundary segments in 2D BEM are split into 2 son segments of half length.

Besides the Galerkin error $\|u - u_\ell\|$, we plot the natural $(h - h/2)$ -error estimators η_ℓ as well as the (problem dependent) error estimator $\tilde{\mu}_\ell$. We empirically compare uniform mesh-refinement with $\tilde{\mu}_\ell$ -steered mesh-refinement. Throughout, Algorithm 1 turns out to be effective in the sense that even the optimal rate of convergence is recovered.

4.1 Finite element method

In the first experiment, we consider the numerical solution of (11). The domain $\Omega \subset \mathbb{R}^2$, visualized in Fig. 4, is a rotated Z-shape which is obtained by rotation of $(-0.25, 0.25)^2 \setminus \text{conv}\{(0, 0), (-0.25, -0.25), (0, -0.25)\}$ with angle $-\pi/8$. The exact solution reads

$$u(x) = r^{4/7} \cos(4\varphi/7) \quad \text{in polar coordinates } x = r(\cos \varphi, \sin \varphi) \quad (56)$$

and has a generic singularity at the re-entrant corner $(0, 0)$. We stress that the rotation of the Z-shape is done in a way that provides homogeneous Dirichlet data at the two diagonal boundary edges, cf. Fig. 4. The initial mesh \mathcal{T}_0 consisted of $N = 14$ rectangular triangles, where the longest edges are the reference edges.

Recall that, for a piecewise H^2 solution, the optimal order of convergence in this experiment is $\mathcal{O}(N^{-1/2})$, where $N = \#\mathcal{T}$ is the number of elements. Figure 5 provides the experimental results for uniform and adaptive mesh-refinement, where we use

$$\tilde{\xi}_\ell = (\tilde{\mu}_\ell^2 + \text{osc}_\ell^2)^{1/2}$$

and $\theta = 0.25$ for marking in the adaptive algorithm. Adaptive mesh-refinement is performed as described in Sect. 2.2. We stress that Theorem 4 predicts convergence of this adaptive scheme. Uniform mesh-refinement leads to a suboptimal order of con-




Fig. 4 Initial mesh in FE Example 4.1 with 14 rectangular elements (left) and initial mesh in BE Example 4.2 with 10 affine boundary segments (right). The reference edges for the newest vertex bisection in the FE mesh are indicated by double lines, and the nodes on the Dirichlet boundary are indicated by red bullets (left) (color figure online)




Fig. 5 Error $\|u - u_\ell\| = \|\nabla(u - u_\ell)\|_{L^2(\Omega)}$ and error estimators η_ℓ and $\tilde{\zeta}_\ell = (\tilde{\mu}_\ell^2 + \text{osc}_\ell^2)^{1/2}$ for uniform and adaptive FEM in Example 4.1, where we use $\theta = 0.25$ and $\tilde{\zeta}_\ell$ for marking in Algorithm 1. Note that the adaptive strategy yields the optimal order of convergence

vergence $\mathcal{O}(N^{-2/7})$. The optimal order of convergence $\mathcal{O}(N^{-1/2})$ is recovered by use of the adaptive algorithm.

4.2 Boundary element method in 2D

In this experiment, we consider Symm’s integral equation

$$Vu = (K + 1/2)g \tag{57}$$

which is equivalent to the Dirichlet problem

$$\begin{aligned} -\Delta U &= 0 \quad \text{in } \Omega, \\ U &= g \quad \text{on } \Gamma. \end{aligned} \tag{58}$$

In this case, the exact solution of (57) is the normal derivative $u = \partial_n U$ of $U \in H^1(\Omega)$, and U can be obtained from (u, g) by use of Green’s third formula [18]. Kg denotes the double-layer potential, which is formally defined as the Cauchy principal value




Fig. 6 Error $\|u - u_\ell\| \sim \|u - u_\ell\|_{H^{-1/2}(\Gamma)}$ and error estimators η_ℓ and $\tilde{\mu}_\ell$ for uniform and adaptive BEM in Example 4.2, where we use $\theta = 0.25$ and $\tilde{\mu}_\ell$ for marking in Algorithm 1. Note that the adaptive strategy yields the optimal order of convergence




Fig. 7 Initial mesh with $N = 96$ rectangular triangles (left) and $\tilde{\mu}_\ell$ -adaptively generated mesh with $N = 20.806$ triangles (right) in Example 4.3

$$(Kg)(x) = \int_{\Gamma} C \partial_{n(y)} G(s, y) g(y) \, ds_y \quad \text{for } x \in \Gamma$$

and then extended to a linear operator on $H^{1/2}(\Gamma)$ by continuity. We consider the rotated Z-shape Ω of Experiment 4.1, and the exact solution of (58) is given by (56). We stress that Ω is scaled in a way that guarantees $\text{diam}(\Omega) < 1$ to ensure ellipticity




Fig. 8 Error $\|u - u_\ell\| \sim \|u - u_\ell\|_{H^{-1/2}(\Gamma)}$ and error estimators η_ℓ and $\tilde{\mu}_\ell$ for uniform and isotropic adaptive BEM in Example 4.3, where we use $\theta = 0.25$ and $\tilde{\mu}_\ell$ for marking in Algorithm 1

of the single-layer potential. The initial BE mesh \mathcal{T}_0 with $N = 10$ affine boundary segments is shown in Fig. 4.

If the normal derivative $u = \partial U / \partial n$ of U is a piecewise H^1 function then the optimal order of convergence in this experiment is $\mathcal{O}(N^{-3/2})$, in terms of the number $N = \#\mathcal{T}$ of elements [20]. The numerical results are shown in Fig. 6, where we again use $\theta = 0.25$ and $\tilde{\mu}_\ell$ for marking in the adaptive algorithm. The adaptive mesh-refinement is performed as described in Sect. 3.2. Uniform mesh-refinement leads to a suboptimal order of convergence $\mathcal{O}(N^{-4/7})$. This is cured by the adaptive algorithm in the sense that the optimal order of convergence is recovered.

For both, uniform and adaptive mesh-refinement, we observe that the error estimators η_ℓ and μ_ℓ remain parallel with $\|u - u_\ell\|$. This strongly indicates reliability of η_ℓ and thus numerically verifies the saturation assumption (5) for this experiment.

4.3 Boundary element method in 3D

Finally, we compute the discrete solution of Symm’s integral equation

$$Vu = 1 \quad \text{on } \Gamma$$

with Γ the boundary of the Fichera cube $\Omega = [-1, 1]^3 \setminus ([0, 1] \times [-1, 0] \times [0, 1])$. Figure 7 shows the uniform initial mesh \mathcal{T}_0 consisting of $N = 96$ rectangular triangles.

To illustrate the performance of Algorithm 1, Fig. 8 shows the error $\|u - u_\ell\|$ and the estimators η_ℓ and $\tilde{\mu}_\ell$ for uniform and adaptive mesh-refinements. For adaptive mesh-refinement, we use $\theta = 0.25$ and $\tilde{\mu}_\ell$ in Algorithm 1. Moreover, we use the biseq_5 -based mesh-refinement from Sect. 2.2. We observe that uniform mesh-refinement leads to a suboptimal order of convergence of approximately $\mathcal{O}(N^{-1/4})$. The isotropic adaptive strategy leads to an improved order of approximately $\mathcal{O}(N^{-1/2})$. This, as expected, is not optimal since isotropic mesh-refinement does not resolve the generic edge-singularities efficiently [14]. For this fact, the reader is also referred to some heuristics from [8]. As in the previous example, both adaptive and uniform mesh-refinement suggest reliability of η_ℓ . This gives empirical evidence for the saturation assumption (5), which is required in Theorem 7 to guarantee convergence of the adaptive scheme.

Acknowledgements Parts of the results have been achieved during a research stay of C.O. and D.P. at the Hausdorff Institute for Mathematics in Bonn, which is thankfully acknowledged. S.F. acknowledges a grant of the graduate school *Differential Equations – Models in Science and Engineering*, funded by the Austrian Science Fund (FWF) under grant W800-N05. D.P. is partially supported through the research project *Adaptive Boundary Element Method*, funded by the Austrian Science Fund (FWF) under grant P21732.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

1. Ainsworth, M., Oden, J.T.: *A posteriori error estimation in finite element analysis*. Wiley-Interscience [John Wiley & Sons], New-York (2000)
2. Bänsch, E.: Local mesh refinement in 2 and 3 dimensions. *IMPACT Comput. Sci. Eng.* **3**, 181–191 (1991)
3. Bank, R.: Hierarchical bases and the finite element method. *Acta Numerica* **5**, 1–45 (1996)
4. Bank, R., Smith, R.: A posteriori error-estimates based on hierarchical bases. *SIAM J. Numer. Anal.* **30**, 921–935 (1993)
5. Bank, R., Weiser, A.: Some a posteriori error estimators for elliptic partial differential equations. *Math. Comp.* **44**, 283–301 (1985)
6. Bornemann, F., Erdmann, B., Kornhuber, R.: A-posteriori error-estimates for elliptic problems in 2 and 3 space dimensions. *SIAM J. Numer. Anal.* **33**, 1188–1204 (1996)
7. Carstensen, C., Faermann, B.: Mathematical foundation of a posteriori error estimates and adaptive mesh-refining algorithms for boundary integral equations of the first kind. *Eng. Anal. Bound. Elem.* **25**, 497–509 (2001)
8. Carstensen, C., Maischak, M., Praetorius, D., Stephan, E.P.: Residual-based a posteriori error estimate for hypersingular equation on surfaces. *Numer. Math.* **97**(3), 397–425 (2004)
9. Carstensen, C., Praetorius, D.: Averaging techniques for the effective numerical solution of Symm’s integral equation of the first kind. *SIAM J. Sci. Comput.* **27**, 1226–1260 (2006)
10. Cascon, J., Kreuzer, C., Nochetto, R., Siebert, K.: Quasi-optimal convergence rate for an adaptive finite element method. *SIAM J. Numer. Anal.* **46**, 2524–2550 (2008)
11. Deuffhard, P., Leinen, P., Yserentant, H.: Concepts of an adaptive hierarchical finite element code. *IMPACT Comput. in. Sci. and Eng.* **1**, 3–35 (1989)
12. Dörfler, W.: A convergent adaptive algorithm for Poisson’s equation. *SIAM J. Numer. Anal.* **33**, 1106–1124 (1996)

13. Dörfler, W., Nochetto, R.: Small data oscillation implies the saturation assumption. *Numer. Math.* **91**, 1–12 (2002)
14. Ferraz-Leite, S., Praetorius, D.: Simple a posteriori error estimators for the h-version of the boundary element method. *Computing* **83**, 135–162 (2008)
15. Graham, I., Hackbusch, W., Sauter, S.: Finite elements on degenerate meshes: inverse-type inequalities and applications. *IMA J. Numer. Anal.* **25**, 379–407 (2005)
16. Hairer, E., Nørsett, S., Wanner, G.: Solving ordinary differential equations I. Nonstiff problems. Springer, New York (1987)
17. Kossaczky, I.: A recursive approach to local mesh refinement in two and three dimensions. *J. Comput. Appl. Math.* **55**, 275–288 (1995)
18. McLean, W.: Strongly elliptic systems and boundary integral equations. Cambridge University Press, Cambridge (2000)
19. Morin, P., Nochetto, R., Siebert, K.: Data oscillation and convergence of adaptive FEM. *SIAM J. Numer. Anal.* **38**, 466–488 (2000)
20. Sauter, S., Schwab, C.: Randelementmethoden: Analyse, Numerik und Implementierung schneller Algorithmen. Teubner Verlag, Wiesbaden (2004)
21. Sewell, E.: Automatic generation of triangulations for piecewise polynomial approximations. Ph.D. thesis, Purdue University, West Lafayette (1972)
22. Verfürth, R.: A posteriori error estimation and adaptive mesh refinement techniques. *J. Comput. Appl. Math.* **50**, 67–83 (1994)
23. Verfürth, R.: A review of a posteriori error estimation and adaptive mesh-refinement techniques. Teubner, Stuttgart (1996)