

Research article

Open Access

## Microarray genotyping of key experimental strains of *Neisseria gonorrhoeae* reveals gene complement diversity and five new neisserial genes associated with Minimal Mobile Elements.

Lori AS Snyder<sup>1</sup>, John K Davies<sup>2</sup> and Nigel J Saunders\*<sup>1</sup>

Address: <sup>1</sup>Bacterial Pathogenesis and Functional Genomics Group, Sir William Dunn School of Pathology, University of Oxford, South Parks Road, Oxford OX1 3RE, United Kingdom and <sup>2</sup>Bacterial Pathogenesis Research Group, Department of Microbiology, Monash University, VIC 3800, Australia

Email: Lori AS Snyder - [Lori.Snyder@pathology.oxford.ac.uk](mailto:Lori.Snyder@pathology.oxford.ac.uk); John K Davies - [John.Davies@med.monash.edu.au](mailto:John.Davies@med.monash.edu.au); Nigel J Saunders\* - [Nigel.Saunders@pathology.oxford.ac.uk](mailto:Nigel.Saunders@pathology.oxford.ac.uk)

\* Corresponding author

Published: 13 April 2004

Received: 10 November 2003

*BMC Genomics* 2004, **5**:23

Accepted: 13 April 2004

This article is available from: <http://www.biomedcentral.com/1471-2164/5/23>

© 2004 Snyder et al; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

### Abstract

**Background:** There are four widely used experimental strains of *N. gonorrhoeae*, one of which has been sequenced and used as the basis for the construction of a multi-strain, mutli-species pan-neisserial microarray. Although the *N. gonorrhoeae* population structure is thought to be less diverse than *N. meningitidis*, there are some recognized gene-complement differences between strains, including the 59 genes of the Gonococcal Genetic Island. In this study we have investigated the three experimental strains that have not been sequenced to determine the extent and nature of their similarities and differences.

**Results:** Using the Pan-*Neisseria* microarray, three commonly used gonococcal laboratory experimental strains were investigated (F62, MS11, & FA19). Genes absent from these strains, but present in strain FA1090, were assessed as is possible with typical microarrays. Due to the design of this microarray, additional genes were also identified. Differences were associated with Minimal Mobile Elements (MMEs) or known divergences. Genotyping indicates the presence of genes previously only described in meningococci and shows the presence of the complete Gonococcal Genetic Island in *N. gonorrhoeae* strain FA19. Five new neisserial genes were identified through microarray genotyping and subsequent sequencing of two divergent MMEs in *N. gonorrhoeae* strain MS11 and four MMEs in *N. gonorrhoeae* strain FA19. No differences were identified between *N. gonorrhoeae* strains FA1090 and F62, indicating that these strains are very similar.

**Conclusion:** This study shows extensive similarity between the experimental strains, associated with a varying number of strain-specific genes. This provides a framework for those working with these strains to refer to the available gonococcal genome sequence, and is the first detailed comparison of gene complements between gonococcal strains.

### Background

The *Neisseria gonorrhoeae* population structure is not clonal [1-5], its panmictic structure being the result of

horizontal genetic exchange [6]. The pathogenic *Neisseria* spp. are naturally transformable [7] and chromosomal changes are largely mediated by homologous

recombination. Natural competence for transformation and homologous recombination allow the generation of mosaic genes, such as the divergent *opa* genes [8,9], antigenic variation through recombination with silent cassettes, such as in the *pilE/pilS* system [10,11], and the horizontal exchange of whole genes or groups of genes in Minimal Mobile Elements (MMEs) [12]. A Minimal Mobile Element (MME) is defined as a region between two conserved genes within which different genes are found in different strains. These elements are described as 'minimal' reflecting the fact that they do not have features suggesting mobilization by associated transposases, nor do they currently appear to have features by which remotely encoded systems would facilitate mobilization. These were initially identified through comparisons of the available *Neisseria* spp. genome sequences, on which basis the MME*pheS-pheT* was the first to be studied in an extended set of strains [12]. In this region, flanked by the highly conserved *pheS* and *pheT* genes, eight different intergenic regions were found in the *Neisseria* strains that were investigated. Polymorphisms within the flanking gene sequences are indicative of the points of homologous recombination between the imported DNA and the native DNA. Further, assessments of five additional MMEs in the genome sequences show these to be present in different combinations in different strains, supporting the model that these are horizontally, rather than vertically, transferred elements [12].

*N. gonorrhoeae* may show less diversity than *Neisseria meningitidis* because it is more likely to be isolated away from other *Neisseria* spp. in its primary host niche [1]. The exchange of highly similar sequences between strains would prevent sequence analysis revealing the full extent of horizontal exchange. It is quite possible that recombination within genes coding for surface exposed proteins, or those under direct selection, may reveal a higher degree of recombinatorial diversity than is currently established from the predominantly housekeeping gene sequences investigated to date [10,13-15].

There are four commonly used laboratory strains of *N. gonorrhoeae*. The complete genome sequence is available for *N. gonorrhoeae* strain FA1090 [16], for which there is also a physical map [17,18]. This strain was originally recovered from a patient with disseminated gonococcal infection (DGI) in 1983 [19]. There is also a physical map of the *N. gonorrhoeae* strain MS11 chromosome [20]. It is from strain MS11 that the Gonococcal Genetic Island (GGI) has been sequenced [21]. The GGI is composed of 59 predicted coding regions, including genes believed to be involved in the secretion of DNA. This strain was isolated from a patient with uncomplicated gonorrhoea [22]. *N. gonorrhoeae* strain F62 was also isolated from a patient with uncomplicated gonorrhoea [23], but it does not

contain the GGI [21]. The last of the four commonly used laboratory strains of *N. gonorrhoeae* is strain FA19, which was isolated from a patient with both gonorrhoea and DGI. Additionally, it has caused DGI in a laboratory worker, following a mouth-pipetting accident [24]. While other strains are certainly used experimentally, these four represent those most commonly used and for which there is the greatest body of existing literature available.

Since these strains have been widely used in experimental studies it is important to be able to compare them, and to provide a means of relating each to the available genome sequence of *N. gonorrhoeae* strain FA1090. Some differences between these four commonly used laboratory strains, in addition to the presence/absence of the GGI [21], have been noted previously. The expression of ICAM-1 by human mucosal epithelial cells is upregulated in response to strain FA1090, but not to strain MS11 [25]. Truncations of the proteoglycan PG-1 showed different potencies against strains FA19 and F62 [26]. There are two immunological classes of the major outer membrane protein Protein I (encoded by *porB*), designated PIA and PIB [2]. Strains F62 and FA19 have PIA, while strains FA1090 and MS11 have PIB [2,27,28]. While this is not an exhaustive list, it does indicate that some difference in gene complement between these strains may be expected, and defining the presence and absence of gene-complement differences may provide keys for the community to address their significant functional differences.

The Pan-*Neisseria* microarray was constructed by an international consortium of neisseriologists (Saunders, Davies, et al – in preparation). The probes printed onto this microarray represent the coding sequences of three complete neisserial genomes (*N. gonorrhoeae* strain FA1090 [16], *N. meningitidis* strain MC58 [29], *N. meningitidis* strain Z2491 [30]) as well as the coding regions from the GGI [21]. For those coding regions of sufficient length, each probe is complementary to 150–450 bp of the most conserved regions of the CDS between the different genome sequences, that are also best able to discriminate between paralogous genes. Basing the design upon the most conserved regions between the different strains and species makes the array minimally strain specific facilitating the use of the Pan-*Neisseria* microarray in analyzing strains to which it was not designed.

This paper addresses the gene complement differences between the four most commonly used laboratory strains, utilizing the Pan-*Neisseria* microarray, comparative genomic analysis, and sequencing of the newly identified divergent MMEs revealed by the microarray data. This is the first use of a multi-species, multi-genome microarray to conduct gene complement studies. Additionally, this

**Table 1: Genomotyping results of *N. gonorrhoeae* strain FA1090 versus *N. gonorrhoeae* strain MS11.**

<b>Gene probes hybridizing to FA1090 but not MS11</b>	
<p><b>Genes known to be associated with divergence:</b>  XNG0049 (<i>pilC1</i>)  XNG0210 (hypothetical)  XNG0211 (hypothetical)  XNG0212 (hypothetical)  XNG0213 (hypothetical)  XNG1280 (hypothetical)  XNG0442 (hypothetical)  XNG1509 (hypothetical)</p> <p><b>Genes associated with strain-specific regions:</b>  XNG0080 (hypothetical)  XNG0081 (hypothetical)  XNG0528 (hypothetical)  XNG0529 (hypothetical)  XNG0669 (hypothetical)  XNG0670 (type I restriction enzyme M protein)  XNG0671 (<i>ribA-B</i>)  XNG0741 (hypothetical)  XNG1618 (hypothetical)</p>	<p><b>Lack of hybridization may be due to:</b>  divergence in MS11 <i>pilC</i> sequence (Z50180)  <i>maf</i>-associated divergence  <i>maf</i>-associated divergence  <i>maf</i>-associated divergence  <i>maf</i>-associated divergence  <i>maf</i>-associated divergence  prophage region-associated divergence  prophage region-associated divergence</p> <p><b>Associated strain-specific region:</b>  hypo-<i>pglB</i> MME  hypo-<i>pglB</i> MME  hypo-<i>aspA</i> MME  hypo-<i>aspA</i> MME  <i>ribAB</i>-hypo MME  <i>ribAB</i>-hypo MME  <i>ribAB</i>-hypo MME  <i>ung</i>-hypo MME  <i>nuoL-nuoM</i> MME</p>
<b>Gene probes hybridizing to MS11 but not FA1090</b>	
<p><b>Genes known to be associated with divergence</b>  NMA1797 (<i>tspB</i>)  NMB1119 (hypothetical)  NMB1120 (hypothetical)  GCJDS60 (suppressor)  GCJDS61 (regulator)</p> <p><b>Genes associated with strain-specific regions:</b>  59 Gonococcal Genetic Island genes</p>	<p><b>Hybridization to non-FA1090 probes may be due to:</b>  divergence in MS11 <i>tspB</i> sequence  prophage region-associated divergence  prophage region-associated divergence  prophage region-associated divergence  prophage region-associated divergence</p> <p><b>Associated strain-specific region:</b>  <i>ung</i>-hypo MME</p>

work supports and expands our observation of Minimal Mobile Elements in the *Neisseria* spp. [12].

## Results

### **Comparison of *N. gonorrhoeae* strains F62, MS11, and FA19 to the sequenced strain FA1090 using the Pan-*Neisseria* microarray**

Chromosomal DNA from *N. gonorrhoeae* strain FA1090 was fluorescently labelled with Cy3-dCTP and mixed with Cy5-dCTP labelled chromosomal DNA from either *N. gonorrhoeae* strain F62, MS11, or FA19, and hybridized to the Pan-*Neisseria* microarray slide. Analysis of labelled DNA hybridization indicated that there are gene complement differences between strains FA1090 and MS11 (Table 1), and strains FA1090 and FA19 (Table 2). No differences in gene complement were noted in microarray data comparison of *N. gonorrhoeae* strains FA1090 and F62. To confirm the original results for strains FA1090 and F62 these strains were re-sourced from an additional laboratory and the experiments repeated. The repeat experiment confirmed the absence of any differences.

### **Differences between *N. gonorrhoeae* strains FA1090 and MS11**

The results of hybridization with strains FA1090 and MS11 are shown in Table 1. One of the two strain MS11 *pilC* genes [31] is sufficiently divergent not to hybridize to any of the four *pilC* probes on the microarray (X0049, X1797, B0049, B1847). The T-cell stimulating antigen *tspB* [32] is present in multiple copies in the sequenced neisserial genomes [16,29,30], and the sequences are often divergent. Hybridization of strain MS11 with a meningococcal *tspB* probe (A1797) is a reflection of that divergence and possible exchange between the species. Additionally, probes to five *maf*-associated hypothetical genes hybridized to strain FA1090 labelled chromosomal DNA, but not to that from strain MS11. Through analysis of the complete neisserial genome sequences, it has been suggested that the region downstream of *mafB*, which is itself present in multiple copies, contains silent cassettes of *mafB* akin to the silent *pilS* cassettes that are capable of homologous recombination into the expression locus *pilE* [30]. The regions containing the *mafA* and *mafB* genes and


**Table 2: Genomotyping results of *N. gonorrhoeae* strain FA1090 versus *N. gonorrhoeae* strain FA19.**

<b>Gene probes hybridizing to FA1090 but not FA19</b>	
<p><b>Genes known to be associated with divergence:</b>  XNG0049 (<i>pilC1</i>)  XNG0211 (hypothetical)  XNG0212 (hypothetical)  XNG0213 (hypothetical)  XNG1848 (hypothetical)  XNG0458 (hypothetical)  XNG0459 (regulator)  XNG0460 (hypothetical)  XNG0461 (regulator)  XNG0462 (hypothetical)  XNG1006 (hypothetical)  XNG1509 (hypothetical)</p> <p><b>Genes associated with strain-specific regions:</b>  XNG0438 (hypothetical)  XNG0741 (hypothetical)  XNG0528 (hypothetical)  XNG0529 (hypothetical)  XNG1114 (hypothetical)  XNG1115 (methylase)  XNG1441 (hypothetical)</p>	<p><b>Lack of hybridization may be due to:</b>  divergence in FA19 <i>pilC</i> sequence  <i>maf</i>-associated divergence  <i>maf</i>-associated divergence  <i>maf</i>-associated divergence  <i>maf</i>-associated divergence  prophage region-associated divergence  prophage region-associated divergence  prophage region-associated divergence  prophage region-associated divergence  prophage region-associated divergence  prophage region-associated divergence  prophage region-associated divergence</p> <p><b>Associated strain-specific region:</b>  pilin-AzIC MME  ung-hypo MME  hypo-aspA MME  hypo-aspA MME  uvrA-hypo MME  uvrA-hypo MME  nadC-xthA MME</p>
<b>Gene probes hybridizing to FA19 but not FA1090</b>	
<p><b>Genes known to be associated with divergence</b>  NMB1541 (<i>lbpB</i>)  NMA0036 (hypothetical)  NMB2109 (hypothetical)  NMB2110 (hypothetical)  NMA1360 (hypothetical)  NMB1119 (hypothetical)  NMB1120 (hypothetical)  GCJDS60 (suppressor)  GCJDS61 (regulator)</p> <p><b>Genes associated with strain-specific regions:</b>  NMA2121 (hypothetical)  59 Gonococcal Genetic Island genes</p>	<p><b>Hybridization to non-FA1090 probes may be due to:</b>  <i>lbpB</i> is known to be dead in FA1090 (AF072890)  <i>maf</i>-associated divergence  <i>maf</i>-associated divergence  <i>maf</i>-associated divergence  the presence of a sequenced gene fragment in FA19  prophage region-associated divergence  prophage region-associated divergence  prophage region-associated divergence  prophage region-associated divergence</p> <p><b>Associated strain-specific region:</b>  hypo-hypo MME  ung-hypo MME</p>

the proposed silent *mafB* cassettes also contain many hypothetical genes. These features of the *maf* regions, present in the neisserial genomes in up to four locations [16], are not conserved, and are composed largely of short hypothetical genes of low %G+C separating what are believed to be the silent *mafB* recombination exchange cassettes [30]. It is these *maf*-associated low %G+C potential short coding sequences that appear to be divergent between the gonococcal strains, which is also the case between the four sequenced genome strains. Also present in strain FA1090, but absent from strain MS11, are two prophage-associated hypothetical genes. Additionally, there are four prophage-associated genes present in strain MS11, which are absent from strain FA1090. As prophages and prophage-associated genes are horizontally trans-

ferred and as each of the four neisserial genome sequences has quite different prophage and prophage-associated gene complements, it is not unexpected that differences in hybridization to these sequences would be found between the gonococcal strains.


All of the other differences between *N. gonorrhoeae* strain FA1090 and *N. gonorrhoeae* strain MS11 are associated with MMEs [12], the sites of which could be determined by comparison of the three complete neisserial genome sequences [16,29,30]. One of these MME regions is the GGI (Figure 1), which has been sequenced previously from strain MS11 [21]. The four remaining MMEs showing differences in the presence of strain FA1090 genes



**Figure 1**  
**The MME flanked by *ung* and a hypothetical protein.**  
 This MME contains the 59 predicted coding regions of the Gonococcal Genetic Island (GGI) in *N. gonorrhoeae* strains MS11 and FA19, one hypothetical gene in *N. gonorrhoeae* strain FA1090 [16], a putative methylase and a hypothetical gene in *N. meningitidis* strains MC58 and Z2491 [29,30], and two hypothetical genes, *piv*, and a putative regulator in *N. meningitidis* strain FAM18. Similar genes are indicated in the same colour.

were investigated further through PCR amplification and sequencing.

The MME flanked by *nuoL* and *nuoM* is shown in Figure 2, from which four newly described neisserial genes were sequenced from strain MS11 (Accession AY386266). These lack any significant similarity to sequences in the public databases. Although these genes are not represented on the Pan-*Neisseria* microarray, it was possible using the microarray data to identify the *nuoL-nuoM* region as a target for sequencing due to the lack of hybridization of the labelled strain MS11 DNA to the strain




**Figure 2**  
**The MME flanked by *nuoL* and *nuoM*.** This MME contains a hypothetical gene in *N. gonorrhoeae* strain FA1090 [16], short intergenic sequences in *N. meningitidis* strains MC58 [29] and FAM18, two hypothetical genes in *N. meningitidis* strain Z2491 [30], and four newly described neisserial genes of unknown function in *N. gonorrhoeae* strain MS11 (AY386266).

FA1090 probe for XNG1618 or to the alternate *N. meningitidis* strain Z2491 probes for NMA0001 and NMA2230.

The MME flanked by a putative lipopolysaccharide biosynthesis protein and *pglB* is shown in Figure 3. In *N. gonorrhoeae* strain FA1090 and *N. meningitidis* strains Z2491 and FAM18, there are two hypothetical genes (*NMA0640* & *NMA0641*) in the intervening region. In *N. gonorrhoeae* strain MS11, however, there is a short (167 bp) intergenic region (Accession AY386267), similar to that found in *N. meningitidis* strain MC58.

The remaining two MMEs, identified through absence of hybridization of microarray probes with gonococcal strain MS11, failed to amplify either from strain MS11 or from control strain FA1090. The first is flanked by a hypothetical gene and *aspA*, containing three hypothetical genes in *N. gonorrhoeae* strain FA1090, spanning a region of over 7 kb. Of these three genes, two do not hybridize with the FA1090-derived probes, and appear to be absent in strain MS11. The second of these remaining MMEs is flanked by a hypothetical gene and *ribA-B* and in *N. gonorrhoeae* strain FA1090 this region, of approximately 11.5 kb, contains genes encoding a restriction-modification system. The size of these MMEs in strain FA1090 (7 & 11.5 kb), may account for the failure to amplify these regions.



**Figure 3**  
**The MME flanked by a putative LPS biosynthesis gene and *pgIB*.** This MME contains two hypothetical genes in *N. gonorrhoeae* strain FA1090 [16], *N. meningitidis* strain Z2491 [30], and *N. meningitidis* strain FAM18; and a short intergenic region in *N. meningitidis* strain MC58 [29] and *N. gonorrhoeae* strain MS11 (AY386267). Similar genes are indicated in the same colour.

#### Differences between *N. gonorrhoeae* strains FA1090 and FA19

As with gonococcal strain MS11, divergence is indicated by the genotyping results for the strain FA19 *pilC* gene, *maf*-associated hypothetical genes, and prophage-associated genes (Table 2). Additional points of variation between *N. gonorrhoeae* strain FA1090 and strain FA19 have been confirmed, by sequencing, to be due to hybridization to sequences not present in strain FA1090.

#### Cross hybridizations to divergent sequences revealed through sequencing

Probe A0036 was designed to *NMA0036*, which is within an MME. In *N. gonorrhoeae* strain FA1090 *XNG1638* is in this MME location, probed by X1638. Strain FA19 genomic DNA hybridized both with probe A0036 and X1638, suggesting that both genes, which were believed to be mutually exclusive in this MME, are present in strain FA19. This MME was therefore amplified and sequenced from strain FA19 (data not presented). This sequence shows that the MME region is similar to that present in strain FA1090 and did not contain the MME-associated gene *NMA0036*, therefore the hybridization of strain FA19 to probe A0036 must be to a sequence outside of this MME. The probe A0036 cross-hybridizes in *N. meningitidis* strain MC58 at 86% over 132 bp with *NMB0656*, which is a *maf*-associated hypothetical gene. In *N. gonor-*

*rhoeae* strain FA19, therefore, this probe may be hybridizing to a *maf*-associated region, rather than to the MME as initially suspected.

A second probe designed against *N. meningitidis* strain Z2491, A1360, was found to hybridize the FA19 labelled genomic DNA. It was known that this probe cross-hybridizes slightly to fragments of the meningococcal gene in strain FA1090 (28 bp at 96% and 37 bp at 89%), but not at a level that would be expected to be detected on the Pan-*Neisseria* microarray and which did not, in fact, generate a microarray hybridization signal. Sequencing of this region from strain FA19 revealed that an additional 106 bp is present in this strain (data not presented), generating a region in strain FA19 that is 91% similar to the 181 bp length of the A1360 probe. Although this sequence homology is to a longer sequence than is in strain FA1090, it is still a fragment and not the complete meningococcal gene in strain FA19. Therefore, this result is due to sequence diversity and cross-hybridization of the probe to a gene fragment, rather than the presence of an entire gene.

#### Presence of the Gonococcal Genetic Island in strain FA19


Labelled chromosomal DNA from *N. gonorrhoeae* strain FA19 hybridized to all 59 probes for the GGI, indicating for the first time, the presence of the complete island in this strain (Figure 1).

#### MME-associated differences between strains FA1090 and FA19


The MME-associated region detected in strain MS11, flanked by a hypothetical gene and *aspA*, was also detected in strain FA19, although PCR amplification of this region from strain FA19 failed as it did with strains FA1090 and MS11 (see above). The remaining four strain FA19 MMEs, indicated to be divergent through genotyping, revealed additional differences between the gonococcal laboratory strains (Figures 4, 5, 6, 7), including a large new neisserial gene (Figure 6).

Hypothetical gene *XNG1441* is not present in strain FA19. This gene is located between *nadC* and *xthA* in strain FA1090, where it disrupts a hypothetical gene identified in *N. meningitidis* strain MC58, *NMB0397* (Figure 4). This meningococcal hypothetical gene is disrupted by different means in all other strains from which this region has been sequenced, including in *N. gonorrhoeae* strain FA19 (Accession AY386268). Therefore the absence of *XNG1441* in strain FA19 is confirmed by sequencing and an additional means of disrupting *NMB0397* has been identified.

Hypothetical gene *XNG0438* is also missing from strain FA19. In this case the MME containing this gene has conserved features in the meningococcal strains, in the form



**Figure 4**  
**The MME flanked by *nadC* and *xthA*.** This MME contains a hypothetical gene (*NMB0397*) in *N. meningitidis* strain MC58 [29]. This gene is disrupted in a different way in all other strains from which it has been sequenced. In *N. meningitidis* strain Z2491 [30], it is interrupted by a CREE, which also disrupts the next gene, in *N. meningitidis* strain FAM18 it is disrupted by a 61 bp indel that causes a frame-shift, in *N. gonorrhoeae* strain FA1090 [16] it is disrupted by a hypothetical gene, and in *N. gonorrhoeae* strain FA19 (AY386268) it is disrupted by a different CREE from that in this location in *N. meningitidis* strain Z2491. Similar genes or fragments thereof are indicated in the same colour.




**Figure 5**  
**The MME flanked by a pilin-related gene and an *AzIC*-related gene.** This MME contains a hypothetical gene in *N. gonorrhoeae* strain FA1090 [16], a compound CREE structure in *N. meningitidis* strains MC58 [29], Z2491 [30], and FAM18, in which two nearly identical CREE flank a variable intergenic region composed largely of tandem repeats and which disrupts the 3' end of the *AzIC*-related gene, and a single CREE in *N. gonorrhoeae* strain FA19 (AY386269), where the 3' end of the *AzIC*-related gene is disrupted by other intergenic sequence.


of two nearly identical Correia Repeat Enclosed Elements (CREE) [33-35] flanking a variable length intergenic region composed largely of tandem repeats (Figure 5). The Correia Repeat [33,34] is an intergenic sequence element commonly found as an inverted repeat with a characteristic core region, most often being 105 or 153 bp in length and present in over 100 copies in the genome [35]. Although it has not been determined how the CREE is mobilized, it is present in different locations between the genomes and the target site is TA, which is copied upon insertion [35]. The insertion of a CREE can generate a promoter at either its the 5' [36] or 3' end [37] and can disrupt the sequence into which it has been inserted. In this case the CREE disrupts the 3' end of the *AzIC*-related gene that flanks the MME, generating a termination codon for this coding region within one of the CREE. Sequencing from strain FA19 revealed only one CREE in this region, although the *AzIC*-related gene was still disrupted at the 3' end by divergent sequence (Accession AY386269). Therefore the absence of *XNG0438* in strain FA19 has been confirmed by sequencing and an additional variation in the intergenic region of this MME has been identified.

In *N. gonorrhoeae* strain FA1090, the MME flanked by *uvrA* and a hypothetical gene contains another hypothetical gene and a cytosine-specific methylase gene (Figure 6). In *N. meningitidis* strains Z2491 and FAM18, there are three and two hypothetical genes in this location, respectively. The region 3' of *uvrA* is also the site of chromosomal rearrangement in *N. meningitidis* strain MC58, relative to the other strains. DNA from strain FA19 does not hybridize to any of these genes within the *uvrA*-associated MME. PCR and sequencing of this region from *N. gonorrhoeae* strain FA19 revealed one new coding sequence within this MME (Figure 6; Accession AY386270). This sequence has no homology to complete coding sequences within the public databases, although the first 137 bp (100% DNA identity) of this CDS, 3' of *uvrA*, is included as unannotated sequence at the end of the GenBank entry for *uvrA* from library clone pJKD909, derived from *N. gonorrhoeae* (U34760).

The hybridization of the probe for *N. meningitidis* strain Z2491 CDS *NMA2121*, suggested that this gene is present in strain FA19 in an MME flanked by two hypothetical genes (Figure 7). Annotated as a hypothetical protein, this CDS is not present in the other two neisserial genome



**Figure 6**  
**The MME flanked by *uvrA* and a hypothetical gene.**  
 This MME contains a gene encoding a putative methylase and a hypothetical gene in *N. gonorrhoeae* strain FA1090 [16], three hypothetical genes in *N. meningitidis* strain Z2491 [30], two hypothetical genes in *N. meningitidis* strain FAM18, and one new neisserial gene of unknown function in *N. gonorrhoeae* strain FA19 (AY386270). This is the point of a chromosomal rearrangement in *N. meningitidis* strain MC58 [29].



**Figure 7**  
**The MME flanked by two hypothetical genes.** This MME contains an intergenic sequence in *N. gonorrhoeae* strain FA1090 [16], and a hypothetical gene in *N. meningitidis* strain Z2491 [30]. In *N. gonorrhoeae* strain FA19 (AY386271) there is a homologue of the meningococcal CDS, which has been disrupted by a CREE 44 bp into the gene. The CREE in this location introduces a termination codon and generates a frame-shift. However, it also generates a potential promoter 5' of a potential ribosomal binding site and secondary initiation codon internal to *NMA2121*. This is the point of a chromosomal rearrangement in *N. meningitidis* strains MC58 [29] and FAM18. Similar genes are indicated in the same colour.

sequences, nor in the recently completed *N. meningitidis* strain FAM18 genome sequence. This MME was sequenced from *N. gonorrhoeae* strain FA19 to determine if a complete version of *NMA2121* was present in this location (Accession AY386271). Sequencing demonstrated that although this gonococcal strain does contain a CDS with homology to the sequence of the meningococcal CDS *NMA2121*, it is disrupted. A 107 bp CREE is present 44 bp into the CDS, which introduces a termination codon shortly thereafter and generates a frame-shift (Figure 7). The presence of this CREE also generates a potential promoter at the 3' end of the element, similar to those described previously and identified 5' of *uvrB* [37]. This, coupled with a potential ribosome binding site and secondary translational initiation codon, suggests that the *NMA2121* homologue may be expressed in *N. gonorrhoeae* strain FA19, despite the apparent interruption of the gene by a CREE.

## Discussion

The Pan-*Neisseria* microarray, coupled with MME analysis of sites of divergence indicated by the microarray analysis, has revealed five new neisserial genes not represented on the microarray, as well as identifying the 59 genes of the GGI in *N. gonorrhoeae* strain FA19. There are also 36 genes that are present within strain FA1090 that are not universally represented in the other main experimental strains, 16 of which are located within MMEs that are identifiable from comparison of the complete genome sequences of gonococcal strain FA1090 and the available complete meningococcal genome sequences. The ability to identify this range of differences is in part due to the unique nature of the Pan-*Neisseria* microarray (Saunders, Davies, et al. – in preparation). Its design incorporates probes to the most conserved regions of genes, having been evaluated through the comparison of *N. gonorrhoeae* strain FA1090 sequences [16] with those of *N. meningitidis* strains MC58 [29] and Z2491 [30]. Genes from the two meningococcal genomes that were not optimally probed by the gonococcal probes ( $\geq 90\%$  homology over 150 bp) or where not represented by the gonococcal probes, had additional probes added to the microarray. The 59 predicted coding sequences of the Gonococcal Genetic Island, as sequenced from *N. gonorrhoeae* strain MS11 [21], are also represented on the microarray. This microarray composition meant that for the three gonococcal laboratory strains evaluated here, it was possible to determine, solely from the microarray results, that 36 strain FA1090 genes are absent from the other gonococcal strains, while 130 genes are present in these strains that are absent from strain FA1090. This second group of genes could not have been identified with a single-genome microarray based upon strain FA1090 alone, emphasizing the strength of this multi-species, multi-genome design strategy. The microarray results additionally indicated regions of the chromosomes likely



to contain divergent sequence, analysis of which revealed five new genes.

In strain MS11, 17 probes designed to strain FA1090 genes did not hybridize, including nine genes associated with five MMEs. Sequencing of two of these MMEs confirmed the absence of the strain FA1090 genes and identified four new genes in strain MS11. One of the five MMEs is the Gonococcal Genetic Island, the sequence of which is known from strain MS11. The remaining two MMEs (*hypo-aspA* and *ribAB-hypo*) may simply be too large to amplify using the techniques described here, but may therefore represent the location of additional genes present in strain MS11 and absent in FA1090, and vice versa. There are 19 strain FA1090 probes that do not hybridize to strain FA19 DNA, with seven of these being contained in five MMEs, one of which is the Gonococcal Genetic Island. Sequencing of three of the MMEs confirmed the absence of the strain FA1090 genes and identified one new neisserial gene in strain FA19. The remaining MME (*hypo-aspA*) was also identified in strain MS11. No differences were observed between *N. gonorrhoeae* strains FA1090 and F62. Discounting those genes that are likely to display differences due to divergence (*pilC*, *lbpB*, *tspB*, *maf*-associated, prophage-associated, and the *NMA1360* gene fragment) a total of 80 genes were found to contribute to inter-strain gene-complement differences, of which 59 comprise the Gonococcal Genetic Island and one represents the hypothetical strain FA1090 gene that is present in its place (*XNG0741*; Figure 1). This does not take into account those genes that may be found in strains MS11 and FA19 in the large *hypo-aspA* MME and in strain MS11 in the large *ribAB-hypo* MME.

Although this approach to strain comparison, utilizing a multi-species, multi-genome microarray is novel, powerful, and has identified differences between these strains, there are some additional forms of divergence that this approach cannot address. This means that although we have an indication of the significant similarity of strains FA1090 and F62, which will facilitate the use of the available genome sequence in this strain, they cannot be assumed to be identical. There may be other differences between these and the other strains such as: (a) the presence of additional genes that are not currently on the microarray, although these cannot be associated with the absence of MME-associated genes as they are in strains MS11 and FA19; (b) divergence outside of the probe regions used for construction of the microarray, which were selected due to their maximal conservation across the species ( $\geq 90\%$  ID over a probe of 150–450 bp), therefore the divergence in the *fbp* iron acquisition genes (strain F62 sequence: U33937) is not detected because the probes hybridize to the conserved regions of the genes at  $\geq 98\%$  over the length of the probe; (c) differences in tran-

scriptional regulation, post-transcriptional regulation, and phase variation, for example as in *pgtA*, which is phase variable in strain FA1090 but not in strain F62 where the gene lacks the phase variation-mediating poly-G tract [38]. Despite these caveats this study clearly shows some of the significant advantages and power of this microarray design approach in its application to genotyping, and provides a new basis upon which to inter-relate the data obtained from the many studies performed using these strains.

## Conclusions

The multi-strain, multi-species microarray design strategy can be used to identify the presence and absence of genes in a more informative fashion than arrays based upon single genomes, or even single species. This approach can be used to identify those genes that are trans-species, rather than species associated, as demonstrated from limited complete genome sequenced examples. This approach is also able to identify chromosomal discontinuities which when coupled to MME analysis can identify new sequences and genes not present in any of the sequences used for microarray design. This study identified many differences, which are likely to account for at least some of their recorded differences in behaviour between the gonococcal experimental strains. This included the identification of five new neisserial genes. The recently completed *N. meningitidis* serogroup C strain FAM18 complete genome sequence <ftp://ftp.sanger.ac.uk/pub/pathogens/nm/> identified 35 new neisserial genes, other than those associated with a large prophage region. This array-based approach is therefore a comparatively cheap and efficient way of finding new genes within a species for which existing genomes are available. The range of differences that have been identified in this study emphasize the utility of a pan-species microarray, with the design characteristics used, in the investigation of diverse bacterial populations.

## Methods

### Bacterial strains and growth conditions

*N. gonorrhoeae* strains FA1090, F62, MS11, and FA19, were propagated on GC agar (Difco Laboratories) containing the Kellogg supplement and ferric nitrate [39] at 37°C under 5% (v/v) CO<sub>2</sub>. These strains were kindly supplied by Prof. William M. Shafer (FA1090, F62, FA19) and Prof. Richard F. Rest (FA1090, F62, MS11).

### DNA extraction, labelling, and microarray hybridization

Chromosomal DNA was extracted using the AquaPure Genomic DNA Isolation Kit (BIORAD). FluoroLink™ Cy3-dCTP and FluoroLink™ Cy5-dCTP (Amersham Pharmacia Biotech) were incorporated into 10 µg of chromosomal DNA using random hexamer primers (Invitrogen) and DNA polymerase I, Klenow fragment (Bioline, UK). Labelled DNA:DNA probe microarray hybridizations

were conducted in  $4 \times$  SSC, 0.2875% SDS at  $65^\circ\text{C}$  overnight. The pan-*Neisseria* microarray was designed, generated and printed as will be described fully elsewhere (Saunders, Davies, et al. – in preparation). This microarray design contains 2704 probes directed to the most conserved regions of the 6294 annotated potential coding sequences from *N. gonorrhoeae* strain FA1090 [16], *N. meningitidis* strains MC58 [29] and Z2491 [30], and additional sequence from *N. gonorrhoeae* strain MS11 including the Gonococcal Genetic Island [21]. Each probe of 150–450 bp was selected to target within the most conserved ( $\geq 90\%$  identical) sequence region of each coding sequence between the genomes against which it was designed. In this way, the pan-*Neisseria* microarray achieves maximal hybridization to orthologous genes in different strains and species. Each probe was also assessed for cross-hybridization within the same genome. If any significant cross-hybridization was detected, being assessed through sequence homology of 70% over 40 bp, the probe was redesigned to exclude the potential for cross-hybridization with other genes. In this way, the pan-*Neisseria* microarray eliminates false positive results due to the hybridization to non-representative probes. As an example of the design detail and probe hybridization fidelity, the *pilC* gene from *N. gonorrhoeae* strain FA1090 XNG0049 does not hybridize to either of the *pilC* probes designed against *N. meningitidis* strain MC58 *pilC* genes NMB0049 and NMB1847. While the gonococcal gene is 78.8% (over 3300 bp) and 82.3% (over 3293) identical, with respect to each of the meningococcal genes, the probes for the meningococcal genes fall below the 70% over 40 bp cross-hybridization design threshold and no signal from these probes is detected in control experiments when the microarray is hybridized with *N. gonorrhoeae* strain FA1090 DNA. Microarray slides were scanned on an Axon GenePix 4000 B scanner using GenePix Pro v 3.0.6.90 and analyzed using GACK [40] downloaded from <http://falkow.stanford.edu/whatwedo/software/>.

#### PCR amplification and sequencing

PCR amplification from chromosomal DNA was performed using BIO-X-ACT™ Long DNA Polymerase (BIO-LINE) or *Taq* DNA polymerase (Invitrogen) according to the manufacturers' instructions. Primers used are listed in additional file 1 Table S1. PCR amplification of two regions, *hypo-aspA* and *ribAB-hypo*, failed even after alternate primers were designed. Automated sequencing used ABI Prism® BigDye™ Terminator Cycle Sequencing version 3.0 (Applied Biosystems) and was resolved on an ABI Prism® 3100 DNA Sequencer (Applied Biosystems).

#### Bioinformatics

ACEDB [41] was used to analyze the complete genome sequences of the *Neisseria* spp., as described previously

[42]. MMEs were identified through comparison of the available genomes in the regions of the chromosome indicated as divergent from the microarray data. Sequence traces were analyzed using Trev [43]. Sequences were aligned and assembled using the Wisconsin Package from GCG (Accelrys). Homology searches were performed using BLAST [44] against the EMBL databases, accessed through the Oxford University Bioinformatics Centre.

#### Accession numbers for sequence data

The sequence of the MME region containing four new gonococcal genes between *nuoL* and *nuoM* from *N. gonorrhoeae* strain MS11, accession number AY386266, and that containing one new gonococcal gene adjacent to *uvrA* from *N. gonorrhoeae* strain FA19, accession number AY386270, have been submitted to GenBank. Additional MME region sequences submitted to GenBank are: that adjacent to *pglB* from *N. gonorrhoeae* strain MS11 (AY386267); between *nadC* and *xthA* from *N. gonorrhoeae* strain FA19 (AY386268); between a type IV pilin-related genes and an AzlC-related protein gene from *N. gonorrhoeae* strain FA19 (AY386269); and between two hypothetical protein genes from *N. gonorrhoeae* strain FA19 (AY386271).

#### Authors' contributions

LS is responsible for all experimental manipulations, analysis of microarray and sequencing data, identification of MMEs, writing of the manuscript, and assisted in probe design for the pan-*Neisseria* microarray. JD is responsible for the production of the pan-*Neisseria* microarray slides used in this study and provided support in the preparation of the manuscript. NS conceived of the pan-*Neisseria* microarray and its use in this kind of study, was the primary designer of the probes for the microarray, supervised this work, and provided support in the preparation of this manuscript.

#### Additional material

##### Additional File 1

Table S1. Oligonucleotide primers used in this study. A Table of the oligonucleotide primers used in this study for the amplification and sequencing of the MMEs presented from *N. gonorrhoeae* strain MS11 and *N. gonorrhoeae* strain FA19.

Click here for file

[<http://www.biomedcentral.com/content/supplementary/1471-2164-5-23-S1.pdf>]

#### Acknowledgements

The authors would like to thank Dr. Sarah Butcher for continued bioinformatics support and Julian Robinson from the Sir William Dunn School of Pathology sequencing service. NJS was supported by a Wellcome Advanced Research Fellowship. LASS is supported by a Wellcome Trust Project

Grant. JKD is the recipient of a Project Grant from the Australian National Health and Medical Research Council. The meningococcal genome sequence data were produced by the *N. meningitidis* serogroup C strain FAM18 Sequencing Group at the Wellcome Trust Sanger Institute, and can be obtained from <ftp://ftp.sanger.ac.uk/pub/pathogens/nm/>. The *N. gonorrhoeae* genome sequence was obtained from the University of Oklahoma, <http://www.genome.ou.edu/gono.html>, the Gonococcal Genome Sequencing Project which was supported by USPHS/NIH grant #AI-38399, GenBank accession number AE004969 [16].

## References

- Spratt BG, Smith NH, Zhou J, O'Rourke M, Feil E: **The population genetics of the pathogenic *Neisseria***. *Society for General Microbiology Volume 52*. Edited by: Baumberg S, Young JPW, Wellington EMH and Saunders JR. SGM Press; 1995:143-160.
- Carbonetti NH, Simnad VI, Seifert HS, So M, Sparling PF: **Genetics of protein I of *Neisseria gonorrhoeae*: construction of hybrid porins**. *Proc Natl Acad Sci U S A* 1988, **85**:6841-6845.
- O'Rourke M, Stevens E: **Genetic structure of *Neisseria gonorrhoeae* populations: a non-clonal pathogen**. *J Gen Microbiol* 1993, **139 (Pt 11)**:2603-2611.
- O'Rourke M, Spratt BG: **Further evidence for the non-clonal population structure of *Neisseria gonorrhoeae*: extensive genetic diversity within isolates of the same electrophoretic type**. *Microbiology* 1994, **140 (Pt 6)**:1285-1290.
- De La Fuente L, Vazquez JA: **Genetic structures of non-penicillinase-producing *Neisseria gonorrhoeae* strains in relation to auxotype and serovar class**. *J Infect Dis* 1994, **170**:696-700.
- Saunders NJ, Hood DW, Moxon ER: **Bacterial evolution: bacteria play pass the gene**. *Curr Biol* 1999, **9**:R180-3.
- Biswas GD, Thompson SA, Sparling PF: **Gene transfer in *Neisseria gonorrhoeae***. *Clin Microbiol Rev* 1989, **2 Suppl**:S24-8.
- Aho EL, Dempsey JA, Hobbs MM, Klapper DG, Cannon JG: **Characterization of the opa (class 5) gene family of *Neisseria meningitidis***. *Molecular Microbiology* 1991, **5**:1429-1437.
- Hobbs MM, Seiler A, Achtman M, Cannon JG: **Microevolution within a clonal population of pathogenic bacteria: recombination, gene duplication and horizontal genetic exchange in the opa gene family of *Neisseria meningitidis***. *Mol Microbiol* 1994, **12**:171-180.
- Seifert HS, Ajioka RS, Marchal C, Sparling PF, So M: **DNA transformation leads to pilin antigenic variation in *Neisseria gonorrhoeae***. *Nature* 1988, **336**:392-395.
- Mehr IJ, Seifert HS: **Differential roles of homologous recombination pathways in *Neisseria gonorrhoeae* pilin antigenic variation, DNA transformation and DNA repair**. *Mol Microbiol* 1998, **30**:697-710.
- Saunders NJ, Snyder LAS: **The minimal mobile element**. *Microbiology* 2002, **148**:3756-3760.
- Halter R, Pohlner J, Meyer TF: **IgA protease of *Neisseria gonorrhoeae*: isolation and characterization of the gene and its extracellular product**. *EMBO J* 1984, **3**:1595-1601.
- Feavers IM, Heath AB, Bygraves JA, Maiden MC: **Role of horizontal genetic exchange in the antigenic variation of the class I outer membrane protein of *Neisseria meningitidis***. *Mol Microbiol* 1992, **6**:489-495.
- Connell TD, Black VJ, Kawula TH, Barritt DS, Dempsey JA, Kverneland K, Stephenson A, Schepart BS, Murphy GL, Cannon JG: **Recombination among protein II genes of *Neisseria gonorrhoeae* generates new coding sequences and increases structural variability in the protein II family**. *Mol Microbiol* 1988, **2**:227-236.
- Lewis LA, Gillaspay AF, McLaughlin RE, Gipson M, Ducey T, Ownbey T, Hartman K, Nydick C, Carson M, Vaughn J, Thomson C, Song L, Lin S, Yuan X, Najjar F, Zhan M, Ren Q, Zhu H, Qi S, Kenton SM, Lai H, White JD, Clifton S, Roe BA, Dyer DW: **The gonococcal genome sequencing project**. *unpublished*.
- Dempsey JA, Cannon JG: **Locations of genetic markers on the physical map of the chromosome of *Neisseria gonorrhoeae* FA1090**. *J Bacteriol* 1994, **176**:2055-2060.
- Dempsey JA, Litaker W, Madhure A, Snodgrass TL, Cannon JG: **Physical map of the chromosome of *Neisseria gonorrhoeae* FA1090 with locations of genetic markers, including opa and pil genes**. *J Bacteriol* 1991, **173**:5476-5486.
- Nachamkin I, Cannon JG, Mittler RS: **Monoclonal antibodies against *Neisseria gonorrhoeae*: production of antibodies directed against a strain-specific cell surface antigen**. *Infect Immun* 1981, **32**:641-648.
- Bihimaier A, Romling U, Meyer TF, Tummeler B, Gibbs CP: **Physical and genetic map of the *Neisseria gonorrhoeae* strain MS11-N198 chromosome**. *Mol Microbiol* 1991, **5**:2529-2539.
- Dillard JP, Seifert HS: **A variable genetic island specific for *Neisseria gonorrhoeae* is involved in providing DNA for natural transformation and is found more often in disseminated infection isolates**. *Mol Microbiol* 2001, **41**:263-277.
- Swanson J: **Studies on gonococcus infection. IV. Pili: their role in attachment of gonococci to tissue culture cells**. *J Exp Med* 1973, **137**:571-589.
- La Scolea LJ, Dul MJ, Young FE: **Stability of pathogenic colony types of *Neisseria gonorrhoeae* in liquid culture by using the parameters of colonial morphology and deoxyribonucleic acid transformation**. *J Clin Microbiol* 1975, **1**:165-170.
- Daly JA, Lee TJ, Spitznagel JK, Sparling PF: **Gonococci with mutations to low-level penicillin resistance exhibit increased sensitivity to the oxygen-independent bactericidal activity of human polymorphonuclear leukocyte granule extracts**. *Infect Immun* 1982, **35**:826-833.
- Jarvis GA, Li J, Swanson KV: **Invasion of human mucosal epithelial cells by *Neisseria gonorrhoeae* upregulates expression of intercellular adhesion molecule 1 (ICAM-1)**. *Infect Immun* 1999, **67**:1149-1156.
- Qu XD, Harwig SS, Shafer WM, Lehrer RI: **Protegrin structure and activity against *Neisseria gonorrhoeae***. *Infect Immun* 1997, **65**:636-639.
- Joiner KA, Puentes SM, Warren KA, Scales RA, Judd RC: **Complement binding on serum-sensitive and serum-resistant transformants of *Neisseria gonorrhoeae*: effect of presensitization with a non-bactericidal monoclonal antibody**. *Microb Pathog* 1989, **6**:343-350.
- Olesky M, Hobbs M, Nicholas RA: **Identification and analysis of amino acid mutations in porin IB that mediate intermediate-level resistance to penicillin and tetracycline in *Neisseria gonorrhoeae***. *Antimicrob Agents Chemother* 2002, **46**:2811-2820.
- Tettelin H, Saunders NJ, Heidelberg J, Jeffries AC, Nelson KE, Eisen JA, Ketchum KA, Hood DW, Peden JF, Dodson RJ, Nelson WC, Gwinn ML, DeBoy R, Peterson JD, Hickey EK, Haft DH, Salzberg SL, White O, Fleischmann RD, Dougherty BA, Mason T, Ciecko A, Parksey DS, Blair E, Cittone H, Clark EB, Cotton MD, Utterback TR, Khouri H, Qin H, Vamathevan J, Gill J, Scarlato V, Masignani V, Pizza M, Grandi G, Sun L, Smith HO, Fraser CM, Moxon ER, Rappuoli R, Venter JC: **Complete genome sequence of *Neisseria meningitidis* serogroup B strain MC58**. *Science* 2000, **287**:1809-1815.
- Parkhill J, Achtman M, James KD, Bentley SD, Churcher C, Klee SR, Morelli G, Basham D, Brown D, Chillingworth T, Davies RM, Davis P, Devlin K, Feltham T, Hamlin N, Holroyd S, Jagels K, Leather S, Moule S, Mungall K, Quail MA, Rajandream MA, Rutherford KM, Simmonds M, Skelton J, Whitehead S, Spratt BG, Barrell BG: **Complete DNA sequence of a serogroup A strain of *Neisseria meningitidis* Z2491**. *Nature* 2000, **404**:502-506.
- Jonsson AB, Rahman M, Normark S: **Pilus biogenesis gene, pilC, of *Neisseria gonorrhoeae*: pilC1 and pilC2 are each part of a larger duplication of the gonococcal genome and share upstream and downstream homologous sequences with opa and pil loci**. *Microbiology* 1995, **141 (Pt 10)**:2367-2377.
- Woolridge KG, Kizil G, Jones M, De-Netto LA, Todd I, Ala'Aldeen DAA: **Identification and analysis of TspB, a potent T-cell and B-cell stimulating *Neisseria*-specific antigen**. *Eleventh International Pathogenic *Neisseria* Conference* Edited by: Nassif X, Quentin-Millet M-J and Taha M-K. Nice, France, Editions E.D.K.; 1998:197.
- Correia FF, Inouye S, Inouye M: **A family of small repeated elements with some transposon-like properties in the genome of *Neisseria gonorrhoeae***. *J Biol Chem* 1988, **263**:12194-12198.
- Correia FF, Inouye S, Inouye M: **A 26-base-pair repetitive sequence specific for *Neisseria gonorrhoeae* and *Neisseria meningitidis* genomic DNA**. *J Bacteriol* 1986, **167**:1009-1015.
- Liu SV, Saunders NJ, Jeffries A, Rest RF: **Genome analysis and strain comparison of *Correia* Repeats and *Correia* Repeat-Enclosed Elements in pathogenic *Neisseria***. *J Bacteriol* 2002, **184**:6163-6173.

36. Snyder LAS, Shafer WM, Saunders NJ: **Divergence and transcriptional analysis of the division cell wall (dcw) gene cluster in *Neisseria* spp.** *Mol Microbiol* 2003, **47**:431-442.
37. Black CG, Fyfe JA, Davies JK: **A promoter associated with the neisserial repeat can be used to transcribe the *uvrB* gene from *Neisseria gonorrhoeae*.** *J Bacteriol* 1995, **177**:1952-1958.
38. Banerjee A, Wang R, Supernavage SL, Ghosh SK, Parker J, Ganesh NF, Wang PG, Gulati S, Rice PA: **Implications of phase variation of a gene (*pgtA*) encoding a pilin galactosyl transferase in gonococcal pathogenesis.** *J Exp Med* 2002, **196**:147-162.
39. Kellogg D.S., Jr., Peacock W.L., Jr., Deacon WE, Brown L, Pirkle CI: ***Neisseria gonorrhoeae*. I. Virulence genetically linked to clonal variation.** *Journal of Bacteriology* 1963, **85**:1274-1279.
40. Kim CC, Joyce EA, Chan K, Falkow S: **Improved analytical methods for microarray-based genome-composition analysis.** *Genome Biol* 2002, **3**:RESEARCH0065.
41. Durbin R, Thierry-Mieg JT: **A *C. elegans* DataBase. Documentation, code and data available from <http://www.acedb.org>.** 1991.
42. Snyder LA, Butcher SA, Saunders NJ: **Comparative whole-genome analyses reveal over 100 putative phase-variable genes in the pathogenic *Neisseria* spp.** *Microbiology* 2001, **147**:2321-2332.
43. Bonfield JK, Beal KF, Betts MJ, Staden R: **Trev: a DNA trace editor and viewer.** *Bioinformatics* 2002, **18**:194-195.
44. Altschul SF, Gish W, Miller W, Myers EV, Lipman DJ: **Basic local alignment search tool.** *J Mol Biol* 1990, **215**:403-410.

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

