

Dopamine modulates striato-frontal functioning during temporal processing

Catherine R. G. Jones^{1*} and Marjan Jahanshahi²

¹ Department of Psychology, University of Essex, Colchester, UK

² Sobell Department of Motor Neuroscience and Movement Disorders, UCL Institute of Neurology, London, UK

*Correspondence: crgjones@essex.ac.uk

With genesis in the ventral tegmental area and the substantia nigra pars compacta (SNc), the neurotransmitter dopamine influences brain function through three distinct pathways: the nigrostriatal, mesolimbic, and mesocortical. Dopamine plays a crucial role in a range of functions, for example: reward-related processing (Schultz et al., 1997) and reinforcement learning (Montague et al., 1996; Frank and Claus, 2006; Frank et al., 2007), working memory (Brozoski et al., 1979; Kimberg et al., 1997; Lustig et al., 2005; Cools et al., 2008), and motor function, including determining the vigor of actions (e.g., Niv et al., 2007; Smith and Villalba, 2008). Here our focus is on the role of dopamine in temporal processing, a less commonly recognized function (Meck, 1996; Buhusi and Meck, 2005; Meck et al., 2008; Jones and Jahanshahi, 2009; Allman and Meck, 2011; Coull et al., 2011 – but see Hata, 2011).

A growing body of research has sought to characterize the role of dopamine in interval timing, which can be broadly thought of as motor and perceptual timing in the milliseconds- and seconds-range. The influence of dopamine on interval timing has been demonstrated in pharmacological studies both with animals (Drew et al., 2003; Matell et al., 2006; Cheng et al., 2007; Meck et al., 2011) and humans (Rammsayer, 1993, 1997, for reviews see Meck et al., 2008; Jones and Jahanshahi, 2009; Coull et al., 2011). Using the peak-interval procedure, in which a learnt temporal interval is reproduced, animal research has established that dopamine agonists lead to the interval being underestimated, whereas dopamine antagonists lead to overestimation (e.g., Drew et al., 2003; Matell et al., 2004, 2006; MacDonald and Meck, 2005). These results have been interpreted as the effect of dopamine agonists and antagonists on speeding and slowing an “internal clock,” respectively. Lesions to the SNc and the caudate-putamen (CPu) both impair temporal control on the task, while

rats with lesions to the nucleus accumbens show no evidence of disrupted temporal performance (Meck, 2006a). This pattern implicates the nigrostriatal (substantia nigra–dorsal striatum) dopamine pathway in interval timing. Further, levodopa, a precursor to dopamine that is commonly used to treat Parkinson’s disease (PD), restores timing performance in rats with lesions to the SNc but not in those with lesions to the CPu, which may reflect the distinct roles of these structures in temporal calculation (Meck, 2006a). Work on healthy humans has established that haloperidol, a non-specific D2 receptor antagonist, attenuates both milliseconds- and seconds-range perceptual timing (comparing the length of two stimuli), whereas remoxipride, which blocks D2 receptors in the mesolimbic and mesocortical tracts, only impairs seconds-range timing (Rammsayer, 1997). These results were considered to support the role of the nigrostriatal system in milliseconds-range timing. More recently, Wiener et al. (2011) were able to apply a more targeted investigation by studying the effect of different single-nucleotide genetic polymorphisms on perceptual timing. Participants with a polymorphism affecting the density of striatal D2 receptors showed increased variability for milliseconds- but not seconds-range perceptual timing. Conversely, participants with a polymorphism that affects an enzyme (COMT) influencing prefrontal dopamine showed greater variability only in the seconds-range. Thus, these data suggest a double dissociation, with the nigrostriatal pathway being important for milliseconds-range perceptual timing and the mesocortical pathway being important for seconds-range perceptual timing.

Parkinson’s disease is a movement disorder associated with degeneration of dopaminergic neurons in the SNc. There is now a body of evidence that motor and perceptual timing within the milliseconds- and seconds-range are impaired in PD (see Jones

and Jahanshahi, 2009; Allman and Meck, 2011; Coull et al., 2011). Dopaminergic medication often improves performance on perceptual (e.g., Pastor et al., 1992a; Malapani et al., 1998) and motor timing tasks (e.g., Pastor et al., 1992b; O’Boyle et al., 1996) in patients with PD; although the pattern of effects (e.g., whether accuracy or variability is affected) is variable and significant effects are not always found (e.g., Pastor et al., 1992b; O’Boyle et al., 1996; Jones et al., 2008; Harrington et al., 2011b). The variable findings might relate to the insufficiency of dopaminergic medication for fully restoring striato-frontal function (Harrington et al., 2011b), or to the inadequacy of medication withdrawal and the lingering effects of long-acting medication in patients tested “off” medication. Difficulties with interval timing have also been observed in other disorders that are associated with dopaminergic dysfunction, including schizophrenia and attention-deficit/hyperactivity disorder (ADHD; see Jones and Jahanshahi, 2009; Allman and Meck, 2011).

There is consensus from imaging studies that the basal ganglia, particularly the dorsal striatum, are engaged during interval timing (e.g., Rao et al., 2001; Harrington et al., 2004, 2010, 2011a,b; Jahanshahi et al., 2006, 2010; Coull et al., 2008, for reviews see Meck et al., 2008; Jones and Jahanshahi, 2009; Coull et al., 2011). The basal ganglia are closely connected to the cortex through a series of striato-cortical loops (Alexander et al., 1986). Recording from neural ensembles in animal studies supports the role of both striatal and cortical regions in encoding temporal intervals (e.g., Matell et al., 2003, 2011; Lebedev et al., 2008; Höhn et al., 2011), whilst cortical lesions in rats attenuate temporal performance (Meck, 2006b). Further, patients with cortical lesions and healthy individuals with short-lasting TMS-induced disruption to cortical function demonstrate difficulties on temporal

tasks (Jones et al., 2004; Coslett et al., 2009). Thus there has been increasing interest in explaining and exploring how interval timing might be distributed across key cortical and subcortical regions. The obvious mediator for this relationship is dopamine. To better examine the effects of dopamine on striato-frontal function in PD, we investigated the neural correlates of motor timing in a repetitive tapping task, in patients tested both “on” and “off” apomorphine, a non-selective dopamine receptor agonist (Jahanshahi et al., 2010). Using effective connectivity analysis, we established that task-related coupling between the left head of the caudate nucleus and the prefrontal cortex was increased when the patients were “on” medication compared to the “off” state. The data support the proposal that dopamine modulates the coupling between frontal and striatal regions during motor timing. Fronto-striatal connectivity in PD has also been investigated during a perceptual timing task where the durations of two stimuli were compared (Harrington et al., 2011b). Consistent with the previous study, greater activation of the putamen and superior frontal gyrus was observed “on” compared to “off” medication during the decision phase of the perceptual timing task. However, the dominant finding was that connectivity between the basal ganglia and cortex was greater “off” than “on” medication during this task.

A challenge for the field has been to develop a biologically plausible model of interval timing. The striatal beat frequency model (Matell and Meck, 2000, 2004) proposes that the striatum receives cortical and thalamic oscillatory activity that serves as a clock signal. Using a process of coincidence detection, striatal spiny neurons fire when a criterion number of neuronal inputs oscillate with the same beat frequency. It is suggested that nigrostriatal phasic dopamine signals the onset and offset of a timed interval, whereas tonic dopamine alters the frequencies of the cortical oscillations, thus directly modulating the speed of the “internal clock.” According to this model, dopamine is a critical mediator of temporal calculations (see Allman and Meck, 2011; Oprisan and Buhusi, 2011).

Interval timing fits under the broad umbrella of temporal processing, which includes circadian rhythms through to the psychological relationship between time and

memory. The different types of temporal experience are poorly defined and understood (Grondin, 2010), and the relationship between these various types of temporal processing has received little attention. A complete account of interval timing should seek to position this process within the wider context of temporal processing. Particularly, an unexplored question is whether the role of dopamine in interval timing can be integrated with ostensibly distinct areas of dopamine-focused research that have a temporal component. The temporal difference model of learning proposes that the phasic activity of midbrain dopamine neurons code the time discrepancy between the expected and actual delivery of reward, calculating a reward prediction error (e.g., Montague et al., 1996; Schultz et al., 1997; Hollerman and Schultz, 1998). Dopamine neurons in the substantia nigra have also been implicated in temporal discounting, which is the waning of reward value with increasing delay, and in the temporal uncertainty inherent in delayed rewards (Kobayashi and Schultz, 2008). Related to this, administration of dopamine antagonists to healthy participants increases the degree of temporal discounting, i.e., the relative value of delayed rewards decreases, leading to impulsive behavior (Pine et al., 2010). Other work proposes that tonic levels of dopamine in the nucleus accumbens play an important role in response vigor (i.e., the rate of responding) by calculating the average rate of reward (Niv et al., 2007), a process that has an intrinsic temporal component. Finally, mesocortical dopamine is critical for working memory (Brozoski et al., 1979; Kimberg et al., 1997; Cools et al., 2008), which is essentially the process of maintaining information “on line” over a time interval. To date, the investigation of the role of dopamine in reward prediction error, reinforcement learning, temporal discounting, working memory, response vigor, and interval timing have remained largely distinct fields. There may be a value in exploring whether the temporal components of these tangential fields can be integrated with research that seeks to formulate a role for dopamine in modulating interval timing.

REFERENCES

Alexander, G. E., DeLong, M. R., and Strick, P. L. (1986). Parallel organization of functionally segregated circuits linking basal ganglia and cortex. *Annu. Rev. Neurosci.* 9, 357–381.

Allman, M. J., and Meck, W. H. (2011). Pathophysiological distortions in time perception and timed performance. *Brain*. doi: 10.1093/brain/awr210. [Epub ahead of print].

Brozoski, T. J., Brown, R. M., Rosvold, H. E., and Goldman, P. S. (1979). Cognitive deficit caused by regional depletion of dopamine in prefrontal cortex of rhesus monkey. *Science* 205, 929–932.

Buhusi, C. V., and Meck, W. H. (2005). What makes us tick? Functional and neural mechanisms of interval timing. *Nat. Rev. Neurosci.* 6, 755–765.

Cheng, R. K., Ali, Y. M., and Meck, W. H. (2007). Ketamine “unlocks” the reduced clock-speed effect of cocaine following extended training: evidence for dopamine-glutamate interactions in timing and time perception. *Neurobiol. Learn. Mem.* 88, 149–159.

Cools, R., Gibbs, S. E., Miyakawa, A., Jaquist, W., and D’Esposito, M. (2008). Working memory capacity predicts dopamine synthesis capacity in the human striatum. *J. Neurosci.* 28, 1208–1212.

Coslett, H. B., Shenton, J., Dyer, T., and Wiener, M. (2009). Cognitive timing: neuropsychology and anatomic basis. *Brain Res.* 1254, 38–48.

Coull, J. T., Cheng, R. K., and Meck, W. H. (2011). Neuroanatomical and neurochemical substrates of timing. *Neuropsychopharmacology* 36, 3–25.

Coull, J. T., Nazarian, B., and Vidal, F. (2008). Timing, storage, and comparison of stimulus duration engage discrete anatomical components of a perceptual timing network. *J. Cogn. Neurosci.* 12, 2185–2197.

Drew, M. R., Fairhurst, S., Malapani, C., Horvitz, J. C., and Balsam, P. D. (2003). Effects of dopamine antagonists on the timing of two intervals. *Pharmacol. Biochem. Behav.* 75, 9–15.

Frank, M. J., and Claus, E. D. (2006). Anatomy of a decision: striatal-orbitofrontal interactions in reinforcement learning, decision making, and reversal. *Psychol. Rev.* 113, 300–326.

Frank, M. J., Moustafa, A. A., Haughey, H. M., Curran, T., and Hutchinson, K. E. (2007). Genetic triple dissociation reveals multiple roles for dopamine in reinforcement learning. *Proc. Natl. Acad. Sci. U.S.A.* 9, 16311–16316.

Grondin, S. (2010). Timing and time perception: a review of recent behavioral and neuroscience findings and theoretical directions. *Atten. Percept. Psychophys.* 72, 561–582.

Harrington, D. L., Boyd, L. A., Mayer, A. R., Sheltraw, D. M., Lee, R. R., Huang, M., and Rao, S. M. (2004). Neural representation of interval encoding and decision making. *Brain Res. Cogn. Brain Res.* 21, 193–205.

Harrington, D. L., Castillo, G. N., Fong, C. H., and Reed, J. D. (2011a). Neural underpinnings of distortions in the experience of time across senses. *Front. Integr. Neurosci.* 5:32. doi: 10.3389/fnint.2011.00032

Harrington, D. L., Castillo, G. N., Greenberg, P. A., Song, D. D., Lessig, S., Lee, R. R., and Rao, S. M. (2011b). Neurobehavioral mechanisms of temporal processing deficits in Parkinson’s disease. *PLoS ONE* 6, e17461. doi: 10.1371/journal.pone.0017461

Harrington, D. L., Zimelman, J. L., Hinton, S. C., and Rao, S. M. (2010). Neural modulation of temporal encoding, maintenance, and decision processes. *Cereb. Cortex* 20, 1274–1285.

Hata, T. (2011). Glutamate – a forgotten target for interval timing. *Front. Integr. Neurosci.* 5:27. doi: 10.3389/fnint.2011.00027

Höhn, S., Dallérac, G., Faure, A., Urbach, Y., Nguyen, H. P., Riess, O., von Hörsten, S., Le Blanc, P., Desvignes, N., El Massioui, N., Brown, B. L., and Doyère, V. (2011).

- Behavioral and in vivo electrophysiological evidence for presymptomatic alteration of prefronto-striatal processing in the transgenic rat model for Huntington disease. *J. Neurosci.* 31, 8986–8997.
- Hollerman, J. R., and Schultz, W. (1998). Dopamine neurons report and error in the temporal prediction of reward during learning. *Nat. Neurosci.* 1, 304–309.
- Jahanshahi, M., Jones, C. R., Dirnberger, G., and Frith, C. D. (2006). The substantia nigra pars compacta and temporal processing. *J. Neurosci.* 26, 12266–12273.
- Jahanshahi, M., Jones, C. R., Zijlmans, J., Katzenschlager, R., Lee, L., Quinn, N., Frith, C. D., and Lees, A. J. (2010). Dopaminergic modulation of striato-frontal connectivity during motor timing in Parkinson's disease. *Brain* 133, 727–745.
- Jones, C. R., and Jahanshahi, M. (2009). The substantia nigra, the basal ganglia, dopamine and temporal processing. *J. Neural Transm. Suppl.* 73, 161–171.
- Jones, C. R., Rosenkranz, K., Rothwell, J. C., and Jahanshahi, M. (2004). The right dorsolateral prefrontal cortex is essential in time reproduction: an investigation with repetitive transcranial magnetic stimulation. *Exp. Brain Res.* 158, 66–372.
- Jones, C. R. G., Malone, T. J., Dirnberger, G., Edwards, M., and Jahanshahi, M. (2008). Basal ganglia, dopamine and temporal processing: performance on three timing tasks on and off medication in Parkinson's disease. *Brain Cogn.* 68, 30–41.
- Kimberg, D. Y., D'Esposito, M., and Farah, M. J. (1997). Effects of bromocriptine on human subjects depend on working memory capacity. *Neuroreport* 8, 3581–3585.
- Kobayashi, S., and Schultz, W. (2008). Influence of reward delays on responses of dopamine neurons. *J. Neurosci.* 28, 7837–7846.
- Lebedev, M. A., O'Doherty, J. E., and Nicolelis, M. A. (2008). Decoding of temporal intervals from cortical ensemble activity. *J. Neurophysiol.* 99, 166–186.
- Lustig, C., Matell, M. S., and Meck, W. H. (2005). Not “just” a coincidence: frontal-striatal synchronization in working memory and interval timing. *Memory* 13, 441–448.
- MacDonald, C. J., and Meck, W. H. (2005). Differential effects of clozapine and haloperidol on interval timing in the supraseconds range. *Psychopharmacology (Berl.)* 182, 232–244.
- Malapani, C., Rakitin, B., Meck, W. H., Deweer, B., Dubois, B., and Gibbon, J. (1998). Coupled temporal memories in Parkinson's disease: a dopamine-related dysfunction. *J. Cogn. Neurosci.* 10, 316–331.
- Matell, M. S., Bateson, M., and Meck, W. H. (2006). Single-trials analyses demonstrate that increases in clock speed contribute to the methamphetamine-induced horizontal shifts in peak-interval timing functions. *Psychopharmacology (Berl.)* 188, 201–212.
- Matell, M. S., King, G. R., and Meck, W. H. (2004). Differential adjustment of interval timing by the chronic administration of intermittent or continuous cocaine. *Behav. Neurosci.* 118, 150–156.
- Matell, M. S., and Meck, W. H. (2000). Neuropsychological mechanisms of interval timing behavior. *Bioessays* 22, 94–103.
- Matell, M. S., and Meck, W. H. (2004). Cortico-striatal circuits and interval timing: coincidence detection of oscillatory processes. *Brain Res. Cogn. Brain Res.* 21, 139–170.
- Matell, M. S., Meck, W. H., and Nicolelis, M. A. (2003). Interval timing and the encoding of signal duration by ensembles of cortical and striatal neurons. *Behav. Neurosci.* 117, 760–773.
- Matell, M. S., Shea-Brown, E., Gooch, C., Wilson, A. G., and Rinzel, J. A. (2011). A heterogeneous population code for elapsed time in rat medial agranular cortex. *Behav. Neurosci.* 125, 54–73.
- Meck, W. H. (1996). Neuropharmacology of timing and time perception. *Brain Res. Cogn. Brain Res.* 3, 227–242.
- Meck, W. H. (2006a). Neuroanatomical localization of an internal clock: a functional link between mesolimbic, nigrostriatal, and mesocortical dopaminergic systems. *Brain Res.* 1109, 93–107.
- Meck, W. H. (2006b). Frontal cortex lesions eliminate the clock speed effect of dopaminergic drugs on interval timing. *Brain Res.* 1108, 157–167.
- Meck, W. H., Cheng, R. K., MacDonald, C. J., Gainetdinov, R. R., Caron, M. G., and Çevik, M. Ö. (2011). Genodose dependent effects of methamphetamine on interval timing in dopamine-transporter knockout mice. *Neuropharmacology*. doi: 10.1016/j.neuropharm.2011.01.042. [Epub ahead of print].
- Meck, W. H., Penney, T. B., and Pouthas, V. (2008). Cortico-striatal representation of time in animals and humans. *Curr. Opin. Neurobiol.* 18, 145–152.
- Montague, P. R., Dayan, P., and Sejnowski, T. J. (1996). A framework for mesencephalic dopamine systems based on predictive Hebbian learning. *J. Neurosci.* 16, 1936–1947.
- Niv, Y., Daw, N. D., Joel, D., and Dayan, P. (2007). Tonic dopamine: opportunity costs and the control of response vigor. *Psychopharmacology (Berl.)* 191, 507–520.
- O'Boyle, D. J., Freeman, J. S., and Cody, F. W. (1996). The accuracy and precision of timing of self-paced, repetitive movements in subjects with Parkinson's disease. *Brain* 119, 51–70.
- Oprisan, S. A., and Buhusi, C. V. (2011). Modeling pharmacological clock and memory patterns of interval timing in a striatal beat-frequency model with realistic, noisy neurons. *Front. Integr. Neurosci.* 5:52. doi: 10.3389/fnint.2011.00052
- Pastor, M. A., Jahanshahi, M., Artieda, J., and Obeso, J. A. (1992a). Time estimation and reproduction is abnormal in Parkinson's disease. *Brain* 115, 875–891.
- Pastor, M. A., Jahanshahi, M., Artieda, J., and Obeso, J. A. (1992b). Performance of repetitive wrist movements in Parkinson's disease. *Brain* 115, 211–225.
- Pine, A., Shiner, T., Seymour, B., and Dolan, R. J. (2010). Dopamine, time, and impulsivity in humans. *J. Neurosci.* 30, 8888–8896.
- Rammesayer, T. (1993). On dopaminergic modulation of temporal information processing. *Biol. Psychol.* 36, 209–222.
- Rammesayer, T. H. (1997). Are there dissociable roles of the mesostriatal and mesolimbocortical dopamine systems on temporal information processing in humans? *Neuropsychobiology* 35, 36–45.
- Rao, S. M., Mayer, A. R., and Harrington, D. L. (2001). The evolution of brain activation during temporal processing. *Nat. Neurosci.* 4, 317–323.
- Schultz, W., Dayan, P., and Montague, P. R. (1997). A neural substrate of prediction and reward. *Science* 275, 1593–1599.
- Smith, Y., and Villalba, R. (2008). Striatal and extrastriatal dopamine in the basal ganglia: an overview of its anatomical organization in normal and Parkinsonian brains. *Mov. Disord.* 23, 534–547.
- Wiener, M., Lohoff, F. W., and Coslett, H. B. (2011). Double dissociation of dopamine genes and timing in humans. *J. Cogn. Neurosci.* 23, 2811–2821.

Received: 07 October 2011; accepted: 07 October 2011; published online: 25 October 2011.

Citation: Jones CRG and Jahanshahi M (2011) Dopamine modulates striato-frontal functioning during temporal processing. *Front. Integr. Neurosci.* 5:70. doi: 10.3389/fnint.2011.00070

Copyright © 2011 Jones and Jahanshahi. This is an open-access article subject to a non-exclusive license between the authors and Frontiers Media SA, which permits use, distribution and reproduction in other forums, provided the original authors and source are credited and other Frontiers conditions are complied with.