

## Orientations, lattice polytopes, and group arrangements II: Modular and integral flow polynomials of graphs

Beifang Chen and Richard Stanley

ABSTRACT. We study modular and integral flow polynomials of graphs by means of subgroup arrangements and lattice polytopes. We introduce an Eulerian equivalence relation on orientations, flow arrangements, and flow polytopes; and we apply the theory of Ehrhart polynomials to obtain properties of modular and integral flow polynomials. The emphasis is on the geometrical treatment through subgroup arrangements and Ehrhart polynomials. Such viewpoint leads to a reciprocity law for the modular flow polynomial, which gives rise to an interpretation on the values of the modular flow polynomial at negative integers, and answers a question by Beck and Zaslavsky.

### 1. Introduction

The flow polynomial  $\varphi(G, t)$  of a graph  $G$  was introduced by Tutte [24] as a conceptual dual to the chromatic (or tension) polynomial of  $G$ . When  $G$  is a planar graph,  $\varphi(G, t)$  is essentially the chromatic polynomial  $\chi(G^*, t)$  of the dual graph  $G^*$  in the sense that  $\chi(G^*, t) = t^{c(G)}\varphi(G, t)$ , where  $c(G)$  is the number of connected components of  $G$ . The historic Four-Color Conjecture of the time was made by Tutte into the Five-Flow Conjecture: any bridgeless graph admits a nowhere-zero integer 5-flow. Both conjectures are still open and stimulate studies on chromatic and flow polynomials. In a seminal paper [19], Rota introduced characteristic polynomial for posets and observed that  $\varphi(G, t)$  is the characteristic polynomial of the circuit lattice of  $G$ . Greene and Zaslavsky [15] made Rota's observation transparent between the flow polynomial and the characteristic polynomial by using hyperplane arrangements. As a special case of Zaslavsky's formula [25], the absolute value  $|\varphi(G, -1)|$  counts the number of totally cyclic orientations of  $G$ , which is a dual analog of Stanley's result on chromatic polynomials:  $|\chi(G, -1)|$  counts the number of acyclic orientations of  $G$ . However, Stanley's result [20] includes an interpretation of the values of  $\chi(G, t)$  at negative integers, known as the Reciprocity Law of chromatic polynomials. More recently, Kochol [17] showed that the number

---

1991 *Mathematics Subject Classification.* 05A99, 05B35, 05C15, 06A07, 52B20, 52B40.

*Key words and phrases.* Subgroup arrangement, hyperplane arrangement, orientation, totally cyclic orientation, directed Eulerian subgraph, Eulerian equivalence relation, flow polytope, integer flow, modular flow, flow polynomial, dual flow polynomial, characteristic polynomial, Tutte polynomial, reciprocity law.

Research of the first author is supported by RGC Competitive Earmarked Research Grants 600703, 600506, and 600608.

of integer-valued  $q$ -flows is a polynomial function of  $q$  and introduced the integral flow polynomial  $\varphi_{\mathbb{Z}}(G, t)$ ; Beck and Zaslavsky [1] studied the modular and integral flow polynomials for graphs and signed graphs, using Ehrhart polynomials of lattice polytopes. The present paper, as a continuation of [11], is to associate flow group arrangements with graphs, and to obtain a clear picture of the relation between the integral flow polynomial and the modular flow polynomial. The byproduct of this association is a generalization of the Reciprocity Law of chromatic and tension polynomials to modular and integral flow polynomials, and the interpretation of the values of the modular and integral flow polynomials at zero and negative integers. The geometric method of our exposition may be modified to obtain analogous results on Tutte polynomials.

Let  $G = (V, E)$  be a finite graph with possible loops and multiple edges. We write  $V = V(G)$ ,  $E = E(G)$ . For each subset  $X \subseteq E$ , denote by  $\langle X \rangle$  the induced subgraph  $(V, X)$ . An *orientation* on  $G$  is a (multivalued) function  $\varepsilon : V \times E \rightarrow \{-1, 0, 1\}$  such that (i)  $\varepsilon(v, e)$  has the ordered double-value  $\pm 1$  or  $\mp 1$  if  $e$  is a loop at a vertex  $v$  and has a single-value otherwise, (ii)  $\varepsilon(v, e) = 0$  if  $v$  is not an end-vertex of  $e$ , and (iii)  $\varepsilon(u, e)\varepsilon(v, e) = -1$  if  $e$  has two distinct end-vertices  $u, v$ . Pictorially, if  $e$  is a non-loop edge with distinct end-vertices  $u, v$ , then  $\varepsilon(u, e) = -\varepsilon(v, e) = 1$  (or  $\varepsilon(v, e) = -\varepsilon(u, e) = 1$ ), and it means that  $e$  is assigned an arrow from  $u$  to  $v$ , which contributes exactly one out-degree at  $u$  and one in-degree at  $v$ . If  $e$  is a loop at a vertex  $v$ , then  $\varepsilon(v, e) = \pm 1$  or  $\mp 1$ , and it means that the loop  $e$  is assigned an arrow, pointing away and to the vertex  $v$ , which contributes exactly one out-degree and one in-degree at  $v$ . We assume  $-(\pm 1) = \mp 1$ ,  $-(\mp 1) = \pm 1$ . A graph  $G$  together with an orientation  $\varepsilon$  is called a *digraph*, denoted  $(G, \varepsilon)$ . A digraph is said to be *directed Eulerian* if its in-degree equals its out-degree at every vertex.

Let  $(G, \varepsilon)$  be a digraph throughout the whole paper. Associated with  $(G, \varepsilon)$  is the *incidence matrix*  $\mathbf{M} = \mathbf{M}(G) := [\mathbf{m}_{v,e}]_{V \times E}$ , where  $\mathbf{m}_{v,e} = 0$  if the edge  $e$  is a loop and  $\mathbf{m}_{v,e} = \varepsilon(v, e)$  if  $e$  is not a loop. Let  $A$  be an abelian group. A *flow* of  $(G, \varepsilon)$  with values in  $A$ , or an *A-flow*, is a function  $f : E \rightarrow A$ , satisfying the *Conservation Law*:

$$\sum_{e \in E} \mathbf{m}_{v,e}(v, e)f(e) = 0 \quad \text{or} \quad \sum_{e \in E} \varepsilon(v, e)f(e) = 0, \quad v \in V, \quad (1.1)$$

where  $\varepsilon(v, e)$  is counted twice in the second sum as  $-1$  and  $1$  if  $e$  is a loop at its unique end-vertex  $v$ . A flow  $f$  is said to be *nowhere-zero* if  $f(e) \neq 0$  for all  $e \in E$ . We denote by  $F(G, \varepsilon; A)$  the abelian group of all  $A$ -flows of  $(G, \varepsilon)$ . The *flow arrangement* of  $(G, \varepsilon)$  with the abelian group  $A$  is the group arrangement  $\mathcal{A}_{\text{FL}}(G, \varepsilon; A)$  of  $F(G, \varepsilon; A)$ , consisting of the subgroups

$$F_e := \{f \in F(G, \varepsilon; A) \mid f(e) = 0\}, \quad e \in E. \quad (1.2)$$

Coincidentally, we shall see that the characteristic polynomial  $\chi(\mathcal{A}_{\text{FL}}(G, \varepsilon; A), t)$  is equal to the (*modular*) *flow polynomial*  $\varphi(G, t)$ , defined for  $t = q$  as the number of nowhere-zero flows of  $(G, \varepsilon)$  with values in an abelian group of order  $q$ . The polynomial  $\varphi$  is independent of the chosen orientation  $\varepsilon$  and the abelian group structure; see Rota [19] and Tutte [23]. For a complete information about modular and integral flows, we refer to the book of Zhang [27].

Recall that a *cut* of  $G$  is a nonempty edge subset of the form  $[S, S^c]$ , where  $S \subseteq V$  is a nonempty proper subset,  $S^c := V - S$  is the complement of  $S$ , and  $[S, S^c]$  is the set of all edges between the vertices of  $S$  and the vertices of  $S^c$ . Let

$U = [S, S^c]$  be a cut. A *direction* of  $U$  is an *orientation*  $\varepsilon_U$  on the induced subgraph  $(V, U)$  such that the arrows of the edges in  $U$  are either all from  $S$  to  $S^c$  or all from  $S^c$  to  $S$ ;  $U$  together with a direction  $\varepsilon_U$  is called a *directed cut*, denoted  $(U, \varepsilon_U)$ . If  $(U, \varepsilon)$  is a directed cut, we call  $(U, \varepsilon)$  a directed cut of both  $(G, \varepsilon)$  and  $\varepsilon$ , and say that the cut  $U$  is *directed* in  $(G, \varepsilon)$ . Let  $\mathcal{O}(G)$  denote the set of all orientations on  $G$ . We denote by  $\mathcal{O}_{\text{TC}}(G)$  the set of all orientations without directed cut (also known as *totally cyclic orientations*, as they are the orientations in which every edge belongs to a directed circuit).

Let  $\varphi_{\mathbb{z}}(G, q)$  denote the number of nowhere-zero integer-valued flows  $f$  of  $(G, \varepsilon)$  such that  $0 < |f(e)| < q$  for all  $e \in E$ . As pointed out by Beck and Zaslavsky [1], the function  $\varphi_{\mathbb{z}}(G, q)$  was never mentioned to be a polynomial until Kochol [17]. If  $\varepsilon$  is totally cyclic, we introduce the counting functions

$$\varphi_{\varepsilon}(G, q) := \#\{f \in F(G, \varepsilon; \mathbb{Z}) \mid 0 < f(e) < q, e \in E\}, \quad (1.3)$$

$$\bar{\varphi}_{\varepsilon}(G, q) := \#\{f \in F(G, \varepsilon; \mathbb{Z}) \mid 0 \leq f(e) \leq q, e \in E\}, \quad (1.4)$$

and the relatively open 0-1 polytope

$$\Delta_{\text{FL}}^+(G, \varepsilon) := \{f \in F(G, \varepsilon; \mathbb{R}) \mid 0 < f(e) < 1, e \in E\}. \quad (1.5)$$

The closure  $\bar{\Delta}_{\text{FL}}^+(G, \varepsilon)$  is a 0-1 polytope (whose vertices are 0-1 vectors), and is the convex hull of all 0-1 flows of  $(G, \varepsilon)$  in  $\mathbb{R}^E$  (flows whose values are either 0 or 1), called the *flow polytope* of  $(G, \varepsilon)$ . We introduce the following counting function

$$\bar{\varphi}_{\mathbb{z}}(G, q) := \#\{(\rho, f) \mid \rho \in \mathcal{O}_{\text{TC}}(G), f \in F(G, \rho; \mathbb{Z}), 0 \leq f(e) \leq q, e \in E\}. \quad (1.6)$$

We shall see that  $\varphi_{\mathbb{z}}(G, q)$ ,  $\varphi_{\varepsilon}(G, q)$  are polynomial functions of positive integers  $q$ , and  $\bar{\varphi}_{\mathbb{z}}(G, q)$ ,  $\bar{\varphi}_{\varepsilon}(G, q)$  are polynomial functions of nonnegative integers  $q$ , and that  $\varphi_{\mathbb{z}}(G, q)$  is independent of the chosen orientation  $\varepsilon$ . The corresponding polynomial  $\varphi_{\mathbb{z}}(G, t)$  ( $\bar{\varphi}_{\mathbb{z}}(G, t)$ ) is called the (*dual*) *integral flow polynomial* of  $G$ , and  $\varphi_{\varepsilon}(G, t)$  ( $\bar{\varphi}_{\varepsilon}(G, t)$ ) the *local (dual) flow polynomial with respect to the orientation*  $\varepsilon$ . The names and notations are so selected in order to easily recognize these polynomials.

We first reproduce a result due to Kochol [17] about Equation (1.8), and due to Beck and Zaslavsky [1] about the combinatorial interpretation of the values of  $\varphi_{\mathbb{z}}(G, t)$  at nonpositive integers.

**Theorem 1.1** (Kochol [17], Beck and Zaslavsky [1]). *Let  $G = (V, E)$  be a finite bridgeless graph with possible loops and multiple edges.*

- (a) *If the orientation  $\varepsilon$  is totally cyclic, then  $\Delta_{\text{FL}}^+(G, \varepsilon)$  is a relatively open 0-1 polytope in  $\mathbb{R}^E$  of dimension  $n(G)$ ;  $\varphi_{\varepsilon}(G, t)$  and  $\bar{\varphi}_{\varepsilon}(G, t)$  are Ehrhart polynomials of  $\Delta_{\text{FL}}^+(G, \varepsilon)$  and  $\bar{\Delta}_{\text{FL}}^+(G, \varepsilon)$  respectively, and satisfy the Reciprocity Law:*

$$\varphi_{\varepsilon}(G, -t) = (-1)^{n(G)} \bar{\varphi}_{\varepsilon}(G, t), \quad (1.7)$$

where  $n(G) = |E| - r(G)$  and  $r(G)$  is the number of edges of a maximal spanning forest of  $G$ . Moreover,

$$\varphi_{\varepsilon}(G, 0) = (-1)^{n(G)}, \quad \bar{\varphi}_{\varepsilon}(G, 0) = 1.$$

(b) The integral flow polynomials  $\varphi_{\mathbb{Z}}(G, t)$  and  $\bar{\varphi}_{\mathbb{Z}}(G, t)$  can be written as

$$\varphi_{\mathbb{Z}}(G, t) = \sum_{\rho \in \mathcal{O}_{\text{TC}}(G)} \varphi_{\rho}(G, t), \quad (1.8)$$

$$\bar{\varphi}_{\mathbb{Z}}(G, t) = \sum_{\rho \in \mathcal{O}_{\text{TC}}(G)} \bar{\varphi}_{\rho}(G, t), \quad (1.9)$$

and satisfy the Reciprocity Law:

$$\varphi_{\mathbb{Z}}(G, -t) = (-1)^{n(G)} \bar{\varphi}_{\mathbb{Z}}(G, t). \quad (1.10)$$

In particular,  $|\varphi_{\mathbb{Z}}(G, 0)|$  counts the number of totally cyclic orientations on  $G$ .

There are analogous results on the modular flow polynomial  $\varphi(G, t)$ . To do this we need to introduce an equivalence relation on the set  $\mathcal{O}(G)$  of orientations on  $G$ . Two orientations  $\varepsilon_1, \varepsilon_2$  on  $G$  are said to be *Eulerian equivalent*, written  $\varepsilon_1 \sim \varepsilon_2$ , if the spanning subgraph induced by the edge subset  $\{e \in E \mid \varepsilon_1(v, e) \neq \varepsilon_2(v, e)\}$  is a directed Eulerian subgraph with respect to the orientation either  $\varepsilon_1$  or  $\varepsilon_2$ . We shall see that  $\sim$  is indeed an equivalence relation on  $\mathcal{O}(G)$ . Moreover, if an Eulerian equivalence class intersects  $\mathcal{O}_{\text{TC}}(G)$ , the whole equivalence class is contained in  $\mathcal{O}_{\text{TC}}(G)$ . So  $\sim$  induces an equivalence relation on the set  $\mathcal{O}_{\text{TC}}(G)$  of totally cyclic orientations. Let  $[\mathcal{O}_{\text{TC}}(G)]$  denote a set of distinct representatives, exact one representative from each equivalence class of  $\sim$  on  $\mathcal{O}_{\text{TC}}(G)$ . We introduce the following counting function

$$\bar{\varphi}(G, q) := \#\{(\rho, f) \mid \rho \in [\mathcal{O}_{\text{TC}}(G)], f \in F(G, \rho; \mathbb{Z}), 0 \leq f(e) \leq q, e \in E\}. \quad (1.11)$$

We next produce the following Theorem 1.2, which answers a question by Beck and Zaslavsky [1] about the combinatorial interpretation of the values of the modular flow polynomial  $\varphi(G, t)$  at zero and negative integers.

**Theorem 1.2.** *Let  $G = (V, E)$  be a finite bridgeless graph with possible loops and multiple edges. Then  $\varphi(G, q)$  ( $\bar{\varphi}(G, q)$ ) is a polynomial function of degree  $n(G)$  of positive (nonnegative) integers  $q$ , and satisfy the Reciprocity Law:*

$$\varphi(G, -t) = (-1)^{n(G)} \bar{\varphi}(G, t). \quad (1.12)$$

Moreover,

$$\varphi(G, t) = \sum_{\rho \in [\mathcal{O}_{\text{TC}}(G)]} \varphi_{\rho}(G, t), \quad (1.13)$$

$$\bar{\varphi}(G, t) = \sum_{\rho \in [\mathcal{O}_{\text{TC}}(G)]} \bar{\varphi}_{\rho}(G, t). \quad (1.14)$$

In particular,  $|\varphi(G, -1)|$  counts the number of totally cyclic orientations on  $G$ , and  $|\varphi(G, 0)|$  counts the number of Eulerian equivalence classes of totally cyclic orientations.

Equation (1.13) is recently obtained by Kochol [17] with a formal proof in different form. The combinatorial interpretation of  $|\varphi(G, -1)| = T_G(0, 2)$  is due to Las Vergnas [18], see also Brylawski and Oxley [6]. At the moment of this revising, we noticed a paper by Breuer and Sanyal [5] on modular flow reciprocity, which is quite different from our Reciprocity Law (1.12). The difference lies in that the result of [5] on  $\varphi(G, -q)$  for a positive integer  $q$  involves the counting of flows modulo  $q$ , our result only involves nonnegative integer flows bounded by  $q$ , and

that the bijection between the two counting sets is nontrivial; see Section 5 for the detailed discussion. The proof of Theorem 1.2 is rigorous and self-contained. The following corollary is an immediate consequence of Theorem 1.2.

**Corollary 1.3.** *The value  $T_G(0,1)$  of the Tutte polynomial  $T_G(x,y)$  counts the number of Eulerian equivalence classes of totally cyclic orientations on  $G$ .*

## 2. Characteristic polynomials of group arrangements

Let  $\Omega$  be a finitely generated abelian group. By a *flat* of  $\Omega$  we mean a coset of a subgroup of  $\Omega$ . For a subgroup  $\Gamma \subseteq \Omega$ , we denote by  $\text{Tor}(\Gamma)$  the torsion subgroup of  $\Gamma$  and write  $|\Gamma| := |\text{Tor}(\Gamma)| t^{\text{rank}(\Gamma)}$ . By a *subgroup arrangement* (or just *arrangement*) of  $\Omega$  we mean a finite collection of flats of  $\Omega$ . Associated with a subgroup arrangement  $\mathcal{A}$  is the semilattice  $\mathcal{L}(\mathcal{A})$ , whose members are nonempty sets obtained from all possible intersections of flats in  $\mathcal{A}$ . The *characteristic polynomial* of  $\mathcal{A}$  is defined as

$$\chi(\mathcal{A}, t) = \sum_{X \in \mathcal{L}(\mathcal{A})} \frac{|\text{Tor}(\Omega)|}{|\text{Tor}(\Omega/\langle X \rangle)|} \mu(X, \Omega) t^{\text{rank}(X)}, \quad (2.1)$$

where  $\mu$  is the Möbius function of the poset  $\mathcal{L}(\mathcal{A})$ , whose partial order is the set inclusion,  $\langle X \rangle := \{x - y \mid x, y \in X\}$ .

Let  $\mathcal{B}(\Omega)$  be the Boolean algebra generated by cosets of all subgroups of  $\Omega$ , i.e., every member of  $\mathcal{B}(\Omega)$  is obtained from cosets of subgroups of  $\Omega$  by taking unions, intersections, and complements finitely many times. A *valuation* on  $\Omega$  with values in an abelian group  $A$  is a map  $\nu : \mathcal{B}(\Omega) \rightarrow A$  such that

$$\begin{aligned} \nu(\emptyset) &= 0, \\ \nu(X \cup Y) &= \nu(X) + \nu(Y) - \nu(X \cap Y) \end{aligned}$$

for  $X, Y \in \mathcal{B}(\Omega)$ . A valuation  $\nu$  is said to be *translation invariant* if

$$\nu(S + x) = \nu(S)$$

for  $S \in \mathcal{B}(\Omega)$  and any  $x \in \Omega$ ; and  $\nu$  is said to satisfy *multiplicativity* if

$$\nu(A + B) = \nu(A) \nu(B)$$

for subgroups  $A, B \subseteq \Omega$  such that  $A + B$  is a direct sum of  $A$  and  $B$ , and the subgroup  $A + B$  is a direct summand of  $\Omega$ .

**Theorem 2.1** (Chen [11]). *For any finitely generated abelian group  $\Omega$ , there exists a unique translation invariant valuation  $\lambda : \mathcal{B}(\Omega) \rightarrow \mathbb{Q}[t]$  such that the multiplicativity is satisfied and*

$$\lambda(\Omega) = |\text{Tor}(\Omega)| t^{\text{rank}(\Omega)} = |\Omega|.$$

*In particular,  $\lambda(\Gamma) = \frac{|\Omega|}{|\Omega/\Gamma|}$  for any subgroup  $\Gamma \subseteq \Omega$ , and for any subgroup arrangement  $\mathcal{A}$  of  $\Omega$ ,*

$$\lambda\left(\Omega - \bigcup_{X \in \mathcal{A}} X\right) = \chi(\mathcal{A}, t).$$

The analogue of Theorem 2.1 for vector spaces was obtained by Ehrenborg and Readdy [13]. Let  $V$  be a vector space over an infinite field. Let  $\mathcal{L}(V)$  be the lattice of all affine subspaces of  $V$ . We denote by  $\mathcal{B}(V)$  the Boolean algebra generated by  $\mathcal{L}(V)$ . A *subspace arrangement* of  $V$  is a finite collection  $\mathcal{A}$  of affine subspaces of  $V$ . The *torsion* of any subspace is just the zero space. Then *characteristic polynomial*

$\chi(\mathcal{A}, t)$  of a subspace arrangement  $\mathcal{A}$  can be defined by the same formula (2.1) for subgroup arrangement.

**Theorem 2.2** (Ehrenborg and Readdy [13]). *For any finite-dimensional vector space  $V$  over an infinite field  $\mathbb{K}$ , there exists a unique translation invariant valuation  $\lambda : \mathcal{B}(V) \rightarrow \mathbb{Z}[t]$  such that  $\lambda(W) = t^{\dim W}$  for subspaces  $W \subseteq V$ . Moreover, for a subspace arrangement  $\mathcal{A}$  of  $V$ ,*

$$\lambda\left(V - \bigcup_{X \in \mathcal{A}} X\right) = \chi(\mathcal{A}, t).$$

One may combine Theorems 2.1 and 2.2 by considering arrangements of affine submodules. Let  $M$  be a finitely generated left  $R$ -module over a commutative ring  $R$ ; we restrict  $R$  to the cases of  $\mathbb{R}$ ,  $\mathbb{Z}$ , and  $\mathbb{Z}/q\mathbb{Z}$ . By a *flat* of  $M$  we mean a subset of the form  $a + N = \{a + x \mid x \in N\}$ , where  $N$  is a submodule of  $M$ . Let  $\mathcal{L}(M)$  be the lattice of all flats of  $M$ , and  $\mathcal{B}(M)$  the Boolean algebra generated by  $\mathcal{L}(M)$ . For each subset  $S \subseteq M$ , we denote by  $1_S$  the characteristic function of  $S$ .

Let  $\mathcal{A}$  be a finite collection of flats in  $M$ , called a *submodule arrangement* of  $M$ . Let  $\mathcal{L}(\mathcal{A})$  be the poset whose members are nonempty sets obtained by taking all possible intersections of flats in  $\mathcal{A}$ , and whose partial order  $\leq$  is the set inclusion. For each  $X \in \mathcal{L}(\mathcal{A})$ , we define

$$X^\circ := X - \bigcup_{Y \in \mathcal{L}(\mathcal{A}), Y < X} Y.$$

Clearly,  $\{X^\circ \mid X \in \mathcal{L}(\mathcal{A})\}$  is a family of disjoint subsets of  $M$ . Then for each  $X \in \mathcal{L}(\mathcal{A})$ ,

$$1_X = \sum_{Y \in \mathcal{L}(\mathcal{A}), Y \leq X} 1_{Y^\circ}.$$

By the Möbius inversion, for each  $X \in \mathcal{L}(\mathcal{A})$ ,

$$1_{X^\circ} = \sum_{Y \in \mathcal{L}(\mathcal{A}), Y \leq X} \mu(Y, X) 1_Y.$$

In particular,  $M^\circ = M - \bigcup \mathcal{A} = M - \bigcup_{X \in \mathcal{A}} X$  and

$$1_{M - \bigcup \mathcal{A}} = \sum_{Y \in \mathcal{L}(\mathcal{A})} \mu(Y, M) 1_Y. \quad (2.2)$$

Thus for any valuation  $\nu$  on  $\mathcal{B}(M)$ , we have the Inclusion-Exclusion Formula:

$$\nu(M - \bigcup \mathcal{A}) = \sum_{X \in \mathcal{L}(\mathcal{A})} \mu(X, M) \nu(X). \quad (2.3)$$

This is a prototype of many existing formulas when  $\nu$  is taken to be various valuations; see [8, 13, 26].

Let  $M$  be the Euclidean  $n$ -space  $\mathbb{R}^n$ . One has half-spaces  $\{x \in \mathbb{R}^n \mid f(x) \leq c\}$  (with linear functionals  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  and constant real numbers  $c$ ), convex polyhedra (intersections of half-spaces), and the Boolean algebra  $\mathcal{B}(\mathcal{P}^n)$  (generated by half-spaces by taking intersections, unions, and relatively complements finitely many times). There are two valuations  $\chi$  and  $\bar{\chi}$  on  $\mathcal{B}(\mathcal{P}^n)$ , both are referred to the *Euler characteristic* (see [7, 22, 26], for example), such that for any relatively open convex polyhedron  $P$ ,

$$\chi(P) = (-1)^{\dim P}, \quad \bar{\chi}(P) = \lim_{r \rightarrow \infty} \chi(P \cap [-r, r]^n).$$

By Groemer's extension theorem [16],  $\chi$  and  $\bar{\chi}$  can be extended to be linear functionals on the functional space spanned by characteristic functions of convex polyhedra. Now let  $\mathcal{A}$  be a hyperplane arrangement of  $\mathbb{R}^n$ . Evaluating  $\chi$  and  $\bar{\chi}$  on both sides of (2.3), one obtains Zaslavsky's first and second counting formulas (see [13, 25, 26]):

$$|\chi(\mathcal{A}, -1)| = \text{number of regions of } \mathbb{R}^n - \bigcup \mathcal{A}, \quad (2.4)$$

$$|\chi(\mathcal{A}, 1)| = \text{number of relatively bounded regions of } \mathbb{R}^n - \bigcup \mathcal{A}. \quad (2.5)$$

### 3. Modular flow polynomials

Let  $(H_i, \varepsilon_i)$  be subdigraphs of the graph  $G = (V, E)$ ,  $i = 1, 2$ . The *coupling* of  $\varepsilon_1$  and  $\varepsilon_2$  is a function  $[\varepsilon_1, \varepsilon_2] : E \rightarrow \{-1, 0, 1\}$ , defined for each edge  $e \in E$  (at its one end-vertex  $v$ ) by

$$[\varepsilon_1, \varepsilon_2](e) = \begin{cases} 1 & \text{if } e \in E(H_1) \cap E(H_2), \varepsilon_1(v, e) = \varepsilon_2(v, e), \\ -1 & \text{if } e \in E(H_1) \cap E(H_2), \varepsilon_1(v, e) \neq \varepsilon_2(v, e), \\ 0 & \text{otherwise.} \end{cases} \quad (3.1)$$

The following proposition is straightforward.

**Proposition 3.1** (Berge [2]). (a) *A function  $f : E(G) \rightarrow A$  is a flow of  $(G, \varepsilon)$  if and only if for any directed cut  $(U, \varepsilon_U)$ ,*

$$\sum_{e \in U} [\varepsilon, \varepsilon_U](e) f(e) = 0.$$

(b) *In particular, the digraph  $(G, \varepsilon)$  is directed Eulerian if and only if for any directed cut  $(U, \varepsilon_U)$ ,*

$$\sum_{e \in U} [\varepsilon, \varepsilon_U](e) = 0.$$

(c) *The graph  $G$  is Eulerian if and only if every cut  $U$  contains even number of edges.*

Let  $F_{\text{nz}}(G, \varepsilon; A)$  denote the set of all nowhere-zero flows with values in  $A$ , i.e.,

$$F_{\text{nz}}(G, \varepsilon; A) := \{f \in F(G, \varepsilon; A) \mid f(e) \neq 0, e \in E\}.$$

If  $|A| = q$  is finite, it is well-known (see [27]) that the counting function

$$\varphi(G, q) := |F_{\text{nz}}(G, \varepsilon; A)| \quad (3.2)$$

is a polynomial function of  $q$ , depending only on the order  $|A|$ , but not on the chosen orientation  $\varepsilon$  and the group structure of  $A$ . The polynomial  $\varphi(G, t)$  is called the *modular flow polynomial* of  $G$ .

For two orientations  $\rho, \sigma \in \mathcal{O}(G)$ , there is an involution  $P_{\rho, \sigma} : A^E \rightarrow A^E$ , defined by

$$(P_{\rho, \sigma} f)(e) = \begin{cases} f(e) & \text{if } \rho(v, e) = \sigma(v, e), \\ -f(e) & \text{if } \rho(v, e) \neq \sigma(v, e). \end{cases} \quad (3.3)$$

In fact,  $P_{\rho, \varepsilon} f = [\rho, \varepsilon]f$ . Obviously,  $P_{\rho, \rho}$  is the identity map,  $P_{\rho, \sigma} P_{\sigma, \varepsilon} = P_{\rho, \varepsilon}$ .

**Lemma 3.2.** *The involution  $P_{\rho, \varepsilon}$  is a group isomorphism. Moreover,*

$$\begin{aligned} P_{\rho, \varepsilon} F(G, \varepsilon; A) &= F(G, \rho; A), \\ P_{\rho, \varepsilon} F_{\text{nz}}(G, \varepsilon; A) &= F_{\text{nz}}(G, \rho; A). \end{aligned}$$

PROOF. It is clear that  $P_{\rho,\varepsilon}$  is invertible and  $P_{\rho,\varepsilon}^{-1} = P_{\rho,\varepsilon}$ . Let  $f \in A^E$ . The group isomorphism follows from the fact that at each vertex  $v$ ,

$$\sum_{e \in E} \rho(v, e)(P_{\rho,\varepsilon} f)(e) = \sum_{e \in E} \rho(v, e)\rho(v, e)\varepsilon(v, e)f(e) = \sum_{e \in E} \varepsilon(v, e)f(e).$$

Since  $P_{\rho,\varepsilon} f(e) \neq 0$  is equivalent to  $f(e) \neq 0$ , it follows that  $P_{\rho,\varepsilon} F_{\text{nz}}(G, \varepsilon; A) = F_{\text{nz}}(G, \rho; A)$ .  $\square$

Let  $T$  be a *maximal forest* of  $G$  in the sense that every component of  $T$  is a spanning tree of a component of  $G$ . For each edge  $e$  of the complement  $T^c := E - E(T)$ , let  $C_e$  denote the unique circuit contained in  $T \cup e$ , and let  $\rho_e$  be a direction of  $C_e$  (i.e.  $(C_e, \rho_e)$  is directed Eulerian) such that  $\rho_e(e) = \varepsilon(e)$ . It is easy to see that  $[\varepsilon, \rho_e]$  is a flow of  $(G, \varepsilon)$ .

**Lemma 3.3** (Berge [2]). *Let  $T$  be a maximal spanning forest of  $G$ . Then each flow  $f$  of the digraph  $(G, \varepsilon)$  can be expressed as a unique linear combination*

$$f = \sum_{e \in T^c} f(e)[\varepsilon, \rho_e]. \quad (3.4)$$

The system (1.1) for a flow  $f$ , whose equations are indexed by vertices  $v \in V$ , is equivalent to the system (3.4), which can be written as

$$f(x) = \sum_{e \in T^c} f(e)[\varepsilon, \rho_e](x), \quad x \in T, \quad (3.5)$$

whose equations are indexed by edges  $x \in T$ . In other words, the values  $f(e)$  for  $e \in T^c$  can be arbitrarily specified, and  $f(x)$  for  $x \in T$  are determined by (3.5).

Let  $n(G)$  denote the cycle rank of  $G$ . If  $T$  is a maximal spanning forest of  $G$ , then  $n(G)$  is the number of edges of  $T^c$ . Note that

$$n(G) = |E| - |V| + c(G),$$

where  $c(G)$  is the number of connected components of  $G$ . Lemma 3.3 shows that the abelian group  $F(G, \varepsilon; A)$  is of rank  $n(G)$ . The *flow arrangement* of the digraph  $(G, \varepsilon)$  with the abelian group  $A$  is the subgroup arrangement  $\mathcal{A}_{\text{FL}}(G, \varepsilon; A)$  of  $F(G, \varepsilon; A)$ , consisting of the subgroups

$$F_e \equiv F_e(G, \varepsilon; A) := \{f \in F(G, \varepsilon; A) \mid f(e) = 0\}, \quad e \in E. \quad (3.6)$$

The semilattice  $\mathcal{L}(\mathcal{A}_{\text{FL}}(G, \varepsilon; A))$  consists of the subgroups

$$F_X \equiv F_X(G, \varepsilon; A) := \{f \in F(G, \varepsilon; A) \mid f(x) = 0, x \in X\}, \quad X \subseteq E.$$

Then  $F \equiv F_\emptyset = F(G, \varepsilon; A)$ . Using (3.5), it is easy to see that the arrangement  $\mathcal{A}_{\text{FL}}(G, \varepsilon; A)$  is isomorphic to the arrangement  $\mathcal{A}_{\text{FL}}(G, T, \varepsilon; A)$  of  $A^{E(T^c)}$ , consisting of the subgroups

$$H_e := \begin{cases} \{f \in A^{E(T^c)} \mid f(e) = 0\} & \text{if } e \in T^c, \\ \{f \in A^{E(T^c)} \mid \sum_{x \in T^c} [\varepsilon, \rho_x](e)f(x) = 0\} & \text{if } e \in T. \end{cases}$$

The isomorphism is given by the restriction  $f \mapsto f|_{E(T^c)}$ , sending  $F_e$  to  $H_e$ .

**Lemma 3.4.** *The abelian group  $F(G, \varepsilon; A)$  is isomorphic to the product group  $A^{n(G)}$ . Moreover, for any subset  $X \subseteq E$ , the subgroup  $F_X(G, \varepsilon; A)$  is isomorphic to the product group  $A^{n(E-X)}$ . In particular, if  $|A| = q$  is finite, then*

$$|F(G, \varepsilon; A)| = q^{n(G)}, \quad |F_X(G, \varepsilon; A)| = q^{n(E-X)}. \quad (3.7)$$



PROOF. Trivial □

**Theorem 3.5.** *Let  $A$  be an abelian group such that either  $|A| = q$  is finite, or  $A = \mathbb{Z}$ , or  $A$  is an infinite field. Then*

$$\varphi(G, q) = \lambda \left( F(G, \varepsilon; A) - \bigcup_{e \in E} F_e \right) \Big|_{t=q} = \chi(\mathcal{A}_{\text{FL}}(G, \varepsilon; A), q). \quad (3.8)$$

Moreover,  $|\varphi(G, -1)|$  counts the number of totally cyclic orientations on  $G$ .

PROOF. Let  $X, Y \subseteq E$ . If  $|A| \geq 2$  (including  $|A| = \infty$ ), then  $F_X(G, \varepsilon; A) \subseteq F_Y(G, \varepsilon; A)$  is equivalent to that circuits of  $\langle E - X \rangle$  are contained in  $\langle E - Y \rangle$ . Thus the map  $F_X(G, \varepsilon; A) \mapsto F_X(G, \varepsilon; \mathbb{R})$  is an isomorphism from  $\mathcal{L}(\mathcal{A}_{\text{FL}}(G, \varepsilon; A))$  to  $\mathcal{L}(\mathcal{A}_{\text{FL}}(G, \varepsilon; \mathbb{R}))$ . Consequently, we have the same Möbius function

$$\mu(F_X(G, \varepsilon; A), F_Y(G, \varepsilon; A)) = \mu(F_X(G, \varepsilon; \mathbb{R}), F_Y(G, \varepsilon; \mathbb{R})).$$

If  $A$  is infinite, applying the valuation  $\lambda$  to both sides of (2.2), we have

$$\lambda \left( F - \bigcup_{e \in E} F_e \right) = \sum_{F_X \in \mathcal{L}(\mathcal{A}_{\text{FL}}(G, \varepsilon; A))} \mu(F_X, F) t^{n\langle E - X \rangle}.$$

If  $A$  is finite and  $|A| = q$ , applying the counting measure  $\#$  to both sides of (2.2), we have

$$\varphi(G, q) = \sum_{F_X \in \mathcal{L}(\mathcal{A}_{\text{FL}}(G, \varepsilon; A))} \mu(F_X, F) q^{n\langle E - X \rangle}.$$

The identity (3.8) follows immediately for positive integers  $q$ .

For  $q = 1$ , we have  $A = \{0\}$  and  $F_{\text{nz}}(G, \varepsilon; A) = \emptyset$ . Hence  $\varphi(G, 1) = 0$ . Since the hyperplane arrangement  $\mathcal{A}_{\text{FL}}(G, \varepsilon; \mathbb{R})$  is central (all hyperplanes pass through the origin), there is no relatively bounded region. Zaslavsky's second counting formula (2.5) confirms that  $\chi(\mathcal{A}_{\text{FL}}(G, \varepsilon; \mathbb{R}), 1) = 0$ .

Finally, by Zaslavsky's counting formula (2.4),  $(-1)^{n(G)}\varphi(G, -1)$  counts the number of regions of the complement  $F - \bigcup_{e \in E} F_e$ . By Lemmas 4.2 and 4.3, the regions of the complement  $F(G, \varepsilon; \mathbb{R}) - \bigcup_{e \in E} F_e(G, \varepsilon; \mathbb{R})$  correspond bijectively to the totally cyclic orientations on  $G$ . □

#### 4. Integral flow polynomials

In this section we apply the Ehrhart polynomial theory to study integral flow polynomials. Let us recall some well-known facts about lattice polytopes and Ehrhart polynomials. Let  $P$  be a relatively open lattice polytope of  $\mathbb{R}^n$ , i.e.,  $P$  is open in the flat that  $P$  spans, and the vertices of  $P$  are lattice points of  $\mathbb{Z}^n$ . The closure of  $P$  is denoted by  $\bar{P}$ . A bounded *lattice polyhedron* is a disjoint union of finitely many relatively open lattice polytopes. Let  $X$  be a bounded lattice polyhedron and  $q$  a positive integer. The *dilatation* of  $X$  by  $q$  is the polyhedron  $qX := \{qx \mid x \in X\}$ . Let

$$L(X, q) := \#(qX \cap \mathbb{Z}^n).$$

It is known that  $L(X, q)$  is a polynomial function of degree  $\dim X$  in the positive integer variable  $q$ , called the *Ehrhart polynomial* of  $X$ . Moreover, the leading coefficient of  $L(X, t)$  is the volume of  $X$ ; the constant term  $L(X, 0)$  is the Euler characteristic  $\chi(X)$ . In particular, if  $X = P$  is a relatively open lattice polytope, then  $L(P, q)$  and  $L(\bar{P}, q)$  satisfy the *Reciprocity Law*:

$$L(P, -t) = (-1)^{\dim P} L(\bar{P}, t);$$

the constant term of  $L(\bar{P}, t)$  is 1, and the constant term of  $L(P, t)$  is  $(-1)^{\dim P}$ . All these and other related properties about Ehrhart polynomials can be found in [9, 10, 21].

Flows with values in  $\mathbb{R}$  are called *real flows*; and flows with values in  $\mathbb{Z}$  are called *integer flows*. A flow  $f \in F(G, \varepsilon; \mathbb{R})$  is called a *q-flow* if  $|f(e)| < q$  for all  $e \in E$ . We define the set of all real *q*-flows of  $(G, \varepsilon)$  as

$$F(G, \varepsilon; q) := \{f \in F(G, \varepsilon; \mathbb{R}) : |f(e)| < q, e \in E\}.$$

We denote by  $F_{\mathbb{Z}}(G, \varepsilon; q)$  the set of all integer *q*-flows of  $(G, \varepsilon)$ , and by  $F_{\text{nz}}(G, \varepsilon; q)$  the set of all nowhere-zero integer *q*-flows, i.e.,

$$F_{\text{nz}}(G, \varepsilon; q) := \{f \in F_{\mathbb{Z}}(G, \varepsilon; q) \mid f(e) \neq 0, e \in E\}.$$

Clearly,  $F_{\text{nz}}(G, \varepsilon; q)$  is the set of lattice points of the dilatation  $q\Delta_{\text{FL}}(G, \varepsilon)$  (dilated by  $q$ ) of the nonconvex polyhedron

$$\Delta_{\text{FL}}(G, \varepsilon) := \{f \in F(G, \varepsilon; \mathbb{R}) : 0 < |f(e)| < 1, e \in E\}.$$

It follows that the counting function

$$\varphi_{\mathbb{Z}}(G, q) := |F_{\text{nz}}(G, \varepsilon; q)| = L(\Delta_{\text{FL}}(G, \varepsilon), q) \quad (4.1)$$

is an Ehrhart polynomial of degree  $\dim \Delta_{\text{FL}}(G, \varepsilon)$  in the positive integer variable  $q$ . In fact, we shall see that  $|F_{\text{nz}}(G, \varepsilon; q)|$  is independent of the chosen orientation  $\varepsilon$ . We call  $\varphi_{\mathbb{Z}}(G, t)$  the *integral flow polynomial* of  $G$ .

**Lemma 4.1.** *The involution  $P_{\rho, \varepsilon}$  is a group isomorphism from  $\mathbb{R}^E$  to itself. Moreover,*

$$\begin{aligned} P_{\rho, \varepsilon} \Delta_{\text{FL}}(G, \varepsilon) &= \Delta_{\text{FL}}(G, \rho), \\ P_{\rho, \varepsilon} F_{\text{nz}}(G, \varepsilon; q) &= F_{\text{nz}}(G, \rho; q). \end{aligned}$$

PROOF. Let  $A = \mathbb{R}$ ,  $f \in \mathbb{R}^E$ ,  $e \in E$ . Note that  $0 < |P_{\rho, \varepsilon} f(e)| < 1$  is equivalent to  $0 < |f(e)| < 1$ . Hence  $P_{\rho, \varepsilon} \Delta_{\text{FL}}(G, \varepsilon) = \Delta_{\text{FL}}(G, \rho)$ . Similarly,  $P_{\rho, \varepsilon} f(e) \in \mathbb{Z}$  is equivalent to  $f(e) \in \mathbb{Z}$ ; and  $0 < |P_{\rho, \varepsilon} f(e)| < q$  is equivalent to  $0 < |f(e)| < q$ . Thus  $P_{\rho, \varepsilon} F_{\text{nz}}(G, \varepsilon; q) = F_{\text{nz}}(G, \rho; q)$ .  $\square$

Let  $A = \mathbb{R}$ . The subgroup arrangement  $\mathcal{A}_{\text{FL}}(G, \varepsilon; \mathbb{R})$  is a hyperplane arrangement of  $F(G, \varepsilon; \mathbb{R})$ . The complement of  $\mathcal{A}_{\text{FL}}(G, \varepsilon; \mathbb{R})$  is the set

$$F(G, \varepsilon; \mathbb{R}) - \bigcup \mathcal{A}_{\text{FL}}(G, \varepsilon; \mathbb{R}) = \{f \in F(G, \varepsilon; \mathbb{R}) \mid f(e) \neq 0, e \in E\}.$$

For each edge  $e \in E$  with end-vertices  $u, v$ , the nonzero condition  $f(e) \neq 0$  can be split into two inequalities:

$$f(e) > 0 \quad \text{and} \quad f(e) < 0;$$

the former can be interpreted as an orientation of  $e$  agreeing with  $\varepsilon(u, e)$ , and the latter is interpreted as an orientation of  $e$  opposite to  $\varepsilon(u, e)$ .

For each orientation  $\rho \in \mathcal{O}(G)$ , we introduce the open convex cone

$$C^{\rho}(G, \varepsilon) := \{f \in F(G, \varepsilon; \mathbb{R}) : [\rho, \varepsilon](e)f(e) > 0, e \in E\}.$$

The complement  $F(G, \varepsilon; \mathbb{R}) - \bigcup \mathcal{A}_{\text{FL}}(G, \varepsilon; \mathbb{R})$  is a disjoint union of these open convex cones, some of them may be empty. By Lemma 4.2 below, the cone  $C^{\rho}(G, \varepsilon)$  is isomorphic to the open convex cone

$$C^{+}(G, \rho) := \{f \in F(G, \rho; \mathbb{R}) \mid f(e) > 0, e \in E\}.$$

We introduce the relatively open polytopes

$$\Delta_{\text{FL}}^+(G, \rho) := \{f \in F(G, \rho; \mathbb{R}) \mid 0 < f(e) < 1, e \in E\},$$

$$\Delta_{\text{FL}}^\rho(G, \varepsilon) := \{f \in F(G, \varepsilon; \mathbb{R}) \mid 0 < [\rho, \varepsilon](e)f(e) < 1, e \in E\}.$$

If  $\Delta_{\text{FL}}^+(G, \rho) \neq \emptyset$  (equivalent to that  $\rho$  is totally cyclic), then the closure of  $\Delta_{\text{FL}}^+(G, \rho)$  is the polytope

$$\bar{\Delta}_{\text{FL}}^+(G, \rho) := \{f \in F(G, \rho; \mathbb{R}) \mid 0 \leq f(e) \leq 1, e \in E\}. \quad (4.2)$$

Whether the orientation  $\rho$  is totally cyclic or not, the set  $\bar{\Delta}_{\text{FL}}^+(G, \rho)$  is always a polytope, and is called a *flow polytope of  $G$  with respect to  $\rho$* .

**Lemma 4.2.**  $P_{\rho, \varepsilon} \Delta_{\text{FL}}^\rho(G, \varepsilon) = \Delta_{\text{FL}}^+(G, \rho)$ ; and disjoint decomposition

$$\Delta_{\text{FL}}(G, \varepsilon) = \bigsqcup_{\rho \in \mathcal{O}(G)} \Delta_{\text{FL}}^\rho(G, \varepsilon). \quad (4.3)$$

PROOF. Since  $P_{\rho, \varepsilon} f = [\rho, \varepsilon]f$  for  $f \in \mathbb{R}^E$ , then the first identity is trivial by definition of  $\Delta_{\text{FL}}^\rho(G, \varepsilon)$ ,  $\Delta_{\text{FL}}^+(G, \rho)$ , and Lemma 4.1.

Let  $f \in \Delta_{\text{FL}}(G, \varepsilon)$ . We define an orientation  $\rho$  on  $G$  as follows: for each edge  $e$  at its one end-vertex  $v$ , set  $\rho(v, e) = \varepsilon(v, e)$  if  $f(e) > 0$  and  $\rho(v, e) = -\varepsilon(v, e)$  if  $f(e) < 0$ . Then  $f \in \Delta_{\text{FL}}^\rho(G, \varepsilon)$ . Conversely, each  $\Delta_{\text{FL}}^\rho(G, \varepsilon)$  is obviously contained in  $\Delta_{\text{FL}}(G, \varepsilon)$ . The union is clearly disjoint.  $\square$

Notice that the open convex cone  $C^+(G, \varepsilon)$  may be empty for the given orientation  $\varepsilon$ . If  $(G, \varepsilon)$  contains a directed cut, then it is impossible to have positive real flows by Proposition 3.1(a), thus  $C^+(G, \varepsilon) = \emptyset$ . To have  $C^+(G, \varepsilon) \neq \emptyset$ , the orientation  $\varepsilon$  must be totally cyclic.

**Lemma 4.3.** (a)  $\Delta_{\text{FL}}^+(G, \varepsilon) \neq \emptyset$  if and only if  $\varepsilon \in \mathcal{O}_{\text{TC}}(G)$ .

(b) If  $\varepsilon \in \mathcal{O}_{\text{TC}}(G)$ , then  $\bar{\Delta}_{\text{FL}}^+(G, \varepsilon)$  is a 0-1 polytope, i.e., all its vertices are flows of  $(G, \varepsilon)$  with values in  $\{0, 1\}$ .

(c) If  $\varepsilon \in \mathcal{O}_{\text{TC}}(G)$ , then every flow of  $(G, \varepsilon)$  with values in  $\{0, 1\}$  is a vertex of  $\bar{\Delta}_{\text{FL}}^+(G, \varepsilon)$ .

PROOF. (a) Trivial.

(b) Let  $f$  be a vertex of  $\bar{\Delta}_{\text{FL}}^+(G, \varepsilon)$ . It is enough to show that  $f$  is integer-valued. By Linear Programming the vertex  $(x_e = f(e) : e \in E)$  is a unique solution of a linear system of the form

$$\begin{aligned} \sum_{e \in E} \mathbf{m}_{v, e} x_e &= 0, & v \in V, \\ x_e &= a_e, & e \in E_f, \end{aligned}$$

where  $a_e = 0$  or  $1$ ,  $E_f$  is an edge subset,  $|E_f| = |E| - n(G)$ . The system is equivalent to the linear system

$$\sum_{e \in E'_f} \mathbf{m}_{v, e} x_e = b_v, \quad v \in V,$$

where  $E'_f := E - E_f$ ,  $b_v \in \mathbb{Z}$ , and the rank of the matrix  $[\mathbf{m}_{v, e}]_{V \times E'_f}$  is  $n(G)$ . Since the incidence matrix  $\mathbf{M} = [\mathbf{m}_{v, e}]_{V \times E}$  is totally unimodular (see [4], p.35), the submatrix  $[\mathbf{m}_{v, e}]_{V \times E'_f}$  is row equivalent to the matrix  $\begin{bmatrix} I \\ 0 \end{bmatrix}$  over  $\mathbb{Z}$ , where  $I$  is the identity matrix when  $V$  is linearly labeled. It then follows that the solution  $(x_e = f(e) : e \in E)$  is an integer vector.

(c) Notice that a flow of  $(G, \varepsilon)$  with values in  $\{0, 1\}$  is just the characteristic function of the edge set of a directed Eulerian subgraph of  $(G, \varepsilon)$ . Let  $f$  be a

flow with values in  $\{0, 1\}$ . Suppose  $f$  is not a vertex of  $\bar{\Delta}_{\text{FL}}^+(G, \varepsilon)$ . Then there are distinct flows  $f_i$  of  $(G, \varepsilon)$  with values in  $\{0, 1\}$  such that  $f = \sum_{i=1}^k a_i f_i$ , where  $a_i > 0$ ,  $\sum_{i=1}^k a_i = 1$ ,  $k \geq 2$ . Let  $e$  be an edge such that  $f(e) = 1$ . Then  $f_i(e) = 1$  for all  $i$ ; otherwise, say,  $f_1(e) = 0$ , then  $f(e) = \sum_{i=1}^k a_i f_i(e) \leq \sum_{i=2}^k a_i < 1$ , which contradicts  $f(e) = 1$ . Thus  $f_i = f$  for all  $i$ ; this is contradictory to the distinctness of  $f_i$ .  $\square$

Recall that  $\varphi_\varepsilon(G, q)$  for a positive integer  $q$  is the number of integer flows of  $(G, \varepsilon)$  with values in  $\{1, 2, \dots, q-1\}$ . In other words,  $\varphi_\varepsilon(G, q)$  is the number of integer flows of  $(G, \varepsilon)$  with values in the open interval  $(0, q)$ . Clearly,  $\varphi_\varepsilon(G, q)$  counts the number of lattice points of the dilatation  $q\Delta_{\text{FL}}^+(G, \varepsilon)$ , i.e.,

$$\varphi_\varepsilon(G, q) = L(\Delta_{\text{FL}}^+(G, \varepsilon), q). \quad (4.4)$$

We call  $\varphi_\varepsilon(G, q)$  the *local flow polynomial of  $G$  with respect to  $\varepsilon$* . Analogously, let  $\bar{\varphi}_\varepsilon(G, q)$  denote the number of integer flows of  $(G, \varepsilon)$  with values in  $\{0, 1, \dots, q\}$ . In other words,  $\bar{\varphi}_\varepsilon(G, q)$  is the number of integer flows of  $(G, \varepsilon)$  with values in the closed interval  $[0, q]$ . Then  $\bar{\varphi}_\varepsilon(G, q)$  counts the number of lattice points of  $q\bar{\Delta}_{\text{FL}}^+(G, \varepsilon)$ , i.e.,

$$\bar{\varphi}_\varepsilon(G, q) = L(\bar{\Delta}_{\text{FL}}^+(G, \varepsilon), q). \quad (4.5)$$

We call  $\bar{\varphi}_\varepsilon(G, q)$  the *local dual flow polynomial of  $G$  with respect to  $\varepsilon$* . Now we denote by  $\bar{\varphi}_z(G, q)$  the number of pairs  $(\rho, f)$ , where  $\rho$  is a totally cyclic orientation on  $G$  and  $f$  is an integer flow of  $(G, \rho)$  with values in  $\{0, 1, \dots, q\}$ . We call  $\bar{\varphi}_z(G, q)$  the *dual integral flow polynomial of  $G$* .

#### PROOF OF THEOREM 1.1.

(a) By Lemma 4.3, the closure  $\bar{\Delta}_{\text{FL}}^+(G, \varepsilon)$  of the open polytope  $\Delta_{\text{FL}}^+(G, \varepsilon)$  is the convex hull of the lattice points  $f \in \mathbb{Z}^E$  such that  $f(e) \in \{0, 1\}$  for all  $e \in E$  and satisfying (1.1). The Reciprocity Law and the interpretation of the constant term follow from the Reciprocity Law and the properties of Ehrhart polynomials.

(b) Note that  $F_{\text{Izz}}(G, \varepsilon; q) = q\Delta_{\text{FL}}(G, \varepsilon)$ . By Lemma 4.2, we have a disjoint union

$$q\Delta_{\text{FL}}(G, \varepsilon) = \bigsqcup_{\rho \in \mathcal{O}_{\text{TC}}(G)} q\Delta_{\text{FL}}^\rho(G, \varepsilon), \quad (4.6)$$

where each lattice open polytope  $\Delta_{\text{FL}}^\rho(G, \varepsilon)$  is isomorphic to the 0-1 open polytope  $\Delta_{\text{FL}}^+(G, \rho)$  by the unimodular transformation  $P_{\rho, \varepsilon}$ . Then (1.8) follows immediately from (4.6); (1.9) follows from definition of  $\bar{\varphi}_z(G, q)$ . The Reciprocity Law (1.10) follows from (1.7)-(1.9). The interpretation of the constant term  $\varphi_z(G, 0)$  follows from (1.8) and  $\varphi_\varepsilon(G, 0) = (-1)^{n(G)}$ .  $\square$

### 5. Interpretation of modular flow polynomial

This section is devoted to interpreting the values of the modular flow polynomial in a way similar to how the modular tension polynomial was interpreted in [11]. For the graph  $G = (V, E)$  and a positive integer  $q$ , there is a modulo  $q$  map

$$\text{Mod}_q : \mathbb{R}^E \rightarrow (\mathbb{R}/q\mathbb{Z})^E, \quad (\text{Mod}_q f)(x) = f(x) \pmod{q}, \quad f \in \mathbb{R}^E.$$

Then  $\text{Mod}_q(\mathbb{Z}^E) = (\mathbb{Z}/q\mathbb{Z})^E$  is a subgroup of the toric group  $(\mathbb{R}/q\mathbb{Z})^E$ , and  $\text{Mod}_q(F(G, \varepsilon; \mathbb{Z}))$  is a subgroup of  $\text{Mod}_q(\mathbb{Z}^E)$ .

Given orientations  $\rho, \sigma \in \mathcal{O}(G)$ ; there is an involution  $Q_{\rho, \sigma} : [0, q]^E \rightarrow [0, q]^E$  defined by

$$(Q_{\rho, \sigma} g)(e) = \begin{cases} g(e) & \text{if } \rho(v, e) = \sigma(v, e), \\ q - g(e) & \text{if } \rho(v, e) \neq \sigma(v, e), \end{cases}$$

where  $g \in [0, q]^E$  and  $v$  is an end-vertex of the edge  $e$ . Clearly,  $Q_{\rho, \sigma}$  is a bijection from  $[0, q]^E$  to  $[0, q]^E$ , and is also a bijection from  $(0, q)^E$  to  $(0, q)^E$ , where  $(0, q) = \{x \in \mathbb{R} \mid 0 < x < q\}$ . Moreover,  $Q_{\rho, \sigma} Q_{\sigma, \varepsilon} = Q_{\rho, \varepsilon}$ .

Recall that two orientations  $\varepsilon_1, \varepsilon_2 \in \mathcal{O}(G)$  are said to be *Eulerian equivalent*, written  $\varepsilon_1 \sim \varepsilon_2$ , if the induced spanning subdigraph by the edge subset

$$E(\varepsilon_1 \neq \varepsilon_2) := \{e \in E \mid \varepsilon_1(v, e) \neq \varepsilon_2(v, e), v \text{ is an end-vertex of } e\}$$

is a directed Eulerian subgraph with the orientation either  $\varepsilon_1$  or  $\varepsilon_2$ .

- Lemma 5.1.** (a) *The relation  $\sim$  is an equivalence relation on  $\mathcal{O}(G)$ .*  
 (b) *Let  $\rho, \sigma \in \mathcal{O}(G)$  be Eulerian equivalent. If  $\rho$  is totally cyclic, so is  $\sigma$ .*  
 (c) *Let  $\rho, \sigma \in \mathcal{O}(G)$  and  $\rho \sim \sigma$ . Then  $Q_{\rho, \sigma} : q\bar{\Delta}_{\text{FL}}^+(G, \sigma) \rightarrow q\bar{\Delta}_{\text{FL}}^+(G, \rho)$  is a bijection, sending lattice points to lattice points. In particular,*

$$\begin{aligned} Q_{\rho, \sigma}(q\Delta_{\text{FL}}^+(G, \sigma)) &= q\Delta_{\text{FL}}^+(G, \rho), \\ \varphi_\rho(G, q) &= \varphi_\sigma(G, q), \\ \bar{\varphi}_\rho(G, \varepsilon; q) &= \bar{\varphi}_\sigma(G, q). \end{aligned}$$

PROOF. (a) The reflexivity and the symmetric property are obvious. Transitivity is a straightforward computation. Note that a digraph  $(H, \rho)$  is Eulerian if and only if for all  $v \in V(H)$ ,

$$\sum_{e \in E(H)} \rho(v, e) = 0.$$

Let  $\varepsilon_i \in \mathcal{O}(G)$  ( $i = 1, 2, 3$ ) be such that  $\varepsilon_1 \sim \varepsilon_2$  and  $\varepsilon_2 \sim \varepsilon_3$ . Then

$$\sum_{e \in E(\varepsilon_1 \neq \varepsilon_2)} \varepsilon_2(v, e) = \sum_{e \in E(\varepsilon_2 \neq \varepsilon_3)} \varepsilon_2(v, e) = 0.$$

Thus

$$\begin{aligned} \sum_{e \in E(\varepsilon_1 \neq \varepsilon_3)} \varepsilon_1(v, e) &= \sum_{e \in E(\varepsilon_1 = \varepsilon_2 \neq \varepsilon_3)} \varepsilon_1(v, e) + \sum_{e \in E(\varepsilon_1 \neq \varepsilon_2 = \varepsilon_3)} \varepsilon_1(v, e) \\ &= \sum_{e \in E(\varepsilon_1 = \varepsilon_2 \neq \varepsilon_3)} \varepsilon_2(v, e) - \sum_{e \in E(\varepsilon_1 \neq \varepsilon_2 = \varepsilon_3)} \varepsilon_2(v, e) \\ &= \sum_{e \in E(\varepsilon_1 = \varepsilon_2 \neq \varepsilon_3) \sqcup E(\varepsilon_1 \neq \varepsilon_2 \neq \varepsilon_3)} \varepsilon_2(v, e) \\ &\quad - \sum_{e \in E(\varepsilon_1 \neq \varepsilon_2 = \varepsilon_3) \sqcup E(\varepsilon_1 \neq \varepsilon_2 \neq \varepsilon_3)} \varepsilon_2(v, e) \\ &= \sum_{e \in E(\varepsilon_2 \neq \varepsilon_3)} \varepsilon_2(v, e) - \sum_{e \in E(\varepsilon_1 \neq \varepsilon_2)} \varepsilon_2(v, e) = 0. \end{aligned}$$

This means that  $\varepsilon_1$  is Eulerian equivalent to  $\varepsilon_3$ .

(b) Suppose  $(G, \rho)$  contains a directed cut  $(U, \varepsilon_U)$ , where  $U = [S, S^c]$  with  $S \subseteq V$ . Since  $E(\rho \neq \sigma)$  is directed Eulerian with the orientation  $\rho$ , then by Proposition 3.1(b),

$$\sum_{x \in U \cap E(\rho \neq \sigma)} [\rho, \varepsilon_U](x) = 0.$$

Since  $[\rho, \varepsilon_U] \equiv 1$ , it follows that  $U \cap E(\rho \neq \sigma) = \emptyset$ . This means that  $\rho(v, e) = \sigma(v, e)$  for all edges  $e \in U$ , where  $v$  is an end-vertex of  $e$  and  $v \in S$ . So  $(U, \varepsilon_U)$  is a directed cut of  $(G, \varepsilon)$ . This is a contradiction.

(c) For a flow  $f \in q\bar{\Delta}_{\text{FL}}^+(G, \sigma)$  ( $f \in q\Delta_{\text{FL}}^+(G, \sigma)$ ), we have

$$\sum_{e \in E(v)} \rho(v, e)(Q_{\rho, \sigma} f)(e) = \sum_{e \in E(v)} \sigma(v, e)f(e) + q \sum_{e \in E(\sigma \neq \rho)} \rho(v, e) = 0.$$

This shows that  $Q_{\rho, \sigma} f \in q\bar{\Delta}_{\text{FL}}^+(G, \rho)$  ( $Q_{\rho, \sigma} f \in q\Delta_{\text{FL}}^+(G, \rho)$ ). Clearly,  $Q_{\rho, \sigma}$  sends lattice points to lattice points by definition. Therefore,  $\bar{\varphi}_\rho(G, q) = \bar{\varphi}_\sigma(G, q)$  and  $\varphi_\rho(G, q) = \varphi_\sigma(G, q)$ .  $\square$

For two Eulerian equivalent orientations  $\rho, \sigma$ , we have seen that the digraph  $(G, \rho)$  contains no directed cut if and only if  $(G, \sigma)$  contains no directed cut. This means that  $\sim$  induces an equivalence relation on  $\mathcal{O}_{\text{TC}}(G)$ ; and each equivalence class of  $\sim$  in  $\mathcal{O}_{\text{TC}}(G)$  is an equivalence class of  $\sim$  in  $\mathcal{O}(G)$ . We denote by  $[\mathcal{O}_{\text{TC}}(G)]$  the quotient set  $\mathcal{O}_{\text{TC}}(G)/\sim$  of Eulerian equivalence classes. For each  $\rho \in \mathcal{O}_{\text{TC}}(G)$ , let  $[\rho] \in [\mathcal{O}_{\text{TC}}(G)]$  denote the equivalence class of  $\rho$ , and define

$$\varphi_{[\rho]}(G, q) = \varphi_\rho(G, q) \quad \text{and} \quad \bar{\varphi}_{[\rho]}(G, q) = \bar{\varphi}_\rho(G, q).$$

The following nontrivial Lemma is due to Tutte. It is crucial to the proof of our main result Theorem 1.2; so we present a proof with our notations.

**Lemma 5.2** (Tutte [23]). *The map  $\text{Mod}_q : F_{\mathbb{Z}}(G, \varepsilon; q) \rightarrow F(G, \varepsilon; \mathbb{Z}/q\mathbb{Z})$  and its restriction  $\text{Mod}_q : F_{\text{nz}}(G, \varepsilon; q) \rightarrow F_{\text{nz}}(G, \varepsilon; \mathbb{Z}/q\mathbb{Z})$  are surjective.*

**PROOF.** The second part of the lemma implies the first part. In fact, every flow  $f \in F(G, \varepsilon; \mathbb{Z}/q\mathbb{Z})$  can be viewed as a nowhere-zero flow  $f|_{E_f}$  on the subdigraph  $(V, E_f, \varepsilon)$ , where  $E_f := \{e \in E \mid f(e) \neq 0\}$ . Let  $g$  be a nowhere-zero integer  $q$ -flow on  $(V, E_f, \varepsilon)$  such that  $\text{Mod}_q(g) = f|_{E_f}$ . Then  $g$  is extended to an integer  $q$ -flow on  $(G, \varepsilon)$  by setting  $g \equiv 1$  on  $E - E_f$ .

Now for each  $f \in F_{\mathbb{Z}}(G, \varepsilon; q)$  we write  $\tilde{f} = \text{Mod}_q f$ . We identify  $\mathbb{Z}/q\mathbb{Z}$  with the set  $\{0, 1, \dots, q-1\}$  and view each modular flow  $g \in F(G, \varepsilon; \mathbb{Z}/q\mathbb{Z})$  as an integer-valued function  $g : E \rightarrow \{0, 1, \dots, q-1\}$ . Then the map  $Q_{\rho, \varepsilon}$  maps  $F(G, \varepsilon; \mathbb{Z}/q\mathbb{Z})$  to  $F(G, \rho; \mathbb{Z}/q\mathbb{Z})$ , and  $F_{\text{nz}}(G, \varepsilon; \mathbb{Z}/q\mathbb{Z})$  to  $F_{\text{nz}}(G, \rho; \mathbb{Z}/q\mathbb{Z})$ . For each  $g \in [0, q]^E$  and  $\rho \in \mathcal{O}(G)$ , we define

$$\eta(g, \rho) := \sum_{v \in V} \left| \sum_{e \in E} \rho(v, e)g(e) \right|.$$

Fix a modular flow  $\tilde{f} \in F_{\text{nz}}(G, \varepsilon; \mathbb{Z}/q\mathbb{Z})$ . Let  $\rho^*$  be a particular orientation on  $G$  such that

$$\eta(Q_{\rho^*, \varepsilon} \tilde{f}, \rho^*) = \min\{\eta(Q_{\rho, \varepsilon} \tilde{f}, \rho) \mid \rho \in \mathcal{O}(G)\}.$$

We write  $f^* := Q_{\rho^*, \varepsilon} \tilde{f}$  and define  $f := P_{\varepsilon, \rho^*} f^*$ .

If  $\eta(f^*, \rho^*) = 0$ , then  $\sum_{e \in E} \rho^*(v, e) f^*(e) = 0$  for all  $v \in V$ . This means that  $f^*$  is an integer  $q$ -flow of  $(G, \rho^*)$ . Whence  $f$  is an integer  $q$ -flow of  $(G, \varepsilon)$ . By definition of  $P_{\varepsilon, \rho^*}$  and  $Q_{\rho^*, \varepsilon}$ , we see that

$$\begin{aligned} f(e) &= \begin{cases} \tilde{f}(e) & \text{if } \varepsilon(v, e) = \rho^*(v, e) \\ \tilde{f}(e) - q & \text{if } \varepsilon(v, e) \neq \rho^*(v, e) \end{cases} \\ &= \tilde{f}(e) \pmod{q} \quad \text{for all } e \in E. \end{aligned}$$

The surjectivity of  $\text{Mod}_q : F_{\text{nz}}(G, \varepsilon; q) \rightarrow F_{\text{nz}}(G, \varepsilon; \mathbb{Z}/q\mathbb{Z})$  follows immediately.

We now claim that  $\eta(f^*, \rho^*) = 0$ . Suppose  $\eta(f^*, \rho^*) > 0$ . Then there exist a vertex  $u \in V$  and an integer  $k$  such that

$$\sum_{e \in E} \rho^*(u, e) f^*(e) = kq > 0 \quad (< 0). \quad (5.1)$$

Note that for any function  $g : E \rightarrow A$ , where  $A$  is an abelian group, and for any orientation  $\rho \in \mathcal{O}(G)$ , we have

$$\sum_{v \in V} \sum_{e \in E} \rho(v, e) g(e) = 0.$$

In particular, for the function  $f^*$  and the orientation  $\rho^*$ , we have

$$\sum_{v \in V} \sum_{e \in E} \rho^*(v, e) f^*(e) = 0.$$

Since (5.1) and  $f^*$  is a modular flow, there exists a vertex  $w$  such that

$$\sum_{e \in E} \rho^*(w, e) f^*(e) = -lq < 0 \quad (> 0).$$

Thus there is a path  $P = v_0 e_1 v_1 \cdots e_n v_n$  with  $v_0 = u$  and  $v_n = w$ , such that  $\rho^*(v_0, e_1) = 1$ ,  $\rho^*(v_n, e_n) = -1$ , and  $\rho^*(v_i, e_i) \rho^*(v_i, e_{i+1}) = -1$ , where  $1 \leq i \leq n-1$ . Let  $\rho^{**}$  be an orientation on  $G$  given by

$$\rho^{**}(v, e) = \begin{cases} -\rho^*(v, e) & \text{if } e = e_i \text{ for some } 1 \leq i \leq n, \\ \rho^*(v, e) & \text{otherwise.} \end{cases}$$

We write  $f^{**} := Q_{\rho^{**}, \rho^*} f^*$ . Then  $f^{**} = Q_{\rho^{**}, \rho^*} Q_{\rho^*, \varepsilon} \tilde{f} = Q_{\rho^{**}, \varepsilon} \tilde{f}$ .

Notice that at each vertex  $v \in V$ , we have

$$\sum_{e \in E} \rho^{**}(v, e) f^{**}(e) = \sum_{e \in E} \rho^*(v, e) f^*(e) - q \sum_{e \in E(\rho^* \neq \rho^{**})} \rho^*(v, e).$$

In particular, for the vertices  $u, w$ , and other vertices  $v$  different from  $u$  and  $w$ , we have

$$\begin{aligned} \sum_{e \in E} \rho^{**}(u, e) f^{**}(e) &= \sum_{e \in E} \rho^*(u, e) f^*(e) - q = (k-1)q, \\ \sum_{e \in E} \rho^{**}(w, e) f^{**}(e) &= \sum_{e \in E} \rho^*(w, e) f^*(e) + q = (1-l)q, \\ \sum_{e \in E} \rho^{**}(v, e) f^{**}(e) &= \sum_{e \in E} \rho^*(v, e) f^*(e). \end{aligned}$$

It follows that

$$\eta(f^{**}, \rho^{**}) = \eta(f^*, \rho^*) - 2q < \eta(f^*, \rho^*).$$

This is contradictory to the selection of  $\rho^*$  that  $\eta(f^*, \rho^*)$  is minimum.  $\square$

For each real-valued function  $f : E \rightarrow \mathbb{R}$  and any orientation  $\rho$  on  $G$ , we associate with  $f$  and  $\rho$  an orientation  $\rho_f$ , defined for each  $(v, e) \in V \times E$  by

$$\rho_f(v, e) = \begin{cases} \rho(v, e) & \text{if } f(e) > 0, \\ -\rho(v, e) & \text{if } f(e) \leq 0. \end{cases} \quad (5.2)$$

For two orientations  $\rho, \sigma \in \mathcal{O}(G)$ , we associate a *symmetric difference function*  $I_{\rho, \sigma} : E \rightarrow \{0, 1\}$ , defined for each edge  $e \in E$  (and its one end-vertex  $v$ ) by

$$I_{\rho, \sigma}(e) = \begin{cases} 0 & \text{if } \rho(v, e) = \sigma(v, e), \\ 1 & \text{if } \rho(v, e) \neq \sigma(v, e). \end{cases} \quad (5.3)$$

**Lemma 5.3.** *Let  $f_1, f_2 \in F(G, \varepsilon; q)$ . If  $f_1(e) \equiv f_2(e) \pmod{q}$  for all edges  $e \in E$ , then  $\varepsilon_{f_1}$  and  $\varepsilon_{f_2}$  are Eulerian equivalent.*

PROOF. Let us simply write  $\varepsilon_i := \varepsilon_{f_i}$ ,  $i = 1, 2$ . It suffices to show that at each vertex  $v$ ,

$$\sum_{e \in E(\varepsilon_1 \neq \varepsilon_2)} \varepsilon_2(v, e) = 0. \quad (5.4)$$

Since  $f_1(e) \equiv f_2(e) \pmod{q}$  for all  $e \in E$ , then

$$f_1(e) = f_2(e) + a_e q, \quad e \in E,$$

where  $a_e \in \{-1, 0, 1\}$ . More precisely, for each edge  $e$  at its one end-vertex  $v$ ,

$$f_1(e) = \begin{cases} f_2(e) & \text{if } \varepsilon_1(v, e) = \varepsilon_2(v, e), \\ f_2(e) - q & \text{if } \varepsilon_1(v, e) \neq \varepsilon_2(v, e) = \varepsilon(v, e), \\ f_2(e) + q & \text{if } \varepsilon_1(v, e) \neq \varepsilon_2(v, e) \neq \varepsilon(v, e). \end{cases}$$

Let  $g_i = P_{\varepsilon_i, \varepsilon} f_i$ ,  $i = 1, 2$ . Then  $g_i$  are real  $q$ -flows of  $(G, \varepsilon_i)$ , and for an edge  $e$  at its one end-vertex  $v$ ,

$$g_1(e) = \begin{cases} g_2(e) & \text{if } \varepsilon_1(v, e) = \varepsilon_2(v, e), \\ q - g_2(e) & \text{if } \varepsilon_1(v, e) \neq \varepsilon_2(v, e). \end{cases}$$

Note that at each vertex  $v \in V$ ,

$$\sum_{e \in E} \varepsilon_i(v, e) g_i(e) = 0, \quad i = 1, 2. \quad (5.5)$$

Now consider the case of  $i = 1$  in (5.5). Replace  $g_1(e)$  by  $g_2(e)$  if  $\varepsilon_1(v, e) = \varepsilon_2(v, e)$  and by  $q - g_2(e)$  if  $\varepsilon_1(v, e) \neq \varepsilon_2(v, e)$ ; we obtain

$$\begin{aligned} & \sum_{e \in E(\varepsilon_1 = \varepsilon_2)} \varepsilon_2(v, e) g_2(e) - \sum_{e \in E(\varepsilon_1 \neq \varepsilon_2)} \varepsilon_2(v, e) (q - g_2(e)) \\ &= \sum_{e \in E} \varepsilon_2(v, e) g_2(e) - q \sum_{e \in E(\varepsilon_1 \neq \varepsilon_2)} \varepsilon_2(v, e) = 0. \end{aligned}$$

Applying (5.5) for  $i = 2$ , we see that (5.4) is true.  $\square$

**Lemma 5.4.** *Let  $\rho, \sigma, \omega \in \mathcal{O}_{\text{TC}}(G)$  be Eulerian equivalent orientations, and let  $f \in q\Delta_{\text{FL}}^\rho(G, \varepsilon)$  be a real  $q$ -flow. Then*

- (a)  $\varepsilon_f = \rho$ .
- (b)  $P_{\varepsilon, \sigma} Q_{\sigma, \rho} P_{\rho, \varepsilon} (q\Delta_{\text{FL}}^\rho(G, \varepsilon)) = q\Delta_{\text{FL}}^\sigma(G, \varepsilon)$ .
- (c)  $P_{\varepsilon, \sigma} Q_{\sigma, \rho} P_{\rho, \varepsilon} f = P_{\varepsilon, \omega} Q_{\omega, \rho} P_{\rho, \varepsilon} f$  if and only if  $\sigma = \omega$ .
- (d)  $F(G, \varepsilon; q) \cap \text{Mod}_q^{-1}(\text{Mod}_q f) = \{P_{\varepsilon, \alpha} Q_{\alpha, \rho} P_{\rho, \varepsilon} f \mid \alpha \sim \rho\}$ .



PROOF. (a) By definition of  $\Delta_{\text{FL}}^\rho(G, \varepsilon)$  and the fact  $f \in q\Delta_{\text{FL}}^\rho(G, \varepsilon)$ , we have  $[\rho, \varepsilon](e)f(e) > 0$  for all  $e \in E$ . So for each edge  $e$  at its one end-vertex  $v$ ,  $\rho(v, e) = \varepsilon(v, e)$  if  $f(e) > 0$ , and  $\rho(v, e) = -\varepsilon(v, e)$  if  $f(e) < 0$ . By definition of  $\varepsilon_f$ , we see that  $\varepsilon_f = \rho$ .

(b) Recall Lemma 4.2 and Lemma 5.1(c) that

$$P_{\rho, \varepsilon}(q\Delta_{\text{FL}}^\rho(G, \varepsilon)) = q\Delta_{\text{FL}}^+(G, \rho) \quad \text{and} \quad Q_{\sigma, \rho}(q\Delta_{\text{FL}}^+(G, \rho)) = q\Delta_{\text{FL}}^+(G, \sigma).$$

Since  $P_{\varepsilon, \sigma}$  is an involution, then  $P_{\varepsilon, \sigma}(q\Delta_{\text{FL}}^+(G, \sigma)) = q\Delta_{\text{FL}}^\sigma(G, \varepsilon)$ . The identity follows immediately by composition.

(c) The sufficiency is trivial. For necessity, we write  $g := P_{\varepsilon, \sigma}Q_{\sigma, \rho}P_{\rho, \varepsilon}f$  and  $h := P_{\varepsilon, \omega}Q_{\omega, \rho}P_{\rho, \varepsilon}f$ . Since  $g \in q\Delta_{\text{FL}}^\sigma(G, \varepsilon)$  and  $h \in q\Delta_{\text{FL}}^\omega(G, \varepsilon)$ , we have  $\varepsilon_g = \sigma$  and  $\varepsilon_h = \omega$ . Clearly, if  $g = h$ , then  $\sigma = \omega$ .

(d) Let us write  $g := P_{\varepsilon, \alpha}Q_{\alpha, \rho}P_{\rho, \varepsilon}f$  for an orientation  $\alpha$  such that  $\alpha \sim \rho$ . By definition of  $P_{\varepsilon, \alpha}, Q_{\alpha, \rho}, P_{\rho, \varepsilon}$ , we have  $g \in \Delta_{\text{FL}}^\alpha(G, \varepsilon)$  and

$$g(e) = f(e) + a_e q, \quad e \in E,$$

where  $a_e \in \{-1, 0, 1\}$ . Clearly,  $g(e) \equiv f(e) \pmod{q}$  for all  $e \in E$ ; namely,  $g \in \text{Mod}_q^{-1}(\text{Mod}_q f)$ . Conversely, let  $h \in F(G, \varepsilon; q)$  be such that  $\text{Mod}_q h = \text{Mod}_q f$ . We have

$$h(e) = f(e) + b_e q, \quad e \in E,$$

where  $b_e \in \{-1, 0, 1\}$ . By definition of  $\varepsilon_f, \varepsilon_h$ , a straightforward calculation shows that for each non-loop edge  $e$  (and its one end-vertex  $v$ ),

$$h(e) = \begin{cases} f(e) & \text{if } h(e) > 0, f(e) > 0 \ (\Leftrightarrow \varepsilon_h(v, e) = \varepsilon_f(v, e) = \varepsilon(v, e)), \\ f(e) & \text{if } h(e) \leq 0, f(e) \leq 0 \ (\Leftrightarrow \varepsilon_h(v, e) = \varepsilon_f(v, e) \neq \varepsilon(v, e)), \\ f(e) - q & \text{if } h(e) \leq 0, f(e) > 0 \ (\Leftrightarrow \varepsilon_h(v, e) \neq \varepsilon_f(v, e) = \varepsilon(v, e)), \\ f(e) + q & \text{if } h(e) > 0, f(e) \leq 0 \ (\Leftrightarrow \varepsilon_h(v, e) \neq \varepsilon_f(v, e) \neq \varepsilon(v, e)). \end{cases}$$

By definition of  $P_{\varepsilon, \varepsilon_h}, Q_{\varepsilon_h, \varepsilon_f}, P_{\varepsilon_f, \varepsilon}$ , another straightforward calculation shows that for each non-loop edge  $e$  (and its one end-vertex  $v$ ),

$$(P_{\varepsilon, \varepsilon_h}Q_{\varepsilon_h, \varepsilon_f}P_{\varepsilon_f, \varepsilon}f)(e) = \begin{cases} f(e) & \text{if } \varepsilon_h(v, e) = \varepsilon_f(v, e) = \varepsilon(v, e), \\ f(e) & \text{if } \varepsilon_h(v, e) = \varepsilon_f(v, e) \neq \varepsilon(v, e), \\ f(e) - q & \text{if } \varepsilon_h(v, e) \neq \varepsilon_f(v, e) = \varepsilon(v, e), \\ f(e) + q & \text{if } \varepsilon_h(v, e) \neq \varepsilon_f(v, e) \neq \varepsilon(v, e). \end{cases}$$

This means that  $h = P_{\varepsilon, \varepsilon_h}Q_{\varepsilon_h, \varepsilon_f}P_{\varepsilon_f, \varepsilon}f$ . Since  $\varepsilon_f = \rho$  by Part (a), thus  $h = P_{\varepsilon, \alpha}Q_{\alpha, \rho}P_{\rho, \varepsilon}f$  with  $\alpha = \varepsilon_h$ .  $\square$

**Proposition 5.5.** *The number of orientations on  $G$  that are Eulerian equivalent to  $\varepsilon$  is the number of 0-1 flows of  $(G, \varepsilon)$ , i.e.,*

$$\#[\varepsilon] = \bar{\varphi}_\varepsilon(G, 1) = |\bar{\Delta}_{\text{FL}}^+(G, \varepsilon) \cap \mathbb{Z}^E|. \quad (5.6)$$

PROOF. It is enough to show that the following map

$$[\varepsilon] = \{\rho \in \mathcal{O}(G) \mid \rho \sim \varepsilon\} \rightarrow \bar{\Delta}_{\text{FL}}^+(G, \varepsilon) \cap \mathbb{Z}^E, \quad \rho \mapsto I_{\rho, \varepsilon},$$

is a bijection. Fix an orientation  $\rho$  that is Eulerian equivalent to  $\varepsilon$ . Note that  $I_{\rho, \varepsilon}$  is the characteristic function of the edge subset  $E(\rho \neq \varepsilon)$ . Since  $E(\rho \neq \varepsilon)$  is directed Eulerian with the orientation  $\varepsilon$ . Then  $I_{\rho, \varepsilon}$  is a flow of  $(G, \varepsilon)$  with values in  $\{0, 1\}$ . So the map  $\rho \mapsto I_{\rho, \varepsilon}$  is well-defined, and is clearly injective. Conversely, given a 0-1 flow  $f$  of  $(G, \varepsilon)$ . Let  $\rho$  be an orientation on  $G$  defined by  $\rho(v, e) = \varepsilon(v, e)$  if  $f(e) = 0$  and  $\rho(v, e) = -\varepsilon(v, e)$  if  $f(e) = 1$ , where  $v$  is an end-vertex of the edge  $e$ . Then

$E(\rho \neq \varepsilon) = \{e \in E(G) \mid f(e) = 1\}$ , which is directed Eulerian with the orientation  $\varepsilon$ . This means that  $\rho \sim \varepsilon$  and  $I_{\rho, \varepsilon} = f$ . Hence the map is surjective.  $\square$

PROOF OF THEOREM 1.2.

It has been shown that  $\varphi(G, q)$  is a polynomial function of degree  $n(G)$ . Fix an orientation  $\rho \in \mathcal{O}_{\text{TC}}(G)$ . For each orientation  $\sigma \sim \rho$  and any  $f \in q\Delta_{\text{FL}}^\sigma(G, \varepsilon)$ , we have

$$\begin{aligned} |F(G, \varepsilon; q) \cap \text{Mod}_q^{-1}(\text{Mod}_q f)| &= \#\{P_{\varepsilon, \alpha} Q_{\alpha, \rho} P_{\rho, \varepsilon} f \mid \alpha \sim \sigma\} \\ &= \#[\sigma] = \#[\rho]. \end{aligned} \quad (5.7)$$

Now apply Parts (b) and (d) of Lemma 5.4; we have the disjoint unions

$$\begin{aligned} \bigsqcup_{\sigma \in [\rho]} q\Delta_{\text{FL}}^\sigma(G, \varepsilon) &= \bigsqcup_{\sigma \in [\rho]} P_{\varepsilon, \sigma} Q_{\sigma, \rho} P_{\rho, \varepsilon} (q\Delta_{\text{FL}}^\rho(G, \varepsilon)) \\ &= \bigsqcup_{\sigma \in [\rho], f \in q\Delta_{\text{FL}}^\rho(G, \varepsilon)} \{P_{\varepsilon, \sigma} Q_{\sigma, \rho} P_{\rho, \varepsilon} f\} \\ &= \bigsqcup_{f \in q\Delta_{\text{FL}}^\rho(G, \varepsilon)} F(G, \varepsilon; q) \cap \text{Mod}_q^{-1}(\text{Mod}_q f) \\ &= F(G, \varepsilon; q) \cap \text{Mod}_q^{-1} \text{Mod}_q (q\Delta_{\text{FL}}^\rho(G, \varepsilon)). \end{aligned}$$

Note that the above orientation  $\rho$  can be replaced by any orientation  $\sigma$  that is Eulerian equivalent to  $\rho$ . We further have

$$\bigsqcup_{\sigma \in [\rho]} q\Delta_{\text{FL}}^\sigma(G, \varepsilon) = F(G, \varepsilon; q) \cap \text{Mod}_q^{-1} \text{Mod}_q \left( \bigsqcup_{\sigma \in [\rho]} q\Delta_{\text{FL}}^\sigma(G, \varepsilon) \right). \quad (5.8)$$

On the one hand, recall  $\varphi_\sigma(G, q) = \varphi_\rho(G, q)$  whenever  $\sigma \sim \rho$  (see Lemma 5.1); then the number of lattice points in the left-hand side of (5.8) is

$$\varphi_\rho(G, q) \cdot \#[\rho].$$

On the other hand, (5.7) implies that the number of lattice points of the right-hand side of (5.8) is

$$\left| \text{Mod}_q \left( \bigsqcup_{\sigma \in [\rho]} q\Delta_{\text{FL}}^\sigma(G, \varepsilon) \cap \mathbb{Z}^E \right) \right| \cdot \#[\rho].$$

It then follows that

$$\left| \text{Mod}_q \left( \bigsqcup_{\sigma \in [\rho]} q\Delta_{\text{FL}}^\sigma(G, \varepsilon) \cap \mathbb{Z}^E \right) \right| = \varphi_\rho(G, q).$$

Note that (4.3) implies the disjoint decomposition

$$F_{\text{NZZ}}(G, \varepsilon; q) = \bigsqcup_{\rho \in [\mathcal{O}_{\text{TC}}(G)]} \bigsqcup_{\sigma \in [\rho]} q\Delta_{\text{FL}}^\sigma(G, \varepsilon) \cap \mathbb{Z}^E.$$

Applying the map  $\text{Mod}_q$ , we obtain the following disjoint decomposition

$$\begin{aligned} F_{\text{Nz}}(G, \varepsilon; \mathbb{Z}/q\mathbb{Z}) &= \text{Mod}_q(F_{\text{NZZ}}(G, \varepsilon; q)) \\ &= \bigsqcup_{[\rho] \in [\mathcal{O}_{\text{TC}}(G)]} \text{Mod}_q \left( \bigsqcup_{\sigma \in [\rho]} q\Delta_{\text{FL}}^\sigma(G, \varepsilon) \cap \mathbb{Z}^E \right). \end{aligned} \quad (5.9)$$

The first equation follows from the surjectivity of  $\text{Mod}_q$  by Lemma 5.2. To see the disjointness of the union in the second equation, suppose the union is not disjoint. This means that there exist integer flows  $f_i \in q\Delta_{\text{Fl}}^{\sigma_i}(G, \varepsilon)$  and orientations  $\rho_i$  such that  $\sigma_i \sim \rho_i$  ( $i = 1, 2$ ),  $[\rho_1] \neq [\rho_2]$ , and  $\text{Mod}_q(f_1) = \text{Mod}_q(f_2)$ . Then Lemma 5.4(a) implies that  $\varepsilon_{f_i} = \sigma_i$ , and Lemma 5.3 implies that  $\varepsilon_{f_1} \sim \varepsilon_{f_2}$ . It follows by transitivity that  $\rho_1 \sim \rho_2$ , i.e.,  $[\rho_1] = [\rho_2]$ . This is a contradiction.

Counting the number of elements of both sides of (5.9), we obtain

$$\varphi(G, q) = \sum_{[\rho] \in [\mathcal{O}_{\text{TC}}(G)]} \varphi_\rho(G, q).$$

The Reciprocity Law (1.12) follows from the Reciprocity Law (1.7) and the definition of  $\bar{\varphi}(G, q)$ .

Let  $q = 0$ . We have  $\varphi(G, 0) = (-1)^{n(G)} \bar{\varphi}(G, 0)$  by (1.12). Since  $\bar{\varphi}_\rho(G, 0) = 1$  for all  $\rho \in \mathcal{O}_{\text{TC}}(G)$ , we see that  $\bar{\varphi}(G, 0) = \#\mathcal{O}_{\text{TC}}(G)$  by (1.14). Hence  $\varphi(G, 0) = (-1)^{n(G)} \#\mathcal{O}_{\text{TC}}(G)$ .  $\square$

Searching online we found a result on modular flow reciprocity by Breuer and Sanyal [5], which states that  $(-1)^{n(G)} \varphi(G, -q)$  counts the numbers of pairs  $(f, \sigma)$ , where  $f$  is a flow of  $(G, \varepsilon)$  modulo  $q$  and  $\sigma$  is a totally cyclic reorientation of the digraph  $(G/\text{supp } f, \varepsilon)$ . The result can be written as the sum

$$\varphi(G, -q) = (-1)^{n(G)} \sum_{X \subseteq E} \varphi(\langle X \rangle, q) |\mathcal{O}_{\text{TC}}(G/X)|, \quad (5.10)$$

where  $G/X$  is the graph obtained from  $G$  by contracting the edges of  $X$ . The term  $\varphi(\langle X \rangle, q) |\mathcal{O}_{\text{TC}}(G/X)|$  in (5.10) is nonzero if and only if the graphs  $\langle X \rangle$  and  $G/X$  are bridgeless. The formula (5.10) can be argued straightforward as follows.

Notice the trivial fact that each flow  $f$  corresponds to a nowhere-zero flow on its support  $\text{supp}(f) := \{e \in E \mid f(e) \neq 0\}$ . This means that

$$t^{n(X)} = \sum_{Y \subseteq X} \varphi(\langle Y \rangle, t), \quad X \subseteq E.$$

The Möbius inversion implies

$$\varphi(\langle X \rangle, t) = \sum_{Y \subseteq X} (-1)^{|X-Y|} t^{n(Y)}, \quad X \subseteq E.$$

In particular, for  $X = E$  and  $t = -q$ , we have

$$\begin{aligned} \varphi(G, -q) &= \sum_{Y \subseteq E} (-1)^{|E-Y|+n(Y)} q^{n(Y)} \\ &= \sum_{Y \subseteq E} (-1)^{|E-Y|+n(Y)} \sum_{X \subseteq Y} \varphi(\langle X \rangle, q) \\ &= \sum_{X \subseteq E} \varphi(\langle X \rangle, q) \sum_{X \subseteq Y \subseteq E} (-1)^{|E-Y|+n(Y)}. \end{aligned}$$

Since  $n\langle Y \rangle = n\langle X \rangle + n\langle Y \rangle / X$  for all edge subsets  $Y$  of  $G$  such that  $X \subseteq Y$ , we see that for each fixed edge subset  $X \subseteq E$ ,

$$\begin{aligned} \sum_{X \subseteq Y \subseteq E} (-1)^{|E-Y|+n\langle Y \rangle} &= \sum_{X \subseteq Y \subseteq E} (-1)^{|E-Y|+n\langle X \rangle+n\langle Y \rangle / X} \\ &= (-1)^{n\langle X \rangle} \sum_{Z \subseteq E(G/X)} (-1)^{|E(G/X)-Z|+n\langle Z \rangle} \\ &= (-1)^{n\langle X \rangle+n\langle G/X \rangle} |\mathcal{O}_{\text{TC}}(G/X)|. \end{aligned}$$

The above last equality follows from the Zaslavsky formula (2.4) about the flow arrangement  $\mathcal{A}_{\text{FL}}(G, \varepsilon; \mathbb{R})$ , i.e.,

$$\varphi(G, -1) = \sum_{Z \subseteq E} (-1)^{|E|-|Z|+n\langle Z \rangle} = (-1)^{n\langle G \rangle} |\mathcal{O}_{\text{TC}}(G)|.$$

Now the identity (5.10) follows immediately.

## 6. Connection with the Tutte polynomial

The Tutte polynomial (see [3], p.337) of a graph  $G = (V, E)$  is a polynomial in two variables, which may be defined as

$$T_G(x, y) = \sum_{A \subseteq E} (x-1)^{r\langle E \rangle - r\langle A \rangle} (y-1)^{n\langle A \rangle}, \quad (6.1)$$

where  $\langle E \rangle = |V| - c(G)$ ,  $r\langle A \rangle = |V| - c(A)$ ,  $n\langle A \rangle = |A| - r\langle A \rangle$ . The polynomial  $T_G(x, y)$  satisfies the Deletion-Contraction Relation:

$$T_G(x, y) = \begin{cases} xT_{G/e}(x, y) & \text{if } e \text{ is a bridge,} \\ yT_{G-e}(x, y) & \text{if } e \text{ is a loop,} \\ T_{G-e}(x, y) + T_{G/e}(x, y) & \text{otherwise.} \end{cases}$$

It is well-known that the flow polynomial  $\varphi(G, t)$  is related to  $T_G(x, y)$  by

$$\varphi(G, t) = (-1)^{n\langle G \rangle} T_G(0, 1-t). \quad (6.2)$$

Thus

$$\bar{\varphi}(G, t) = (-1)^{n\langle G \rangle} \varphi(G, -t) = T_G(0, t+1).$$

We conclude the information as the following proposition.

**Proposition 6.1.** *The Tutte polynomial  $T_G$  is related to  $\varphi$  and  $\bar{\varphi}$  as follows:*

$$T_G(0, t) = \bar{\varphi}(G, t-1) = (-1)^{n\langle G \rangle} \varphi(G, 1-t). \quad (6.3)$$

*In particular,  $T_G(0, 1) = |\varphi(G, 0)| = \bar{\varphi}(G, 0)$  counts the number of Eulerian equivalence classes of totally cyclic orientations on  $G$ .*

**Example 6.2.** Let  $B_n(u, v)$  be a graph with two vertices  $u, v$ , and  $n$  multiple edges  $e_1, \dots, e_n$  between  $u$  and  $v$ . Let  $\varepsilon$  be an orientation of  $B_n(u, v)$  such that all edges have the direction from  $u$  to  $v$ . The number of integer flows  $f$  of  $B_n(u, v)$  such that  $|f| < q$  is equal to the number of integer solutions of the linear inequality system

$$x_1 + \dots + x_n = 0, \quad -(q-1) \leq x_i \leq q-1.$$

Let  $x_i = y_i - (q-1)$ . The above system reduces to

$$y_1 + \dots + y_n = n(q-1), \quad 0 \leq y_i \leq 2q-2. \quad (6.4)$$

Recall that the number of nonnegative integer solutions of  $y_1 + \cdots + y_n = r$  is

$$\left\langle \begin{matrix} n \\ r \end{matrix} \right\rangle := \binom{n+r-1}{r} = \binom{n+r-1}{n-1}.$$

So the number of nonnegative integer solutions of  $y_1 + \cdots + y_n = n(q-1)$  is  $\binom{nq-1}{n-1}$ , which includes the number of integer solutions of (6.4) and the number of nonnegative integer solutions having at least one  $y_i \geq 2q-1$ .

Let  $s_n(q)$  denote the number of integer solutions of (6.4). To figure out  $s_n(q)$ , let  $Y$  be the set of nonnegative integer solutions of  $y_1 + \cdots + y_n = n(q-1)$ ,  $Y_i$  the set of nonnegative integer solutions of  $y_1 + \cdots + y_n = n(q-1)$  with the  $i$ th variable  $y_i \geq 2q-1$ , and  $Y_0$  the set of integer solutions of (6.4). Then  $Y_0 = Y - \bigcup_{i=1}^n Y_i$ . Consider the case where  $j$  variables are greater than or equal to  $2q-1$ , say,  $y_{n-j+1}, \dots, y_n$ ; then  $y_1 + \cdots + y_{n-j} \leq n(q-1) - j(2q-1)$ . The number of such solutions is equal to the number of nonnegative integer solutions of the equation  $y_1 + \cdots + y_{n-j} + y' = n(q-1) - j(2q-1)$ , which is given by

$$\left\langle \begin{matrix} n-j+1 \\ n(q-1) - j(2q-1) \end{matrix} \right\rangle = \binom{(n-2j)q}{n(q-1) - j(2q-1)} = \binom{(n-2j)q}{n-j}.$$

Applying the Inclusion-Exclusion Principle,

$$\begin{aligned} s_n(q) &= \#(Y) + \sum_{\emptyset \neq I \subseteq [n]} (-1)^{|I|} \# \left( \bigcap_{i \in I} Y_i \right) \\ &= \binom{nq-1}{n-1} + \sum_{j=1}^{\lfloor \frac{n(q-1)}{2q-1} \rfloor} (-1)^j \binom{n}{j} \binom{(n-2j)q}{n-j}. \end{aligned}$$

We list  $s_1, s_2, s_3, s_4$  explicitly as follows:

$$\begin{aligned} s_1(q) &= 1, \quad s_2(q) = 2q-1, \quad s_3(q) = 3q^2 - 3q + 1, \\ s_4(q) &= \binom{4q-1}{3} - 4 \binom{2q}{3} = \frac{1}{3}(2q-1)(8q^2 - 8q + 3). \end{aligned}$$

Let  $s_0(q) (\equiv 1)$  denote the number of zero flows of  $B_n(u, v)$ . Then  $\varphi_z(B_4, q)$ , defined as the number of nowhere-zero integer  $q$ -flows of  $B_4(u, v)$ , counts the number of integer flows  $f$  such that  $0 < |f| < q$ , and is given by

$$\begin{aligned} \varphi_z(B_4, q) &= \# \left( [1-q, q-1]^{E(B_4)} \cap F(B_4, \varepsilon; \mathbb{Z}) - \bigcup_{e \in E(B_4)} F_e \right) \quad (6.5) \\ &= s_4(q) - 4s_3(q) + 6s_2(q) - 4s_1(q) + s_0(q) \\ &= \frac{2}{3}(q-1)(8q^2 - 22q + 21). \end{aligned}$$

Likewise, the number of flows of  $B_n(u, v)$  modulo  $q$  is  $q^{n-1}$ ,  $n \geq 1$ . Thus  $\varphi(B_4, q)$ , defined as the number of nowhere-zero flows of  $B_4(u, v)$  modulo  $q$ , is given by

$$\begin{aligned} \varphi(B_4, q) &= \# \left( F(B_4, \varepsilon; \mathbb{Z}/q\mathbb{Z}) - \bigcup_{e \in E(B_4)} F_e \right) \quad (6.6) \\ &= q^3 - 4q^2 + 6q - 4 + 1 \\ &= (q-1)(q^2 - 3q + 3). \end{aligned}$$

In particular,

$$|\varphi_z(B_4, 0)| = |\varphi(B_4, -1)| = 14, \quad |\varphi(B_4, 0)| = 3.$$

There are 14 totally cyclic orientations on  $B_4(u, v)$  by Theorem 1.1. The 14 orientations can be grouped into 3 Eulerian equivalence classes by Theorem 1.2; see Figure 1.




FIGURE 1. The three Eulerian equivalence classes of  $\mathcal{O}_{\text{TC}}(B_4)$ .

Let  $\varepsilon_k$  be an orientation on  $B_n(u, v)$  such that the arrows of the edges  $e_1, \dots, e_k$  point from  $u$  to  $v$ , and the arrows of the edges  $e_{k+1}, \dots, e_n$  point from  $v$  to  $u$ . If a 0-1 flow  $f$  of  $(B_n, \varepsilon_k)$  has value 1 on exact  $j$  edges of  $e_1, \dots, e_k$ , then  $f$  must have value 1 on exact  $j$  edges of  $e_{k+1}, \dots, e_n$ . Thus by Lemma 5.5,  $\bar{\varphi}_{\varepsilon_k}(B_n, 1)$  is the number of 0-1 flows of  $(B_n, \varepsilon_k)$ , and is given by

$$\bar{\varphi}_{\varepsilon_k}(B_n, 1) = \sum_{j=0}^{\min(k, n-k)} \binom{k}{j} \binom{n-k}{j}.$$

For the case  $n = 4$ , we see that  $\bar{\varphi}_{\varepsilon_0}(B_4, 1) = 1$ ,  $\bar{\varphi}_{\varepsilon_1}(B_4, 1) = 4$ ,  $\bar{\varphi}_{\varepsilon_2}(B_4, 1) = 6$ ,  $\bar{\varphi}_{\varepsilon_3}(B_4, 1) = 4$ , and  $\bar{\varphi}_{\varepsilon_4}(B_4, 1) = 1$ . According to Theorem 1.2,  $\bar{\varphi}_{\varepsilon_k}(B_4, 1)$  counts the number of orientations of  $B_4$  that are Eulerian equivalent to  $\varepsilon_k$ . Indeed, there are 4 orientations Eulerian equivalent to  $\varepsilon_1$  and  $\varepsilon_3$  respectively, and 6 orientations Eulerian equivalent to  $\varepsilon_2$ . These orientations are listed in Figure 1. However, the Eulerian equivalence classes for  $\varepsilon_0$  and  $\varepsilon_4$  are singletons; the integral and modular flow polynomials with respect to these orientations are the zero polynomial. We list the two orientations as follows:



FIGURE 2. The other two Eulerian equivalence classes of orientations of  $B_4$  with directed cut.

**Remark.** The coefficients of the integral flow polynomial  $\varphi_{\mathbb{Z}}(G, t)$  are not necessarily integers as shown in Example 6.2. The combinatorial interpretation on the coefficients of  $\varphi_{\mathbb{Z}}(G, t)$  is particularly wanted.

**Acknowledgement.** We thank the referees, in particular the second referee, for carefully reading the manuscript and offering several valuable comments.

## References

- [1] Beck, M.; Zaslavsky, T. The number of nowhere-zero flows of graphs and signed graphs, *J. Combin. Theory Ser. B* **96** (2006), 901–918.
- [2] Berge, C. *Graphs and Hypergraphs*, American Elsevier, New York, 1973.
- [3] Bollobás, B. *Modern Graph Theory*, Springer, 2002.
- [4] Bondy, J.A.; Murty, U.S.R., *Graph Theory*, Springer, 2008.
- [5] Breuer, F.; Sanyal, R. Ehrhart theory, modular flow reciprocity law, and the Tutte polynomial, arXiv:0907.0845v1 [math.CO] 5 Jul 2009.
- [6] Brylawski, T.; Oxley, J.G. The Tutte polynomial and its applications, in: *Matroid Applications*, N. White (ed.), Encyclopedia of Math. & Its Appl., Vol. **40**, Cambridge: Cambridge Univ. Press, 1992.
- [7] Chen, B. On the Euler characteristics of finite unions of convex sets, *Discrete Comput. Geom.* **10** (1993), 79–93.
- [8] Chen, B. On characteristic polynomials of subspace arrangements, *J. Combin. Theory Ser. A* **90** (2000), 347–352.
- [9] Chen, B. Lattice points, Dedekind sums, and Ehrhart polynomials of lattice polyhedra, *Discrete Comput. Geom.* **28** (2002), 175–199.
- [10] Chen, B. Ehrhart polynomials of lattice polyhedral functions, in: *Integer Points in Polyhedra – Geometry, Number Theory, Algebra, Optimization*, pp. 37–63, Contemp. Math., **374**, Amer. Math. Soc., Providence, RI, 2005.
- [11] Chen, B. Orientations, lattice polytopes, and group arrangements I: Chromatic and tension polynomials of graphs, *Ann. Combin.* **12** (2010), 425–452.
- [12] Crapo, H.; Rota, G.-C. *On the Foundations of Combinatorial Theory: Combinatorial Geometries*, MIT Press, Cambridge, MA, 1970.
- [13] Ehrenborg, R.; Readdy, M.A. On valuations, the characteristic polynomial, and complex subspace arrangements, *Adv. Math.* **134** (1998), 32–42.
- [14] Ehrenborg, R.; Readdy, M.A. The Dowling transform of subspace arrangements. in: Memory of Gian-Carlo Rota, *J. Combin. Theory Ser. A* **91** (2000), 322–333.
- [15] Greene, C; Zaslavsky, T. On the interpretation of Whitney numbers through arrangements of hyperplanes, zonotopes, non-Radon partitions, and orientations of graphs, *Trans. Amer. Math. Soc.* **280** (1983), 97–126.
- [16] Groemer, H. On the extension of additive functionals on the classes of convex sets, *Pacific J. Math.* **75** (1978), 397–410.
- [17] Kochol, M. Polynomials associated with nowhere-zero flows, *J. Combin. Theory Ser. B* **84** (2002), 260–269.
- [18] Las Vergnas, M. Extensions normales d’un matroïde, polynôme de Tutte d’un morphisme. *C. R. Acad. Sci. Paris Sr. A-B* **280** (1975), no. 22, 1479–1482.
- [19] Rota, G.-C. On the foundations of combinatorial theory I. Theory of Möbius functions, *Z. Wahrsch. Verw. Gebiete* **2** (1964), 340–368.
- [20] Stanley, R.P. Acyclic orientations of graphs, *Discrete Math.* **5** (1973), 171–178.
- [21] Stanley, R.P. *Enumerative Combinatorics I*, Cambridge Univ. Press, Cambridge, 1997.
- [22] Schanuel, S. Negative sets have Euler characteristic and dimension, in: *Category Theory* (Como, 1990), 379–385, Lecture Notes in Math. **1488**, Springer, Berlin, 1991.
- [23] Tutte, W.T. On the embedding of linear graphs in surfaces, *Proc. London Math. Soc. Ser. 2* **52** (1950), 474–483.
- [24] Tutte, W.T. A contribution to the theory of chromatic polynomials, *Canadian J. Math.* **6** (1954), 80–91.
- [25] Zaslavsky, T. *Facing up to Arrangements: Face-count Formulas for Partitions of Space by Hyperplanes*, Mem. Amer. Math. Soc. No. 154, 1975.

- [26] Zaslavsky, T. A combinatorial analysis of topological dissections, *Adv. in Math.* **25** (1977), 267–285.
- [27] Zhang, C.-Q. *Integer Flows and Circuit Covers of Graphs*, Marcel Dekker, New York, NY, 1997.

DEPARTMENT OF MATHEMATICS, HONG KONG UNIVERSITY OF SCIENCE AND TECHNOLOGY,  
CLEAR WATER BAY, KOWLOON, HONG KONG  
*E-mail address:* `mabfchen@ust.hk`

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY, CAMBRIDGE,  
MA 02139, USA  
*E-mail address:* `rstan@math.mit.edu`



