

# EUR 246.e

REPRINT

ASSOCIATION EURATOM - C.N.E.N.  
(European Atomic Energy Community and Comitato Nazionale  
per l'Energia Nucleare)

## ELECTRIC QUADRUPOLE TRANSITIONS BETWEEN SOME EXCITED STATES IN SPHERICAL EVEN-EVEN NUCLEI

by

G. SARTORIS

(Istituto di Fisica Teorica dell'Università - Napoli)

J. TOUCHARD

(Laboratoire Joliot-Curie de Physique Nucléaire - Orsay)

1963



Work performed by Istituto Nazionale di Fisica Nucleare -  
Sottosezione di Napoli  
under the Euratom contract No. 001-60-12 MPAI

Reprinted from  
IL NUOVO CIMENTO  
Vol. XXV, No. 6 - 1962

## LEGAL NOTICE

This document was prepared under the sponsorship of the Commission of the European Atomic Energy Community (EURATOM).

Neither the Euratom Commission, its contractors nor any person acting on their behalf :

- 1° — Make any warranty or representation, express or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this document, or that the use of any information, apparatus, method, or process disclosed in this document may not infringe privately owned rights; or
- 2° — Assume any liability with respect to the use of, or for damages resulting from the use of any information, apparatus, method or process disclosed in this document.

*This reprint is intended for restricted distribution only. It reproduces, by kind permission of the publisher, an article from "IL NUOVO CIMENTO", Vol. XXV, No. 6 - 1962, 1375-1382. For further copies please apply to Il Nuovo Cimento, Via Irnerio, 46 - Bologna, Italia.*

*Dieser Sonderdruck ist für eine beschränkte Verteilung bestimmt. Die Wiedergabe des vorliegenden in „IL NUOVO CIMENTO“, Vol. XXV, Nr. 6 - 1962, 1375-1382 erschienen Aufsatzes erfolgt mit freundlicher Genehmigung des Herausgebers. Bestellungen weiterer Exemplare sind an Il Nuovo Cimento, Via Irnerio, 46 - Bologna, Italia, zu richten.*

*Ce tiré à part est exclusivement destiné à une diffusion restreinte. Il reprend, avec l'aimable autorisation de l'éditeur, un article publié dans « IL NUOVO CIMENTO », Vol. XXV, N° 6 - 1962, 1375-1382. Tout autre exemplaire de cet article doit être demandé à Il Nuovo Cimento, Via Irnerio, 46 - Bologna, Italia.*

*Questo estratto è destinato esclusivamente ad una diffusione limitata. Esso è stato riprodotto, per gentile concessione dell'Editore, da « IL NUOVO CIMENTO », Vol. XXV, N° 6 - 1962, 1375-1382. Ulteriori copie dell'articolo debbono essere richieste a Il Nuovo Cimento, Via Irnerio, 46 - Bologna, Italia.*

*Deze overdruk is slechts voor beperkte verspreiding bestemd. Het artikel is met welwillende toestemming van de uitgever overgenomen uit „IL NUOVO CIMENTO“, Serie X, Vol. 27 - 1962, 86-92. Meer exemplaren kunnen besteld worden bij Il Nuovo Cimento, Via Irnerio, 46 - Bologna, Italia.*

## EUR 246.e

REPRINT

ELECTRIC QUADRUPOLE TRANSITIONS BETWEEN SOME EXCITED STATES IN SPHERICAL EVEN-EVEN NUCLEI by G. SARTORIS (Istituto di Fisica Teorica dell'Università - Napoli) and J. TOUCHARD (Laboratoire Joliot-Curie de Physique Nucléaire - Orsay).

Association EURATOM - C.N.E.N.

(European Atomic Energy Community and Comitato Nazionale per l'Energia Nucleare).

Work performed by Istituto Nazionale di Fisica Nucleare, Sottosezione di Napoli, under The Euratom contract No. 001-60-12 MPAl.

Reprinted from „Il Nuovo Cimento”, Vol. XXV, N. 6 - 1962, pages 1375-1382.

The branching ratios from the higher excited states in spherical even-even nuclei are calculated using a simplified model, which includes only a single  $j$  shell, and a pairing plus quadrupole residual interaction between nucleons. The method is applied to the branching from the  $4+$  levels: it is found that the  $E2$  transition to the second  $2+$  level can be enhanced against the corresponding one to the first  $2+$  level for more than one  $4+$  level.

---

## EUR 246.e

REPRINT

ELECTRIC QUADRUPOLE TRANSITIONS BETWEEN SOME EXCITED STATES IN SPHERICAL EVEN-EVEN NUCLEI by G. SARTORIS (Istituto di Fisica Teorica dell'Università - Napoli) and J. TOUCHARD (Laboratoire Joliot-Curie de Physique Nucléaire - Orsay).

Association EURATOM - C.N.E.N.

(European Atomic Energy Community and Comitato Nazionale per l'Energia Nucleare).

Work performed by Istituto Nazionale di Fisica Nucleare, Sottosezione di Napoli, under The Euratom contract No. 001-60-12 MPAl.

Reprinted from „Il Nuovo Cimento”, Vol. XXV, N. 6 - 1962, pages 1375-1382.

The branching ratios from the higher excited states in spherical even-even nuclei are calculated using a simplified model, which includes only a single  $j$  shell, and a pairing plus quadrupole residual interaction between nucleons. The method is applied to the branching from the  $4+$  levels: it is found that the  $E2$  transition to the second  $2+$  level can be enhanced against the corresponding one to the first  $2+$  level for more than one  $4+$  level.

---

## EUR 246.e

REPRINT

ELECTRIC QUADRUPOLE TRANSITIONS BETWEEN SOME EXCITED STATES IN SPHERICAL EVEN-EVEN NUCLEI by G. SARTORIS (Istituto di Fisica Teorica dell'Università - Napoli) and J. TOUCHARD (Laboratoire Joliot-Curie de Physique Nucléaire - Orsay).


Association EURATOM - C.N.E.N.

(European Atomic Energy Community and Comitato Nazionale per l'Energia Nucleare).

Work performed by Istituto Nazionale di Fisica Nucleare, Sottosezione di Napoli, under The Euratom contract No. 001-60-12 MPAl.

Reprinted from „Il Nuovo Cimento”, Vol. XXV, N. 6 - 1962, pages 1375-1382.

The branching ratios from the higher excited states in spherical even-even nuclei are calculated using a simplified model, which includes only a single  $j$  shell, and a pairing plus quadrupole residual interaction between nucleons. The method is applied to the branching from the  $4+$  levels: it is found that the  $E2$  transition to the second  $2+$  level can be enhanced against the corresponding one to the first  $2+$  level for more than one  $4+$  level.



## Electric Quadrupole Transitions between Some Excited States in Spherical Even-Even Nuclei<sup>(\*)</sup>.

G. SARTORIS

*Istituto di Fisica Teorica dell'Università - Napoli*  
*Istituto Nazionale di Fisica Nucleare - Sottosezione di Napoli*

J. TOUCHARD

*Laboratoire Joliot-Curie de Physique Nucléaire - Orsay*

(ricevuto il 21 Maggio 1962)

**Summary.** — The branching ratios from the higher excited states in spherical even-even nuclei are calculated using a simplified model, which includes only a single  $j$  shell, and a pairing plus quadrupole residual interaction between nucleons. The method is applied to the branching from the  $4+$  levels: it is found that the  $E2$  transition to the second  $2+$  level can be enhanced against the corresponding one to the first  $2+$  level for more than one  $4+$  level.

### 1. — Introduction.

It has been shown recently by VAN LIESHOUT, RICCI and GIRGIS<sup>(1)</sup> that several spherical even-even nuclei exhibit an interesting regularity in the decay of some higher excited states. If we consider, for instance, the  $E2$  decay of a given  $4+$  level and introduce the branching ratio:

$$F_x = \frac{B(E2)_{x_1}}{B(E2)_{x_2}},$$

(\*) This research was sponsored in part by the Fondazione Beneduce, Napoli. It is also a part of the research programme of the Nuclear Spectroscopy Group of Naples, working under a contract between EURATOM and C.N.E.N.

(1) R. VAN LIESHOUT, R. A. RICCI and R. K. GIRGIS: *Nuovo Cimento*, 21, 379 (1961).

of the decays into the two lower  $2+$  excited states,  $\omega$  labelling the  $4+$  level of interest (Fig. 1), it appears that  $F_{\omega}$  can be  $\gg 1$  for more than one  $4+$  level.




Fig. 1. - Transitions from a  $4+$  level.

A qualitative explanation of these enhancements has been suggested by RICCI, JEAN and VAN LIESHOUT<sup>(2)</sup>, based on a simple model of « pairing plus quadrupole force » coupling<sup>(3,4)</sup>, in the case of a single  $j$  and only one kind of nucleons<sup>(5)</sup>.

Our purpose, in the present note, is to provide support to that explanation by detailed calculations of the  $E2$  branching ratios. We are aware that the model is too drastic and cannot be applied without further refinement to actual nuclei. The more realistic case of various  $j$  shells and two kinds of nucleons<sup>(6)</sup> will be investigated later.

## 2. - The model.

We introduce the following well-known approximations<sup>(7)</sup> (we do not want to discuss here their validity):

i) The quasi-particle approximation. The creation operator of a quasi-particle  $\beta_m^+$  is defined by the Bogoljubov-Valatin<sup>(8)</sup> transformation:

$$\beta_m^+ = u b_m^+ - (-)^{j+m} v b_{-m},$$

$$u^2 + v^2 = 1,$$

where  $b_m^+$  is the creation operator of a particle in the state  $j, m$ . In our single- $j$  case,  $v^2 = N/2\Omega$  and  $E = \frac{1}{2}G\Omega$ , where  $N$  is the number of particles in the shell,  $\Omega = j + \frac{1}{2}$  the pair capacity of the shell,  $G$  the strength of the pairing force and  $E$  the excitation energy of a quasi-particle.

ii) Quasi-boson approximation. We define:

$$B_{\mu}^{(s)+} = \frac{1}{\sqrt{2}} \sum_{mm'} \langle jjmm' | s\mu \rangle \beta_m^+ \beta_{m'}^+, \quad s = 2, 4 \dots 2j-1,$$

<sup>(2)</sup> R. A. RICCI, M. JEAN and R. VAN LIESHOUT: *Nuovo Cimento*, **25**, 1389 (1962).

<sup>(3)</sup> L. S. KISSLINGER and R. A. SØRENSEN: *Mat. Fys. Medd. Dan. Vid. Selsk.*, **32**, no. 9 (1960).

<sup>(4)</sup> R. A. SØRENSEN: *Nucl. Phys.*, **25**, 674 (1961).

<sup>(5)</sup> B. R. MOTTELSON: *The Many Body Problem* (Paris, 1959).

<sup>(6)</sup> T. TAMURA and T. UDAGAWA: *Prog. Theor. Phys.*, **26**, 947 (1961).

<sup>(7)</sup> B. F. BAYMAN: *Lectures on Seniority, Quasi-particles and Collective Vibrations* (1960).

<sup>(8)</sup> S. T. BELYAEV: *Mat. Fys. Medd. Dan. Vid. Selsk.*, **31**, no. 11 (1959).

and suppose

$$[B_{\mu}^{(s)}, B_{\mu'}^{(s')}] = \delta_{ss'} \delta_{\mu\mu'}$$

$B_{\mu}^{(s)+}$  acts then like the creation operator for a boson of spin  $s$ ; the approximation seems reliable only when the number of quasi-particles is small with respect to the capacity  $\Omega$  of the shell.

iii) The quadrupole force is introduced according to the linearization method<sup>(9-11)</sup>, i.e. only quadratic terms like  $B^+B$  and  $BB$  (and Hermitian conjugates) are taken into account. With the approximations i), ii) and iii) the Hamiltonian is written as:

$$H = H_{s \neq 2} + H_2,$$

$$H_{s \neq 2} = 2E \sum_{s \neq 2, \mu} B_{\mu}^{(s)+} B_{\mu}^{(s)},$$

$$H_2 = 2E \sum_{\mu} B_{\mu}^{(2)+} B_{\mu}^{(2)} - \frac{1}{2} \chi Q^2 \sum_{\mu} (-)^{\mu} [B_{\mu}^{(2)+} + (-)^{\mu} B_{-\mu}^{(2)}] [B_{-\mu}^{(2)+} + (-)^{\mu} B_{\mu}^{(2)}],$$

with

$$Q = (-)^l uv \sqrt{\frac{2}{5}} \langle j \| r^2 Y^{(2)} \| j \rangle.$$

The second term in  $H_2$  comes from the quadrupole force between two nucleons:

$$V_{ab} = - \chi r_a^2 Y_a^{(2)} \cdot r_b^2 Y_b^{(2)}.$$

If we define a vector operator  $J$  by:

$$J_x = \frac{1}{4} (S_2^+ - S_2) = -J_x^+,$$

$$J_y = -\frac{i}{4} (S_2^+ + S_2) = -J_y^+,$$

$$J_z = \frac{1}{2} (S_1 + \frac{5}{2}) = J_z^+,$$

with

$$S_1 = \sum_{\mu} B_{\mu}^{(2)+} B_{\mu}^{(2)} \quad \text{and} \quad S_2 = \sum_{\mu} (-)^{\mu} B_{\mu}^{(2)+} B_{-\mu}^{(2)}$$

then


$$H_2 = -5E + 4(E - \frac{1}{2} \chi Q^2) J_z - 2i \chi Q^2 J_y.$$

<sup>(9)</sup> M. BARANGER: *Phys. Rev.*, **120**, 957 (1960).

<sup>(10)</sup> R. ARVIEU and M. VENERONI: *Compt. Rend.*, **280**, 992 (1960).

<sup>(11)</sup> M. KOBAYASI and T. MARUMORI: *Prog. Theor. Phys.*, **23**, 387 (1960).

In order to find the normal modes of the system, we perform a canonical transformation characterized by the operator  $U = \exp[-2\alpha J_x]$ . Expanding  $U$  into a power series, the transformed Hamiltonian can be calculated with the



help of the commutation rule  $J \wedge J = iJ$ , and is found to be diagonal under the condition:

$$\operatorname{ctgh} 2\alpha = 1 - (2E/\chi Q^2).$$

We can then write the new Hamiltonian as:

$$U^+ H_2 U = -\frac{\hbar\omega}{2}(2E - \hbar\omega) + \hbar\omega S_1$$

with

$$\hbar\omega = 2E \sqrt{1 - \frac{\chi Q^2}{E}}$$

Fig. 2. - Energy spectrum resulting from a « pairing plus quadrupole » force coupling scheme.

The following schematic energy spectrum describes therefore our system of  $N$  interacting particles in the  $j$  shell (Fig. 2):

- A) quasi-boson type excitations with energy  $2E$ ; operators  $B_\mu^{(s \neq 1)^+}$ ;
- B) phonon type excitations with energy  $\hbar\omega$ ; operators  $B_\mu^{(2)^+}$ ;
- C) excitations due to the coupling of phonons and quasi-bosons.

It is clear that for large  $j$  several  $4^+$  levels are obtained which lie above an energy of order  $2\hbar\omega$ . In order to simplify the analysis of the various possible cases, we make the assumption that the quadrupole force is strong enough to depress the  $2^+$  two phonon level below the one quasi-boson level ( $\hbar\omega < E$ ).

We have then four types of  $4^+$  levels labelled by  $x = a, b, c$  and  $d$ :

- a) One  $4^+$  quasi-boson (A band).
- b) Three phonons coupled to  $I = 4^+$  (B band).
- c) Two phonons coupled to  $I = 4^+$  (B band).
- d) A phonon and a quasi-boson coupled to  $I = 4^+$  (C band).

We can then calculate the branching ratio  $F_x$  for each type  $x$ .



## 3. - Electromagnetic transitions.

The  $E2$  transition operator  $\mathcal{M}_\mu^{(2)} = \sum_a r_a^2 Y_\mu^{(2)}(a)$  can be written, in second quantized form, after having performed the quasi-particle transformation as:

$$\mathcal{M}_\mu^{(2)} = Q[B_\mu^{(2)+} + (-)^\mu B_{-\mu}^{(2)}] + q[\beta^+\gamma]_\mu^{(2)},$$

with

$$q = (v^2 - u^2) \frac{1}{\sqrt{5}} \langle j \| r^2 Y^{(2)} \| j \rangle,$$

$$[\beta^+\gamma]_\mu^{(2)} = \sum_{mm'} \langle jjmm' | 2\mu \rangle \beta_m^+ \gamma_{m'},$$

$$\gamma_m = (-)^{j-m} \beta_{-m}.$$

By the canonical transformation  $U$ , the operator  $B_\mu^{(2)+}$  becomes

$$U^+ B_\mu^{(2)+} U = \cosh \alpha B_\mu^{(2)+} - (-)^\mu \sinh \alpha B_{-\mu}^{(2)}.$$

One gets therefore:

$$U^+ \mathcal{M}_\mu^{(2)} U = Q(\cosh \alpha - \sinh \alpha)[B_\mu^{(2)+} + C_\mu^{(2)}] - q \mathcal{N}_\mu^{(2)},$$

$$\mathcal{N}_\mu^{(2)} = -[\beta^+\gamma]_\mu^{(2)} + 2(1 - \cosh 2\alpha) M_\mu^{(2)} + \sinh 2\alpha N_\mu^{(2)} + \sinh 2\alpha O_\mu^{(2)} +$$

$$+ 2(1 - \cosh \alpha) P_\mu^{(2)} + 2(1 - \cosh \alpha) Q_\mu^{(2)} + 2 \sinh \alpha R_\mu^{(2)} + 2 \sinh \alpha T_\mu^{(2)},$$

$$M_\mu^{(2)} = 5 \begin{Bmatrix} 2 & 2 & 2 \\ j & j & j \end{Bmatrix} [B_\mu^{(2)+} C_\mu^{(2)}]_\mu^{(2)} \quad P_\mu^{(2)} = \sum_{s \neq 2} \sqrt{5(2s+1)} \begin{Bmatrix} 2 & 2 & s \\ j & j & j \end{Bmatrix} [B_\mu^{(2)+} C_\mu^{(2)}]_\mu^{(2)},$$

$$N_\mu^{(2)} = 5 \begin{Bmatrix} 2 & 2 & 2 \\ j & j & j \end{Bmatrix} [B_\mu^{(2)+} B_\mu^{(2)+}]_\mu^{(2)} \quad Q_\mu^{(2)} = \sum_{s \neq 2} \sqrt{5(2s+1)} \begin{Bmatrix} 2 & 2 & s \\ j & j & j \end{Bmatrix} [C_\mu^{(2)} B_\mu^{(2)+}]_\mu^{(2)},$$

$$O_\mu^{(2)} = 5 \begin{Bmatrix} 2 & 2 & 2 \\ j & j & j \end{Bmatrix} [C_\mu^{(2)} C_\mu^{(2)}]_\mu^{(2)} \quad R_\mu^{(2)} = \sum_{s \neq 2} \sqrt{5(2s+1)} \begin{Bmatrix} 2 & 2 & s \\ j & j & j \end{Bmatrix} [B_\mu^{(2)+} B_\mu^{(2)+}]_\mu^{(2)},$$

$$T_\mu^{(2)} = \sum_{s \neq 2} \sqrt{5(2s+1)} \begin{Bmatrix} 2 & 2 & s \\ j & j & j \end{Bmatrix} [C_\mu^{(2)} C_\mu^{(2)}]_\mu^{(2)},$$

with

$$C_\mu^{(2)} = (-)^\mu B_{-\mu}^{(2)}.$$

The expressions for the operators  $M$ ,  $N \dots T$  have been obtained forming

commutators of  $J_x$  with  $[\beta^+\gamma]_\mu^{(2)}$ , and making use of the angular momentum recoupling rules <sup>(12)</sup>.

One sees that the transitions  $a_1$  and  $d_2$  are strictly forbidden, *i.e.*  $F_a = 0$  and  $F_d = \infty$ . For the  $B$  band transitions ( $b$  and  $c$ ) one must evaluate <sup>(\*)</sup>:

$$B(E2) = \frac{1}{8} |\langle I = 2 \| \mathcal{M}^{(2)} \| I = 4 \rangle|^2.$$

Using known results for matrix elements of product of tensor operators and the selection rules for matrix elements of the  $B^{(2)}$  operator, one finds

$$F_b = \frac{1}{f} \operatorname{tg}^2 \varphi \frac{(\cosh \alpha - \sinh \alpha)^2}{\sinh^2 2\alpha},$$

$$F_c = \frac{16}{49} f \operatorname{ctg}^2 \varphi \frac{\cosh^2 2\alpha}{(\cosh \alpha - \sinh \alpha)^2},$$

with

$$f = 100 \begin{Bmatrix} 2 & 2 & 2 \\ j & j & j \end{Bmatrix}^2,$$

$$u = \cos(\varphi/2),$$

$$v = -(-)^j \sin(\varphi/2).$$

It is also interesting to evaluate the usual branching ratio

$$F_0 = \frac{B(E2)_{2_4 \rightarrow 2_2}}{B(E2)_{2_4 \rightarrow 0_1}},$$

which is found to be exactly  $2F_b$ , together with the value  $\mathcal{F}$  of  $B(E2)_{2_1 \rightarrow 0_1}$  in units of a single particle estimate  $B(E2)_{j_2}$  for  $(j)_{I=2}^2 \rightarrow (j)_{I=0}^2$  transition, which can be expressed as

$$\frac{\mathcal{F}}{\Omega} = \frac{1}{4} \sin^2 \varphi (\cosh \alpha - \sinh \alpha)^2.$$

#### 4. - Discussion.

Our assumption  $\hbar\omega/E < 1$ , or equivalently  $\chi Q^2/G\Omega > \frac{2}{3}$ , leads to  $\chi$  values much too large as compared with the actual values used for instance by KRIS-

<sup>(12)</sup> A. R. EDMONDS: *Angular Momentum in Quantum Mechanics* (Princeton, 1957).

<sup>(\*)</sup> Strictly speaking, the  $2_1^+$  and the  $4_1^+$  two-phonon levels are degenerate, and therefore no transition between them can occur. However, we may always evaluate  $B(E2)$  for the transition  $C_1$ , and suppose that the degeneracy is slightly removed as is the case in actual nuclei (see Fig. 2).

LINGER and SØRENSEN (4). But it is well known that a single  $j$  model is not able to depress strongly the  $2+$  level. Thus our assumption must be considered only as simulating such a strong effect. We choose as a tentative value  $\hbar\omega = 0.8E$ . On the other hand, the quantity  $f$  varies quite smoothly when  $j$  increases, and one can take a mean value  $f = 0.4$  without affecting significantly the order of magnitude of the branching ratios  $F_n$ .

The single parameter left is then:

$$\xi = \frac{1 - (\hbar\omega/2E)^2}{(2N/\Omega)(1 - N/2\Omega)},$$

which describes the occupancy of the shell (a large  $\xi$  occurs at the beginning or at the end of the shell, and  $\xi_{\min} = 0.84$  at the middle of the shell).




Fig. 3. - Branching ratios  $F_n$  and enhancement factor  $\mathcal{F}$  vs. shell occupancy.

In Fig. 3, we have plotted the quantities of interest against the parameter  $\xi$ . If we exclude the region of large  $\xi$  where the model is inadequate, we can point out the following trends:

i) The branching ratio  $F_b$  is of the order of some units (and even larger when approaching the middle of the shell), whereas  $F_c$  remains always  $\ll 1$ .

ii) The quantity  $\mathcal{F}/\Omega$  has a value 0.2 to 0.6, and the factor  $\mathcal{F}$  cannot be large if  $j$  is not large.

In the middle of the shell,  $\mathcal{F}$  has its maximum value (\*),  $F_0 = 0$  and  $F_0$  as  $F_0$  becomes infinite. We approach then the situation of the vibrational model.

### 5. - Conclusion.

It is thus verified that even within the limit of validity of such a simple model there exists among the possible ones a group of  $4+$  higher excited levels, for which the probability of  $E2$  transition to the  $2'+$  level is enhanced with respect to that to the first  $2+$  level (type *a* and *b*). We also find other  $4+$  levels for which the opposite situation occurs (type *a* and *c*).

As has been mentioned above, the model discussed here can only provide qualitative indications. A more realistic treatment should include several particle levels  $j_1, j_2, j_3, \dots$  with reasonable values of the various parameters. Further calculations in this sense are in progress.

\* \* \*

We wish to thank Prof. M. JEAN and Prof. R. A. RICCI for suggesting the subject of this note and for many stimulating discussions.

---

(\*) It can be remarked that for a pure pairing force,  $\mathcal{F}/\Omega$  has its maximum value 0.25 at the middle of the shell, and that our quadrupole force gives an additional enhancement by a factor 2.5.

### RIASSUNTO (\*)

Si calcolano i rapporti di branching dai più alti livelli eccitati nei nuclei sferici pari-pari, facendo uso di un modello semplificato, che comprende un solo strato  $j$ , ed una interazione residua di coppie e di quadrupolo fra i nucleoni. Si applica il metodo al branching dai livelli  $4+$ ; si trova che la transizione  $E2$  al secondo livello  $2+$  può essere esaltata rispetto a quella corrispondente verso il primo livello  $2+$  da più di un livello  $4+$ .

---

(\*) Traduzione a cura della Redazione.



CDNA00246ENC