

ASSESSING THE BIOAVAILABILITY AND TRANSLOCATION EFFICIENCY OF MINERAL ELEMENTS IN *Lycium barbarum* SPECIES FROM R. MACEDONIA AND R. CHINA

Biljana Balabanova¹, Violeta Ivanova-Petropulos¹, Tiberiu Frentiu², Michaela Ponta², Eniko Covaci², Marin Senila³

¹Faculty of Agriculture, Goce Delčev University, Krste Misirkov No. 10-A, 2000 Štip, Republic of Macedonia

²Faculty of Chemistry and Chemical Engineering, Babes-Boyai University, Cluj-Napoca, Romania

³INCDO-INOE 2000 National Institute for Research and Development of Optoelectronics Bucharest, Research Institut for Analytical Instrumentation, Cluj-Napoca, Romania

INTRODUCTION

Goji berries or wolfberries (*Lycium barbarum* L.) have been traditionally used as food and a medicinal plant. The interest in the chemical composition of goji berries has intensified because of an increased awareness of their possible health benefits. On the other hand, the toxic elements can also be interacting in the plant tissue through the bio-available pathway of the root-soil system.

The present work reports the results obtained for metals (Na, Mg, K, Ca, Mn, Fe, Cu and Zn) in goji berries by using inductively coupled plasma–optical emission spectrometry (ICP-OES), following digestion using a diluted oxidant mixture in a closed-vessel microwave oven. Determinations of Cr, As, Pb, Cd and Ni were realized using the graphite furnace atomic absorption spectrometer (GFAAS). Mercury quantification was realized on the solid samples by the TDAAS method, with the automated direct mercury analyzer hydra-C.

WHY IS GOJI BERRY (*Lycium barbarum*) A SUPERFOOD?

Size does not matter!

- Goji Berry contains ~500 times more vitamin C than oranges!
- It packs ~15 times more iron than spinach!
- Contains 22 minerals including Zn, Fe, Ca..... and 11 amino acids!!

Benefits for human health

Certain forms of some metals/semimetals can be TOXIC, even in relatively small amounts, and therefore pose -


As, Cd, Pb, Hg

RISK TO THE HUMAN HEALTH!!!!

DATA SUMMARY/RESULTS

Table 1. Median values for elements contents in soil and plant species (contents are given in mg/kg on dried mass)


| Element | Goji berries (fruit) | Soil /top layer (0-5 cm) |
|---------|----------------------|--------------------------|
| Na | 44.2 | 157 |
| Mg | 1180 | 5016 |
| K | 22100 | 5016 |
| Ca | 841 | 16860 |
| Mn | 19.4 | 729 |
| Fe | 69.4 | 24290 |
| Cu | 11.1 | 53 |
| Zn | 21.4 | 167 |
| Cr | 0.571 | 27.4 |
| As | 0.015 | 7.53 |
| Pb | 0.187 | 12.4 |
| Cd | 0.04 | 0.451 |
| Ni | 2.5 | 9.63 |
| Hg | 0.0063 | 0.065 |




$$BAF = [C_{plant}] / [C_{soil}]$$

Significant bioaccumulations occurs for BAF>1

Significant translocation occurs for TF>1



$$TF = [C_{fruit}] / [C_{root}]$$




$$BCF = [C_{root}] / [C_{soil}]$$

Significant bioconcentrating occurs for BCF>1

T-test for Dependent Samples

| Element | t | p |
|---------|--------|-------|
| Na | 7.849 | 0.001 |
| Mg | 8.617 | 0.001 |
| K | -4.836 | 0.008 |
| Ca | 3.960 | 0.017 |
| Fe | 4.973 | 0.008 |
| Ni | 5.526 | 0.005 |
| Cu | 3.435 | 0.026 |
| Zn | 2.659 | 0.056 |
| Cr | 4.490 | 0.011 |
| As | -1.044 | 0.356 |
| Cd | -5.928 | 0.004 |
| Hg | 0.991 | 0.378 |
| Pb | 3.088 | 0.037 |
| Mn | -1.952 | 0.123 |

Total elements content (mg/kg)



CONCLUSIONS

▪ The obtained data report the multi-element characterization of different plant parts, and variation in multi-elements content between Macedonian and Chinese species. Bioaccumulation and bioconcentration factor scores revealed the translocation efficiency of metals and nonmetals delcve the *Lycium barbarum* plant parts.

▪ Determination of potentially toxic metals concentration in food products is important for health risk assessment during the food consumption. This kind of study can be used as a tool for the farmers in order to adopt strategies to save the population by minimizing the problems related to metal toxicities. Such assessment for the contaminants is required for the well-being of the population.