

Cardiac Surveillance Findings During Adjuvant and Palliative Trastuzumab Therapy in Patients with Breast Cancer

ERZSÉBET VALICSEK¹, RENÁTA KÓSZÓ¹, ÁGNES DOBI¹, GABRIELLA UHERCSÁK¹,
ZOLTÁN VARGA¹, ANDREA VASS², ÉVA JEBELOVSZKY² and ZSUZSANNA KAHÁN¹

¹Department of Oncotherapy, and ²Second Department of Medicine and Cardiology Center,
University of Szeged, Szeged, Hungary

Abstract. *Background/Aim: Trastuzumab therapy, the standard treatment for human epidermal growth factor receptor type-2 (HER2)-positive breast cancer, is associated with possible cardiotoxicity. We set out to retrospectively analyze the cardiac follow-up data of patients with breast cancer receiving trastuzumab treatment. Patients and Methods: The study involved 47 and 31 patients receiving adjuvant or palliative chemotherapy plus trastuzumab, respectively. Cardiovascular system assessments including echocardiography were regularly performed. Results: A significant heart abnormality was detected in 44.7% of the operable and 41.9% of metastatic cases. In the adjuvant setting, left ventricular ejection fraction changes occurred mostly during treatment and less frequently after its completion (40.4% vs. 19.4%), while in the palliative setting, 35.5% and 40% in the first and the second year of therapy. An asymptomatic atrial septum aneurysm was detected in 8.5% and 13% of the patients in the two groups. Conclusion: Trastuzumab-related cardiotoxicity is mostly manifested in an asymptomatic decrease in left ventricular ejection fraction; hypertension, a high body mass index and left-sided irradiation are its predictors.*

Breast cancer is the most common cancer in women in developed countries. The human epidermal growth factor receptor type-2 (HER2)/neu protein is a 185-kDa transmembrane cytoplasmic tyrosine kinase. Amplification of the *HER2* gene or overexpression of the HER2 protein occurs

in 10-34% of breast cancer cases (1), and is associated with an adverse outcome (2). Trastuzumab (Herceptin[®]; Roche) is a recombinant humanized monoclonal antibody that inhibits tumor cell growth, differentiation and survival by binding with high affinity to the extracellular domain, and blocking the normal regulatory functions of the HER2 receptor (3).

The efficacy and safety of trastuzumab was first demonstrated in patients with HER2-overexpressing metastatic breast cancer (MBC) in different treatment lines, and in combination with standard chemotherapeutic regimens (4-7). In the adjuvant setting, the consistent outcome of several large multi-center randomized trials (8-14) led to the use of trastuzumab in HER2-positive breast cancer becoming a new standard (3, 12).

Retrospective analysis of the pivotal phase II and III trials revealed an increased incidence of cardiac dysfunction, usually manifested as congestive heart failure (CHF), a symptomatic or asymptomatic decrease of the left ventricular ejection fraction (LVEF) or, rarely, cardiac arrhythmias or ischemic heart disease (15). The incidence of cardiotoxicity in patients with MBC was found to be highest in those receiving concurrent trastuzumab and anthracyclines, but low when trastuzumab was administered in combination with paclitaxel or alone (16). The majority of these cases with a cardiac dysfunction were symptomatic, but standard treatment for CHF ameliorated the symptoms (17). In large studies in which the use of trastuzumab in early breast cancer was evaluated, the incidence of symptomatic cardiac dysfunction proved to be low (18), even after a long follow-up or radiotherapy (RT) (13, 14).

Unlike the case with anthracyclines, trastuzumab-related cardiac dysfunction constitutes an entity known as a type-2 chemotherapy-related cardiac dysfunction (CRCDD), a reversible abnormality with no structural changes in the myocardium, not dose-dependent and which does not occur in all patients (19-21). The mechanism of type-2 CRCDD has not been completely clarified and although several risk factors for trastuzumab-induced cardiotoxicity have been described, adverse cardiac events cannot be predicted (22). With the aim of identifying risk factors for cardiac morbidity

This article is freely accessible online.

Correspondence to: Zsuzsanna Kahán, Department of Oncotherapy, University of Szeged, 12 Korányi fasor, H-6720 Szeged, Hungary. Tel: +36 62545404, Fax: +36 62545922, e-mail: kahan.zsuzsanna@med.u-szeged.hu

Key Words: Trastuzumab, cardiac dysfunction, risk factor, hypertension, BMI, radiotherapy.

Table I. The chemotherapeutic regimens applied in patients receiving adjuvant trastuzumab treatment.

Chemotherapeutic regimen	Number of patients receiving adjuvant trastuzumab treatment
CMF	7
CEF	10
AC-paclitaxel	14
AC-docetaxel	6
TAC	1
CMF-docetaxel	3
TEX	3
EC	1
Docetaxel-epirubicin	1
Paclitaxel-carboplatin	1

CMF=cyclophosphamide, methotrexate, 5-fluorouracil, CEF=cyclophosphamide, epirubicin, 5-fluorouracil, AC=adriamycin, cytoxan, TAC=taxotere, Adriamycin, cyclophosphamide, TEX=taxotere, epirubicin, Xeloda, EC=epirubicin, cyclophosphamide.

for our everyday practice, we set-out to perform a retrospective study of cardiac follow-up data on patients with breast cancer receiving trastuzumab in the adjuvant or palliative setting.

Patients and Methods

All the procedures followed were in full accordance with the ethical standards of the appropriate committees on human experimentation (institutional and national) and with the Helsinki declaration.

The retrospective analysis included patients with histologically confirmed, HER2-positive breast cancer who received adjuvant or palliative trastuzumab therapy. All the clinical and pathological data were extracted from the patient files.

The HER2 status was evaluated by immunohistochemical staining or with fluorescence *in situ* hybridization (23).

Trastuzumab therapy. Patients received either adjuvant or neoadjuvant chemotherapy with standard regimens (Table I). Trastuzumab was given for 12 months at the standard dose (an 8 mg/kg loading dose and a 6 mg/kg sustaining dose at 3-week intervals), usually after the chemotherapy, but in the case of the sequential use of taxane monotherapy, it was co-administered with either paclitaxel or docetaxel. After neoadjuvant chemotherapy, it was always used as monotherapy. In metastatic cases, trastuzumab was either co-administered with paclitaxel or was applied as monotherapy following it, continued until disease progression.

Heart function surveillance. Heart function surveillance was carried out in accordance with the internationally accepted guidelines (24). Electrocardiography (ECG) and echocardiography were performed prior to the initiation of trastuzumab therapy, at 3-month intervals during the therapy, and at 6-month intervals after its termination; this was carried out more frequently if any abnormality indicating a deteriorating heart function was detected.

The presence of risk factors such as the age, body mass index (BMI), baseline LVEF and echocardiography abnormalities, hypertension, ischemic heart disease, diabetes mellitus, a smoking history, the duration of trastuzumab therapy, the dose of anthracycline chemotherapy and the administration of RT in cases of left-sided breast cancer were extracted from the patient files.

Standard 2-dimensional transthoracic echocardiography was performed by two experienced cardiologists using a standard ultrasound machine (Toshiba PowerVision 8000 with a 2.0-3.8 MHz transducer; Toshiba, Tustin, CA, USA). The LV dimensions and volume parameters were obtained from standard parasternal and apical 4-chamber views. The LVEF was calculated by the modified Simpson's method, according to the guidelines of the American Society of Echocardiography (24).

Study end-points. The reduction of LVEF by an absolute value of 10% or more from the baseline or its decrease below the level of 50% indicated a significant abnormality. The development of other heart changes or heart failure was additionally analyzed. The discontinuation of trastuzumab therapy was decided on an individual basis, in accordance with the guidelines (25, 26).

Statistical analysis. Data are reported as mean±SE or median values. Comparison of the incidence of abnormal LVEF changes between groups was carried out with the χ^2 test or Fisher's exact test. The difference between the mean LVEF changes during the first and the second year was analyzed with the paired samples *t*-test. SPSS 20.0 for Windows (SPSS Inc., Chicago, IL, USA) was used to perform the analysis. A *p*-value of less than 0.05 was considered significant.

Results

Adjuvant trastuzumab therapy. Between September 2006 and April 2012, 47 patients received adjuvant trastuzumab therapy; the median (range) duration of follow-up was 12 (3-36) months. The mean age (±SE) at diagnosis was 53.0±1.7 years. The tumor characteristics are shown in Table II. Two-thirds of the neoplasms were poorly-differentiated and about three-quarters of them were hormone receptor-negative. The average (±SE) number of chemotherapy cycles that the patients received before or during trastuzumab treatment was 5.62±0.27. Most patients completed the anthracycline-based chemotherapy, 18 patients (38.3%) received concomitant taxane and trastuzumab treatment, while 17 of those with left-sided breast cancer received RT (Table III). At least one of the patient-related risk factors for cardiovascular morbidity was present in 68% of the cases (Table IV).

Echocardiography findings are included in Table V. A total of 42 out of 47 patients completed a 1-year course of trastuzumab therapy, while in five cases, trastuzumab treatment was stopped after a mean duration of 5.2 (2.6-7) months (in four cases due to disease progression and in one due to a deterioration of cardiac function). The mean (±SE) duration of trastuzumab therapy was 10.8±0.4 months, with a median value of 11.9 months. The LVEF was normal at baseline in all cases, but decreased by 10% or more in 21 out of the 47 patients (44.7%). In 7/47 (14.9%) cases, these were

Table II. Tumor characteristics in the patient population treated with adjuvant, and neoadjuvant chemotherapy and adjuvant trastuzumab.

Tumor characteristics		Patients (N=47)	
		N	%
Histological type	Invasive ductal carcinoma	46	97.9
	Invasive lobular carcinoma	1	2.1
Grade	1	2	4.3
	2	16	34.0
	3	29	61.7
ER	Negative ($\leq 10\%$)	33	70.2
	Positive ($> 10\%$)	14	29.8
PR	Negative ($\leq 10\%$)	36	76.6
	Positive ($> 10\%$)	11	23.4
(y)pT	(y)pT1 (≤ 20 mm)	29	61.7
	(y)pT2 (> 20 mm)	18	38.3
(y)pN	(y)pN0	20	42.5
	(y)pN1-2	27	57.5
Distribution	Unifocal	42	89.4
	Multifocal	5	10.6

N=Number, ER=estrogen receptor, PR=progesterone receptor, (y)pT=pathological tumor stage (after neoadjuvant treatment), (y)pN=pathological lymph node stage (after neoadjuvant treatment).

merely isolated accidental findings during the treatment or the follow-up period. Abnormal LVEF changes occurred in 19/47 (40.4%) and 7/36 (19.4%) patients in the first and the second year, respectively. Among those who exhibited an LVEF decrease of 10% or more, the mean (\pm SE) changes were $10.58 \pm 1.82\%$ ($n=19$) and $11.79 \pm 1.29\%$ ($n=7$) in the first and the second year, respectively. None of the patients presented an LVEF $< 50\%$ or symptoms of CHF. Four patients (8.5%) developed atrial septum aneurysm during the treatment or the follow-up period.

The presence of hypertension favored LVEF decrease by 10% or more (27.0% among those with hypertension *vs.* 12.4% among those without hypertension, $p=0.004$). Likewise, LVEF decrease was associated with the earlier delivery of left-sided RT (27.2% among those who received RT *vs.* 15.2% among those who did not, $p=0.023$). No statistically significant difference was found with regard to the presence of diabetes or ischemic heart disease; however, the numbers of such cases were low. With the median BMI taken as threshold, patients with a BMI > 26.6 kg/m² demonstrated a higher incidence of abnormal LVEF changes than that for patients with a BMI less than this, at 28.0% *vs.*

Table III. Selected oncological therapies in the patient population treated with trastuzumab in the adjuvant and palliative settings.

Previous/concomitant oncological therapy	Patients receiving adjuvant trastuzumab treatment (N=47)		Patients receiving palliative trastuzumab treatment (N=31)	
	N	%	N	%
Anthracycline chemotherapy before trastuzumab therapy	41	87.2	12	38.7
Irradiated left-sided breast cancer cases	17	36.2	11	35.5
Whole breast	8	17.0	5	16.1
Chest wall	9	19.1	6	19.4
Regional lymph nodes	9	19.1	9	29.0
Tumor bed boost	8	17.0	3	9.7
Endocrine therapy during trastuzumab monotherapy	14	29.8	0	0.0

N=Number.

10.5% ($p=0.001$). No association was found between abnormal LVEF reduction and treatment with anthracycline-based chemotherapy or its dose. An abnormal LVEF change during the first year correlated significantly with the presence of hypertension and BMI, but not with left-sided RT, while in the second (follow-up) year, only the association with left-sided RT remained significant (Table VI).

Palliative trastuzumab therapy. Between June 2003 and February 2012, 31 patients with MBC received chemotherapy and trastuzumab in the palliative setting, 25 patients in the first-line setting, and six in the second-line. Among them, nine patients were maintained on trastuzumab, while 12 received another oncological therapy throughout observation time. The median duration of follow-up was 12 (3-36) months. The mean age (\pm SE) of the patients was 54.7 ± 2.2 years. Most of the carcinomas were poorly differentiated, and hormone receptor-negative (data not shown). The patients received an average (\pm SE) of 7.59 ± 0.87 chemotherapy cycles before or during trastuzumab therapy; 12/31 patients (38.7%) were treated with anthracycline-based chemotherapy before trastuzumab, while 27 patients (87.1%) received concomitant taxane and trastuzumab therapy, and 11 patients left-sided RT (Table III). The mean (\pm SE) duration of trastuzumab therapy was 27.6 ± 7.1 months. In five cases, trastuzumab monotherapy is ongoing.

Cardiovascular co-morbidities and risk factors were present in 80% of the patients (Table IV).

The initial mean (\pm SE) LVEF was $66.2 \pm 1.1\%$; a cardiac wall dysfunction was revealed in 5/31 patients (16.1%); mitral valve regurgitation and tricuspid valve insufficiency of grade 1 or 2 were the most common cardiac abnormalities

Table IV. Patient-related risk factors for cardiac morbidity in patients with operable and metastatic breast cancer.

Patient-related risk factors for cardiac morbidity	Patients receiving adjuvant trastuzumab treatment (N=47)		Patients receiving palliative trastuzumab treatment (N=31)	
	n	%	n	%
Hypertension	23	48.9	19	61.3
Diabetes mellitus	6	12.8	3	9.7
Ischemic heart disease	3	3.4	1	3.2
Smoking	15	31.9	4	11.0
Median BMI (kg/m ²)	26.6		29.5	
≥Median	24	51.1	16	51.6
<Median	23	48.9	15	48.4

N=Number, BMI=Body Mass index, kg=kilogram, m²=quadratmeter.

(83.9% and 45.2%, respectively), while an aortic valve abnormality was reported in three patients (9.7%) (Table V). In the overall population, 31/133 (23.3%) follow-up echocardiographic tests showed an abnormal LVEF change. The mean (±SE) decrease in the LVEF was 7.85±2.11%. The development of an abnormal LVEF was observed at least once in 13/31 (41.9%) patients: in 11/31 (35.5%), in the first year of therapy and in 6/15 (40%), during the second year. However, in 4/31 (12.9%) cases this was merely an isolated accidental finding. The mean (±SE) decrease in the LVEF in the first year in these 11 patients was 8.7±2.3%. For the six patients who showed abnormal LVEF changes in the second year only, the mean (±SE) decrease was 8.1±2.9%. None of the patients developed symptomatic CHF. Three of the patients had an atrial septum aneurysm, while one patient exhibited pericardial fluid during the treatment.

More abnormal LVEF change was detected in patients with than in those without hypertension (Table VII). Likewise, there was a difference in the second year among those who had received RT and those who had not (Table VII). No difference was found concerning the presence of diabetes or ischemic heart disease, although the numbers of such cases were low. No association was detected between the incidence of abnormal LVEF findings and BMI, treatment with or the dose of anthracyclines, the lines or cycles of previous chemotherapies, nor the cumulative dose of trastuzumab.

Cardiac dysfunction detected during the second year was related to previous left-sided RT (Table VII).

Discussion

In our study, the incidence of asymptomatic cardiac dysfunction categorized as significant was higher than reported in large clinical trials (9-11, 27-35). None of our

Table V. Heart function parameters at baseline in patients receiving adjuvant/palliative trastuzumab treatment.

Heart function parameter at baseline	Population treated with trastuzumab in the adjuvant setting (N=47)	Population treated with trastuzumab in the palliative setting (N=31)
LVEF (mean±SE, %)	67.4±0.9	66.2±1.1
LVEF (median, %)	67.0	66.0
Wall dysfunction (N, %)	0 (0.0)	5 (16.1)
Mitral insufficiency (N, %)	20 (42.6)	26 (83.9)
Tricuspidal insufficiency (N, %)	14 (29.8)	14 (45.2)
Aortic valve abnormality (N %)	2 (4.3)	3 (9.7)
Pulmonary valve abnormality (N, %)	2 (4.3)	0 (0.0)

N=Number, SE=standard error.

patients, however, developed symptomatic CHF, and approximately one-third of the cases in both groups exhibited merely an isolated accidental 'significant' LVEF decrease. The differences between our data and the published results may stem from the relatively high proportion of patients with risk factors for cardiovascular morbidity in our study. The data from the different trials are also not fully comparable as the assessment criteria, the definitions of cardiotoxicity and the testing methods sometimes differ somewhat; nonetheless, our methods did not differ in any fundamental way from those in the literature (9, 10, 29, 36).

Cardiac abnormalities not related to pumping function during trastuzumab therapy are under-reported. In our study, an asymptomatic atrial septum aneurysm was detected in 8.5% of the cases in the adjuvant group, and in 13.0% in the palliative group. Piotrowski *et al.* detected rare asymptomatic left and right bundle branch blocks and other ECG abnormalities (29). Olin *et al.* described newly-developed T-wave inversions (37). The most clearly established risk factors of trastuzumab-related cardiac events are the additive effect of concurrent trastuzumab and anthracycline treatment, previous anthracycline exposure and the cumulative dose of anthracyclines received (16). Trastuzumab seems to change the tertiary structure of the cardiac contractile apparatus reversibly by inhibiting the HER2 receptors of cardiomyocytes, and to lead to progressive myocardial cell apoptosis and destruction (38). This potentiates the adverse effects of anthracyclines, *i.e.* the irreversible apoptosis of myocytes induced by reactive oxygen species (22). Hence, trastuzumab is administered following anthracycline treatment. Heavy alcohol abuse (39), co-existing diabetes (40) and older age (11, 32, 36), overweight and obesity, hypertension (11, 36), RT (41) or genetic factors (22) were suggested as predictors of trastuzumab-induced cardiac

Table VI. Incidence of abnormal cardiac tests during and after adjuvant trastuzumab therapy in relation to selected risk factors.

	First year		Second year	
	Number of patients with risk factors (N=47)	Number of abnormal findings/ number of measurements (N=179)	Number of patients with risk factors (N=36)	Number of abnormal findings/ number of measurements (N=53)
Hypertension				
Yes	23	27/88 (30.7%)	15	3/23 (13%)
No	24	9/91 (9.9%)	21	6/30 (20%)
<i>p</i> -Value	<0.001		0.387	
BMI				
>26.6 kg/m ²	23	26/90 (28.9%)	18	7/28 (25%)
≤26.6 kg/m ²	24	10/89 (11.2%)	18	2/25 (8%)
<i>p</i> -Value	0.003		0.099	
Left-sided RT				
Yes	17	15/60 (25%)	12	7/21 (33.3%)
No	30	21/119 (17.6%)	24	2/32 (6.3%)
<i>p</i> -Value	0.168		0.015	

N=Number, BMI=Body Mass index, kg=kilogram, m²=quadratmeter, RT=radiotherapy.

adverse effects. In our study, among patients receiving adjuvant trastuzumab therapy, hypertension and high BMI proved to be significant risk factors during the first year, as did left-sided RT during the second year of observation. In the adjuvant setting, no association was found between trastuzumab-related cardiotoxicity and diabetes mellitus, ischemic heart disease or the use or dose of anthracyclines. In the metastatic group, only left-sided RT was a significant risk factor during the second year of treatment.

Since trastuzumab-related cardiotoxicity is progressive and initially asymptomatic, regular cardiac monitoring (using multiple-gated acquisition scans or echocardiography) is essential; the discontinuation of trastuzumab therapy and specific therapy are justified in rare cases (24, 25). Besides imaging methods, the measurement of plasma markers, such as N-terminal pro-B-type natriuretic peptide as a marker of myocardial strain, troponin-I as a sign of cardiac cell apoptosis and necrosis (42) or high-sensitivity C-reactive protein (43), may be used for the early detection of a cardiac dysfunction.

The limitations of our study include the small size of the population included and the short follow-up period.

In conclusion, given the substantial survival benefits of trastuzumab therapy that have been recorded in both early and MBC, the risk of cardiac complications (among which an LVEF decrease during trastuzumab treatment is the most

Table VII. The incidence of abnormal cardiac tests during and after palliative trastuzumab therapy in relation to selected risk factors.

	First year		Second year	
	Number of patients with risk factors (N=31)	Number of abnormal findings/ number of measurements (N=97)	Number of patients with risk factors (N=14)	Number of abnormal findings/ number of measurements (N=53)
Hypertension				
Yes	19	17/63 (27.0%)	12	9/28 (32.1%)
No	12	4/34 (11.8%)	2	1/8 (12.5%)
<i>p</i> -Value	0.066		0.269	
Left-sided RT				
Yes	12	9/34 (26.5%)	5	7/15 (46.7%)
No	19	12/63 (19.0%)	9	3/21 (14.3%)
<i>p</i> -Value	0.275		0.039	

N=Number, RT=radiotherapy.

common) can generally be justified. Our findings suggest that trastuzumab-related cardiac dysfunction is manageable, and largely reversible; it does not exclude patients from further treatment with trastuzumab. All patients should be evaluated for their cardiovascular status and risk factors for type-2 CRCD prior to the initiation of trastuzumab treatment, and cardiac surveillance should be routinely performed during treatment with trastuzumab.

References

- Ross JS and Fletcher JA: The HER2/neu oncogene in breast cancer: prognostic factor, predictive factor, and target for therapy. *Stem Cell* 16: 413-428, 1998.
- Slamon DJ, Clark GM, Wong SG, Levin WJ, Ullrich A and McGuire WL: Human breast cancer: correlation of relapse and survival with amplification of the HER2/neu oncogene. *Science* 235: 177-182, 1987.
- Hudis CA: Trastuzumab-Mechanism of action and use in clinical practice. *N Engl J Med* 357: 39-51, 2007.
- Vogel CL, Cobleigh MA, Tripathy D, Gutheil JC, Harris LN, Fehrenbacher L, Slamon DJ, Murphy M, Novotny WF, Burchmore M, Shak S, Stewart SJ and Press M: Efficacy and safety of trastuzumab as a single agent in first-line treatment of HER2-overexpressing metastatic breast cancer. *J Clin Oncol* 20: 719-726, 2002.
- Cobleigh MA, Vogel CL, Tripathy D, Robert NJ, Scholl S, Fehrenbacher L, Wolter JM, Paton V, Shak S, Lieberman G and Slamon DJ: Multinational study of the efficacy and safety of humanized anti-HER2 monoclonal antibody in women who have HER2-overexpressing metastatic breast cancer that has progressed after chemotherapy for metastatic disease. *J Clin Oncol* 17: 2639-2648, 1999.

- 6 Slamon DJ, Leyland-Jones B, Shak S, Fuchs H, Paton V, Bajamonde A, Fleming T, Eiermann W, Wolter J, Pegram M, Baselga J and Norton L: Use of chemotherapy plus a monoclonal antibody against HER2 for metastatic breast cancer that overexpresses HER2. *N Engl J Med* 344: 783-792, 2001.
- 7 Marty M, Cognetti F, Maraninchi D, Snyder R, Mauriac L, Tubiana-Hulin M, Chan S, Grimes D, Antón A, Lluch A, Kennedy J, O'Byrne K, Conte P, Green M, Ward C, Mayne K and Extra JM: Randomized phase II trial of the efficacy and safety of trastuzumab combined with docetaxel in patients with human epidermal growth factor receptor 2-positive metastatic breast cancer administered as first-line treatment: the M77001 study group. *J Clin Oncol* 23: 4265-4274, 2005.
- 8 Joensuu H, Kellokumpu-Lehtinen PL, Bono P, Alanko T, Kataja V, Asola R, Utriainen T, Kokko R, Hemminki A, Tarkkanen M, Turpeenniemi-Hujanen T, Jyrkkiö S, Flander M, Helle L, Ingalsuo S, Johansson K, Jääskeläinen AS, Pajunen M, Rauhala M, Kaleva-Kerola J, Salminen T, Leinonen M, Elomaa I, Isola J and FinHer Study Investigators: Adjuvant docetaxel or vinorelbine with or without trastuzumab for breast cancer. *N Engl J Med* 354: 809-820, 2006.
- 9 Piccart-Gebhart MJ, Procter M, Leyland-Jones B, Goldhirsch A, Untch M, Smith I, Gianni L, Baselga J, Bell R, Jackisch C, Cameron D, Dowsett M, Barrios CH, Steger G, Huang CS, Andersson M, Inbar M, Lichinitser M, Láng I, Nitz U, Iwata H, Thomssen C, Lohrisch C, Suter TM, Rüschoff J, Suto T, Gatrexore V, Ward C, Straehle C, McFadden E, Dolci MS, Gelber RD and Herceptin Adjuvant (HERA) Trial Study Team: Trastuzumab after adjuvant chemotherapy in HER2-positive breast cancer. *N Engl J Med* 353: 1659-1672, 2005.
- 10 Slamon D, Eiermann W, Robert N, Pienkowski T, Martin M, Press M, Mackey J, Glaspy J, Chan A, Pawlicki M, Pinter T, Valero V, Liu MC, Sauter G, von Minckwitz G, Viscio F, Bee V, Buyse M, Bendahmane B, Tabah-Fisch I, Lindsay MA, Riva A, Crown J and Breast Cancer International Research Group: Adjuvant trastuzumab in HER2-positive breast cancer. *N Engl J Med* 365: 1273-1283, 2011.
- 11 Perez EA, Suman VJ, Davidson NE, Sledge GW, Kaufman PA, Hudis CA, Martino S, Gralow JR, Dakhil SR, Ingle JN, Winer EP, Gelmon KA, Gersh BJ, Jaffe AS and Rodeheffer RJ: Cardiac safety analysis of doxorubicin and cyclophosphamide followed by paclitaxel with or without trastuzumab in the North Central Cancer Treatment Group N9831 adjuvant breast cancer trial. *J Clin Oncol* 26: 1231-1238, 2008.
- 12 Moja L, Tagliabue L, Balduzzi S, Parmelli E, Pistotti V, Guarneri V and D'Amico R: Trastuzumab containing regimens for early breast cancer. *Cochrane Database Syst Rev* 4: CD006243, 2012.
- 13 Perez EA, Romond EH, Suman VJ, Jeong JH, Davidson NE, Geyer CE Jr, Martino S, Mamounas EP, Kaufman PA and Wolmark N: Four-year follow-up of trastuzumab plus adjuvant chemotherapy for operable human epidermal growth factor receptor 2-positive breast cancer: joint analysis of data from NCCTG N9831 and NSABP B-31. *J Clin Oncol* 29: 3366-3373, 2011.
- 14 de Azambuja E, Procter MJ, van Veldhuisen DJ, Agbor-Tarh D, Metzger-Filho O, Steinseifer J, Untch M, Smith IE, Gianni L, Baselga J, Jackisch C, Cameron DA, Bell R, Leyland-Jones B, Dowsett M, Gelber RD, Piccart-Gebhart MJ and Suter TM: Trastuzumab-associated cardiac events at 8 years of median follow-up in the Herceptin Adjuvant Trial (BIG 1-01). *J Clin Oncol* 32: 2159-2165, 2014.
- 15 Bria E, Cuppone F, Fornier M, Nisticò C, Carlini P, Milella M, Sperduti I, Terzoli E, Cognetti F and Giannarelli D: Cardiotoxicity and incidence of brain metastases after adjuvant trastuzumab for early breast cancer: the dark side of the moon? A meta-analysis of the randomized trials. *Breast Cancer Res Treat* 109: 231-239, 2008.
- 16 Seidman A, Hudis C, Pierri MK, Shak S, Paton V, Ashby M, Murphy M, Stewart SJ and Keefe D: Cardiac dysfunction in the trastuzumab clinical trials experience. *J Clin Oncol* 20: 1215-1221, 2002.
- 17 Suter TM, Cook-Bruns N and Barton C: Cardiotoxicity associated with trastuzumab (Herceptin) therapy in the treatment of metastatic breast cancer. *Breast* 13: 173-183, 2004.
- 18 Smith I, Procter M, Gelber RD, Guillaume S, Feyereislova A, Dowsett M, Goldhirsch A, Untch M, Mariani G, Baselga J, Kaufmann M, Cameron D, Bell R, Bergh J, Coleman R, Wardley A, Harbeck N, Lopez RI, Mallmann P, Gelmon K, Wilcken N, Wist E, Sánchez Rovira P, Piccart-Gebhart MJ and HERA study team: 2-year follow-up of trastuzumab after adjuvant chemotherapy in HER2-positive breast cancer: a randomised controlled trial. *Lancet* 369: 29-36, 2007.
- 19 Ewer MS, Voelletich MT, Durand JB, Woods ML, Davis JR, Valero V and Lenihan DJ: Reversibility of trastuzumab-related cardiotoxicity: new insights based on clinical course and response to medical treatment. *J Clin Oncol* 23: 7820-7826, 2005.
- 20 Perez EA and Rodeheffer R: Clinical cardiac tolerability of trastuzumab. *J Clin Oncol* 22: 322-329, 2004.
- 21 Ewer SM and Ewer MS: Cardiotoxicity profile of trastuzumab. *Drug Saf* 31: 459-467, 2008.
- 22 Onitilo AA, Engel JM and Stankowski RV: Cardiovascular toxicity associated with adjuvant trastuzumab therapy: prevalence, patient characteristics, and risk factors. *Ther Adv Drug Saf* 5: 154-166, 2014.
- 23 Wolff AC, Hammond ME, Schwartz JN, Hagerty KL, Allred DC, Cote RJ, Dowsett M, Fitzgibbons PL, Hanna WM, Langer A, McShane LM, Paik S, Pegram MD, Perez EA, Press MF, Rhodes A, Sturgeon C, Taube SE, Tubbs R, Vance GH, van de Vijver M, Wheeler TM, Hayes DF; American Society of Clinical Oncology and College of American Pathologists: American Society of Clinical Oncology/College of American Pathologists guideline recommendations for human epidermal growth factor receptor 2 testing in breast cancer. *J Clin Oncol* 25: 118-145, 2007.
- 24 Lang RM, Bierig M, Devereux RB, Flachskampf FA, Foster E, Pellikka PA, Picard MH, Roman MJ, Seward J, Shanewise JS, Solomon SD, Spencer KT, Sutton MS, Stewart WJ; Chamber Quantification Writing Group; American Society of Echocardiography's Guidelines and Standards Committee and European Association of Echocardiography: Recommendations for chamber quantification: a report from the American Society for Echocardiography's Guidelines and Standards Committee and the Chamber Quantification Writing Group, developed in conjunction with the European Association of Echocardiography, a branch of the European Society of Cardiology. *J Am Soc Echocardiography* 18: 1440-1446, 2005.
- 25 Mackey JR, Clemons M, Côté MA, Delgado D, Dent S, Paterson A, Provencher L, Sawyer MB and Verma S: Cardiac management adjuvant trastuzumab therapy: recommendation of the Canadian Working Group. *Current Oncol* 15: 24-35, 2008.

- 26 Jones AL, Barlow M, Barrett-Lee PJ, Canney PA, Gilmour IM, Robb SD, Plummer CJ, Wardley AM and Verrill MW: Management of cardiac health in trastuzumab-treated patients with breast cancer: updated United Kingdom National Cancer Research Institute recommendations for monitoring. *Br J Cancer* *100*: 684-692, 2009.
- 27 Chang HR: Trastuzumab-based neoadjuvant therapy in patients with HER2-positive breast cancer. *Cancer* *116*: 2856-2867, 2010.
- 28 Láng I, Bell R, Feng FY, Lopez RI, Jassem J, Semiglazov V, Al-Sakaff N, Heinzmann D and Chang J: Trastuzumab retreatment after relapse on adjuvant trastuzumab therapy for human epidermal growth factor receptor 2-positive breast cancer: final results of the Retreatment after HERceptin Adjuvant trial. *Clin Oncol (R Coll Radiol)* *26*: 81-89, 2014.
- 29 Piotrowski G, Gawor R, Stasiak A, Gawor Z, Potemski P and Banach M: Cardiac complications associated with trastuzumab in the setting of adjuvant chemotherapy for breast cancer overexpressing human epidermal growth factor receptor type 2 - a prospective study. *Arch Med Sci* *8*: 227-235, 2012.
- 30 Kelly H, Kimmick G, Dees EC, Collichio F, Gatti L, Sawyer L, Ivanova A, Dressler L, Graham ML and Carey LA: Response and cardiac toxicity of trastuzumab given in conjunction with weekly paclitaxel after doxorubicin/cyclophosphamide. *Clin Breast Cancer* *7*: 237-243, 2006.
- 31 McArthur HL and Chia S: Cardiotoxicity of trastuzumab in clinical practice. *N Engl J Med* *357*: 94-95, 2007.
- 32 Wadhwa D, Fallah-Rad N, Grenier D, Krahn M, Fang T, Ahmadi R, Walker JR, Lister D, Arora RC, Barac I, Morris A and Jassal DS: Trastuzumab mediated cardiotoxicity in the setting of adjuvant chemotherapy for breast cancer: a retrospective study. *Breast Cancer Res Treat* *117*: 357-364, 2009.
- 33 Tarantini L, Cioffi G, Gori S, Tuccia F, Boccardi L, Bovelli D, Lestuzzi C, Maurea N, Oliva S, Russo G, Faggiano P and Italian Cardio-Oncologic Network: Trastuzumab adjuvant chemotherapy and cardiotoxicity in real-world women with breast cancer. *J Card Fail* *18*: 113-119, 2012.
- 34 Gori S, Colozza M, Mosconi AM, Franceschi E, Basurto C, Cherubini R, Sidoni A, Rulli A, Bisacci C, De Angelis V, Crinò L and Tonato M: Phase II study of weekly paclitaxel and trastuzumab in anthracycline- and taxane-pretreated patients with HER2-overexpressing metastatic breast cancer. *Br J Cancer* *90*: 36-40, 2004.
- 35 Esteva FJ, Valero V, Booser D, Guerra LT, Murray JL, Pusztai L, Cristofanilli M, Arun B, Esmaeli B, Fritsche HA, Sneige N, Smith TL and Hortobagyi GN: Phase II study of weekly docetaxel and trastuzumab for patients with HER2-overexpressing metastatic breast cancer. *J Clin Oncol* *20*: 1800-1808, 2002.
- 36 Tan-Chiu E, Yothers G, Romond E, Geyer CE Jr, Ewer M, Keefe D, Shannon RP, Swain SM, Brown A, Fehrenbacher L, Vogel VG, Seay TE, Rastogi P, Mamounas EP, Wolmark N and Bryant J: Assessment of cardiac dysfunction in a randomized trial comparing doxorubicin and cyclophosphamide followed by paclitaxel, with or without trastuzumab as adjuvant therapy in node-positive, human epidermal growth factor receptor 2-overexpressing breast cancer: NSABP B-31. *J Clin Oncol* *23*: 7811-7819, 2005.
- 37 Olin RL, Desai SS, Fox K and Davidson R: Non-myopathic cardiac events in two patients treated with trastuzumab. *Breast J* *13*: 211-212, 2007.
- 38 Singh KK, Shukla PC, Quan A, Lovren F, Pan Y, Wolfstadt JJ, Gupta M, Al-Omran M, Leong-Poi H, Teoh H and Verma S: Herceptin, a recombinant humanized anti-ERBB2 monoclonal antibody, induces cardiomyocyte death. *Biochem Biophys Res Commun* *411*: 421-426, 2011.
- 39 Lemieux J, Diorio C, Côté MA, Provencher L, Barabé F, Jacob S, St-Pierre C, Demers E, Tremblay-Lemay R, Nadeau-Larochelle C, Michaud A and Laflamme C: Alcohol and HER2 polymorphisms as risk factor for cardiotoxicity in breast cancer treated with trastuzumab. *Anticancer Res* *33*: 2569-2576, 2013.
- 40 Serrano C, Cortés J, De Mattos-Arruda L, Bellet M, Gómez P, Saura C, Pérez J, Vidal M, Muñoz-Couselo E, Carreras MJ, Sánchez-Ollé G, Tabernero J, Baselga J and Di Cosimo S: Trastuzumab-related cardiotoxicity in the elderly: a role for cardiovascular risk factors. *Ann Oncol* *23*: 897-902, 2012.
- 41 Cao L, Hu WG, Kirova YM, Yang ZZ, Cai G, Yu XL, Ma JL, Guo XM, Shao ZM and Chen JY: Potential impact of cardiac dose-volume on acute cardiac toxicity following concurrent trastuzumab and radiotherapy. *Cancer Radiother* *18*: 119-124, 2014.
- 42 Goel S, Simes RJ and Beith JM: Exploratory analysis of cardiac biomarkers in women with normal cardiac function receiving trastuzumab for breast cancer. *Asia Pac J Clin Oncol* *7*: 276-280, 2011.
- 43 Onitilo AA, Engel JM, Stankowski RV, Liang H, Berg RL and Doi SA: High-sensitivity C-reactive protein (hs-CRP) as a biomarker for trastuzumab-induced cardiotoxicity in HER2-positive early-stage breast cancer: a pilot study. *Breast Cancer Res Treat* *134*: 291-298, 2012.

Received April 21, 2015

Revised May 25, 2015

Accepted May 27, 2015