

# SUPPLEMENTARY INFORMATION


1

2       **Arctic sea ice melt leads to atmospheric**  
3                   **new particle formation**


4

5       **M. Dall’Osto, D. Beddows, P. Tunved, R. Krejci, J. Ström, H-C**  
6       **Hansson, Y.J. Yoon, Ki-Tae Park, S. Becagli, R. Udisti, T.**  
7       **Onasch, C.D. O’Dowd, R. Simo and Roy M. Harrison**


8



**Supplementary Figure S1.** Plot of Dunn Index and Silhouette Width against cluster number to identify the main cluster groups within the data. Cluster number 9 was been identified as the optimum number of clusters within the daily DMPS average number size distributions. Dashed line marks the optimum (9<sup>th</sup>) clusters.




(a)




(b)

**Supplementary Figure S2.** Real time concentrations ( $\text{cm}^{-3}$ ) data examples of (a) Nucleation event with growth and (b) nascent ultrafine particles with no subsequent growth. The total particle number concentration ( $N_{10-500\text{nm}}$ ) was found correlated with concurrent measurements taken of total particle number concentrations with CPC3010 and CPC2025, allowing to obtain the  $N_{3-10\text{nm}}$  concentrations.



**Supplementary Figure S3.** Air mass back trajectory cluster analysis results (left) and annual variation of the percentage of the monthly cluster count (right). This plot was created using the R software (R Development Core Team, R i386 3.3.2; [www.r-project.org](http://www.r-project.org)).



**Supplementary Figure S4.** Calculation of Condensational Sink ( $s^{-1}$ ) for the five aerosol categories

SMPS cluster	Air mass classification					Geographical region			
	1	2	3	4	5	sea	sea ice	snow	land
	N	N	SW	E	S				
<b>Nucleation</b>	14	7	25	15	38	43	39	14	4
<b>Bursting</b>	10	6	28	19	37	43	37	17	3
<b>Nascent</b>	10	6	22	20	42	51	29	13	7
<b>Accumulation I</b>	18	24	12	31	14	14	62	23	1
<b>Accumulation II</b>	11	15	19	23	30	32	45	20	3

**Supplementary Table S1.** Relation of aerosol SMPS cluster analysis (Figure 1) with air mass back trajectories cluster analysis (cluster 1-5: main origin N: North, SW: South West, E: East and S: South; Figure S3) and geographical region (sea, sea ice, snow, land). Percentages for air mass classification (Figure S3) are normalized over aerosol cluster category (Figure 1). Percentages of geographical region are normalised by the period of passage over land, snow, open water and sea ice .