

SCIENTIFIC REPORTS

OPEN

Phylogenetic and functional potential links pH and N₂O emissions in pasture soils

Md Sainur Samad¹, Ambarish Biswas¹, Lars R. Bakken², Timothy J. Clough³, Cecile A. M. de Klein⁴, Karl G. Richards⁵, Gary J. Lanigan⁵ & Sergio E. Morales¹

Received: 16 August 2016

Accepted: 10 October 2016

Published: 26 October 2016

Denitrification is mediated by microbial, and physicochemical, processes leading to nitrogen loss via N₂O and N₂ emissions. Soil pH regulates the reduction of N₂O to N₂, however, it can also affect microbial community composition and functional potential. Here we simultaneously test the link between pH, community composition, and the N₂O emission ratio (N₂O/(NO + N₂O + N₂)) in 13 temperate pasture soils. Physicochemical analysis, gas kinetics, 16S rRNA amplicon sequencing, metagenomic and quantitative PCR (of denitrifier genes: *nirS*, *nirK*, *nosZI* and *nosZII*) analysis were carried out to characterize each soil. We found strong evidence linking pH to both N₂O emission ratio and community changes. Soil pH was negatively associated with N₂O emission ratio, while being positively associated with both community diversity and total denitrification gene (*nir* & *nos*) abundance. Abundance of *nosZII* was positively linked to pH, and negatively linked to N₂O emissions. Our results confirm that pH imposes a general selective pressure on the entire community and that this results in changes in emission potential. Our data also support the general model that with increased microbial diversity efficiency increases, demonstrated in this study with lowered N₂O emission ratio through more efficient conversion of N₂O to N₂.

The Anthropocene has resulted in a loss of global biodiversity and enhanced greenhouse gas emissions¹. A major driver of change has been the transformation of land for agriculture purposes, needed to sustain the expanding global populations². These changes are expected to drive further reductions in biodiversity and the loss of associated ecosystem services³. Of the greenhouse gases associated with agriculture, nitrous oxide (N₂O) is of particular concern due to its global warming potential (>300 times more powerful as CO₂) and ozone-depleting capabilities⁴⁻⁷.

The mechanisms that control N₂O production and loss from soils are still being debated, with identified regulators comprising physical, chemical and biological factors⁸. Soil pH has been identified as a master regulator of gaseous N emissions, with the propensity of soils to release N₂O over N₂ tightly linked to this⁹. Two mechanisms have been proposed for explaining the role of pH: (i) a distal impact on the genetic potential in soils through re-arrangements of the microbial community and (ii) a proximal impact driven by modulation of the direct reactions catalysing the conversion of N₂O to N₂ by microbial enzymes¹⁰. However, emissions of N₂O are controlled at multiple levels: (i) the available genetic potential within the soil microbial community (genotype)¹¹, (ii) the activation or de-activation of the potential in response to an environmental signal (transcriptional regulation controlling expression of genotype)^{12,13}, (iii) the translation of transcripts leading to an immature or apoprotein (translational regulation)¹⁴, (iv) maturation of a protein resulting in an active enzyme (post-translational regulation)¹⁴, (v) export of enzymes when activity is not cytoplasmic (e.g. *sec/tat* dependent secretion as is the case for *NosZ*)¹⁵⁻¹⁷, and (vi) degradation or turnover rate of enzymes once active¹⁸. These controls cover both the production of N₂O and the consumption, or turnover, into N₂ by a different process. As a result emissions are limited by what may be summarized as: (i) genetic potential, (ii) transcriptional regulation, and (iii) enzymatic activity. The outcome is a complex array of regulators and processes that are likely to change across time and space.

¹Department of Microbiology and Immunology, Otago School of Medical Sciences, University of Otago, Dunedin, New Zealand. ²Department of Environmental Sciences, Norwegian University of Life Sciences, Ås, Norway. ³Department of Soil and Physical Sciences, Lincoln University, Lincoln, New Zealand. ⁴AgResearch Invermay, Mosgiel, New Zealand. ⁵Teagasc, Environmental Research Centre, Johnstown Castle, Wexford, Ireland. Correspondence and requests for materials should be addressed to S.E.M. (email: sergio.morales@otago.ac.nz)




Figure 1. Relationship between soil pH, N_2O emission ratio, community phylogenetic and functional potential. Relationships of $N_2O/(NO + N_2O + N_2)$ with pH (A), Shannon diversity based on 16S OTUs clustered at 97% sequence similarity (B), and total gene abundance (gene abundance per 5 ng soil DNA) for denitrification genes (*nirS*, *nirK*, *nosZI* and *nosZII*) based on qPCR (C). Changes in community composition at phylum level for Irish (IR) and New Zealand (NZ) soils ranked by country (a–c: IR; Ireland soils, d–m: NZ: New Zealand soils) and decreasing N_2O emission ratio (D). Microbial community dissimilarities of soils with different emission profiles as determined using NMDS (Bray–Curtis) ordination (E).

Despite the complexity, observations support the role of both distal and proximal regulators^{19,20}. Distal impacts by pH are proposed to be driven by selecting for community shifts at both functional and phylogenetic levels²¹ with shifts in available potential (functional gene abundances) resulting in shifts in phenotypes (observed emissions)^{22,23}. Proximal impacts by pH provide a clearer mechanism. Low pH causes a shift in active organisms²⁴, but more importantly pH disrupts the activity of the N_2O reductase by interfering with assembly^{25–27}. Although evidence supports the role of pH in regulating emissions and community structure^{28–31} studies linking all three remain sparse.

An additional consideration is the role of biodiversity in supporting ecosystem processes like N (nitrogen) cycling. It has been proposed that biodiversity is a universal regulator of ecosystem processes³². Although microbial studies that support the role of microbial diversity in controlling productivity^{33,34}, N cycling^{35–37} and even N_2O emissions³⁸ exist, these rely on single manipulated soils or small sample sizes. However, such studies serve to establish a hypothesis that aligns with ecological theory. That is, with increasing diversity there is increased redundancy and efficiency of ecosystem processes^{39,40}. This has been observed in other microbial studies^{35,41}, including those associated with N_2O emissions⁴². However, a detailed study linking gaseous emissions (NO , N_2O and N_2), pH and microbial diversity, over soils with varying parent materials and climates, is lacking.

In this study we aimed to link phenotypes (emission potential) to genotypes (functional potential and community composition) across 13 soils with varying pH (5.57–7.03) representing both Northern and Southern Hemisphere soils. These soils were selected as they represent the normally observed pH range in agronomic grasslands (recommended pH optima = 6.2–6.5). Using this dataset our goal was to simultaneously explore the relationship between pH, diversity and emissions. We hypothesized that the effect of pH on emissions would be linked to changes in whole communities, and not solely to denitrification functional potential. To test this, we quantified the abundance of genes involved in denitrification using quantitative PCR and metagenomic analysis, and examined their relationship with the emissions potential (N_2O ratio = $N_2O/(NO + N_2O + N_2)$). We also determined the microbial community composition and diversity of each soil and identified patterns linked to both changes in pH and emissions.

Results

pH dependent changes in emissions linked to denitrifier community size as well as to total community diversity and composition. The preferential loss of N from soils as N_2O , or alternatively the efficiency of conversion of N_2O to N_2 , as determined using the N_2O ratio ($N_2O/(NO + N_2O + N_2)$) was negatively associated with soil pH ($R^2 = 0.83$, $p < 0.001$) (Fig. 1A). However, when individual gases produced during denitrification were considered, pH was only strongly and inversely associated with emissions of N_2O ($R^2 = 0.62$, $p < 0.01$), with other gases showing no clear pattern (NO [$R^2 = 0.12$, $p = 0.25$], N_2 [$R^2 = 0.21$, $p = 0.11$]) (Supplementary Fig. S1). The N_2O ratio was negatively, and pH was positively, associated with microbial diversity ($R^2 = 0.57$, $p < 0.01$; $R^2 = 0.49$, $p < 0.01$), as well as to total denitrification gene (*nir* & *nos*) abundance ($R^2 = 0.57$, $p < 0.01$) (Fig. 1B,C and Supplementary Fig. S2). Across all soils the Proteobacteria, Actinobacteria

and Firmicutes phyla were the dominant phyla, and represented >75% of total microbial populations in pasture soils (Fig. 1D). Comparison of samples based on 16S rRNA community composition visualised with a non-metric multidimensional scaling (NMDS) plot, using a Bray-Curtis dissimilarity matrix, also displayed a significant link to the N₂O emission ratio and pH (Fig. 1E and Supplementary Fig. S3–S4). A Mantel test, however, supported the correlation between microbial community structure and both the N₂O ratio ($r = 0.57$, $p < 0.001$) and pH ($r = 0.61$, $p < 0.001$). A pvclust analysis (hierarchical clustering with p-values calculated via multiscale bootstrap resampling, Supplementary Fig. S5) demonstrated that while at a 95% confidence level the clusters formed represented replicates for the same site, at lower confidence levels (< 95%) soils could be clustered geographically (4 clusters: 1 Ireland; 3 New Zealand: Otago, Canterbury and North Island).

pH and the N₂O ratio correlate to distinct microbial populations. Operational taxonomic units (OTUs at 97% sequence similarity) significantly associated to changes in emissions, or pH, were identified using Spearman's rank correlation (Fig. 2). A total of 590 OTUs displaying both a statistically significant result ($p < 0.05$) and a strong effect ($r \geq 0.5$ or $r \leq -0.5$), based separately on either variable, were analysed. The number of detected OTUs was 2.5-fold larger for pH (554 OTUs) than for N₂O ratio (224 OTUs) (Fig. 2). Surprisingly, the number of OTUs either positively or negatively correlated, to either variable, was relatively conserved indicating an almost 1:1 replacement of OTUs along the gradient. For pH, 49.2% of detected OTUs were positively and 50.7% were negatively correlated, whereas for the N₂O ratio 47.8% were positively and 52.2% were negatively correlated. As a general trend, taxa showed a strongly conserved antiparallelism in relationship to pH and N₂O ratio consistent with prior trends (Fig. 1). While certain phyla displayed conserved patterns (e.g. Chloroflexi and Bacteroidetes), all phyla had examples of contrasting responses suggesting diverse life strategies. However, certain lineages at lower taxonomic levels did present consistent patterns (e.g. class Ktedonobacteria within the Chloroflexi, Subgroup 1 & 2 of the Acidobacteria, and Frankiales within the Actinobacteria). Lineages with known functional roles associated to N cycling like the Nitrospirae (positive correlation to pH and a negative correlation to N₂O ratio) and the Thaumarchaeota (mostly negative correlation to pH and a positive correlation to N₂O ratio) showed clear responses. It is also worth noting that candidate phyla (WD272, WS3) as well as other poorly studied phyla (e.g. Armatimonadetes) showed strong correlations with the N₂O ratio. For full taxonomic lineages and corresponding response to pH and emissions see Supplementary Table S1.

Linking denitrifying genes with pH and N₂O emissions. To determine the effect of varying pH on the genetic potential for denitrification, qPCR analysis was performed for key denitrification genes. Results confirmed a link between pH and the denitrification potential of soils (total [sum] abundance of all measured denitrification genes [*nirS*, *nirK*, *nosZI*, *nosZII*]). A positive association with pH ($R^2 = 0.41$, $p < 0.05$) was observed, with an inverse response observed based on emissions (negative association with N₂O ratio [$R^2 = 0.57$, $p < 0.01$]) (Fig. 3). To confirm observations, and to account for potential biases associated with primers and PCR, we determined the total abundance (per 2.63 million reads per sample) of denitrification genes in metagenomes created from 6 soils (Fig. 3 and Supplementary Fig. S6). Trends based on total denitrification gene abundance were conserved between approaches ($R^2 = 0.66$, $p < 0.05$), however, discrepancies were observed when clade specific *nosZ* gene correlations were performed. For Clade I trends were similar based on either qPCR of metagenome, although these were not statistically significant ($R^2 = 0.44$). However, results for Clade II based on metagenomic data showed a strong and statistically significant link to both pH ($R^2 = 0.69$, $p < 0.05$) and N₂O ratio ($R^2 = 0.63$, $p = 0.059$) that was not consistent with qPCR results. Despite low PCR efficiencies (average 66%), the abundance of *nosZ* genes belonging to Clade II were consistently higher than Clade I for both methods (~5-fold based on metagenome and 1.02-fold based on qPCR) (Figs 3–4). Irish soils had significantly higher numbers (1.9-fold, $p < 0.05$, Welch's *t*-test on metagenome data) of *nosZ* genes compared to New Zealand. It was also observed that taxonomic richness and diversity for Clade II was approximately 3-fold higher than for Clade I. A total of 11 different phyla (Bacteroidetes, Firmicutes Verrucomicrobia, Gemmatimonadetes, Thermomicrobia, Proteobacteria [Alpha, Beta, Delta and Gamma], Spirochaetes, Aquificae, Euryarchaeota, Crenarchaeota, and Chloroflexi) were identified based on *nosZ* sequences. The Bacteroidetes dominated those belonging to Clade II (*nosZ*) while the Alphaproteobacteria dominated within Clade I (Fig. 4 and Supplementary Fig. S7). We also examined the *nirS* and *nirK* genes individually, and found a positive association with pH ($R^2 = 0.53$, $p < 0.05$) and negative association with N₂O ratio ($R^2 = 0.38$, $p < 0.05$) for *nirS* (Supplementary Fig. S8). However, no significant associations were observed for the *nirK* gene.

Linking functional richness with pH and N₂O emissions. To account for changes in community metabolic potential outside of those previously explored, trait (function) specific patterns, associated to pH and emissions, were explored by determining the functional richness at two different levels: general N metabolism (all N cycling related genes detected) and total functional potential (total number of different genes detected). No pattern was observed between functional richness (total functional richness as well as functional richness of N-metabolism) and pH or N₂O emission ratio in the soil (Supplementary Fig. S9).

Discussion

Results support the role of native soil pH in shaping community composition and diversity. Microbial community changes were associated to both geographic changes (country and region) as well as to N₂O emissions potential, as has been described previously^{21,43}. It is important to note that N₂O emissions potential, or ratio, as defined in this study (N₂O/(NO + N₂O + N₂)) refers to the propensity of soils to emit N₂O over other denitrification gas intermediates. Here this is accomplished using a controlled environment where all other factors were held constant. While this does not reflect the absolute (total amount) of N lost through the process, it is possibly the best predictor of the propensity of the soils to emit N₂O^{8,9}. However, this potential, and the observed phenotype,




Figure 2. Taxonomic summary of OTUs significantly associated ($p < 0.05$ after BH correction; $r \geq 0.5$ [Red] or ≤ -0.5 [Green]) to either pH or N_2O emissions ratio. The graph represents a cladogram of 590 OTUs. Nodes on the tree (moving outwards from centre) correspond to taxonomic level [Domain, Phylum, Class, Order, Family, Genus and OTUs]. Shaded areas of branches delineate defined taxonomic groups. Abbreviations: S, Subgroup-22; H, Holophagae; SG, 7, 10 and 17 denotes Acidobacterial orders (subgroups); Rhodo., Rhodospirillales; Sphing., Sphingomonadales; Xantho., Xanthomonadales; Burk., Burkholderiales; Nit., Nitrosomonadales; Frank., Frankiales; Mic., Micrococcales; Thermo., Thermoleophila; Acid., Acidimicrobiia; KD4, KD4-96; An., Anaerolineae; Sphingobac., Sphingobacteriia; Cyto., Cytophagia; Flavo., Flavobacteriia; Spa., Spartobacteria; Ver., Verrucomicrobiae; Plancto., Planctomycetes; Planc., Planctomycetacia; Gemma., Gemmatimonadetes; SB, Solirubrobacterales; CO, Comamonadaceae. See Supplementary file (Table S1) for full classification.

can be modulated by fluctuating factors and require observations at the denitrification level through expression profiling (transcriptional/translational level) to identify real time drivers of N_2O emissions^{24,26,27}. Despite these limitations our observations highlight a conserved response to pH in both Northern and Southern Hemisphere soils. This suggests pH is part of a universally conserved mechanism selecting for both emissions and microbial communities. The range of pH observed in our soils (5.57–7.03) was sufficient to capture the range at which the N_2O reductase and N_2O emissions fluctuate in response to pH^{26,44–46}. Soil pH controls not only the assembly of the N_2O reductase^{26,27}, but also alters general expression patterns²⁴ and selects for shifts in microbial community composition³¹ indirectly influencing the abundance and type of functional genes in soils. Thus pH can have confounding effects due to its role in shaping the genotype, expression and eventual phenotype associated with denitrification.




Figure 3. Relationship between abundance of denitrification genes (based on absolute quantification of metagenome & qPCR abundance of *nirS*, *nirK*, *nosZI*, *nosZII*), $N_2O/(NO + N_2O + N_2)$ and pH. (A–C) Comparison of gene abundances based on either metagenomic (i.e. gene abundance per 2.63 million reads) or qPCR analysis (gene abundance per 5 ng soil DNA) for 6 soils. (D–F) Response of total denitrification genes, *nosZ* Clade I and II abundances based on metagenomic analysis for 6 soils against $N_2O/(NO + N_2O + N_2)$ (gray) and pH (black). (G–I) Response of total denitrification genes, *nosZ* Clade I and II abundances based on qPCR analysis for all 13 soils against $N_2O/(NO + N_2O + N_2)$ (gray) and pH (black).




Figure 4. Abundance (genes per 2.63 million reads) and predicted taxonomy of nitrous oxide reductase (*nosZ*) genes by soil (3 New Zealand [HT, Horotiu; LM, Lismore; TP, Templeton] and 3 Ireland soils [JT, Johnstown; SH, Solohead; MP, Moorepark]). (A), and summarized by Clade (B), based on metagenomics analysis. Clade I: Total abundance (150), Richness (4), Shannon Diversity (0.68), Evenness (0.49). Clade II: Total abundance (824), Richness (14), Shannon Diversity (1.87), Evenness (0.46).

While our findings support prior work, we show that of all the three measured gases only N_2O had a significant association with pH when compared to maximum emission levels, with maximum observed N_2O emissions decreasing with higher pH (Fig. S1). This was consistent with a lack of correlation between pH and individual denitrification genes. This is potentially due to the modular nature of denitrification^{19,47,48} where different steps within the pathway are encoded in distinct operons which do not necessarily depend on nor are associated with each other. Despite no strong correlations between emissions and denitrification specific genes, we found that of the two clades of *nosZ* gene one was dominant. Both qPCR and metagenome results show that Clade II are highly abundant, despite amplification efficiencies being poor (66%) for Clade II primers. Further, trends between metagenomic and qPCR data did not match and suggested that Clade II primers do not provide an accurate view of the abundance within our soils. Despite an apparent under representation (based on qPCR) for *nosZII*, the average Clade II/Clade I abundance ratio was >1 both for PCR-based and metagenomics analysis and is in line with prior observations of their dominance in certain soils⁴⁹. It also aligns with reports linking the abundance of Clade II with the emissions potential of soils⁵⁰. Our results also support the predicted diversity based on clade,

with Clade II being represented in almost 3-times more phyla (Fig. 4)⁵¹. Despite evidence supporting the taxonomic conservation for the two clades (different *nosZ* types are found restricted to certain microbial groups)^{50–52} our data shows that these organisms can be associated with soils displaying contrasting pH and emissions ratios.

Despite the lack of correlation between specific denitrification genes and pH, we did observe a trend of decreasing abundance of denitrification genes and overall diversity (based on 16S analysis) with decreasing pH. The role of diversity in regulating ecosystem processes has been long debated^{39,40}. The significance of microorganisms in this debate has only vaguely been addressed, relative to their predicted diversity⁵³, despite their expected importance^{54,55}. Available studies suggest that when specific microbial functional groups (i.e. methanotrophy vs respiration) are used to test diversity/ecosystem process relationships, significant trends can be uncovered^{35–37,41}. For N₂O, studies suggest that diversity plays a role, with decreases in diversity leading to increases in emissions^{38,56}. Our results support and expand on those observations indicating a role for diversity-mediated responses at multiple levels (from whole community, to specific populations linked to denitrification). Though our data do not allow a mechanism to be determined, we hypothesize that an increase in diversity ensures a steady population of microbes that are capable of sustaining a process (e.g. N₂O reduction) over a range of conditions. This diversity is still under the proximal control of regulators thus it can be modulated based on spatially and temporally controlled factors.

Identification of specific organisms responding to either pH or emissions highlighted co-varying trends. For example, while many organisms associated to changes in pH were identified as being associated to changes in emissions, not all organisms were. This implies that while certain organisms are selected by pH, they may not play a role in controlling emissions. Alternatively, some organisms that do play a role, might not be selected for by pH alone. While such correlations allow for development of new hypotheses they serve only as a first step in identifying the mechanisms controlling emissions and the role individual organisms may play. Our study also does not address the role or contributions other pathways (like nitrification) might play in regulating N₂O emissions.

Methods

Sample collection and processing. Soil samples used in this study and their physio-chemical properties have been described previously⁹. Soils were selected to represent intensive agricultural grasslands with a representative pH range close to the agronomic optimum of 6.5. Briefly, soil samples were collected from 13 permanent grassland (managed agricultural) sites in Ireland (Johnstown, Moorepark, Solohead) and New Zealand (Horotiu, Lismore, Manawatu, Mayfield, Otokia, Te Kowhai, Templeton, Tokomairiro, Warepa, Wingatui), representing Northern and Southern hemisphere sites. Soil cores (n > 3) were collected randomly from each site using a corer (25 mm diameter by 100 mm long), and excluded the grass layer. For each site, replicate cores were sieved to <4 mm, composited and immediately shipped to the Norwegian University of Life Sciences, Norway for analysis. Soil samples for kinetics were stored at 4 °C in the lab until analyzed (within one week). Soils for DNA extraction were immediately frozen and stored at –20 °C until extracted. Three separate DNA extractions were performed from 0.25 g of soil material from each site (total 39) with the PowerLyzer® PowerSoil® DNA Isolation Kit (MoBio, Carlsbad, CA) as per manufacturer's instructions. DNA concentration, purity and contamination with humics were assessed with a Nanodrop Spectrophotometer, ND-1000 (Thermo Scientific). DNA yields ranged between 8–21 ng/μl (median = 13; standard error = 0.6) with no detection of humic acids (median absorbance at 320 nm = 0.008; standard error = 0.0010) indicating high quality extractions.

Gas kinetics. Gas kinetics methods were described in detail in Samad *et al.*⁹. Briefly, soils (100 g dry weight) were provided with nitrate (2 mM NH₄NO₃) by flooding in 500 ml filter funnels (Millipore) with 4.5 cm diameter (0.2 μm) Millipore filters at least three times for 10 minutes. To obtain a homogeneous distribution of NO₃[–] and to remove excess liquid from soils a vacuum was applied. After NO₃[–] adjustment, 20 g (dry weight equivalent) of each soil was transferred to a 120 ml serum vial and sealed with an air-tight butyl-rubber septa and an aluminum crimp cap. For each site triplicate vials were prepared and incubated at 20 °C using an automated GC system⁵⁷. The soils were first incubated for 40 h under oxic conditions and then incubated under anoxic conditions for over 200 h. The emission of NO, N₂O and N₂ were measured at 5 h intervals under anoxic conditions. The product ratio of N₂O (i.e. N₂O/(NO + N₂O + N₂)) was calculated and the maximum value observed during incubation for each soil was used. The maximum value represents the highest potential of each soil to emit N₂O. While NO₃[–] concentrations are likely to see a small increase due to nitrification of the added NH₄⁺ during oxic incubation, resulting in soil-to-soil differences in available NO₃[–] at the beginning of the anoxic incubations, these differences are unlikely to affect the kinetics of denitrification (and the product ratios) since the NO₃[–] concentration applied (2 mM) was 2–3 orders of magnitude higher than *K*s for NO₃[–] reductases⁵⁸. Further, wetting of soils did not result in emissions with kinetics only measurable in the presence of exogenously added N.

Quantification of bacterial community and functional gene abundance. Quantitative PCR (qPCR) was performed on all 39 extractions to determine total bacterial abundance and the abundance of four denitrification functional marker genes (*nirS*, *nirK*, *nosZ* (Clade I) & *nosZ* (Clade II)) in each soil. Reactions were performed in 96-well plates using the ViiA7 real-time PCR system (Applied Biosystems, Carlsbad, CA). Standards for qPCR were generated using a 10-fold serial dilution (10⁸ to 10¹) of known copy numbers of pGEM-T easy (Promega, Madison, Wisconsin, USA) cloned template. All quantifications were performed using 4 technical replicates for each DNA sample loaded into the same plate, with each plate containing replicated standards and no template controls (PCR efficiencies shown in Supplementary Table S2. Amplification of *nosZ* Clade II and *nirK* targets was not possible with multiple tested polymerase brands even after optimization. As a result, two different master mixes (ABI and Thermo Scientific) were used as specific below. All reactions were performed in 20 μl volumes containing: 1 × Master Mix (ABI for *nirS* & *nosZI* or Thermo Scientific for *nirK* & *nosZII*), 0.5–1 μM of each primer (0.5 μM for *nirS* & *nosZI* and 1 μM for *nirK* & *nosZII*), 5 ng of template DNA and autoclaved Milli-Q H₂O

to a final volume of 20 μ l. Primers and qPCR conditions are summarized in Supplementary Table S2. A melt curve analysis (95 °C for 15 s, 60 °C for 1 min then increasing 0.05 °C/s (data acquisition) until 95 °C) was performed at the end of reactions to test for specificity and to confirm no amplification in the negative control. No inhibition was observed and all samples tested amplified.

Analysis of 16S rRNA gene by amplicon sequencing. 16S rRNA gene libraries were created for each DNA extraction using bacterial/archaeal primers 515F/806R targeting the V4 region of the 16S rRNA gene. Library preparation and sequencing were conducted according to the standard protocol (Version 4_13) of the Earth Microbiome Project⁵⁹ and libraries were paired-end sequenced using the Illumina MiSeq platform. Preliminary processing was carried out in Qiime (version 1.9.0) using default parameters⁶⁰. Sequences were clustered into Operational Taxonomic Units (OTUs) at 97% sequence similarity using the SILVA version 119 reference library⁶¹ and UCLUST⁶². Taxonomic classification was assigned using BLAST analysis against the SILVA database⁶³. Samples were then rarified and randomly subsampled 10 times (using the Qiime command 'multiple_rarefactions_even_depth.py') to equal depths (16,000). Samples below that threshold were removed for a total of 38 samples retained. All 10 OTU tables per sample were subsequently merged and exported for processing in R. All downstream analysis were performed in R⁶⁴ and described in detail in supplemental information. The 16S rRNA amplicon sequences were submitted to NCBI, SRA database (SRA accession: SRP080971).

Metagenomic sequence analysis. Six sites (Ireland: Johnstown, Moorepark, Solohead and New Zealand: Horotiu, Lismore, Templeton) representing a range of emission profiles from each country were selected for metagenomic analysis. Libraries for each metagenome were generated using the Illumina Nextera XT library preparation kit. Duplicate MiSeq 2 × 250 base paired end runs were carried out for each of the 6 samples. Sequences were submitted to and annotated using the MG-RAST server⁶⁵. Metagenomic data is available through the MG-RAST server (ID numbers 4644147.3 to 4644142.3). Sequence counts ranged from 2,634,050–4,851,047 before quality control. Sequences were classified taxonomically using the SILVA SSU ribosomal databases and functionally using KEGG using default settings.

Metagenome quantification of *nosZI* and *nosZII*. To differentiate between Clade I and II variants of the *nosZ* gene, a total of 1463 sequences annotated as being *nosZ* using the KO (KEGG Orthology) database were retrieved from the metagenomic libraries in our study. In order to classify them based on clade and to provide a taxonomic placement a reference database was generated. NosZ amino acid sequences were downloaded from the FunGene database⁶⁶ and classified as Clade I (*nosZI* [PRK02888;Tat dependent]) or Clade II (*nosZII* [nitrous_nosZ_Gp; Sec dependent]) based on conserved protein domains using CD-Search⁶⁷. Classification was confirmed by detection of signal peptides using the PRED-TAT algorithm⁶⁸. Taxonomy for each reference sequence was retrieved from NCBI using accession numbers associated to reference sequences. Metagenome extracted *nosZ* sequences were annotated by identifying their closest match to the reference database using BLASTX (word_size: 3, E-value:10). Matches with 60% identity and 40 amino acids coverage (cutoff) were retained and classified based on the best match. A total of 974 sequences of the original 1463 were annotated.

Statistical analyses. All statistical analyses were performed in R⁶⁴ using the phyloseq⁶⁹, pvclust⁷⁰ and vegan⁷¹ packages. Detailed descriptions can be found in supplemental methods.

References

- Vitousek, P. M., Mooney, H. A., Lubchenco, J. & Melillo, J. M. Human domination of Earth's ecosystems. *Science* **277**, 494–499 (1997).
- Tilman, D. *et al.* Agricultural sustainability and intensive production practices. *Nature* **418**, 671–677 (2002).
- Tilman, D. *et al.* Forecasting agriculturally driven global environmental change. *Science* **292**, 281–284 (2001).
- Robertson, G. P. Greenhouse gases in intensive agriculture: contributions of individual gases to the radiative forcing of the atmosphere. *Science* **289**, 1922–1925 (2000).
- Ravishankara, A. R., Daniel, J. S. & Portmann, R. W. Nitrous oxide (N₂O): the dominant ozone-depleting substance emitted in the 21st Century. *Science* **326**, 123–125 (2009).
- IPCC Climate Change 2014. *Synthesis Report* (eds Pachauri, R. K. & Meyer, L. A.) (Cambridge Univ. Press, 2014).
- Tian, H. *et al.* The terrestrial biosphere as a net source of greenhouse gases to the atmosphere. *Nature* **531**, 225–228 (2016).
- Saggar, S. *et al.* Denitrification and N₂O:N₂ production in temperate grasslands: Processes, measurements, modelling and mitigating negative impacts. *Sci Total Environ* **465**, 173–195 (2013).
- Samad, M. S. *et al.* High-resolution denitrification kinetics in pasture soils link N₂O emissions to pH, and denitrification to C mineralization. *Plos One* **11**, e0151713–11 (2016).
- Wallenstein, M. D., Myrold, D. D., Firestone, M. & Voytek, M. Environmental controls on denitrifying communities and denitrification rates: insights from molecular methods. *Ecol Appl* **16**, 2143–2152 (2006).
- Braker, G. & Conrad, R. Diversity, structure, and size of N₂O-producing microbial communities in soils—what matters for their functioning? *Adv Appl Microbiol* **75**, 33–70 (2011).
- Kern, M. & Simon, J. Three transcription regulators of the Nss family mediate the adaptive response induced by nitrate, nitric oxide or nitrous oxide in *Wolmella succinogenes*. *Environ Microbiol* **18**, 2899–2912 (2015).
- Qu, Z., Bakken, L. R., Molstad, L., Frostegård, Å. & Bergaust, L. L. Transcriptional and metabolic regulation of denitrification in *Paracoccus denitrificans* allows low but significant activity of nitrous oxide reductase under oxic conditions. *Environ Microbiol* **18**, 2951–2963 (2016).
- Dreusch, A., Bürgisser, D. M., Heizmann, C. W. & Heizmann, C. W. Lack of copper insertion into unprocessed cytoplasmic nitrous oxide reductase generated by an R20D substitution in the arginine consensus motif of the signal peptide. *Biochim Biophys Acta* **1319**, 311–318 (1997).
- Bernhard, M., Friedrich, B. & Siddiqui, R. A. *Ralstonia eutropha* TF93 is blocked in Tat-mediated protein export. *J Bacteriol* **182**, 581–588 (2000).
- Heikkilä, M. P., Honisch, U., Wunsch, P. & Zumft, W. G. Role of the Tat transport system in nitrous oxide reductase translocation and Cytochrome *cdl* biosynthesis in *Pseudomonas stutzeri*. *J Bacteriol* **183**, 1663–1671 (2001).

17. Simon, J., Einsle, O., Kroneck, P. M. H. & Zumft, W. G. The unprecedented nosgene cluster of *Wolinella succinogenes* encodes a novel respiratory electron transfer pathway to cytochrome c nitrous oxide reductase. *FEBS Lett* **569**, 7–12 (2004).
18. Vogel, C. & Marcotte, E. M. Insights into the regulation of protein abundance from proteomic and transcriptomic analyses. *Nat Rev Genet* **13**, 228–232 (2012).
19. Philippot, L., Andert, J., Jones, C. M. & Bru, D. Importance of denitrifiers lacking the genes encoding the nitrous oxide reductase for N₂O emissions from soil. *Glob Change Biol* **17**, 1497–1504 (2011).
20. Bakken, L. R., Bergaust, L. & Liu, B. Regulation of denitrification at the cellular level: a clue to the understanding of N₂O emissions from soils. *Phil Trans R Soc. B* **367**, 1226–1234 (2012).
21. Morales, S. E. & Jha, N. & Saggart, S. Biogeography and biophysicochemical traits link N₂O emissions, N₂O emission potential and microbial communities across New Zealand pasture soils. *Soil Biol Biochem* **82**, 87–98 (2015).
22. Nishizawa, T. *et al.* Inoculation with N₂-generating denitrifier strains mitigates N₂O emission from agricultural soil fertilized with poultry manure. *Biol Fertil Soils* **50**, 1001–1007 (2014).
23. Shiina, Y. *et al.* Relationship between soil type and N₂O reductase genotype (*nosZ*) of indigenous soybean Bradyrhizobia: *nosZ*-minus populations are dominant in Andosols. *Microb Environ* **29**, 420–426 (2014).
24. Brenzinger, K., Dörsch, P. & Braker, G. pH-driven shifts in overall and transcriptionally active denitrifiers control gaseous product stoichiometry in growth experiments with extracted bacteria from soil. *Front Microbiol* **6**, 1226–11 (2015).
25. Bergaust, L., Mao, Y., Bakken, L. R. & Frostegard, Å. Denitrification response patterns during the transition to anoxic respiration and posttranscriptional effects of suboptimal pH on nitrogen oxide reductase in *Paracoccus denitrificans*. *Appl Environ Microb* **76**, 6387–6396 (2010).
26. Liu, B., Mørkved, P. T., Frostegård, Å. & Bakken, L. R. Denitrification gene pools, transcription and kinetics of NO, N₂O and N₂ production as affected by soil pH. *Fems Microbiol Ecol* **72**, 407–417 (2010).
27. Liu, B., Frostegard, Å. & Bakken, L. R. Impaired reduction of N₂O to N₂ in acid soils is due to a posttranscriptional interference with the expression of *nosZ*. *mBio* **5**, e01383–14–e01383–14 (2014).
28. Nicol, G. W., Leininger, S., Schleper, C. & Prosser, J. I. The influence of soil pH on the diversity, abundance and transcriptional activity of ammonia oxidizing archaea and bacteria. *Environ Microbiol* **10**, 2966–2978 (2008).
29. Lauber, C. L., Hamady, M., Knight, R. & Fierer, N. Pyrosequencing-based assessment of soil pH as a predictor of soil bacterial community structure at the continental scale. *Appl Environ Microb* **75**, 5111–5120 (2009).
30. Čuhel, J. *et al.* Insights into the effect of soil pH on N₂O and N₂ emissions and denitrifier community size and activity. *Appl Environ Microb* **76**, 1870–1878 (2010).
31. Rousk, J. *et al.* Soil bacterial and fungal communities across a pH gradient in an arable soil. *ISME J* **4**, 1340–1351 (2010).
32. Tilman, D. The ecological consequences of changes in biodiversity: A search for general principles. *Ecology* **80**, 1455–1474 (1999).
33. Ptačník, R. *et al.* Diversity predicts stability and resource use efficiency in natural phytoplankton communities. *Proc Nat Acad Sci* **105**, 5134–5138 (2008).
34. Schnitzer, S. A. *et al.* Soil microbes drive the classic plant diversity–productivity pattern. *Ecology* **92**, 296–303 (2011).
35. Griffiths, B. S. *et al.* Ecosystem response of pasture soil communities to fumigation-induced microbial diversity reductions: an examination of the biodiversity–ecosystem function relationship. *Oikos* **90**, 279–294 (2000).
36. Wertz, S. *et al.* Maintenance of soil functioning following erosion of microbial diversity. *Environ Microbiol* **8**, 2162–2169 (2006).
37. Wittebolle, L. *et al.* Initial community evenness favours functionality under selective stress. *Nature* **458**, 623–626 (2009).
38. Wagg, C., Bender, S. F., Widmer, F. & van der Heijden, M. G. A. Soil biodiversity and soil community composition determine ecosystem multifunctionality. *Proc Nat Acad Sci* **111**, 5266–5270 (2014).
39. Loreau, M. *et al.* Biodiversity and ecosystem functioning: current knowledge and future challenges. *Science* **294**, 804–808 (2001).
40. Tilman, D., Isbell, F. & Cowles, J. M. Biodiversity and ecosystem functioning. *Annu Rev Ecol Evol Syst* **45**, 471–493 (2014).
41. Levine, U. Y., Teal, T. K., Robertson, G. P. & Schmidt, T. M. Agriculture's impact on microbial diversity and associated fluxes of carbon dioxide and methane. *ISME J* **5**, 1683–1691 (2011).
42. Domeignoz-Horta, L. A. *et al.* The diversity of the N₂O reducers matters for the N₂O:N₂ denitrification end-product ratio across an annual and a perennial cropping system. *Front Microbiol* **6**, 698–10 (2015).
43. Morales, S. E., Cosart, T. & Holben, W. E. Bacterial gene abundances as indicators of greenhouse gas emission in soils. *ISME J* **4**, 799–808 (2010).
44. Obia, A., Cornelissen, G., Mulder, J. & Dörsch, P. Effect of soil pH increase by biochar on NO, N₂O and N₂ Production during Denitrification in Acid Soils. *Plos One* **10**, e0138781–19 (2015).
45. McMillan, A. M. S. *et al.* Can pH amendments in grazed pastures help reduce N₂O emissions from denitrification? - The effects of liming and urine addition on the completion of denitrification in fluvial and volcanic soils. *Soil Biol Biochem* **93**, 90–104 (2016).
46. Russenes, A. L., Korsaaeth, A., Bakken, L. R. & rsch, P. D. Spatial variation in soil pH controls off-season N₂O emission in an agricultural soil. *Soil Biol Biochem* **99**, 36–46 (2016).
47. Zumft, W. G. Cell biology and molecular basis of denitrification. *Microbiol Mol Biol R* **61**, 533–616 (1997).
48. Philippot, L. Denitrifying genes in bacterial and Archaeal genomes. *Biochim Biophys Acta* 355–376 (2002).
49. Orellana, L. H. *et al.* Detecting nitrous oxide reductase (*nosZ*) genes in soil metagenomes: method development and implications for the nitrogen cycle. *mBio* **5**, e01193–14–e01193–14 (2014).
50. Jones, C. M., Spor, A., Brennan, F. P. & Breuil, M. C. Recently identified microbial guild mediates soil N₂O sink capacity. *Nature Climate Change* **4**, 801–805 (2014).
51. Jones, C. M., Graf, D., Bru, D., Philippot, L. & Hallin, S. The unaccounted yet abundant nitrous oxide-reducing microbial community: a potential nitrous oxide sink. *ISME J* **7**, 417–426 (2013).
52. Sanford, R. A. *et al.* Unexpected nondenitrifier nitrous oxide reductase gene diversity and abundance in soils. *Proc Nat Acad Sci* **109**, 19709–19714 (2012).
53. Locey, K. J. & Lennon, J. T. Scaling laws predict global microbial diversity. *Proc Nat Acad Sci* **113**, 5970–5975 (2016).
54. Van Der Heijden, M. G. A., Bardgett, R. D. & van Straalen, N. M. The unseen majority: soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems. *Ecol Letters* **11**, 296–310 (2008).
55. Graham, E. B. *et al.* Do we need to understand microbial communities to predict ecosystem function? A comparison of statistical models of nitrogen cycling processes. *Soil Biol Biochem* **68**, 279–282 (2014).
56. Philippot, L. *et al.* Loss in microbial diversity affects nitrogen cycling in soil. *7*, 1609–1619 (2013).
57. Molstad, L., Dörsch, P. & Bakken, L. R. Robotized incubation system for monitoring gases (O₂, NO, N₂O, N₂) in denitrifying cultures. *J Microbiol Meth* **71**, 202–211 (2007).
58. Hassan, J., Qu, Z., Bergaust, L. L. & Bakken, L. R. Transient Accumulation of NO₂⁻ and N₂O during Denitrification Explained by Assuming Cell Diversification by Stochastic Transcription of Denitrification Genes. *PLoS Comput Biol* **12**, e1004621–e1004624 (2016).
59. Caporaso, J. G. *et al.* Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq platforms. *ISME J* **6**, 1621–1624 (2012).
60. Caporaso, J. G. *et al.* QIIME allows analysis of high-throughput community sequencing data. *Nat Meth* **7**, 335–336 (2010).
61. Quast, C. *et al.* The SILVA ribosomal RNA gene database project: improved data processing and web-based tools. *Nucleic Acid Res* **41**, D590–D596 (2012).
62. Edgar, R. C. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **26**, 2460–2461 (2010).

63. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search tool. *J Mol Biol* **215**, 403–410 (1990).
64. R Development Core Team. R: A language and environment for statistical computing. <https://www.r-project.org/> (R Foundation for Statistical Computing, Vienna, Austria, 2008).
65. Meyer, F. *et al.* The metagenomics RAST server – a public resource for the automatic phylogenetic and functional analysis of metagenomes. *BMC Bioinformatics* **9**, 386 (2008).
66. Fish, J. A. *et al.* FunGene: the functional gene pipeline and repository. *Front Microbiol* **4**, 291 (2013).
67. Marchler-Bauer, A. & Bryant, S. H. CD-Search: protein domain annotations on the fly. *Nucleic Acid Res* **32**, W327–W331 (2004).
68. Bagos, P. G., Nikolaou, E. P., Liakopoulos, T. D. & Tsigirigos, K. D. Combined prediction of Tat and Sec signal peptides with hidden Markov models. *Bioinformatics* **26**, 2811–2817 (2010).
69. McMurdie, P. J. & Holmes, S. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data. *Plos One* (2013).
70. Suzuki, R. & Shimodaira, H. Pvcust: an R package for assessing the uncertainty in hierarchical clustering. *Bioinformatics* **22**, 1540–1542 (2006).
71. Oksanen, J., Blanchet, F. G., Kindt, R. & Legendre, P. *Vegan: Community Ecology Package* R package version 2.0-7 <http://CRAN.R-project.org/package=vegan> (2013).

Acknowledgements

We thank all members of the Nitrogen group at the Norwegian University of Life Sciences (<http://www.nmbu.no/en/research/groups/nitrogen>) for technical advice in gas kinetics work. We also thank Dr Xochitl Morgan for her technical assistance with data visualization. This work was funded by the New Zealand Government through the New Zealand Fund for Global Partnerships in Livestock Emissions Research to support the objectives of the Livestock Research Group of the Global Research Alliance on Agricultural Greenhouse Gases (Agreement number: 16084) awarded to SEM and the University of Otago.

Author Contributions

S.E.M., L.R.B., C.A.M.d.K., T.J.C., K.G.R. and G.J.L. designed the experiments. M.S.S., C.A.M.d.K., T.J.C., K.G.R. and G.J.L. collected samples and processed. M.S.S., A.B., S.E.M. and L.R.B. analysed the data. All authors were involved in the paper writing and revision process.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Samad, M. S. *et al.* Phylogenetic and functional potential links pH and N₂O emissions in pasture soils. *Sci. Rep.* **6**, 35990; doi: 10.1038/srep35990 (2016).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016

SCIENTIFIC REPORTS

OPEN

Erratum: Phylogenetic and functional potential links pH and N₂O emissions in pasture soils

Md Sainur Samad, Ambarish Biswas, Lars R. Bakken, Timothy J. Clough, Cecile A. M. de Klein, Karl G. Richards, Gary J. Lanigan & Sergio E. Morales

Scientific Reports 6:35990; doi: 10.1038/srep35990; published online 26 October 2016; updated on 06 December 2016

The original version of this Article contained errors in the spelling of the author Md Sainur Samad, which was incorrectly given as M.d. Sainur Samad.

These errors have now been corrected in the PDF and HTML versions of the Article.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016