

ATLAS Sensitivity to Leptoquarks, W_R and Heavy Majorana Neutrinos in Final States with High- p_T Dileptons and Jets with Early LHC Data at 14 TeV proton-proton collisions

Vikas Bansal (ATLAS Collaboration)

University of Pittsburgh, 3941 O'Hara St., Pittsburgh, PA 15260, USA

Dilepton-jet final states are used to study physical phenomena not predicted by the standard model. The ATLAS discovery potential for leptoquarks and Majorana Neutrinos is presented using a full simulation of the ATLAS detector at the Large Hadron Collider. The study is motivated by the role of the leptoquark in the Grand Unification of fundamental forces and the see-saw mechanism that could explain the masses of the observed neutrinos. The analysis algorithms are presented, background sources are discussed and estimates of sensitivity and the discovery potential for these processes are reported.

1. Introduction

The Large Hadron Collider (LHC) will soon open up a new energy scale that will directly probe for physical phenomena outside the framework of the Standard Model (SM). Many SM extensions inspired by Grand Unification introduce new, very heavy particles such as leptoquarks. Extending the SM to a larger gauge group that includes, e.g. Left-Right Symmetry (LRS) [1], could also explain neutrino masses via the see-saw mechanism. The LRS-based Left-Right Symmetric Model (LRSM) [10] used as a guide for presented studies, extends the electroweak gauge group of the SM from $SU(2)_L \times U(1)_Y$ to $SU(2)_L \times SU(2)_R \times U(1)_{B-L}$ and thereby introduces Z^0 and right-handed W bosons. If the LRS breaking in nature is such that all neutrinos become Majoranas, the LRSM predicts the see-saw mechanism [2] that elegantly explains the masses of the three light neutrinos.




Figure 1: Feynman diagrams for the pair production of scalar leptoquarks via gluon-gluon fusion (a-d) and quark anti-quark annihilation (e-f).

2. Search for scalar leptoquarks

Leptoquarks (LQ) are hypothetical bosons carrying both quark and lepton quantum numbers, as well as fractional electric charge [3, 4]. Leptoquarks could, in principle, decay into any combination of any flavor lepton and any flavor quark. Experimental limits on lepton number violation, flavor-changing neutral currents, and proton decay favor three generations of leptoquarks. In this scenario, each leptoquark couples to a lepton and a quark from the same SM generation [5]. Leptoquarks can either be produced in pairs by the strong interaction or in association with a lepton via the leptoquark-quark-lepton coupling. Figure 1 shows Feynman diagrams for the pair production of leptoquarks at the LHC.

This contribution describes the search strategy for leptoquarks decaying to either an electron and a quark or a muon and a quark leading to final states with two leptons and at least two jets. The branching fraction

of a leptoquark to a charged lepton and a quark is denoted as β^1 .

MC-simulated signal events have been studied [6] using Monte Carlo (MC) samples for first generation (1st gen.) and second generation (2nd gen.) scalar leptoquarks simulated at four masses of 300 GeV, 400 GeV, 600 GeV, and 800 GeV with the MC generator Pythia [7] at 14 TeV pp center-of-mass energy. The next to leading order (NLO) cross section [8] for the above simulated signal decreases with leptoquark mass from a few pb to a few fb with mass point of 400 GeV at (2.24 ± 0.38) pb.

¹ $\beta = 1$ would mean that leptoquarks do not decay into quarks and neutrinos.

Table I Partial cross-section (pb) that remains after each selection criterion for the first generation leptoquark channel. Baseline selection corresponds to the first selection of two electron candidates and two jets from the reconstructed objects. The Z/DY cross section is for the region $M(ee) < 60 \text{ GeV}$. VB pairs corresponds to the diboson processes of WW, WZ, and ZZ.

Physics sample	Before selection	Baseline selection	S_T 490 GeV	M_{ee} 120 GeV	$M_{lj}^1 - M_{lj}^2$ mass window [320-480] - [320-480] [GeV]
LQ (400 GeV)	2.24	1.12	1.07	1.00	0.534
Z/DY 60 GeV	1808.	49.77	0.722	0.0664	0.0036
tt	450.	3.23	0.298	0.215	0.0144
VB pairs	60.94	0.583	0.0154	0.0036	0.00048
Multijet	10^8	20.51	0.229	0.184	0.0

Table II Partial cross-section (pb) that remains after each selection criterion for the second generation leptoquark channel. Baseline selection corresponds to the first selection of two muon candidates and two jets from the reconstructed objects. The Z/DY cross section is for the region $M(\mu\mu) < 60 \text{ GeV}$. VB pairs corresponds to the diboson processes of WW, WZ, and ZZ.

Physics sample	Before selection	Baseline selection	p_T 60 GeV p_T^{jet} 25 GeV	S_T 600 GeV	$M(\mu\mu)$ 110 GeV	M_{lj} mass window [300 - 500] [GeV]
LQ (400 GeV)	2.24	1.70	1.53	1.27	1.23	0.974
Z/DY 60 GeV	1808.	79.99	2.975	0.338	0.0611	0.021
tt	450.	4.17	0.698	0.0791	0.0758	0.0271
VB pairs	60.94	0.824	0.0628	0.00846	0.00308	0.00205
Multijet	10^8	0.0	0.0	0.0	0.0	0.0

2.1. Reconstruction and objects selection

Signal reconstruction requires selection of two high quality leptons and at least two jets. Each signal jet and lepton candidate is required to have transverse momentum (p_T) $> 20 \text{ GeV}$. This helps to suppress low p_T background predicted by the SM. Leptons are required to have pseudorapidity $|\eta|$ below 2.5, which is the inner detector's acceptance, whereas jets are restricted to $|\eta| < 4.5$ to suppress backgrounds from underlying event and minimum bias events that dominate in the forward region of the detector. In addition, leptons are required to pass identification criteria, which, in case of electrons, are based on electromagnetic-shower shape variables in the calorimeter and, in the case of muons, are based on finding a common track in the muon spectrometer and the inner detector together with a muon isolation² requirement in the calorimeter. Electron candidates are also required to have a matching track in the

inner detector. Furthermore, it is required that signal jet candidates are spatially separated from energy clusters in the electromagnetic calorimeter that satisfy electron identification criteria. Finally, a pair of leptoquark candidates are reconstructed from lepton-jet combinations. Given the fact that these four objects can be combined to give two pairs, the pair that has minimum mass difference between the two leptoquark candidates is assumed to be the signal.

2.2. Background Studies

The main backgrounds to the signal come from tt, and Z=DY + jets production processes. Multijet production where two jets are misidentified as electrons, represents another background to the dielectron(1st gen.) channel. In addition, minor contributions arise from diboson production. Other potential background sources, such as single-top production, were also studied and found to be insignificant.

The backgrounds are suppressed and the signal significance is improved by taking advantage of the fact that the final state particles in signal-like events have relatively large p_T . A scalar sum of transverse momenta of signal jets and lepton candidates, denoted by S_T , helps in reducing the backgrounds while retaining most of the signal. The other variable used to increase the signal significance is the invariant mass of

² $E_T^{\text{iso}} = p_T > 0.3$, where p_T is muon candidate's transverse momentum and E_T^{iso} is energy detected in the calorimeters in a cone of $R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} = 0.2$ around muon candidate's reconstructed trajectory.

the two leptons, M_{ll} . The distributions of these two variables for the first generation channel are shown in Fig. 2.




Figure 2: S_T (top) and M_{ee} of the selected electron pair after S_T requirement (bottom) in 1st gen. leptoquark MC events ($m_{LQ} = 400$ GeV). Both distributions are normalized to 100 pb^{-1} of integrated pp luminosity.

After applying optimized selection on these two variables, S_T and M_{ll} , relative contributions from the background processes from $t\bar{t}$, $Z=DY$, diboson and multijet are 22%, 7%, 0.4% and 18%, respectively. Partial cross-section for the signal and the background processes passing the selection criteria are shown in tables I and II for the first and second generation channels, respectively. Figure 3 shows the invariant masses³ of the reconstructed leptoquark candidates before and after background suppression criteria are applied to the MC data.

2.3. Sensitivity and Discovery Potential

ATLAS's sensitivity to leptoquark signal for a 400 GeV mass hypothesis and with an integrated pp luminosity of 100 pb^{-1} is summarized in Fig. 4. The




Figure 3: Reconstructed electron-jet invariant mass for 1st gen. leptoquark ($m_{LQ} = 400$ GeV) in signal and background MC events after baseline selection (top) and after additional selection criteria based on S_T and m_{ll} (bottom) have been applied. Both distributions are normalized to 100 pb^{-1} of integrated pp luminosity.

cross-sections include systematic uncertainties of 50%. Leptoquark-like events in the ATLAS detector are triggered by single leptons with an efficiency of 97%. ATLAS is sensitive to leptoquark masses of about 565 GeV and 575 GeV for 1st and 2nd generations, respectively, at the given luminosity of 100 pb^{-1} provided the predicted cross-sections for the pair production of leptoquarks are correct.

3. Search for W_R bosons and heavy Majorana neutrinos

W_R bosons are the right-handed counterpart of the SM W bosons. These right-handed intermediate vector bosons are predicted in LRSMs and can be produced at the LHC in the same processes as the SM W 's and Z . They decay into heavy Majorana neutrinos. The Feynman diagram for W_R production and subsequent decay to Majorana neutrino is shown in Fig. 5.

This section describes the analysis of W_R production and its decays $W_R \rightarrow eN_e$ and $W_R \rightarrow N_e \nu_e$, fol-

³These distributions contain two entries per event corresponding to the two reconstructed leptoquark candidates.

Table III LR SM dielectron analysis. Partial cross-section (pb) that remains after each selection criterion for the dielectron channel.

Physics sample	Before selection	Baseline selection	M (ejj) 100 GeV	M (eejj) 1000 GeV	M (ee) 300 GeV	S_T 700 GeV
LR SM _18_3	0.248	0.0882	0.0882	0.0861	0.0828	0.0786
LR SM _15_5	0.470	0.220	0.220	0.215	0.196	0.184
Z/DY 60 GeV	1808.	49.77	43.36	0.801	0.0132	0.0064
tt	450.	3.23	3.13	0.215	0.0422	0.0165
VB pairs	60.94	0.583	0.522	0.0160	0.0016	0.0002
Multijet	10^8	20.51	19.67	0.0490	0.0444	0.0444

Table IV LR SM dimuon analysis. Partial cross-section (pb) that remains after each selection criterion for the dimuon channel.

Physics sample	Before selection	Baseline selection	M (jj) 100 GeV	M (jj) 1000 GeV	M () 300 GeV	S_T 700 GeV
LR SM _18_3	0.248	0.145	0.145	0.141	0.136	0.128
LR SM _15_5	0.470	0.328	0.328	0.319	0.295	0.274
Z/DY 60 GeV	1808.	79.99	69.13	1.46	0.0231	0.0127
tt	450.	4.17	4.11	0.275	0.0527	0.0161
VB pairs	60.94	0.824	0.775	0.0242	0.0044	0.0014
Multijet	10^8	0.0	0.0	0.0	0.0	0.0

lowed by the decays $N_e \rightarrow e^0 q$ and $N \rightarrow q^0 q$, which are detected in final states with (at least) two leptons and two jets. The two leptons can be of either same-sign or opposite-sign charge due to the Majorana nature of neutrinos. This analysis in both the dielectron and the dimuon channels has been performed without separating dileptons into same-sign and opposite-sign samples.

Studies [6] of the discovery potential for W_R and Majorana neutrinos N_e and N have been performed using MC samples where $M(N_1) = 300$ GeV; $M(W_R) = 1800$ GeV (referred to as LR SM _18_3) and $M(N_1) = 500$ GeV; $M(W_R) = 1500$ GeV (referred to as LR SM _15_5), simulated with PYTHIA according to a particular implementation [9] of LR SM [10]. The production cross-sections $(pp(14 \text{ TeV}) \rightarrow W_R X)$ times the branching fractions $(W_R \rightarrow N_1 \rightarrow lljj)$ are 24.8 pb and 47 pb for LR SM _18_3 and LR SM _15_5, respectively.

3.1. Reconstruction and objects selection

Signal event candidates are reconstructed using two electron or muon candidates and two jets that pass the standard selection criteria as discussed in section 2.1. The two signal jet candidates are combined with each of the signal leptons and the combination that gives the smaller invariant mass is assumed to be the new

heavy neutrino candidate. The other remaining lepton is assumed to come directly from the decay of the W_R boson. If signal electrons and signal jets overlap in R within 0.4 then, to avoid double counting, only the two signal jets are used to reconstruct the invariant masses of the heavy neutrino candidate and W_R .

3.2. Background Studies

The main backgrounds to the LR SM analyses studied here are the same as mentioned in section 2.2. The same background suppression criteria as in the leptoquark analyses are also effective here, namely S_T and m_{ll} . The distributions of these two variables for the dimuon channel are shown in Fig. 6. Partial cross-section for the signal and the background processes passing the selection criteria are shown in tables III and IV for the dielectron and dimuon channels, respectively. Figure 7 shows the invariant mass of the reconstructed W_R candidates before and after background suppression criteria are applied to the MC data.

3.3. Sensitivity and Discovery Potential

Signal significance for W_R analyses in the dielectron and dimuon channels as a function of integrated luminosity at 14 TeV is summarized in Fig. 8. The results include systematic uncertainty of 45% and 40% for dielectron and dimuon channel, respectively. The




Figure 4: 5 discovery potential for 1st and 2nd gen. 400 GeV scalar leptoquark versus β^2 , with and without background systematic uncertainty (top). M in μm^2 of scalar leptoquark versus leptoquark mass at 100 pb^{-1} of integrated pp luminosity at 5 (background systematic uncertainty is included) (bottom).




Figure 5: Feynman diagram for W_R production and its decay to the Majorana neutrino N at the LHC.

events in this analysis are also triggered by single leptons with an efficiency of 97%.

4. Conclusions

Dilepton-jet based final states have been discussed in both electron and muon channels. Discovery potential for leptoquarks and LRSM with early LHC data have been investigated with the predicted cross-sections for these models. Assuming $\alpha = 1$, both 1st




Figure 6: LRSM analysis. The distributions of S_T (top) and $M(\mu\mu)$ (bottom) for signals and backgrounds normalized to 100 pb^{-1} of integrated pp luminosity after baseline selection in the dimuon analysis.

and 2nd generations leptoquarks could be discovered with masses up to 550 GeV with 100 pb^{-1} of data. Two LRSM mass points LRSM_18_3 and LRSM_15_5 for the W_R bosons and heavy Majorana neutrinos have been studied. The discovery of these new particles with such masses would require integrated luminosities of 150 pb^{-1} and 40 pb^{-1} , respectively.




Figure 7: LRS M analysis. The distributions of the reconstructed invariant masses for $W_R \rightarrow N$ candidates in background and signal (LRS M_18_3 and LRS M_15_5) events before (top) and after (bottom) background suppression is performed in dimuon channel analysis. Both distributions are normalized to 100 pb^{-1} of integrated pp luminosity.

References

- [1] Pati, J. C. and Mohapatra, R. N., Phys. Rev. D 11, 566 (1975).
- [2] Mohapatra, R. N. and Senjanovic, G., Phys. Rev. D 23, 165 (1981).
- [3] Buchmuller, W. and Wyler, D., Phys. Lett. B 177, 377 (1986).
- [4] Georgi, H. and Glashow, S. L., Phys. Rev. Lett. 32, 438 (1974).

- [5] Leurer, M., Phys. Rev. D 49, 333 (1994).
- [6] Aad, G. et al. [The ATLAS Collaboration], arXiv:0901.0512 [hep-ex].
- [7] T. Sjöstrand, et al., JHEP 05 (2006) 026."
- [8] Kramer, M. et al., Phys. Rev. D 71, 057503 (2005).
- [9] Ferrari, A. et al., Phys. Rev. D 62, 013001 (2000).
- [10] Huitu, K. et al., Nuc. Phys. B 487, 27 (1997).




Figure 8: LRS M analysis. Expected signal significances versus integrated pp luminosity for N_e, N and W_R mass hypotheses, according to signal MC samples LRS M_18_3 and LRS M_15_5. Open symbols show sensitivities without systematic uncertainties. Sensitivities shown with closed symbols include an overall relative uncertainty of 45% (40%), estimated for background contributions in the dielectron (dimuon) analysis.