

1 Metal bioaccumulation and cellular fractionation in an epigeic  
2 earthworm (*Lumbricus rubellus*): the interactive influences of  
3 population exposure histories, site-specific geochemistry and  
4 mitochondrial genotype.

5

6 **Jane Andre<sup>ab\*</sup>**, **Stephen R. Stürzenbaum<sup>c</sup>**, **Peter Kille<sup>a</sup>**, **A. John Morgan<sup>a</sup>** and  
7 **Mark E. Hodson<sup>b</sup>**.

8

9 <sup>a</sup> Cardiff School of Biosciences, Cardiff University, Park Place, Cardiff, CF10 3US.

10 <sup>b</sup> Department of Soil Science, School of Human and Environmental Sciences, University of  
11 Reading, Whiteknights, Reading, RG6 6DW.

12 <sup>c</sup> King's College London, School of Biomedical & Health Sciences, Department of  
13 Biochemistry, Pharmaceutical Sciences Research Division, London, SE1 9NH

14 \* Author for correspondence. Current address: Dr. Jane Andre, School of Health and  
15 Medicine, Division of Biomedical and Life Sciences, Lancaster University, Lancaster, LA1  
16 4YQ

17

18

19

20

21

22

23

24 Abstract

25 Subcellular fractionation techniques were used to describe temporal changes (at  
26 intervals from T<sub>0</sub> to T<sub>70</sub> days) in the Pb, Zn and P partitioning profiles of *Lumbricus*  
27 *rubellus* populations from one calcareous (M<sub>DH</sub>) and one acidic (M<sub>CS</sub>) geographically  
28 isolated Pb/Zn-mine sites and one reference site (C<sub>PF</sub>). M<sub>DH</sub> and M<sub>CS</sub> individuals were  
29 laboratory maintained on their native field soils; C<sub>PF</sub> worms were exposed to both  
30 M<sub>DH</sub> and M<sub>CS</sub> soils. Site-specific differences in metal partitioning were found:  
31 notably, the putatively metal-adapted populations, M<sub>DH</sub> and M<sub>CS</sub>, preferentially  
32 partitioned higher proportions of their accumulated tissue metal burdens into insoluble  
33 CaPO<sub>4</sub>-rich organelles compared with naive counterparts, C<sub>PF</sub>. Thus, it is plausible  
34 that efficient metal immobilization is a phenotypic trait characterising metal tolerant  
35 ecotypes. Mitochondrial cytochrome oxidase II (COII) genotyping revealed that the  
36 populations indigenous to mine and reference soils belong to distinct genetic lineages,  
37 differentiated by ~13%, with 7 haplotypes within the reference site lineage but fewer  
38 (3 and 4, respectively) in the lineage common to the two mine sites. Collectively,  
39 these observations raise the possibility that site-related genotype differences could  
40 influence the toxico-availability of metals and, thus, represent a potential confounding  
41 variable in field-based eco-toxicological assessments.

42

43 Keywords: earthworms, Pb & Zn, subcellular fractionation, field & lab exposures,  
44 genotyping

45

46

47

48

49 Introduction

50 Direct toxic effects arise in metal-exposed organisms not as a consequence of the total  
51 accumulated tissue metal burden *per se* but when the rate of uptake overtakes the  
52 combined rates of excretion and detoxification, such that the internal metal-specific  
53 concentration threshold of metabolically-available metal is exceeded (Rainbow, 2007;  
54 Pan and Wang, 2008). The threshold concentration denoting the transition from no  
55 adverse effect to an observable adverse effect for a given metal is referred to as the  
56 critical body residue (CBR) (McCarthy and Mackay, 1993; Péry, *et al.*, 2005). Thus,  
57 only a fraction of the body burden is toxicologically (re)active or available (Rainbow,  
58 2002; Vijver, *et al.*, 2004). Organisms have evolved mechanisms to regulate the  
59 bioreactivities of essential and non-essential metals (Campbell, *et al.*, 2006). In  
60 general these initially entail binding and trafficking by chaperone molecules. Essential  
61 cations may subsequently be delivered to physiologically labile intracellular storage  
62 sites, classically exemplified by Ca<sup>2+</sup>-storing endoplasmic reticulum regions, whilst  
63 excess essential and non-essential cations can also either be excreted directly or  
64 immobilized as insoluble products in specialized organelles often with long half-lives.  
65 These structures possess diverse morphologies and matrix compositions (Hopkin,  
66 1989) that are generically referred to as ‘metal-rich granules’ or ‘concretions’  
67 (Campbell, *et al.*, 2006).

68

69 Improved toxic effects prediction and ecological risk assessment would be likely  
70 outcomes of a better knowledge of the fate and speciation of metal within sentinel  
71 organisms (Vijver, *et al.*, 2006; Huang, *et al.*, 2009; Jones, *et al.*, 2009). Although  
72 there is some evidence from studies on aquatic invertebrates that the toxico-available  
73 metals are associated with the cytosolic (soluble) fraction (Perceval, *et al.*, 2006; Péry,  
74 *et al.*, 2008), it is generally the case that the relationship between metal induced

75 toxicity and accumulated burden is difficult to evaluate due to the cellular  
76 compartmentalization of metals (Campbell, *et al.*, 2006; Vijver, *et al.*, 2006).  
77 Techniques such as analytical electron microscopy and synchrotron-based X-ray  
78 absorption spectroscopy have been used to some extent to characterize the ligand-  
79 binding speciation of metals and metalloids in invertebrate tissues (Cotter-Howells, *et*  
80 *al.*, 2005; Langdon, *et al.*, 2005; Arnold, *et al.*, 2008; Andre, *et al.*, 2009). However, a  
81 much more widely used method for segregating invertebrate metal burdens into  
82 operationally defined detoxified- and non-detoxified subcellular metal compartments  
83 is to differentially centrifuge tissue homogenates. To date, such studies have mainly  
84 concentrated on aquatic animals (Honeycutt, *et al.*, 1995; Wallace and Lopez, 1997;  
85 Conder, *et al.*, 2002; Wallace, *et al.*, 2003; Cain, *et al.*, 2004; Vijver, *et al.*, 2004), but  
86 there is a burgeoning body of publications on the assessment of metal partitioning in  
87 earthworms (Arnold *et al.*, 2008; Andre *et al.*, 2009; Huang, *et al.*, 2009; Vijver, *et*  
88 *al.*, 2006; Li, *et al.*, 2008; Jones, *et al.*, 2009).

89

90 Voets, *et al.*, (2009) reviewed some of the literature demonstrating that the cellular  
91 metal distribution patterns in indigenous invertebrate and vertebrate populations often  
92 differ from the distribution patterns observed in naive counterpart organisms exposed  
93 to metals in laboratory or field-based transplant experiments. Evidently both exposure  
94 history and genetic differentiation are biotic variables that can lead to modifications of  
95 the efficiency of metal detoxification by invertebrates (Wallace, *et al.*, 2003) as well  
96 as vertebrates (Knapen, *et al.*, 2004). Morgan, *et al.* (2007) also noted that the genetic  
97 background of a population can confound biomarker assays, a further indication that  
98 the balance between the sensitive and detoxified metal pools can be altered by micro-  
99 evolutionary events. Given that comprehensive phylogenetic studies on earthworms

100 using mitochondrial and nuclear markers have recently revealed high intra-species  
101 genetic diversity (Velavan, *et al.*, 2007; Novo, *et al.*, 2008) and deeply divergent  
102 genetic lineages, possibly in some cases corresponding with cryptic species (King, *et*  
103 *al.*, 2008; Shepeleva, *et al.*, 2008; Pérez-Losada, *et al.*, 2009), it is a major omission  
104 that, to the best of our knowledge, no studies hitherto have explicitly attempted to  
105 describe the cellular partitioning of metals in field populations of earthworms with  
106 respect to exposure history and genotype. A recent report (Langdon, *et al.*, 2009) that  
107 populations of the species *Lumbricus rubellus* inhabiting abandoned arsenic mine  
108 sites have evolved resistance to the metalloid brings the omission into sharp focus.

109

110 The present study had two main aims. First, to investigate the interactive influences of  
111 population exposure history and site-specific geochemistry on subcellular metal (Pb,  
112 Zn) and P partitioning by comparing two putative adapted *L. rubellus* populations  
113 sampled from geochemically contrasting disused Pb/Zn mines (one acidic and one  
114 calcareous, respectively) and maintained on their native soils with each other and with  
115 reference earthworms transferred experimentally to both polluted soils. Phosphorus  
116 partitioning was monitored because phosphate is recognised as the predominant  
117 counter-ion in earthworm Pb- and Zn-sequestering cellular compartments (Cotter-  
118 Howells, *et al.*, 2005). The second study aim was to use mitochondrial cytochrome  
119 oxidase II (COII) to genotype the three field populations. Andre, *et al.* (2010)  
120 observed site-specific differences in the tissue and subcellular partitioning profiles of  
121 *L. rubellus* populations indigenous to calcareous and acidic sites, respectively.  
122 Moreover, the authors reported that the two identified genetically distinct *L. rubellus*  
123 lineages were differentially distributed across a heterogeneous polluted landscape,  
124 with lineage 'A' predominating within a calcareous Pb/Zn-polluted 'island' and

125 lineage 'B' predominating in an adjacent acidic polluted location. The present study  
126 extended these previous observations through the novel combination of cell  
127 fractionation and genotype analyses applied to geographically isolated populations.

128

## 129 Materials and Methods

### 130 Soil and earthworm collection and preparation

131 Soil and earthworms (mature, *L. rubellus*) were collected from one control site,  
132 Pontcanna Fields ( $C_{PF}$ ) ST 165779 (GPS: 51:29.63122N 3:12.24983W) and two  
133 contaminated disused, metalliferous mine sites, Draethen Hollow ( $M_{DH}$ ) ST 217877  
134 (GPS: 51:34.96185N 3:7.88760W) and Cwmystwyth Stream ( $M_{CS}$ ) SN 803748 (GPS:  
135 52:21.48890N 3:45.54702W). At least ten soil samples (excluding the litter layer),  
136 taken from a 0-5cm depth, were randomly collected from the sampling areas,  
137 combined and mixed. The pH of all soils were measured in deionised H<sub>2</sub>O (Boisson,  
138 *et al.*, 1998) prior to them being oven dried at 30°C overnight, sieved to <2mm, then  
139 digested in boiling 16N HNO<sub>3</sub> (Morgan and Morgan, 1990) and analysed for major  
140 inorganic constituents by inductively coupled plasma - optical emission spectroscopy  
141 (ICP-OES; Perkin-Elmer Optima 3000). Analysis of an in house certified reference  
142 material (a sewage sludge amended soil) indicated that that the overall analytical error  
143 did not exceed 5.2%. In addition the calibration accuracy of the instrument was assessed  
144 through the analysis of an in-house matrix-matched standard and was within 10%. To  
145 provide an indication of the organic matter content, loss on ignition (LOI) was  
146 determined for each soil sample. 10g (dry weight) of each soil was weighed in a glass  
147 crucible and heated to 500°C overnight. The percentage weight reduction was then  
148 recorded.

149

**Comment [MEH1]:** Have to include this as otherwise the reviewer would expect a reference number to see which commercially available CRM was used. If you have the name or number of the CRM you used (normally something like SXXX I can add in the certified composition.

**Comment [MEH2]:** This reads correctly

150 Sub-cellular fractionation

151 Thirty boxes containing 300g of contaminated M<sub>CS</sub> soil and 30 boxes containing 300g  
152 of contaminated M<sub>DH</sub> soil were established. The soils were wetted to 70% of their  
153 water-holding capacity and reference site (C<sub>PF</sub>) earthworms were placed into 15 boxes  
154 (three individuals per box) of M<sub>CS</sub> and 15 boxes of M<sub>DH</sub> soil. Similarly, 3 native M<sub>DH</sub>  
155 earthworms were placed into each of the remaining 15 boxes containing M<sub>DH</sub> soil and  
156 3 native M<sub>CS</sub> earthworms into each of the remaining 15 boxes containing M<sub>CS</sub> soil. At  
157 1, 3, 14, 28 and 70 days of soil exposure, 3 boxes of each soil-earthworm combination  
158 (i.e. maximum 'n' per 'treatment' = nine) were selected at random and the  
159 earthworms depurated prior to freezing. They were depurated for an initial period of 48  
160 hours on moistened filter paper (which was changed daily to prevent coprophagy),  
161 followed by 24 hours in a filter-paper free petri-dish (with de-ionised H<sub>2</sub>O) to allow  
162 exudation of any filter paper consumed (Arnold and Hodson, 2007). The total  
163 exposure period extended to 10 weeks in order to allow sufficient time for the toxico-  
164 available fraction to stabilize, as was shown to be the case in the lumbricid species  
165 *Eisenia fetida* with no previous history of metal exposure (Jones *et al.*, 2009). When  
166 required the earthworms were defrosted, weighed, homogenized in 0.01M Tris-HCl,  
167 pH 7.5, and fractionated as described in Arnold, *et al.*, (2008) into a soluble "C  
168 fraction" (cytosolic fraction including soluble proteins such as metallothionein and  
169 heat shock proteins) and separate insoluble "D" (metal-rich granules) and "E  
170 fractions" (tissue fragments, mitochondrial and gut contents) which for the purposes  
171 of this study were combined (see supplementary Figure 1). Individual fractions were  
172 digested in boiling 16N HNO<sub>3</sub> (Morgan and Morgan, 1990). Samples were made up to  
173 volume with ultra-pure water and analysed for major inorganic constituents by ICP-  
174 OES with resulting concentrations expressed as mg of metal per kg (wet weight) of

175 earthworm. Blanks were included for each analyses and detection limits were calculated as  
176 800  $\mu\text{g L}^{-1}$ , 200  $\mu\text{g L}^{-1}$  and 400  $\mu\text{g L}^{-1}$  for P, Pb and Zn respectively. No certified reference  
177 materials exist for use with this fractionation method but previous analysis of standard  
178 additions were within range (10%), indicating good recovery in the matrix (Arnold et al.,  
179 2008) and precision, calculated from repeat analyses of samples was < 5 %. Fresh,  
180 depurated, weights of the worms sacrificed at each time interval were recorded prior  
181 to fractionation; weight change over the exposure periods were taken as estimates of  
182 physiological condition.

183

#### 184 Mitochondrial Cytochrome Oxidase II (COII) genotyping

185 *L. rubellus* earthworms from the M<sub>DH</sub> (n=22), M<sub>CS</sub> (n=32) and C<sub>PF</sub> (n=29) sites, and  
186 single *L. castaneus* and *L. eiseni* (from uncontaminated sites in Lancashire, England)  
187 were collected by digging and hand-sorting. The animals were transported back to the  
188 laboratory in their native soil and depurated (Arnold and Hodson, 2007). A short  
189 length (approximately 3cm) of posterior segments was clipped from all *L. rubellus*  
190 individuals, and genomic DNA was extracted using DNAzol reagent (Invitrogen Ltd.,  
191 Paisley, UK). Forward (5'-TAGCTCACTTAGATGCCA) and reverse (5'-  
192 GTATGCGGATTTCTAATTGT) *L. rubellus*-specific primers were used to amplify  
193 the cytochrome oxidase II (COII) gene, prior to an Exo-SAP-IT PCR clean-up and  
194 sequencing using ABI PRISM<sup>®</sup> BigDye v3.1 Terminator Sequencing technology  
195 (Applied Biosystems, USA) as described by Andre, *et al.*, (2010). Raw sequence  
196 traces were confirmed using Finch TV before being imported into Mega v3.1 (Kumar,  
197 *et al.*, 2004) for alignment and tree construction. The distance-based neighbour  
198 joining (NJ) algorithm (Saitou and Nei, 1987), using p-distance, was used to estimate  
199 tree topology and calculate branch lengths.

200

**Comment [MEH3]:** If you want to convert these into an equivalent mg per kg wet weight so that they are the same units as the results you report you need to assume a mass of earthworm digested and use this, together with the volume of digestate to calculate them

201 Results

202 Soil analysis

203 Tables 1 and 2 show the concentration of Pb, Zn and P in soil and earthworms  
204 sampled from the metalliferous M<sub>CS</sub> and M<sub>DH</sub> and reference C<sub>PF</sub> sites, as well as the  
205 percentage body weight change over the full extent of the exposure period. The Pb  
206 and Zn soil concentrations were highest at the calcareous M<sub>DH</sub> site; acidic M<sub>CS</sub> soil  
207 was only mildly contaminated, but contained significantly higher Pb and Zn  
208 concentrations than the reference C<sub>PF</sub> soil. Phosphorus concentration was significantly  
209 higher in C<sub>PF</sub> reference soil than in the two metalliferous soils. C<sub>PF</sub> earthworms  
210 maintained higher whole body P concentrations after 70 days of exposure to both  
211 metalliferous soils when compared with their M<sub>DH</sub> and M<sub>CS</sub> counterparts. Mean total  
212 earthworm tissue Pb and Zn levels to some extent reflected the corresponding soil Pb  
213 and Zn concentrations, although it is noteworthy that the worms indigenous to the  
214 acidic M<sub>CS</sub> site had a Pb bioaccumulation factor of greater than 1 (based upon dry-  
215 weight values, data not shown).

216

217 Body mass changes

218 Mortality was evident across all treatment groups and the mean fresh weights of  
219 earthworms, including C<sub>PF</sub> worms on their 'own' unpolluted reference soil, decreased  
220 considerably over the exposure period. These observations indicate that a degree of  
221 stress mediated by dietary restriction and/or metal toxicity was experienced by all  
222 earthworms in our experimental regime.

223

224 Sub-cellular fractionation

225 *Lead*: Following a ten-week exposure period, the *ex-situ* partitioning profiles were  
226 similar for both indigenous and naive introduced earthworms exposed to the same soil  
227 (Figure 1). Significant increases in Pb concentration were seen in the soluble (C) and  
228 insoluble (D+E) fractions of all worms exposed to M<sub>DH</sub> soil, and C<sub>PF</sub> individuals  
229 exposed to M<sub>CS</sub> soil. M<sub>CS</sub> individuals only demonstrated a slight increase in insoluble  
230 Pb. Destroying the physical integrity of the field soils by indiscriminate sampling,  
231 drying, sieving, homogenisation, and re-hydrating appears to have released more  
232 metal for uptake into earthworm tissues above the corresponding equilibrated field  
233 levels. Pb was found to preferentially partition into the non-soluble or detoxified  
234 (D+E) fraction in all earthworm/soil combinations (Figure 1B). Plotting the time  
235 course partitioning data with the soluble fraction Pb values expressed as a percentage  
236 of the whole body Pb concentration values (Figure 2) revealed differences in the  
237 efficiencies of incorporating Pb into the detoxified fraction between indigenous  
238 worms and naive worms introduced into the metalliferous soils. Specifically, and  
239 consistently over the entire exposure period, the proportion of Pb within the sensitive  
240 soluble fraction of M<sub>CS</sub> earthworms was proportionately less than that in C<sub>PF</sub>  
241 earthworms maintained on M<sub>CS</sub> soil (Figure 2A). A similar efficiency difference was  
242 found between M<sub>DH</sub> and C<sub>PF</sub> earthworms, but only after 10-weeks of exposure (Figure  
243 2B); at earlier intervals no difference was apparent in Pb partitioning between these  
244 two populations. Naïve earthworms accumulated Pb linearly in all three fractions over  
245 the duration of the exposure period. In contrast, after 28 days M<sub>DH</sub> earthworms  
246 preferentially partitioned the majority of accumulated metal into the insoluble  
247 (detoxified) fraction. As the concentrations of Cu, Ni, and Sr did not change  
248 appreciably over the 10 week exposure period (data not shown), this implies that the

249 temporal changes in the concentrations of Pb and Zn in the subcellular fractions were  
250 not directly linked to the loss of whole-worm weight over this period.

251

252 *Zinc:* The temporal partitioning profiles of Zn resemble those of Pb, with indigenous  
253 worms with multi-generational histories of metal exposure ( $M_{DH}$  and  $M_{CS}$ ) and naive  
254 worms with no previous field history of exposure ( $C_{PF}$ ) each sequestering Zn  
255 primarily in the insoluble (D+E) fraction, and restricting the cytosolic soluble Zn  
256 fraction within relatively narrow limits (Figure 3). Zn uptake in both the soluble and  
257 insoluble fractions by naive  $C_{PF}$  earthworms exposed to  $M_{DH}$  soil occurred in a linear  
258 fashion during the entire exposure period, whereas after 28 days of exposure  $M_{DH}$   
259 earthworms appeared to preferentially partition Zn into the insoluble (detoxified)  
260 fraction. The similarities between Pb and Zn partitioning extended to the comparative  
261 efficiency of restricting the metals to the detoxified compartment in indigenous versus  
262 introduced populations (Figure 4): the proportion of Zn present in the soluble fraction  
263 was appreciably lower in earthworms from the heavily polluted  $M_{DH}$  site at all time  
264 points compared with that in  $C_{PF}$  worms introduced to the  $M_{DH}$  soil; the proportion of  
265 soluble fraction Zn in  $M_{CS}$  worms was appreciably lower than in  $C_{PF}$  worms  
266 maintained on  $M_{CS}$  soil at three time points (3, 14, and 70 days).

267

268 *Phosphorus:* In both indigenous mine-site populations maintained on their 'own'  
269 soils, and in naive worms introduced to the metalliferous soils, a fairly steady  
270 redistribution of P from the soluble cytosolic phase to the insoluble compartment  
271 occurred during the ten-week exposure period (cf. Figures 5A and 5B). A  
272 considerably higher insoluble P concentration was measured in indigenous and

273 introduced earthworms exposed to M<sub>DH</sub> soil compared with the two treatment groups  
274 exposed to the significantly less polluted M<sub>CS</sub> soil (Figure 5).

275

#### 276 Cytochrome oxidase II (COII) genotyping

277 The phylogenetic structure of the study populations was assessed using the  
278 mitochondrial cytochrome oxidase II (mtDNA COII) gene sequence data of  
279 individuals sampled from the three sites. Good quality COII nucleotide sequences  
280 (304bp) were aligned from 85 *L. rubellus* earthworms and from individuals of *L.*  
281 *castaneus* and *L. eiseni* to rule out the possibility of misidentification. Only functional  
282 COII sequences, with no stop or nonsense codons in the reading frame, were used.  
283 Intra- and inter-site evolutionary relationships were phylogenetically analysed (Figure  
284 6), with the tree constructed using the distance-based neighbour joining (NJ)  
285 algorithm based upon p-distance. Only one representative of each site and haplotype  
286 are shown and the resulting tree topology was well supported by bootstrap analyses.  
287 The sampled *L. rubellus* individuals could be resolved into two distinct genetic  
288 lineages (lineage A and B, respectively), with a mean inter-lineage mtDNA sequence  
289 divergence of 13%.

290

291 C<sub>PF</sub> earthworms grouped exclusively within lineage A, and comprised 7 distinct  
292 haplotypes, with a between-haplotype diversity of 1 to 4% (Figure 6). In comparison,  
293 the M<sub>DH</sub> and M<sub>CS</sub> populations derived from mine-associated soils belonged  
294 predominantly to the lineage B genotype, and comprised 3 (M<sub>DH</sub>) and 4 haplotypes  
295 (M<sub>CS</sub>) exhibiting between-haplotype diversity of 1 to 2%, respectively. Only one  
296 individual from each mine site had lineage A genotype signatures. The genetic  
297 distance between *L. rubellus* and two other *Lumbricus* species (*L. castaneus* and *L.*

298 *eiseni*) was calculated as 18.1%, thus indicating that the *L. rubellus* field populations  
299 were correctly assigned.

300

### 301 Discussion

302 Abandoned metal mine soils in the UK and elsewhere harbour locally adapted  
303 earthworm populations with innate abilities to tolerate phenomenally high internal  
304 body loads of certain metals. For example, earthworms evidently thrive in field soils  
305 contaminated to degrees exceeding by an order of magnitude the exposure level that  
306 severely compromises reproduction in spiked laboratory soils (Spurgeon, *et al.*, 1994).  
307 That these are residents and not immigrants from less-contaminated surrounding soil  
308 is one way of interpreting the 'patchy' pattern of genotype distributions observed in *L.*  
309 *rubellus* across geochemically heterogeneous metalliferous landscapes (Andre, *et al.*,  
310 2010). A number of published studies provide mechanistic insights concerning the  
311 modes of metal detoxification within discrete subcellular compartments in these  
312 chronically exposed natural populations (Morgan and Morris, 1982; Morris and  
313 Morgan, 1986; Morgan and Morgan, 1989a; 1998; Sturzenbaum, *et al.*, 2001).  
314 However, evidence of phenotypic differences at the behavioural, physiological and  
315 molecular levels between populations that have undergone multiple generations of  
316 exposure and their counterparts with no comparable metal exposure history in their  
317 native habitat remain sparse. Therefore, by comparing the subcellular partitioning  
318 profiles amongst earthworm populations native to contaminated and clean sites,  
319 further inferences into the metal management strategies of putatively adapted  
320 ecotypes may be gained.

321

322 A Cd-resistant ecotype of the freshwater oligochaete *Lumbriculus hoffmeisteri* has  
323 been shown to possess enhanced Cd accumulation efficiency (Klerks and  
324 Bartholomew, 1991) and a concomitant reduction in the amount of trophically-  
325 available Cd (Wallace and Lopez, 1997). Such integrated duality is also expressed for  
326 Pb in at least one of the earthworm populations,  $M_{DH}$ , examined in the present study.  
327 Specifically, Pb partitioning profiles for  $M_{DH}$  individuals showed a much lower  
328 absolute level of soluble Pb (approximately 57% after 70 days exposure), when  
329 compared to naive  $C_{PF}$  earthworms exposed to the same  $M_{DH}$  polluted soil. This  
330 population appears to have evolved a capability to limit Pb toxico-availability  
331 possibly through modifications of components of  $Ca^{2+}$  transporting pathway, such as  
332 the sarcoplasmic/endoplasmic reticulum calcium ATPase (SERCA) (Andre, *et al.*,  
333 2010). The notion is supported by plotting the soluble Pb fraction of  $M_{DH}$  worms as a  
334 percentage of the total body load; although the total body burden increases over the 10  
335 week exposure period on their native soil, there is a significantly lower proportion of  
336 the Pb burden distributed in the soluble fraction compared with that found in  $C_{PF}$   
337 reference site worms maintained in the laboratory for the same period on  $M_{DH}$   
338 metalliferous soil. A similar differential was also recorded in the proportional Pb  
339 content of the soluble fractions of  $M_{CS}$  and  $C_{PF}$  worms at all time intervals of  
340 laboratory exposure to  $M_{CS}$  soil. This phenomenon of increasing metal concentrations  
341 in earthworms from metal-contaminated soils maintained on their 'own' soils in the  
342 laboratory has been reported by others (e.g. Corp and Morgan, 1991). It is not easily  
343 explained in the cases of metal that are not especially redox active other than as a  
344 consequence of the destruction of the physical structure of soil, with a release of  
345 previously stabilized metal into the 'bioavailable' pool, i.e. a partial reversal of the  
346 'ageing' process. The indication that Pb bioreactivity is reduced in earthworms

347 indigenous to Pb-contaminated soils was reinforced by the finding that the weight loss  
348 experienced by reference C<sub>PF</sub> earthworms maintained on the two studied metalliferous  
349 soils exceeded that experienced by earthworms native to the field soils. These data  
350 highlight the crucial role that intracellular components and machinery play in  
351 facilitating the efficient delivery of metals to intracellular compartments where they  
352 are sequestered in insoluble states.

353

354 Several publications have shown that lysosome-like chloragosomes within the  
355 chlorogogenous tissue (possessing some functional similarities to vertebrate  
356 hepatocytes) are the main metal-sequestering organelles. Chloragosomes represent  
357 phosphate-rich storage compartments for group A, O-seeking, metals (Morgan and  
358 Morris, 1982; Morgan and Morgan, 1989a; b; 1998). Andre, *et al.*, (2009) investigated  
359 the ligand speciation of Pb within whole earthworms using synchrotron- based XAS  
360 analysis, and obtained XANES spectra that unambiguously revealed that *L. rubellus*  
361 with a protracted population history of Pb exposure preferentially sequester the metal  
362 as insoluble pyromorphite [Pb<sub>5</sub>(PO<sub>4</sub>)<sub>3</sub>Cl] and Pb<sub>3</sub>(PO<sub>4</sub>)<sub>2</sub>. Given this fact, the  
363 observation in the present study that intracellular P speciation shifts appreciably in all  
364 exposures over the entire 10 week period from a relatively soluble to less soluble state  
365 presumably to associate with intruding Pb is functionally logical.

366

367 Due to its biological essentiality it is predictable that invertebrates are able to regulate  
368 intracellular Zn levels to a considerable degree. Chromatographic observations  
369 demonstrate that this may be achieved through Zn binding to a variety of low- and  
370 high-molecular weight molecules (Susuki, *et al.*, 1988; Cain and Luoma, 1998; Lock

371 and Janssen, 2001). Homeostatic systems operate to not only sequester and detoxify  
372 excess Zn but, when needed, to release Zn in order to meet the cells physiological  
373 requirements. This system is undoubtedly at work in earthworms from M<sub>DH</sub> as,  
374 despite considerable increases in total body load, they demonstrate the ability to  
375 maintain their intracellular soluble Zn content within relatively narrow limits. Again,  
376 the phosphate-rich chloragosomes are implicated in Zn storage and detoxification  
377 alongside a less well characterised sulphur-rich organelle, the cadmosome  
378 (Sturzenbaum, *et al.*, 1998). The involvement of chloragosomes and cadmosomes in  
379 excess Zn sequestration has been corroborated by XAS analyses, with XANES  
380 spectra indicating that Zn binds to both O- and S-donating ligands (Andre, *et al.*,  
381 2009).

382

383 Cryptic or sibling species are typically found in taxa that thrive in complex,  
384 heterogeneous, environments and have been discovered by genotyping fauna  
385 inhabiting diverse marine, freshwater, and terrestrial habitats (Sturmbauer, *et al.*,  
386 1999; Pinceel, *et al.*, 2004; Mathews, 2006; Pfenninger and Schwenk, 2007). The *L.*  
387 *rubellus* population indigenous to the unpolluted field site, C<sub>PF</sub>, belongs exclusively to  
388 lineage A and can be resolved into 7 haplotypes, whilst the two geographically  
389 isolated mine-site populations both belong to the genotypically distinct lineage B  
390 comprised of 3 and 4, respectively, distinct haplotypes. The number of *L. rubellus*  
391 populations examined was too restricted to draw firm conclusions regarding  
392 microevolutionary genealogies, but the higher intra-lineage diversity of the C<sub>PF</sub>  
393 sample is indicative of a relatively stationary population that has undergone multiple  
394 introductions and bottleneck episodes during its evolutionary history (Harpending,  
395 1994). It is tempting to interpret the comparatively narrow genetic diversity within the

396 lineage B inhabitants of the mine sites as a hallmark of stress-driven genetic erosion  
397 processes (natural selection, genetic drift, inbreeding) having acted upon these  
398 populations. Genetic erosion can certainly accompany small fragmented populations  
399 (Buza *et al.*, 2000) such as those found inhabiting the ‘islands of toxicity’ that typify  
400 abandoned metal mine sites. However, the genetic erosion notion as an explanation of  
401 the genetic structure of mine-associated earthworm populations should be tempered  
402 with the knowledge that calculations from genetic parameters lead to the conclusion  
403 that lineage A (with an inter-stadial expansion time of ~250 000 years BP) is  
404 appreciably ‘older’ than lineage B (expansion time of ~17000 years BP) (Andre, *et*  
405 *al.*, 2010) and may have had the opportunity to evolve more genetic richness. Peles, *et*  
406 *al.* (2003) suggested that certain alleles and genotypes in *L. rubellus* may be more  
407 sensitive to the effects of heavy metals because the frequency of both differed  
408 significantly at polymorphic loci between populations inhabiting sewage  
409 contaminated and reference soils. Conversely, Haimi, *et al.* (2007) reported that metal  
410 contamination did not significantly impact upon clonal diversity in the earthworm  
411 *Dendrobaena octaedra*. Analogous inter- or intra-lineage conclusions cannot firmly  
412 be drawn from the present study on *L. rubellus*. Whether or not the two deeply  
413 divergent *L. rubellus* lineages warrant the status of (cryptic) species must await  
414 further genetic and breeding evidence. Nevertheless, it is noteworthy that Lentzsch  
415 and Gollmack (2006) observed that species richness not ambient soil conditions was  
416 the overriding factor affecting intraspecific diversity and genotype abundance in the  
417 earthworm *Aporrectodea caliginosa*, thus ostensibly supporting the hypothesis that  
418 ecological niches are colonised at a species level prior to local population-level  
419 adaption. The shallow soils often associated with abandoned metal mines usually

420 harbour impoverished earthworm communities, in many instances no more than two  
421 taxonomically accepted representatives of the epigeic ecophysiological group.

422

423 In conclusion, the ability of an adapted population to tolerate the prevailing stress-  
424 evoking conditions in severely polluted habitats most probably involve heritable and  
425 integrated combinations of physiological, morphological and behavioural  
426 modifications. Thus, it is plausible to hypothesise that a metal tolerant earthworm  
427 population has evolved efficient mechanisms of detoxification that feature an  
428 enhanced immobilisation capacity coupled to a relative reduction in the metal  
429 sensitive fraction as an important component of their holistic adaptive arsenal.  
430 However, inferences about population-specific adaptation based on subcellular metal  
431 partitioning profiles should be drawn with a measure of caution because of the  
432 possibility that they could be attributable to lineage-specific traits that are independent  
433 of chronic metal exposure. This is illustrated by the findings of (Heethoff, *et al.*,  
434 2004) that the parthenogenic earthworm *Octolasion tyrtaeum* is differentiated into two  
435 lineages differing significantly in body size. Such findings, together with those arising  
436 from the present study, raise the spectre that field-based eco-toxicological assessments  
437 that utilise earthworms, particularly those emanating from discriminating ‘omics’  
438 measurements, might benefit from the elimination of a potential confounding biotic  
439 variable through prior genotyping of all individuals to establish that they possess  
440 some genetic background equivalence.

441

442 Acknowledgments. This work was performed during the tenure of a Natural  
443 Environment Research Council studentship (JA) registered at Reading University, and  
444 a research grant (F0 0407/AI) awarded by The Leverhulme Trust to AJM. We thank

445 Dr Trevor Pearce of Lancaster University for providing us with specimens of  
446 *Lumbricus eiseni* for genotyping.

447

448 References

449 Andre, J., Charnock, J., Stürzenbaum, S.R., Kille, P., Morgan, A.J., Hodson, M.E.,  
450 2009. Accumulated Metal Speciation in Earthworm Populations with  
451 Multigenerational Exposure to Metalliferous Soils: Cell Fractionation and High-  
452 Energy Synchrotron Analyses. *Environmental Science and Technology* 43, 6822-  
453 6829.

454 Andre, J., King, R.A., Stürzenbaum, S.R., Kille, P., Hodson, M.E., Morgan, A.J.,  
455 2010. Molecular genetic differentiation in earthworms inhabiting a heterogeneous Pb-  
456 polluted landscape. *Environmental Pollution* 158, 883-890.

457 Arnold, R.E., Hodson, M.E., 2007. Effect of time and mode of depuration on tissue  
458 copper concentrations of the earthworms *Eisenia andrei*, *Lumbricus rubellus* and  
459 *Lumbricus terrestris*. *Environmental Pollution* 148, 21-30.

460 Arnold, R.E., Hodson, M.E., Charnock, J.M., Peijnenburg, W.J.G.M., 2008.  
461 Comparison of the subcellular partitioning, distribution and internal speciation of Cu  
462 between Cu-tolerant and naive populations of *Dendrodrilus rubidus* Savigny.  
463 *Environmental Science and Technology* 42, 3900-3905.

464 Boisson, J., Mench, M., Sappin-Didier, V., Sold, P., Vangronsveld, J., 1998. Short-  
465 term in situ immobilization of Cd and Ni by beringite and steel shots application to  
466 long-term sludged plots. *Agronomie* 18, 347-359.

467 Buza, L., Young, A., Thrall, P. (2000). Genetic erosion, inbreeding and reduced  
468 fitness in fragmented populations of the endangered tetraploid pea *Swainsona recta*.  
469 *Biological Conservation* 93, 177-186.

470 Cain, D.J., Luoma, S.N., 1998. Metal exposures to native populations of the caddisfly  
471 *Hydropsyche* (Trichoptera: Hydropsychidae) determined from cytosolic and whole  
472 body metal concentrations. *Hydrobiologia* 386, 103-117.

473 Cain, D.J., Luoma, S.N., Wallace, W.G., 2004. Linking metal bioaccumulation of  
474 aquatic insects to their distribution patterns in a mining- impacted river.  
475 *Environmental Toxicology and Chemistry* 23, 1463-1473.

476 Campbell, P.G.C., Chapman, P.M., Hale, B.A., 2006. Risk assessment of metals in the  
477 environment. *Issues in Environmental Science and Technology* 22, 101-131.

478 Conder, J.M., Seals, L.D., Lanno, R.P., 2002. Method for determining toxicologically  
479 relevant cadmium residues in the earthworm *Eisenia fetida*. *Chemosphere* 49, 1-7.

480 Corp, N., Morgan, A.J., 1991. Accumulation of heavy metals from polluted soils by  
481 the earthworm *Lumbricus rubellus*: Can laboratory exposure of 'control' worms reduce  
482 biomonitoring problems? *Environmental Pollution* 74, 39-52

483 Cotter-Howells, J., Charnock, J.M., Winters, C., Kille, P., Fry, J.C., Morgan, A.J.,  
484 2005. Metal Compartmentation and Speciation in a Soil Sentinel: The Earthworm,  
485 *Dendrodrilus rubidus*. *Environmental Science and Technology* 39, 7731-7740.

486 Haimi, J., Knott, K.E., Yliniemi, I., 2007. Does metal contamination affect clonal  
487 diversity of the parthenogenetic earthworm *Dendrobaena octaedra*? *European Journal*  
488 *of Soil Biology* 43, 268-272.

489 Harpending, H.C., 1994. Signature of Ancient Population-Growth in a Low-  
490 Resolution Mitochondrial-DNA Mismatch Distribution. *Human Biology* 66, 591-600.

491 Heethoff, M., Etzold, K., Scheu, S., 2004. Mitochondrial COII sequences indicate that  
492 the parthenogenetic earthworm *Octolasion tyrtaeum* (Savigny 1826) constitutes of  
493 two lineages differing in body size and genotype. *Pedobiologia* 48, 9-13.

494 Honeycutt, M.E., Roberts, B.L., Roane, D.S., 1995. Cadmium deposition in the  
495 earthworm *Eisenia fetida*. *Ecotoxicology and Environmental Safety* 30, 143-150.

496 Hopkin, S.P., 1989. *Ecophysiology of metals in terrestrial invertebrates*. Elsevier  
497 Applied Science, Barking, UK.

498 Huang, R., Wen, B., Pei, Z., Shan, X.Q., Zhang, S., Williams, P.N., 2009.  
499 Accumulation, Subcellular Distribution and Toxicity of Copper in Earthworm  
500 (*Eisenia fetida*) in the Presence of Ciprofloxacin. *Environmental Science and*  
501 *Technology* 43, 3688-3693.

502 Jones, R.P., Bednar, A.J., Inouye, L.S., 2009. Subcellular compartmentalization of  
503 lead in the earthworm, *Eisenia fetida*: Relationship to survival and reproduction.  
504 *Ecotoxicology and Environmental Safety* 72, 1045-1052.

505 King, R.A., Tibble, A., Symondson, W.O.C., 2008. Opening a can of worms:  
506 unprecedented sympatric cryptic diversity within British lumbricid earthworms.  
507 *Molecular Ecology* 17, 4684-4698.

508 Klerks, P.L., Bartholomew, P.R., 1991. Cadmium accumulation and detoxification in  
509 a Cd-resistant population of the oligochaete *Limnodrilus hoffmeisteri*. *Aquatic*  
510 *Toxicology* 19, 97-112

511 Knapen, D., Bervoets, L., Verheyen, E., Blust, R., 2004. Resistance to water pollution  
512 in natural gudgeon (*Gobio gobio*) populations may be due to genetic adaptation.  
513 *Aquatic Toxicology* 67, 155-165.

514 Kumar, S., Tamura, K., Nei, M., 2004. MEGA3: Integrated software for Molecular  
515 Evolutionary Genetics Analysis and sequence alignment. *Briefings in Bioinformatics*  
516 5, 150-163.

517 Langdon, C.J., Hodson, M.E., Arnold, R.E., Black, S., 2005. Survival, Pb-uptake and  
518 behaviour of three species of earthworm in Pb treated soils determined using an  
519 OECD-style toxicity test and a soil avoidance test. *Environmental Pollution* 138.

520 Langdon, C.J., Morgan, A.J., Charnock, J.M., Semple, K.T., Lowe, C.N., 2009. As-  
521 resistance in laboratory-reared F1, F2 and F3 generation offspring of the earthworm  
522 *Lumbricus rubellus* inhabiting an As-contaminated mine soil. *Environmental*  
523 *Pollution* 157, 3114-3119.

524 Lentzsch, P., Gollack, J., 2006. Genetic diversity of *Aporrectodea caliginosa* from  
525 agricultural sites in Northeast Brandenburg, Germany. *Pedobiologia* 50, 369-376.

526 Li, L.Z., Zhou, D.M., Wang, P., Luo, X.S., 2008. Subcellular distribution of Cd and  
527 Pb in earthworm *Eisenia fetida* as affected by Ca<sup>2+</sup> ions and Cd-Pb interaction.  
528 *Ecotoxicology and Environmental Safety* 71, 632-637.

529 Lock, K., Janssen, C.R., 2001. Zinc and cadmium body burdens in terrestrial  
530 oligochaetes: Use and significance in environmental risk assessment. *Environmental*  
531 *Toxicology and Chemistry* 20, 2067-2072.

532 Mathews, L.M., 2006. Cryptic biodiversity and phylogeographical patterns in a  
533 snapping shrimp species complex. *Molecular Ecology* 15, 4049-4063.

534 McCarthy, L.S., Mackay, D., 1993. Enhancing ecotoxicological modeling and  
535 assessment. *Environmental Science and Technology* 27, 1719-1728.

536 Morgan, A.J., Kille, P., Sturzenbaum, S.R., 2007. Microevolution and ecotoxicology  
537 of metals in invertebrates. *Environmental Science and Technology* 41, 1085-1096.

538 Morgan, A.J., Morris, B., 1982. The accumulation and intracellular compartmentation  
539 of cadmium, lead, zinc and calcium in two earthworm species (*Dendrobaena rubida*  
540 and *Lumbricus rubellus*) living in highly contaminated soil. *Histochemistry and Cell*  
541 *Biology* 75, 269-285.

542 Morgan, J.E., Morgan, A.J., 1989a. The effect of lead incorporation on the elemental  
543 composition of earthworm (Annelida, Oligochaeta) chloragosome granules.  
544 *Histochemistry* 92, 237-241.

545 Morgan, J.E., Morgan, A.J., 1989b. Zinc sequestration by earthworm (Annelida:  
546 Oligochaeta) chloragocytes. *Histochemistry and Cell Biology* 90, 405-411.

547 Morgan, J.E., Morgan, A.J., 1990. The distribution of cadmium, copper, lead, zinc  
548 and calcium in the tissues of the earthworm *Lumbricus rubellus* sampled from one  
549 uncontaminated and four polluted soils. *Oecologia* 84, 559-566.

550 Morgan, J.E., Morgan, A.J., 1998. The distribution and intracellular  
551 compartmentalisation of metals in the endogeic earthworm *Aporrectodea caliginosa*  
552 sampled from an unpolluted and a metal contaminated site. *Environmental Pollution*  
553 99, 167-175.

554 Morris, B., Morgan, A.J., 1986. Calcium-lead interactions in earthworms:  
555 Observations on *Lumbricus terrestris* sampled from a calcareous abandoned lead  
556 mine site. Bulletin of Environmental Contamination and Toxicology 37, 226-233.

557 Novo, M., Velavan, T.P., Almodovar, A., Schulenburg, H., Diaz-Cosin, D.J.,  
558 Michiels, N.K., 2008. Microsatellite markers for the drought-resistant earthworm  
559 *Hormogaster elisae*. Molecular Ecology Resources 8, 901-903.

560 Pan, K., Wang, W.X., 2008. The subcellular fate of cadmium and zinc in the scallop  
561 *Chlamys nobilis* during waterborne and dietary metal exposure. Aquatic Toxicology  
562 90, 253-260.

563 Peles, J.D., Towler, W.I., Guttman, S.I., 2003. Population genetic structure of  
564 earthworms (*Lumbricus rubellus*) in soils contaminated by heavy metals.  
565 Ecotoxicology 12, 379-386.

566 Perceval, O., Couillard, Y., Pinel-Alloul, B., Campbell, P.G.C., 2006. Linking  
567 changes in subcellular cadmium distribution to growth and mortality rates in  
568 transplanted freshwater bivalves (*Pyganodon grandis*). Aquatic Toxicology 79, 87-98.

569 Pérez-Losada, M., Ricoy, M., Marshall, J.C., Domínguez, J., 2009. Phylogenetic  
570 assessment of the earthworm *Aporrectodea caliginosa* species complex (Oligochaeta:  
571 Lumbricidae) based on mitochondrial and nuclear DNA sequences. Molecular  
572 Phylogenetics and Evolution 52, 293-302.

573 Péry, A.R.R., Béthune, A., Gahou, J., Mons, R., Garric, J., 2005. Body residues: a key  
574 variable to analyze toxicity tests with *Chironomus riparius* exposed to copper-spiked  
575 sediments. Ecotoxicology and Environmental Safety 61, 160-167.

576 Péry, A.R.R., Geffard, A., Conrad, A., Mons, R., Garric, J., 2008. Assessing the risk  
577 of metal mixtures in contaminated sediments on *Chironomus riparius* based on  
578 cytosolic accumulation. *Ecotoxicology and Environmental Safety* 71, 869-873.

579 Pfenninger, M., Schwenk, K., 2007. Cryptic animal species are homogeneously  
580 distributed among taxa and biogeographical regions. *BMC Evolutionary Biology* 7,  
581 121.

582 Pinceel, J., Jordaens, K., Van Houtte, N., De Winter, A.J., Backeljau, T., 2004.  
583 Molecular and morphological data reveal cryptic taxonomic diversity in the terrestrial  
584 slug complex *Arion subfuscus/fuscus* (Mollusca, Pulmonata, Arionidae) in continental  
585 north-west Europe. *Biological Journal of the Linnean Society* 83, 23-38.

586 Rainbow, P.S., 2002. Trace metal concentrations in aquatic invertebrates: why and so  
587 what? *Environmental Pollution* 120, 497-507.

588 Rainbow, P.S., 2007. Trace metal bioaccumulation: Models, metabolic availability  
589 and toxicity. *Environment International* 33, 576-582.

590 Saitou, N., Nei, M., 1987. The Neighbor-Joining Method - a New Method for  
591 Reconstructing Phylogenetic Trees. *Molecular Biology and Evolution* 4, 406-425.

592 Shepeleva, O.A., Kodolova, O.P., Zhukovskaya, E.A., Striganova, B.R., 2008.  
593 Genetic diversity of populations of the earthworm *Lumbricus rubellus*  
594 (Hoffm.)(Oligochaeta, Lumbricidae). *Biology Bulletin* 35, 170-177.

595 Spurgeon, D.J., Hopkin, S.P., Jones, D.T., 1994. Effects of cadmium, copper, lead and  
596 zinc on growth, reproduction and survival of the earthworm *Eisenia fetida* (Savigny):

597 Assessing the environmental impact of point-source metal contamination in terrestrial  
598 ecosystems. *Environmental Pollution* 84, 123-130.

599 Sturmbauer, C., Opadiya, G.B., Niederstatter, H., Riedmann, A., Dallinger, R., 1999.  
600 Mitochondrial DNA reveals cryptic oligochaete species differing in cadmium  
601 resistance. *Molecular Biology and Evolution* 16, 967-974.

602 Sturzenbaum, S.R., Kille, P., Morgan, A.J., 1998. The identification, cloning and  
603 characterisation of earthworm metallothionein. *FEBS Letters* 431, 437-442.

604 Sturzenbaum, S.R., Winters, C., Galay, M., Morgan, A.J., Kille, P., 2001. Metal ion  
605 trafficking in earthworms. Identification of a cadmium-specific metallothionein. *The*  
606 *Journal of Biological Chemistry* 276, 34013-34018.

607 Susuki, K.T., Sunaga, H., Aoki, Y., Hatakeyama, S., Sugaya, Y., Sumi, Y., Suzuki,  
608 T., 1988. Binding of cadmium and copper in the mayfly *Baetis Thermicus* that inhabit  
609 a river polluted with heavy metals. *Comparative Biochemistry and Physiology C*  
610 *Comparative Pharmacology* 91C, 487-492.

611 Velavan, T.P., Schulenburg, H., Michiels, N., 2007. Development and  
612 characterization of novel microsatellite markers for the common earthworm  
613 (*Lumbricus terrestris L.*). *Host-Parasite Interactions* 7, 24.

614 Vijver, M.G., Van Gestel, C.A.M., Lanno, R.P., Van Straalen, N.M., Peijnenburg,  
615 W.J.G.M., 2004. Internal metal sequestration and its ecotoxicological relevance: a  
616 review. *Environmental Science and Technology*. 38, 4705-4712.

617 Vijver, M.G., van Gestel, C.A.M., van Straalen, N.M., Lanno, R.P., Peijnenburg,  
618 W.J.G.M., 2006. Biological significance of metals partitioned to subcellular fractions

619 within earthworms (*Aporrectodea Caliginosa*). Environmental Toxicology and  
620 Chemistry 25, 807-814.

621 Voets, J., Redeker, E.S., Blust, R., Bervoets, L., 2009. Differences in metal  
622 sequestration between zebra mussels from clean and polluted field locations. Aquatic  
623 Toxicology 93, 53-60.

624 Wallace, W.G., Lee, B.G., Luoma, S.N., 2003. Subcellular compartmentalisation of Cd  
625 and Zn in two bivalves. 1. Significance of metal-sensitive fractions (MSF) and  
626 biologically detoxified metal (BDM). Marine Ecology Progress Series 249, 183-197.

627 Wallace, W.G., Lopez, G.R., 1997. Bioavailability of biologically sequestered  
628 cadmium and the implications of metal detoxification. Marine Ecology Progress  
629 Series 1997, 149-157.

630

### 631 Figure Legends

#### 632 Figure 1

633 Time course of Pb partitioning into soluble (A) and non-soluble (B) cellular fractions  
634 in *Lumbricus rubellus* sampled from three populations and maintained on two mine-  
635 associated metal contaminated soils. Pb concentrations (per unit wet weight of  
636 earthworm) are presented as mean  $\pm$ S.E. (maximum 'n' =9). C<sub>PF</sub> refers to the  
637 reference site at Pontcanna Fields, M<sub>CS</sub> the metalliferous acidic site Cwmystwyth  
638 Stream and M<sub>DH</sub> the metalliferous calcareous site, Draethen Hollow.

639

#### 640 Figure 2

641 Time course of subcellular Pb distribution in *Lumbricus rubellus* sampled from three  
642 populations and maintained on two mine-associated metal contaminated soils, with

643 the soluble metal concentration expressed as a percentage of the total body  
644 concentration. The Pb levels shown represent fractions extracted from M<sub>CS</sub> and C<sub>PF</sub>  
645 earthworms in M<sub>CS</sub> soil (A) and M<sub>DH</sub> and C<sub>PF</sub> earthworms in M<sub>DH</sub> soil (B). [See Fig.  
646 1 for dataset error bars and Fig. 1 legend for site identifiers.]

647

648 Figure 3

649 Time course of Zn partitioning into soluble (A) and non-soluble (B) cellular fractions  
650 in *Lumbricus rubellus* sampled from three populations and maintained on two mine-  
651 associated metal contaminated soils. Zn concentrations (per unit wet weight of  
652 earthworm) are presented as mean  $\pm$ S.E. (maximum 'n' =9). [See Fig. 1 legend for  
653 site identifiers.]

654

655 Figure 4

656 Time course of subcellular Zn distribution in *Lumbricus rubellus* sampled from three  
657 populations and maintained on two field mine-associated metal contaminated soils,  
658 with the soluble metal concentration expressed as a percentage of the total body  
659 concentration. The Zn levels shown represent fractions extracted from M<sub>CS</sub> and C<sub>PF</sub>  
660 earthworms in M<sub>CS</sub> soil (A) and M<sub>DH</sub> and C<sub>PF</sub> earthworms in M<sub>DH</sub> soil (B). [See Fig. 1  
661 for dataset error bars and Fig. 1 legend for site identifiers.]

662

663 Figure 5

664 Time course of P partitioning into soluble (A) and non-soluble (B) cellular fractions in  
665 *Lumbricus rubellus* sampled from three populations and maintained on two field-  
666 derived metal contaminated soils. P concentrations (per unit wet weight of

667 earthworm) are presented as mean  $\pm$ S.E. (maximum 'n' = 9). [See Fig. 1 legend for  
668 site identifiers.]

669

670 Figure 6

671 Phylogenetic tree based on p-distance of the cytochrome oxidase II mitochondrial  
672 gene of 87 *Lumbricus rubellus* individuals from the contaminated sites M<sub>DH</sub>  
673 (triangles) and M<sub>CS</sub> (circles) and the reference site C<sub>PF</sub> (squares), *L. castaneus* and *L.*  
674 *eiseni* individuals (open squares) are included. Two *L. rubellus* lineages are apparent,  
675 termed lineage 'A' (light grey) and 'B' (dark grey), respectively, with a mean inter-  
676 lineage mtDNA sequence divergence of approximately 13%. Tree topology was well  
677 supported by bootstrap analyses. Only one representative of each haplotype at the  
678 particular sites and are shown; numbers in parentheses indicate the numbers of  
679 individuals of a given haplotype.