



SHARPENS YOUR THINKING

The contribution of injury severity, executive and implicit functions to awareness of deficits after traumatic brain injury (TBI)

MORTON, Nicholas and BARKER, Lynne

Available from Sheffield Hallam University Research Archive (SHURA) at:

<http://shura.shu.ac.uk/6061/>

This document is the author deposited version. You are advised to consult the publisher's version if you wish to cite from it.

Published version

MORTON, Nicholas and BARKER, Lynne (2010). The contribution of injury severity, executive and implicit functions to awareness of deficits after traumatic brain injury (TBI). *Journal of the International Neuropsychological Society*, 16 (06), 1089-1098.

Repository use policy

Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in SHURA to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

Title: The contribution of injury severity, executive and implicit functions to awareness of deficits after Traumatic Brain Injury (TBI).

¹Nicholas Morton, ²Lynne Barker

¹Neurorehabilitation Services Tickhill Road Hospital,
Doncaster, South Yorkshire.

²Dept; of Psychology,
Sheffield Hallam University,
Sheffield, South Yorkshire.

Correspondence concerning this article should be addressed to:
Dr Nicholas Morton
Rotherham, Doncaster and South Humber Mental Health NHS Trust,
Tickhill Road Hospital,
Doncaster, South Yorkshire, DN4 8QL
Phone: +44 (0) 1709 304272
Fax: +44 (0) 1709 307517
E-mail: n.morton@nhs.net

Manuscript length: 4,975 words

ABSTRACT

Deficits in self-awareness are commonly seen after Traumatic Brain Injury (TBI) and adversely affect rehabilitative efforts, independence and quality of life (Ponsford, 2004). Awareness models predict that executive and implicit functions are important cognitive components of awareness though the putative relationship between implicit and awareness processes has not been subject to empirical investigation. (Toglia & Kirk, 2000; Ownsworth, Clare & Morris, 2006; Crosson et al., 1989). Severity of injury, also thought to be a crucial determinant of awareness outcome post-insult, is under-explored in awareness studies (Sherer et al., 1989).

The present study measured the contribution of injury severity, IQ, mood state, executive and implicit functions to awareness in head-injured patients assigned to moderate/severe head-injured groups using several awareness, executive and implicit measures. Severe injuries resulted in greater impairments across most awareness, executive and implicit measures compared to moderate injuries, although deficits were still seen in the moderate group. Hierarchical regression results showed that severity of injury, IQ, mood state, executive and implicit functions made significant unique contributions to selective aspects of awareness. Future models of awareness should account for both implicit and executive contributions to awareness and the possibility that both are vulnerable to disruption after neuropathology.

Keywords: Tacit, conscious, control processes, impairment, neuropathology

INTRODUCTION

Awareness is a complex construct comprising cognitive, psychosocial, and emotional components (Toglia & Kirk, 2000). Neural substrates of awareness are thought to involve diverse brain regions including prefrontal areas, inferior parietal lobe, angular gyrus, supramarginal gyrus, and anterior temporal lobes (Prigatano & Schacter, 1991). Consequently, diffuse bilateral brain pathology seen after severe head injury or advanced dementia is more likely to produce awareness deficits than focal unilateral lesions (Sherer, Hart, Whyte, Todd & Yablon, 2005; Prigatano, 2010). Despite the proposed heterogeneity of neural substrates associated with awareness, frontal pathology is consistently associated with awareness deficits possibly reflecting the functionally integrative role of these brain regions (Rosen et al., 2010, Banks & Weintraub, 2009). Duration of time since injury is also associated with extent of awareness deficits with impairments typically manifest in the post-acute stage several weeks to six months post-injury (Ownsworth, Desbois, Grant, Fleming & Strong 2006; Fleming & Strong, 1999; Hart, Seignourel & Sherer, 1999). Intelligence (Bogod, Mateer & MacDonald, 2003) and emotional state, specifically anxiety and depression, have also been shown to correlate with post-injury awareness (Fleming, Strong & Ashton, 1998). Epidemiological data show that awareness deficits affect approximately 45% of TBI patients (Flashman & McAllister, 2002) and adversely affect rehabilitation compliance and outcome, frequency and severity of socio-behavioral problems, caregiver distress and patient quality of life (King, 1997; Sherer et al., 1998; Wise, Ownsworth & Fleming, 2005).

Awareness models typically incorporate several stages or ‘types’ of awareness (anticipatory, emergent or ‘on-line’ awareness) driven by subsidiary cognitive processes (semantic memory, perceptual, emotional and implicit processes) and integrated by a metacognitive component (metacognitive or intellectual awareness) (Toglia & Kirk, 2000; Morris & Hannesdottir, 2004; Crosson et al., 1989). Executive functions, often termed metacognitive, are higher-order supervisory processes that initiate, maintain or inhibit other cognitive processes to facilitate goal-directed behavior (Miyake et al., 2000; Stuss & Alexander, 2007). Concept formation, self-monitoring and self-appraisal executive functions are considered key processes mediating awareness (Toglia & Kirk, 2000; Crosson et al., 1989). Diverse executive functions have been associated with degree of post-TBI awareness including, planning and mental flexibility, idea generation or fluency, self-regulation, sustained attention, and reasoning ability (Wilson, Alderman, Burgess, Emslie & Evans, 1996; Ownsworth, McFarland & Young, 2000; Ownsworth & Fleming, 2005; Bogod, Mateer & MacDonald 2003; O’Keeffe, Dockree, Moloney, Carton & Robertson, 2007). Some researchers have been unable to replicate earlier findings (Ownsworth & Fleming, 2005), findings reported elsewhere (Bogod, Mateer & MacDonald, 2003), or found no relationship between executive function and awareness (O’Keeffe et al., 2007). Equivocal findings might be explained by use of limited and varied awareness measures across studies, difficulty isolating executive components of executive tasks (Hart, Whyte, Kim & Vaccaro, 2005; Ownsworth & Fleming, 2005), absence of a consistent conceptual framework across studies, and under exploration of neurological variables (injury severity) known to significantly affect executive functions (Mattson & Levin, 1990) and awareness (Sherer et al., 2005).

Awareness models also include an implicit mechanism thought to guide behavioural responses in the absence of conscious awareness of current state (Morris & Hannesdottir, 2004; Schacter, 1990), or conversely contribute to metacognitive awareness by augmenting conscious knowledge (Toglia & Kirk, 2000). Although under-specified the implicit mechanism is an important component of awareness models accounting for tacit awareness (evidenced by task avoidance and behavioral adaptivity) in the absence of conscious awareness of deficits; a pattern of functional outcome well documented in the literature in relation to TBI and Alzheimer's disease (Ownsworth, Clare & Morris, 2006; Trahan, Pépin & Hopps, 2006; Prigatano & Schacter, 1991). It is less clear how implicit processes might augment metacognitive awareness.

Implicit processing refers to the acquisition of information expressed through altered behavior in the absence of subjective awareness of information acquired. There is extensive evidence of the role of implicit processes to social functioning in the experimental literature (see a recent review by Frith & Frith, 2009), and neuroimaging data show that implicit stimuli produce a corresponding neural signature whilst subjective awareness remains at chance levels (Kouider & Dehaene, 2007). Implicit experimental paradigms have also been applied to TBI patients (Barker, Andrade, Romanowski, Morton & Wasti, 2006; Barker, Andrade, Morton, Romanowski & Bowles, 2010; Beldarrain, Grafman, de Valesco, Pascual-Leone & Garcia-Monco, 2002). Barker et al (2006) found that patients with impaired implicit sequence learning had higher behavioral discrepancy scores (indicating impaired awareness) than those with intact implicit learning. These findings hint at a possible contribution of implicit as well as executive processes to awareness post-injury and provide

supporting evidence for an implicit mechanism proposed in awareness models. However, models reviewed here do not account for the possibility that implicit processes are diminished post-injury instead they are generally assumed to be robust to neuropathology (see Reber, 2000 for rationale). Consequently, the integrity of implicit processes post-TBI and possible contribution of both implicit and executive functions has not been subject to empirical scrutiny in awareness studies.

The present study investigated severity of injury, IQ, mood state, implicit and executive contributions to awareness post-injury in a sample of 34 TBI patients using several measures of awareness, executive function and implicit cognition. We hypothesized that executive functions would contribute to metacognitive awareness and that executive and implicit functions would contribute to emergent/anticipatory awareness on the basis of theoretical frameworks. We also expected injury severity to result in greater impaired awareness on executive function and awareness tasks than moderate injury.

METHOD

Participants

Research was conducted in accordance with the declaration of Helsinki and participants gave informed consent. To account for injury severity, thought to be an important determinant of post-injury awareness, 34 participants were assigned to moderate ($n = 11$) and severe ($n = 23$) head-injured groups (Sherer et al., 1998). Injury severity was determined by at least two of the following criteria: (i) GCS score on admission (severe < 9 , moderate ≥ 9 and < 13), (ii) length of coma (severe > 6

hours, moderate < 6 hours > 1 hour), (iii) PTA (severe > 7 days, moderate < 7 days > 1 day), (iv) evidence of focal pathology from routine CT and MRI brain scanning (Lezak, 1995; Mild traumatic brain injury committee, 1993). Participants were a minimum of one year since injury to account for post-acute functional recovery (Fleming, Strong & Ashton, 1998). Lesion site could not be identified for four cases due to absence of imaging records although clinical and medical records indicated anterior neuropathology (see Table 1 for descriptive data).

[Insert table 1 here]

IQ and executive function measures

Intelligence (Wechsler Abbreviated Scale of Intelligence, WASI Wechsler, 1999), pre-morbid IQ (Wechsler Test of Adult Reading, WTAR The Psychological Corporation, 2001), and mood state (Hospital Anxiety and Depression Scale, HADS Zigmond & Snaith, 1983) were measured to account for variables previously shown to affect post-injury awareness (Bogod, Mateer & MacDonald, 2003; Fleming, Strong & Ashton, 1998). Due to the multi-componential nature of awareness, executive and implicit cognitive constructs several measures of each were selected to best capture underlying processes (Miyake et al., 2000; Barker et al., 2006; Andrés & Van Der Linden M, 2002). Awareness measures were chosen to measure metacognitive and emergent/anticipatory aspects of awareness in line with awareness conceptual models (Toglia & Kirk, 2000; Morris & Hannesdottir, 2004; Crosson et al., 1989), and on the recommendation of other researchers (O’Keeffe et al., 2007). Tests of executive function included, the Self-ordered Pointing Test measure of response monitoring (SOPT; Petrides & Milner, 1982), the Sorting Test measure of concept formation (D-

KEFS, Delis, Kaplan & Kramer, 2001), and the Brixton measure of strategy initiation and response inhibition (Hayling and Brixton, Burgess & Shallice, 1997). A Verbal Fluency task (FAS - Benton & Hamsher, 1989) was selected as a modality specific distractor task for the mere exposure effect task to prevent explicit rehearsal of auditory primes and is not considered further here though descriptive data are provided (Table 2 results section). Serial Reaction Time and mere exposure effect implicit tasks were chosen to measure implicit cognition as they are thought to depend on mechanisms governing tacit non-verbal encoding and decoding of contextual cues (Lieberman, 2000), and are sensitive to pathology across patient-based studies (Barker et al., 2006; Barker et al., 2010; Beldarrain et al., 2002).

Metacognitive awareness measures

Awareness Questionnaire (AQ - Sherer et al., 1998; Sherer, Hart & Todd, 2003).

The AQ comprised 17 items comparing individual's pre- and post-injury abilities and consisted of Self and Other versions completed by the participant and family member/significant other respectively. Significant others were selected by patients to complete Other ratings for AQ and DEX questionnaires on the basis that the person knew them well prior to injury and had significant daily contact with them since time of injury. Items were rated on a five-point scale from 1 ("much worse") to 5 ("much better") and summed to give a total score for participant and significant other ratings (range 17-85). Self-awareness score was calculated by subtracting Other ratings from patient ratings to provide an AQ discrepancy score (range -51 to +51). Positive scores indicated that patients underestimated their deficits post-injury. The measure has good internal consistency ($r = 0.88$, Sherer et al., 1998; Sherer, Hart & Todd, 2003).

Dysexecutive Questionnaire (DEX - Wilson et al., 1996).

The DEX is a 20-item, 3 factor (cognitive, emotional and motivational) questionnaire measuring post-TBI deficits and also comprises Self and Other versions. Items are rated on a five-point scale indicating frequency of occurrence ranging from 0 (“never”) to 4 (“very often”). Items were summed to give a total score for participant and Other ratings (range 0-80). Self-ratings were subtracted from Other-ratings to produce a Self-Other (DEX-Discrepancy/Insight) score (-80 to +80). As with the AQ questionnaire positive DEX-Discrepancy scores represented underestimation of deficits by patients. There are no data on the inter-rater reliability of DEX-Other ratings or test-retest reliability for DEX-Self or Other raters. Other raters (one per patient) for both AQ- and DEX-discrepancy were primarily female (85%) and comprised parents (44%), spouse/partner (35%), friends (12%) or other family members (9%).

Self-Awareness of Deficits Interview (SADI - Fleming, Strong & Ashton, 1996).

The SADI is a semi-structured interview measuring three factors, self-awareness of deficits, self-awareness of functional implications of deficits, and ability to set realistic goals. Each section was scored on a four-point scale (range 0-3, total score = 0-9). High scores on the SADI represented low levels of self-awareness across these dimensions. The measure has good inter-rater reliability ($r = 0.82$, Fleming, Strong & Ashton, 1996) and good test-retest reliability ($r = 0.85 - 0.94$, Fleming, Strong & Ashton, 1998).

Emergent/anticipatory awareness measures

Self-Regulation Skills Interview (SRSI - Ownsworth, McFarland & Young, 2000).

The SRSI is a five-item semi-structured interview measuring emergent awareness, anticipatory awareness, strategy generation, strategy-use and strategy effectiveness. The five items were scored on a 10-point scale. Two item scores were summed to provide an Awareness Index score (range 0-20) measuring emergent/anticipatory awareness of a behavioral problem identified by the participant (for example, memory or anger problems) and scored according to standard prompts. The remaining three items were combined to generate a Strategy Index score (range 0-30) measuring participant's awareness of any behavioral strategies they used with the identified problem(s). Again, high scores represented low levels of awareness. The test has good inter-rater ($r = 0.81-0.92$) and test-retest reliability ($r = 0.69-0.91$, Ownsworth, McFarland & Young, 2000).

SADI and SRSI interviews were conducted and scored by the first author. An independent rater scored a random subset of 10 interviews. Results of Pearson's correlations showed a significant degree of inter-rater reliability for SADI total score ($r = 0.95, p < 0.001$), SRSI-Awareness index ($r = 0.96, p < 0.001$), and SRSI-Strategy index scores ($r = 0.97, p < 0.001$) indicating that interview ratings accurately reflected participants' problems.

Implicit Experimental Tasks

Serial Reaction Time task (SRT - Nissen & Bullemer, 1987).

This computer-based task has been used in previous studies with TBI patient and control groups (Barker, Andrade & Romanowski, 2004; Barker et al., 2006; Barker et al., 2010). Participants completed a practice session before beginning the task. In the learning phase participants were told to respond as quickly as possible to a target

(1cm white circle) appearing in a predetermined 10 trial sequence, A B C D B C B D B C, by pressing the corresponding key (v, b, n or m). The circle remained on the screen until the correct key press was made. In random blocks the stimulus circle appeared with the same frequency at screen locations as sequence blocks but did not follow a sequence. The response-stimulus interval was 200 milliseconds and reaction time responses to each trial were recorded. The learning phase consisted of seven blocks of 50 trials comprising an initial random block to discourage participants from explicitly assuming that circles followed a pattern at the outset of the experiment, followed by six sequence blocks. Test phase comprised one sequence block flanked by two random blocks and followed immediately after the learning phase without warning to participants. Self-determined rest breaks appeared after each block of 50 trials. A learning score was calculated by summing reaction time (RT) mean of medians for each ten-trial sequence (5 medians summed to produce a mean RT for each of the three blocks at test). The sequence block mean was subtracted from the random mean (two random block means combined) to produce a sequence learning score. After the task participants completed an explicit knowledge questionnaire with a maximum score of 16 (Seger, 1997).

Mere exposure effect task (Zajonc, 1968; 1980)

This task has been used in previous studies with neuropathological and control groups (Barker et al., 2010; Barker et al., 2006). Participants were instructed to listen to one of two lists of fifteen disyllabic Finnish words, matched for likeability, recorded on compact disc and presented audibly as in previous studies. The word list was presented twice, at a rate of one word per 1.5 seconds. After the acquisition phase, the FAS verbal fluency task was administered as a modality specific distracter to prevent

participants from explicitly rehearsing stimulus words after presentation. Participants then heard a test list containing all 30 words, targets and foils, recorded in random order with a 4 second inter-stimulus interval. For the preference task, participants were asked to guess whether the words meant something good or something bad on the basis of their sound, rating each word as "very nice/good", "slightly nice/good", "slightly nasty/bad" or "very nasty/bad". The mere exposure effect is shown by preference for previously presented words relative to foils. Responses were scored on a four-point scale (0 = very nasty, 3 = very nice). Preference priming scores were calculated by subtracting sum of preference ratings for foil words from sum of preference ratings for target words resulting in a preference score ranging from -30 to +30. Positive scores indicated a mere exposure effect for target words.

Procedure

All measures were administered in counterbalanced order and duration of assessment varied from 2.5 to 4 hours.

RESULTS

We compared scores of moderate and severe groups on neuropsychological measures using the Mann Whitney nonparametric test due to unequal group sizes (Table 1). Groups were not significantly different in duration of time since injury $U = 86.0, p = .34$, and measures of current (WASI-IQ) $U = 98.5, p = .31$, and pre-morbid intelligence (WTAR), $U = 103.0, p = .80$ (Table 2). Mean WASI-IQ scores fell within average ranges and there was no significant reduction in intelligence from premorbid levels as estimated by the WTAR for both groups $F(1,32) = .91, p = .35$. Mean depression ($U = 110.0, p = .60$) and anxiety scores ($U = 116.5, p = .72$) were not different for groups.

[insert table 1 here]

Metacognitive Awareness variables

The severe group had higher total SADI scores than the moderate group indicating greater awareness deficits (Table 1). Self-rating mean was lower than Other mean for both groups on the DEX questionnaire and higher than Other means on the Awareness Questionnaire (where higher scores indicate fewer identified problems) indicating diminished awareness of problems for both patient groups. Self-Other Discrepancy on the Awareness questionnaire was significantly different for the severe group $t(44) = 3.2, p = 0.003$, but not for the moderate group $t(20) = .73, p = .48$. Self-Other discrepancy was also significantly different on the DEX questionnaire for severe $t(44) = -2.3, p = .02$, but not for moderate patients $t(20) = -.17, p = .87$. The severe group had significantly lower DEX Self-ratings than the moderate group $U = 68.5, p = 0.05$, suggesting less awareness of executive/emotional and social problems after severe compared to moderate head injury (Table 2).

[insert table 2 here]

Anticipatory/Emergent measure of awareness

The severe group had larger SRSI-Awareness index scores than the moderate group indicating more impaired awareness (see Table 1 postscript). There was no difference between groups for the SRSI-Strategy measure. Both groups in the current study showed impaired awareness on SRSI subscales compared to previous findings with 38

TBI patients (SRSI-Awareness, $M = 5.6$, SRSI-Strategy $M = 5.4$, Wise, Ownsworth & Fleming, 2005) where larger scores were associated with diminished employment status. To establish whether SRSI-Awareness and SRSI-Strategy subscales shared variance we conducted a Pearson's correlation for total group scores and results showed a moderate correlation between the two measures $r(34) = 0.65, p = .01$.

Executive function tasks

The severe group had significantly lower scores on the Sorting Test measure of concept formation $U = 62.5, p = 0.02$ compared to normal ability in the moderate group (Table 2). Severe participants generated fewer items on FAS verbal fluency $U = 70, p = 0.04$ (although both group mean scores fell within normal ranges), and showed a higher number of errors on the SOPT measure of response monitoring $U = 65.5, p = 0.02$ than the moderate group.

Implicit experimental tasks

Both groups showed a lack of mere exposure effect (no preference for previously exposed targets compared to foils at test), and were not significantly different in this respect $U = 107.0, p = .47$ (see Table 2). This is an impaired pattern of performance on this task compared to control data with 20 participants ($M = 3.5, SD = 3.2$, Barker et al., 2006) and 16 participants ($M = 3.2, SD = 3.3$, Barker et al., 2010) respectively. The moderate group showed some learning on the SRT (shown by a positive mean score), that differed significantly from the severe group who showed little implicit learning $U = 83, p = .05$. However, moderate group performance ($M = 29.3, SD = 61.7$) also fell in the impaired range when compared to control data with 20 participants ($M = 82.4, SD = 41.1$, Barker et al., 2006) and 16 participants ($M = 96.4,$

SD = 40.5, Barker et al., 2010). Neither group reached the cut-off criterion of 16 on the explicit measure, $M = 6.6$ (SD = 4.9) for the moderate group, and $M = 1.9$ (SD = 3.1) for the severe group.

To summarize, groups were not different in time since injury, IQ or mood state. The severe group had greater awareness deficits on SADI and SRSI-Awareness measures than the moderate group, although both groups showed impaired emergent awareness on SRSI-Awareness and -Strategy indices. Groups had discrepant Self- and Other-ratings on AQ and DEX awareness measures that differed significantly for the severe group. The severe group had greater impairment on the SOPT and Sorting Test measures of executive function than the moderate group. Neither group showed a priming effect for previously exposed stimuli and the severe group had significantly lower learning scores on the SRT task than the moderate group, although both group mean scores were low compared to published control data. Due to the small sample size all further analyses were conducted on the total sample of 34 participants.

We conducted 5 separate hierarchical multiple regressions with each of the criterion variables (DEX and AQ-discrepancy, SADI, SRSI-Awareness and SRSI-Strategy) to establish the unique contribution of predictors to the variance in criterion variables with the contribution of other variables accounted for. We followed the standard convention of introducing demographic variables of IQ, mood state and injury severity at block one to account for any effects and measured the unique and additional contribution of theoretically important predictors in block two. In block one, measures of intelligence (WASI), emotional state (HADS combined anxiety and depression score), and severity of injury were entered into the analysis. Executive function variables, SOPT, Sorting Test, and Brixton scores were entered in block two

for the metacognitive criterion variables (SADI, AQ- and DEX-discrepancy scores). For the emergent/anticipatory awareness criterion variables (SRSI-Awareness and SRSI-Strategy), implicit task predictors (SRT learning score and mere exposure effect preference score) were also entered into the analysis with executive function predictor variables at block two (see Table 3).

[Insert table 3 here]

Regression statistics show that the greatest change to models occurred for the SADI criterion variable with the addition of severity of injury, WASI and HADS at block one, and SRSI-Awareness and SRSI-Strategy criterion variables with the addition of executive and implicit variables in block two (Table 3). Beta values and significance levels of each predictor are shown in table four.

[Insert table 4 here]

The strongest unique predictor of the SADI metacognitive criterion variable was severity of injury followed by mood state measured by the HADS. For AQ- and DEX-discrepancy criterion variables the strongest unique predictor was response monitoring measured by the SOPT, followed by mood state (HADS) for AQ-Discrepancy only. For the emergent/anticipatory criterion variables, the strongest unique predictors of SRSI-Awareness were response monitoring (SOPT), concept formation (Sorting test), IQ, severity of injury and implicit learning on the SRT task. The strongest unique predictor of the SRSI-Strategy emergent awareness measure was concept formation, followed by implicit learning on the SRT task. The Brixton

executive function task and mere exposure effect implicit task made no significant contribution to criterion variables in regression models. It is likely that mere exposure effect made no significant contribution to the regression model because neither patient group showed any priming effects at test and the variance in scores across participants was therefore small for the severe group (mean = -1.36, SD 4.0 and for the moderate group (mean = -0.26, SD 4.9). Normal populations typically show a small but robust priming effect on this task (Barker et al., 2006; Barker et al., 2010).

DISCUSSION

The present study investigated the contribution of injury severity, IQ, mood state, executive and implicit function contributions to metacognitive and emergent/anticipatory awareness post-injury. Results of non-parametric analyses showed that the severe group had greater impairment on metacognitive measures of awareness (SADI, AQ- and DEX-Discrepancy), and emergent/anticipatory awareness measured by the SRSI-Awareness subscale. Metacognitive awareness is thought to depend on executive, processes and emergent/anticipatory awareness is thought to depend upon executive, implicit and other sub-component (episodic and working memory) processes (Toglia & Kirk, 2000; Morris & Hannesdottir, 2004). In line with this assumption, the severe group also had greater impairment than the moderate group on executive measures of concept formation, verbal fluency, response monitoring (indexed by Sorting Test, FAS and SOPT respectively), and implicit cognition measured by the SRT task. We did not find global awareness, executive and implicit deficits after severe compared to moderate injury possibly reflecting the multi-componential nature of these functions.

Groups were not different in time since injury, current and premorbid IQ and mood state and showed similarly impaired scores on the SRSI-Strategy measure of emergent/anticipatory awareness and the mere exposure effect implicit task. The moderate group showed impaired metacognitive (AQ- and DEX-discrepancy scores) and emergent awareness (SRSI-Awareness and SRSI-Strategy scores), and diminished implicit learning and priming compared to normative data (Barker et al., 2006; Barker et al., 2010).

Metacognitive awareness measures

We found no appreciable contribution of executive function to metacognitive awareness measured by the SADI, instead only injury severity and mood state significantly predicted scores on this variable. Injury severity is a relatively non-specific variable and we found that executive and implicit functions decreased as a function of severity of injury so it is possible that the significant contribution made by the severity variable to SADI scores masked the contribution of other variables to this measure. However, further investigation of the SADI is probably warranted to establish which aspects of awareness are captured by this measure. Combined anxiety and depression scores also predicted metacognitive awareness measured by SADI and AQ-discrepancy measures with greater emotional distress associated with lower awareness scores (i.e. greater awareness) suggesting that increased emotional distress might heighten self-reflection/self-evaluation (Godfrey, Partridge, Knight & Bishara, 1993; Forgas, 1998; Alloy & Abramson, 1979). It remains to be established whether improved awareness of deficits increases emotional distress or emotional distress precipitates more accurate self-appraisal, and it is likely that the relationship between awareness of functional deficits and mood state is bidirectional.

The executive function of response monitoring (number of errors made) measured by the SOPT was the largest unique predictor of metacognitive awareness measured by AQ- and DEX-discrepancy measures (see Hart et al., 2005 for similar findings). Errors on the SOPT occur due to failure to effectively monitor previously used responses (Petrides & Milner, 1982). Other findings show that diminished self-monitoring processes deleteriously affect self-report accuracy of cognitive deficits post-injury (Schmitz, Rowley, Kawahara & Johnson, 2006; Oddy, Coughlan, Tyerman & Jenkins, 1985). Our findings support the assumption that self-monitoring is an important executive process mediating metacognitive awareness proposed by theoretical models (Toglia & Kirk, 2000; Ownsworth, Clare & Morris, 2006). Response monitoring is also likely to recruit working memory processes and working memory span has been associated with successful SOPT performance (Rich, Blysm & Brandt, 1996). Future work should incorporate executive function and working memory measures to further extract the constituent cognitive components of awareness post-head injury.

Emergent/anticipatory awareness

The strongest unique predictors of emergent/anticipatory awareness measured by SRSI-Awareness were concept formation (Sorting Test scores) and response monitoring (SOPT error scores) contributing an equal amount of variance to the criterion variable. Current IQ status and severity of injury were the second strongest predictors of SRSI-Awareness also contributing an equal amount of variance to emergent awareness scores. In the present study IQ was assessed using two subtests, one of verbal knowledge/reasoning and the other involving abstract rule detection and

nonverbal reasoning processes, either or both of these aspects of ability could have contributed to scores on awareness measures. Severity of injury also made a significant contribution to emergent awareness, as with the SADI metacognitive criterion variable indicating some general effects of diffuse neuropathology to awareness indexed by this measure.

Implicit learning measured by the SRT task made a significant contribution to emergent awareness. Concept formation and implicit learning also contributed significantly to awareness measured by the SRSI-Strategy awareness subscale. This latter effect is likely to reflect shared variance across SRSI-Awareness and –Strategy subscales shown by the moderate correlation between indices, and that specific executive and implicit predictors captured these shared processes across the two measures. In contrast, response monitoring only significantly contributed to SRSI-Awareness, although the beta was large but non-significant for the contribution of response monitoring to awareness indexed by the SRSI-Strategy subscale.

The contribution of implicit learning to emergent awareness shown in present data raises questions about the mechanisms underpinning sequence detection and learning on this task (Barker et al., 2006; Barker et al., 2010; Lieberman, 2000). One candidate explanation is that learning on the task depends upon the integrity of an implicit anticipatory/ predictive mechanism mediated by frontal circuitry and diminished or abolished after neuropathology (Seidler et al., 2005; Wong, Bernat, Bunce & Shevrin, 1997), that may contribute to other functions including emergent awareness. Typically, participants are faster on sequence compared to random trials on the SRT task because they have learned the sequence and can predict/anticipate subsequent

stimulus locations (Seger, 1997). Although the exact cause of absence of learning on the task remains to be elucidated, slowed responses on sequence compared to random blocks at test indicate that participants are unable to anticipate or predict subsequent stimulus locations. Head-injured participants also usually show an absence of explicit knowledge of the sequence shown by low scores on the explicit measure indicating that participants are not relying on explicit awareness of the sequence to facilitate performance in the absence of implicit learning. This finding is relatively robust and has been replicated across several studies with different head-injured cohorts (Barker et al., 2006; Barker et al., 2010; Beldarrain et al., 2002).

Overall, regression data indicate that failure to implicitly encode contextual regularities and accurately predict/anticipate a response, inability to formulate and monitor response strategies and injury severity is associated with impaired emergent awareness. IQ, mood state, injury severity and response monitoring contribute to metacognitive awareness. It remains to be established if these functions share similar neural substrates (although see Seidler et al., 2005; Schmitz et al., 2006; Sherer et al., 2005), and/or typically operate as a functional system(s) mediating awareness that may be differentially affected by neuropathology.

Considered together our findings might shed some light on anecdotal reports of patients who express awareness of deficits but are unable to read and respond to social cues appropriately (Prigatano & Schacter, 1991), and those who show intact implicit awareness by avoiding tasks they find difficult but who seem unable to verbalize awareness deficits (Schacter, 1990; Kihlstrom & Tobias, 1991; Morris & Hannesdottir, 2004). Although speculative these patterns of functional outcome might

reflect some dissociation of emergent/anticipatory and metacognitive awareness and/or differential impairment to cognitive functions contributing to these aspects of awareness. Although there is a growing body of evidence investigating executive components of awareness (Ownsworth, McFarland & Young, 2000; Ownsworth & Fleming, 2005; Bogod, Mateer & MacDonald 2003; O’Keeffe et al., 2007), less attention has been paid to the empirical measurement of implicit cognition in awareness studies. Our findings go some way to elucidate some of the constituent components of metacognitive and emergent/anticipatory awareness and the differential effects of moderate and severe injuries on these functions and awareness more generally.

REFERENCES

Alloy, L.B., & Abramson, L.Y. (1979). Judgement of contingency in depressed and nondepressed students: sadder but wiser? *Journal of Experimental Psychology*, *108*, 441-485.

Andrés, P., & Van Der Linden, M. (2002). Are central executive functions working in patients with focal frontal lesions? *Neuropsychologia*, *40*, 835-845.

Banks S.J., & Weintraub, S. (2009). Generalized and specific-symptom insight in behavioral variant frontotemporal dementia and primary progressive aphasia. *Journal of Neuropsychiatry and clinical neurosciences*, *21*, 299-306.

Barker, L. A., Andrade, J., Morton, N., Romanowski, C. A. J., & Bowles, D. (2010). Investigating the ‘latent’ deficit hypothesis: Age at time of head injury, implicit and

executive functions and behavioral insight. *Neuropsychologia*, 48, 2550-2563.

Barker, L. A., Andrade, J., Romanowski, C. A. J., Morton, N., & Wasti, A. (2006). Implicit cognition is impaired and dissociable in a head-injured group with executive deficits. *Neuropsychologia*, 44, 1413-1424.

Barker, L. A., Andrade, J., & Romanowski, C. A. J. (2004). Impaired implicit cognition with intact executive function after extensive bilateral prefrontal pathology: A case study. *Neurocase*, 10, 233-248.

Beldarrain, M. G., Grafman, J., de Valesco, I. R., Pascual-Leone, A., & Garcia-Monco, J.C. (2002). Prefrontal lesions impair the implicit and explicit learning of sequences on visuomotor tasks. *Experimental Brain Research*, 142, 529-538.

Benton, A.L., & Hamsher, K. deS. (1989). *Multilingual Aphasia Examination*. Iowa City, Iowa: AJA Associates.

Bogod, N.M., Mateer, C.A., & MacDonald, S.W. S. (2003). Self-awareness after traumatic brain injury: a comparison of measures and their relationship to executive functions. *Journal of the International Neuropsychological Society*, 9, 450-458.

Burgess, P.W. & Shallice, T. (1997). *The Hayling and Brixton tests*. Thurston, Suffolk:Thames Valley Test Company.

Crosson, B. C., Barco, P.P., Velozo, C.A., Bolesta, M.M., Cooper, P.V., Werts, D., &

- Brobeck, T.C. (1989). Awareness and compensation in post-acute head injury rehabilitation. *Journal of Head Trauma Rehabilitation, 4*, 46-54.
- Delis, D.C., Kaplan, E., & Kramer, J.H. (2001). *Delis Kaplan Executive Function System*. The Psychological Corporation.
- Flashman, L.A., & McAllister, T.W. (2002). Lack of awareness and its impact in traumatic brain injury. *Neurorehabilitation, 17*, 285-296.
- Forgas, J.P. (1998). On being happy and mistaken: mood effects on the fundamental attribution error. *Journal of Personality & Social Psychology, 75*, 318-331.
- Fleming, J.M., & Strong, J. (1999). A longitudinal study of self-awareness functional deficits underestimated by persons with brain injury. *Occupational Therapy Journal of Research, 19*, 3-17.
- Fleming, J.M., Strong, J., & Ashton, R. (1998). Cluster analysis of self-awareness levels in adults with traumatic brain injury and relationship to outcome. *Journal of Head Trauma Rehabilitation, 13*, 39-51.
- Fleming, J., Strong, J., & Ashton, R. (1996). Self-awareness of deficits in adults with traumatic brain injury. *Brain Injury, 10*, 1-15.
- Frith, C.D., & Frith, U. (2008). Implicit and explicit processes and social cognition. *Neuron, 60*, 503-510.

Godfrey, H.P.D., Partridge, F.M., Knight, R.G., & Bishara, S. (1993). Course of insight disorder and emotional dysfunction following closed head injury: a controlled cross sectional follow-up study. *Journal of Clinical and Experimental Neuropsychology*, *15*, 503-515.

Hart, T., Whyte, J., Kim, J., & Vaccaro, M. (2005). Executive function and self awareness of “real world” behaviour and attention deficits following traumatic brain injury. *Journal of Head Trauma Rehabilitation*, *20*, 333-347.

Hart, T., Seignourel, P. J., & Sherer, M. (1999). A longitudinal study of awareness of deficits after moderate to severe traumatic brain injury. *Neuropsychological rehabilitation*, *19*, 161-176.

Kihlstrom, J.F., & Tobias, B.A. (1991). Anosognosia, consciousness and the self. In: Prigatano, G.P. and Schacter, D. L. (eds.), *Awareness of Deficit After Brain Injury: Clinical and Theoretical Issues*. New York: Oxford University Press.

King, N. (1997). Mild head injury: Neuropathology, sequelae measurement and recovery. *British Journal of Clinical Psychology*, *36*, 161-184.

Kouider, S., & Dehaene, S. (2007). Levels of processing during non-conscious perception: a critical review of visual masking. *Philosophical Transactions of the Royal Society of London B*, *362*, 857-875.

Lezak, M.D. (1995). *Neuropsychological Assessment* (3rd ed.), New York: Oxford University Press.

Lieberman, M. D. (2000). Intuition: A social cognitive neuroscience approach. *Psychological Bulletin*, 126, 109-137.

Mattson, A.J., & Levin, H.S. (1990). Frontal lobe dysfunction following closed head injury: a review of the literature. *Journal of Nervous Mental Disorders*, 178, 282-312.

Mild traumatic brain injury committee of the head injury interdisciplinary special interest group of the American congress of rehabilitation medicine (1993). Definition of mild traumatic brain injury. *Journal of Head Trauma Rehabilitation*, 8, 86-87.

Miyake, A., Friedman, N. P., Emerson, M. J., Witski, A. H., Howerter, A., & Wager, T. D. (2000). The unity and diversity of executive functions and their contribution to complex 'frontal lobe' tasks: A latent variable analysis. *Cognitive Psychology*, 40, 49-100.

Morris, R.G., & Hannesdottir, K. (2004). Loss of "awareness" in Alzheimers disease. In Morris, R.G. & Becker, J.T. (Eds), *Cognitive Neuropsychology of Alzheimer's Disease*. Oxford: Oxford University Press.

Nissen, M. J., & Bullemer, P. (1987). Attentional requirements of learning: Evidence from performance measures. *Cognitive Psychology*, 19, 1-32.

Oddy, M., Coughlan, T., Tyerman, A., & Jenkins D. (1985). Social adjustment after head injury: a further follow-up seven years after injury. *Journal of Neurology, Neurosurgery and Psychiatry*, 48, 564-568.

O'Keeffe, F., Dockree, P., Moloney, P., Carton, S., & Robertson, I.H. (2007). Awareness of deficits in traumatic brain injury: A multidimensional approach to assessing metacognitive knowledge and online awareness. *Journal of the International Neuropsychological Society*, 13, 38-49.

Owensworth, T., Desbois, J., Grant, E., Fleming, J., & Strong, J. (2006). The associations among self-awareness, emotional well-being, and employment outcome following acquired brain injury: A 12-month longitudinal study. *Rehabilitation Psychology*, 51, 50–59.

Owensworth, T., Clare, L., & Morris, R. (2006). An integrated biopsychosocial approach to understanding awareness deficits in Alzheimer's disease and brain injury. *Neuropsychological Rehabilitation*, 16, 415-438.

Owensworth, T.L., & Fleming, J. (2005). The relative importance of meta-cognitive skills, emotional status, and executive function in psychosocial adjustment following acquired brain injury. *Journal of Head Trauma Rehabilitation*, 20, 315-332.

Owensworth, T. L., McFarland, K., & Young, R. (2000). Development and standardization of the Self-Regulation Skills Interview (SRSI): A new clinical assessment tool for acquired brain injury. *The Clinical Neuropsychologist*, 14, 76-92.

Ponsford, J. (Ed.), (2004). *Cognitive and Behavioral Rehabilitation: From Neurobiology to Clinical Practice*. London: The Guildford Press.

Petrides, M., & Milner, B. (1982). Deficits on subject ordered tasks after frontal and temporal-lobe lesions in man. *Neuropsychologia*, 20, 263-274.

Prigatano, G. P; & Schacter, D. L. (Eds;) (1991). *Awareness of deficit after brain injury: Clinical and theoretical issues*. New York: Oxford University Press.

Prigatano, G. P. (2010). *Advances in the study of anosognosia*. New York: Oxford University Press.

Reber, A. S. (2002). Prologue in R. M. French and A. Cleeremans. *Implicit learning and consciousness: An empirical, philosophical and computational consensus in the making*. London. Psychology Press.

Rich, J.B., Blyma, F.W., & Brandt, J. (1996). Self-ordered pointing performance in Huntington's disease patients. *Neuropsychiatry, Neuropsychology and Behavioural Neurology*, 9, 99-106.

Rosen, H. J., Alcantar, O., Rothlind, J., Sturm, V., Kramer, J. H., Weiner, M., & Miller, B. L. (2010). Neuroanatomical correlates of cognitive self-appraisal in neurodegenerative disease. *Neuroimage*, 49, 3358, 3364.

Schacter, D.L. (1990). Toward a cognitive neuropsychology of awareness: implicit knowledge and anosognosia. *Journal of Clinical and Experimental Neuropsychology*, 12, 155-178.

Schmitz, T. W., Rowley, H. A., Kawahara, T. N., & Johnson, S. C. (2006). Neural correlates of self-evaluative accuracy after traumatic brain injury. *Neuropsychologia*, 44, 762-773.

Seger, C. A. (1997). Two forms of sequential implicit learning. *Consciousness and Cognition*, 6, 108-131.

Seidler, R. D., Puroshotham, A., Kim, S. G., Ugurbil, K., Willingham, D., & Ashe, J. (2005). Neural correlates of encoding and expression in implicit sequence learning. *Experimental Brain Research*, 165, 114-124.

Sherer, M., Hart, T., Whyte, J., Todd, N.G., & Yablon, S.A. (2005). Neuroanatomic basis of impaired self-awareness after traumatic brain injury: findings from early computed tomography. *Journal of Head Trauma Rehabilitation*, 20, 287-300.

Sherer, M., Hart, T., & Todd, N. (2003). Measurement of impaired self-awareness after traumatic brain injury: a comparison of the patient competency rating scale and the awareness questionnaire. *Brain Injury*, 17, 25-37.

Sherer, M., Boake, C., Levin, E., Silver, B.V., Ringholz, G., & Walter, M.H. (1998). Characteristics of impaired awareness after traumatic brain injury. *Journal of the*

International Neuropsychology Society, 4, 380-387.

Stuss, D., & Alexander, M. P. (2007). Is there a dysexecutive syndrome? *Philosophical Transactions of the Royal Society of London B, Biological Science*, 362, 901-915.

The Psychological Corporation (2001). *Wechsler Test of Adult Reading*. San Antonio, TX: Harcourt Assessment.

Toglia, J., & Kirk, U. (2000). Understanding awareness deficits following brain injury. *NeuroRehabilitation*, 15, 57-70.

Trahan, E., Pépin, M., & Hopps, S. (2006). Impaired awareness of deficits and treatment adherence among people with traumatic brain injury or spinal cord injury. *Journal of Head Trauma Rehabilitation*, 21, 226-235.

Wechsler, D. (1999). Wechsler Abbreviated Scale of Intelligence. *Psychological Corporation*.

Wilson, B.A., Alderman, N., Burgess, P.W., Emslie, H., & Evans, J. J. (1996). *Behavioural Assessment of the Dysexecutive Syndrome*. Thames Valley Test Company. Bury St. Edmunds, England:

Wise, K., Ownsworth, T., Fleming, J. (2005). Convergent validity of self-awareness measures and their association with employment outcome in adults following

acquired brain injury. *Brain Injury*, 19, 765-775.

Zajonc, R. B. (1968). Attitudinal effects of mere exposure. *Journal of Personality and Social Psychology Monograph*, 9, 1-28.

Zajonc, R. B. (1980). Feeling and thinking: Preferences need no inferences. *American Psychologist*, 35, 151-175.

Zigmond, A.S., & Snaith, R. P. (1983). The Hospital Anxiety and Depression Scale. *Acta Psychiatrica Scandinavica*, 67, 361-70.

Wong, P. S; Bernat, E; Bunce, S; & Shevrin, H. (1997). Brain indices of nonconscious associative learning. *Consciousness and Cognition*, 6, 519-544.

ACKNOWLEDGEMENTS

The authors would like to thank all head-injured participants who took part in the study. The authors report no conflict of interest or sources of financial support.

Table 1: Demographic, neurological, intelligence, emotion and awareness variables for moderate, severe and total TBI group.

Demographic, neurological and test variables	Moderate Group mean (SD's, range) (n = 11)	Severe Group mean (SD's, range) (n = 23)	Total Group mean (SD's, range) (N = 34)
Sex (male:female)	10:1	22:1	32:2
Age (yrs)	31.6 (10.3, 20-54)	36.6 (10.0, 18-55)	35.0 (10.2, 18-55)
Education (yrs)	11.9 (2.4, 10-18)	11.5 (1.3, 9-15)	11.6 (1.7, 9-18)
TBI severity variables			
Time since injury (months)	47.5 (48.8, 12-149)	74.6 (63.7, 12-240)	66.2 (60.0, 12-240)
GCS score on admission	13.0 (2.7, 9-15)	5.2 (3.4, 3-8)	8.1 (5.0, 3-15)
Length of coma (hours)	1.8 (2.4, 0-6)	396.4 (349.1, 8-1176)	277.3 (343.9, 0-1176)
PTA (hours)	57.9 (54.9, 24-168)	1517.5 (1595.7, 192-7200)	1030.9 (1469.8, 24-7200)
Lesion site CT/MRI scans (Number of participants)			
1. Unilateral frontal region	6	9	15
2. Bilateral frontal and other brain region.	0	7	7
4. Unilateral frontal and other brain region	0	2	2
5. Unilateral Temporal	2	4	6
6. No imaging data available	3	1	4
Type of injury			
Road traffic accident	6	14	20
Assault	2	3	5
Fall	3	6	9

Table 2: Moderate and severe subgroup and total group scores for executive and implicit experimental tasks (mean, SD, and range).

Measure	Moderate Group (n = 11)	Severe Group (n = 23)	Total Group, mean (SD), range (N = 34)
<i>Intelligence</i>			
WASI Full Scale IQ	96.0 (14.5)	91.1 (17.6)	92.7 (16.6, 63-135)
WTAR premorbid IQ	98.3 (8.9)	96.2 (13.5)	96.9 (11.9, 74-122)
<i>Emotional State</i>			
HADS – Anxiety	9.8 (3.5)	9.3 (4)	9.4 (3.8, 2-18)
HADS - Depression	6.0 (3.7)	7.4 (4.6)	6.9 (4.4, 1-17)
<i>Awareness Measures</i>			
Self-Awareness of Deficits Interview total score (SADI)	1.6 (1.7)	4.0 (2.2) [†]	3.2 (2.3, 0-8)
AQ-Self	36.7 (10.4)	38.7 (16.0)	38.1 (14.3, 20-85)
AQ-Other	33.7 (8.8)	27.7 (4.4)*	29.7 (6.7, 21-48)
AQ discrepancy score	3.0 (9.9)	11.0 (15.5)	8.4 (14.3, -7-58)
DEX-Self	42.6 (15.7)	34.1 (18.8)**	36.7 (18.1, 1-73)
DEX-Other	44.1 (23.6)	45.5 (13.0)	45.1 (16.7, 4-77)
DEX-discrepancy score	1.5 (15.9)	11.5 (21.8)	8.3 (20.4, -26-63)
Self-Regulatory Skills Interview – Awareness Index (SRSI-A)	10.0 (5.8)	14.8 (4.2) ^{††}	13.2 (5.2, 0-20)
SRSI – Strategy Index (SRSI-S)	16.6 (8.0)	18.5 (6.1)	17.9 (6.7, 6-30)
<i>Executive function measures</i>			
Sorting Test combined description scaled score	9.2 (2.6)	6.6 (3.7)*	7.4 (3.6, 1-16)
Self Ordered Pointing test (SOPT) – total errors	14.2 (12.2)	24.1 (9.1)*	20.9 (11.1, 0-44)
Brixton task – total errors	6.2 (1.7)	5.4 (2.3)	5.6 (2.1, 2-10)
FAS – total number of words	37.7 (13.0)	27.8 (8.4)*	31.0 (11.1, 14-56)
<i>Implicit Cognitive tasks</i>			
Serial Reaction Time Test (SRT) – Learning Score	29.3 (61.7)	-4.1 (146.6)*	6.7 (125.4, -200.2-554.5)
Mere exposure preference score (the difference between target and foil scores)	-0.26 (4.9)	-1.36 (4.0)	-0.62 (4.6, -14-9)

* $p < 0.05$, ** $p = 0.04$, † $p = 0.001$, †† $p = 0.007$ one-tailed

Table 3: Hierarchical regression model statistics for all predictor and criterion variables.

WASI/HADS/ TBI severity variables (block 1)					
Regression statistics	DEPENDENT VARIABLES				
	SADI	AQ-Discrepancy	DEX-Discrepancy	SRSI – Awareness	SRSI – Strategy
F _{Change}	6.69	3.25	1.74	3.55	1.10
<i>p</i>	0.002**	0.04*	0.18	0.04*	0.36
R ² _{Change}	0.44	0.27	0.28	0.19	0.06
Executive function variables (block 2)				Executive function and Implicit cognition variables (block 2)	
Regression statistics	SADI	AQ-Discrepancy	DEX-Discrepancy	SRSI – Awareness	SRSI – Strategy
F _{Change}	0.77	2.84	3.80	4.62	4.01
<i>p</i>	0.53	0.06	0.02*	0.006**	0.01**
R ² _{Change}	0.05	0.20	0.28	0.33	0.35

p* ≤ .05 *p* ≤ .01

Table 4: Beta values and significance level for all predictor and criterion variables in hierarchical regression models.

Predictor variables	Metacognitive criterion variables			Anticipatory/Emergent criterion variables	
	SADI	AQ-Discrepancy	DEX-Discrepancy	SRSI – Awareness	SRSI – Strategy
General Variables					
WASI- IQ <i>P</i> value	-0.29 (0.09)	-0.05 (0.80)	-0.16 (0.44)	-0.35* (0.04)	-0.32 (0.06)
HADS <i>P</i> value	-0.31* (0.05)	-0.44* (0.02)	-0.28 (0.15)	-0.30 (0.06)	-0.04 (0.85)
TBI severity <i>P</i> value	0.40* (0.02)	0.30 (0.11)	0.19 (0.34)	0.35* (0.04)	0.04 (0.85)
Executive Function tasks					
Sorting Test <i>P</i> value	-0.09 (0.70)	0.38 (0.12)	-0.32 (0.09)	-0.42* (0.03)	-0.59** (0.01)
SOPT (error score) <i>P</i> value	0.28 (0.18)	0.52* (0.02)	0.51* (0.02)	0.42* (0.02)	0.31 (0.07)
Brixton Test <i>P</i> value	-0.01 (0.95)	-0.21 (0.22)	-0.27 (0.06)	-0.24 (0.13)	0.09 (0.63)
Implicit measures					
SRT <i>P</i> value				-0.29* (0.04)	-0.29* (0.03)
Mere exposure effect preference score <i>P</i> value				-0.01 (0.97)	0.10 (0.54)

* $p \leq .05$ ** $p \leq .01$