


INTRODUCTION

- Prostate and bladder cancers are the first and fourth most common noncutaneous cancers in men in developed nations
- Plasma is also called the fourth state of matter
 - Ionized gas composed of reactive species, electrical and excited particles
- Cold plasmas recently came into the attention of medical society due to its non-inflammatory selective ablative effects in tumour cells
 - Their effects result from different interactions between plasma components with specific structural cell elements as well as cell functionalities
 - Their selectivity of action results from the lower antioxidant defenses of tumor cells and the cell cycle block at phase G2/M in rapidly multiplying cells
- Plasma therapy is still a developing field, but it seems to be effective against a wide range of tumors – melanoma, glioblastoma, breast, colon, lung and cervix cancer
 - Little is known about its role in the treatment of urologic neoplasms
- Goal:** The aim of this work was to evaluate the cytotoxicity of cold atmospheric plasma (CAP) in prostate and urinary bladder cancer cell lines

MATERIALS AND METHODS

- Our group developed an electronic device capable of generating high output voltage that can ionize a significant fraction of air particles producing cold atmospheric plasma (CAP)
- Cell lines - plated in a concentration of 50.000-100.000 cells/mL in 200pL of cell culture medium
 - Prostate adenocarcinoma - LNCaP and PC3
 - Urinary bladder carcinoma - HT1376
 - Phenotypically normal human fibroblasts cell line - HFF-1
- The device was designed to expose cell cultures seeded in multiwell plates to short periods of CAP
 - Range from 15 to 120 seconds



- Metabolic activity (MA) and protein content (P) were assessed with colorimetric assays MTT and SRB, respectively

DISCUSSION

- CAP is effective on LNCaP and HT1376 cells – over 90% decrease in metabolic activity and protein content after only 60s of exposure
- Regarding to PC3 prostate cancer cell line, there was a decrease in metabolic activity after 60 seconds similar to the other two tumor lines, but the protein content did not have the same reaction even with 120s of exposure
- For the same exposure conditions, there was less change in metabolic activity or protein content of the normal fibroblast line
- CAP may offer a selective anti-tumour therapy capable of providing ablation of tumours after short time courses, typically in the range of seconds-minutes, without damaging with adjacent normal tissues
- Further studies are needed to determine the utility of CAP in the treatment of bladder and prostate cancer

REFERENCES

- Silva-Teixeira, R. et al. Cold atmospheric plasma as an approach to retinoblastoma. *Eur. J. Cancer* 61, S143 (2016).
- Gay-Mimbrera, J. et al. Clinical and Biological Principles of Cold Atmospheric Plasma Application in Skin Cancer. *Adv. Ther.* 33, 894–909 (2016).
- Hoffmann, C., Berganza, C. & Zhang, J. Cold Atmospheric Plasma: methods of production and application in dentistry and oncology. *Med. Gas Res.* 3, 21 (2013).
- von Woedtke, T., Reuter, S., Masur, K. & Weltmann, K. D. Plasmas for medicine. *Phys. Rep.* 530, 291–320 (2013).
- Ahn, H. J. et al. Atmospheric-pressure plasma jet induces apoptosis involving mitochondria via generation of free radicals. *PLoS One* 6, (2011).

RESULTS




Figure 1 - MTT assay results 24 h after plasma therapy application in different human cell lines: prostate adenocarcinoma (PC3 and LNCaP), urinary bladder carcinoma (HT1376) and fibroblasts (HFF1) at distinct times: 15, 30, 60, 90 and 120 seconds (s). Results are expressed as percentage of metabolic activity normalized to control.




Figure 2 - SRB assay results 24 h after plasma therapy application in different human cell lines: prostate adenocarcinoma (PC3 and LNCaP), urinary bladder carcinoma (HT1376) and fibroblasts (HFF1) at distinct times: 15, 30, 60, 90 and 120 seconds (s). Results are expressed as percentage of protein content normalized to control.