



# Estrogens Promote the Production of Natural Neutralizing Antibodies in Fish through G Protein-Coupled Estrogen Receptor 1

María C. Rodenas<sup>1</sup>, Isabel Cabas<sup>1</sup>, Nuria E. Gómez-González<sup>1</sup>, Marta Arizcun<sup>2</sup>, José Meseguer<sup>1</sup>, Victoriano Mulero<sup>1\*</sup> and Alfonsa García-Ayala<sup>1\*</sup>

<sup>1</sup> Department of Cell Biology and Histology, Faculty of Biology, University of Murcia, IMIB-Arixaca, Murcia, Spain,

<sup>2</sup> Centro Oceanográfico de Murcia, Instituto Español de Oceanografía (IEO), Murcia, Spain

## OPEN ACCESS

### Edited by:

Lluís Tort,  
Universitat Autònoma  
de Barcelona, Spain

### Reviewed by:

Carolina Tafalla,  
Instituto Nacional  
de Investigación y Tecnología  
Agraria y Alimentaria, Spain  
Irene Salinas,  
University of New Mexico,  
United States

### \*Correspondence:

Victoriano Mulero  
vmulero@um.es;  
Alfonsa García-Ayala  
agayala@um.es

### Specialty section:

This article was submitted to  
Comparative Immunology,  
a section of the journal  
Frontiers in Immunology

**Received:** 26 April 2017

**Accepted:** 12 June 2017

**Published:** 29 June 2017

### Citation:

Rodenas MC, Cabas I, Gómez-González NE, Arizcun M, Meseguer J, Mulero V and García-Ayala A (2017) Estrogens Promote the Production of Natural Neutralizing Antibodies in Fish through G Protein-Coupled Estrogen Receptor 1. *Front. Immunol.* 8:736. doi: 10.3389/fimmu.2017.00736

Natural antibodies play crucial roles in pathogen elimination, B-cell survival and homeostasis, and inflammatory and autoimmune diseases. Although estrogens are able to regulate both innate and adaptive immune responses, their role in the production of natural antibodies is unknown. Here, we show that the dietary intake of the synthetic estradiol analog, 17 $\alpha$ -ethinylestradiol (EE<sub>2</sub>), one of the most potent pharmaceutical estrogens and intensively used in human therapeutics as a component of most oral contraceptives, regulates the abundance and proliferation of T and IgM<sup>+</sup> B lymphocytes in the teleost fish gilthead seabream (*Sparus aurata* L.). Furthermore, for the first time in vertebrates, it is shown that estrogen signaling through G protein-coupled estrogen receptor 1 (GPER1) induces the production of polyreactive natural antibodies, which are able to crossreact with unrelated antigens and commensal and pathogenic bacteria. In addition, the serum from fish treated with EE<sub>2</sub> or the GPER1 agonist G1 shows higher complement-dependent bactericidal activity than that from non-treated specimens. These results demonstrate that estrogens and GPER1 are the key regulators of natural antibody production and pathogen clearance in fish, paving the way for future studies in other vertebrate classes.

**Keywords:** G protein-coupled estrogen receptor 1, estrogens, lymphocytes, natural antibodies, evolution, fish

## INTRODUCTION

The impact of estrogens on immune responses is well documented (1–4). For many years, the immune system has been considered as a natural target for estrogen action (5), as corroborated by the clear sex differences observed in autoimmune and inflammatory disorders (4).

Estrogens affect different stages of B-cell development and modify the humoral response (3), while a third generation of selective estrogen receptor (ER) modulators has been found to regulate B development and function (6). Moreover, it has recently been described that the activation of ERs directly influences antibody (Ab) expression by binding to switch sites and regulatory elements in the immunoglobulin heavy chain locus of activated B cells (7). All these effects are possible due to the presence of ER in immune cells (4), including lymphocytes (8).

Classically, the action of estrogens was thought to be mediated by nuclear ER $\alpha$  and ER $\beta$ , which function as hormone-inducible transcription factors by binding to the estrogen-responsive element

located within the promoter region of target genes (9), although it was later confirmed that they are also able to rapidly activate transduction pathways *via* non-genomic mechanisms. These additional effects are mediated by a membrane-anchored receptor called G protein-coupled estrogen receptor 1 (GPER1), which was identified by independent laboratories in the early 2000s (10, 11) and later shown to be activated by 17 $\beta$ -estradiol (E<sub>2</sub>) (12–14). GPER1 activation downstream mechanisms include several signaling pathways involving MAPKs, ERK, PI3K, cAMP, and intracellular calcium (15, 16).

Xenoestrogens or environmental estrogens, known as endocrine disruptor chemicals (EDCs), are synthetic or natural substances of high stability that exert toxicity by mimicking the effects of estrogens (17). They have been described as being involved in autoimmunity (18, 19) and as reducing B-cell precursors in mice (20) and have also been found to activate GPER1 (21). 17 $\alpha$ -Ethinylestradiol (EE<sub>2</sub>) is a synthetic analog of E<sub>2</sub> and one of the most potent pharmaceutical estrogens and is intensively used in human therapeutics as a component of most oral contraceptives. The binding affinity of EE<sub>2</sub> to human ERs is one to two times higher than E<sub>2</sub> (22). However, it has been calculated that 16–68% of the dose is excreted in the urine or feces (23), reaching the waste water treatment plants, where cannot be totally eliminated (24). Consequently, an unquantified load of xenoestrogens are released into the aquatic environment, where they can be absorbed by sediment and persist for long periods, or taken up by animals and concentrated in their tissues (22, 25). It has been found in concentrations up to 21 ng/l in a south-western European river (Mira, Portugal) (26).

The gilthead seabream (*Sparus aurata* L.) is a seasonal marine teleost of great commercial value in the Mediterranean area. Its hermaphrodite character makes it an interesting animal model to analyze the role of estrogens, natural or synthetic, in the immune response. We have previously demonstrated that nuclear ERs and GPER1 are expressed in gilthead seabream head kidney (bone marrow equivalent) leukocytes (27, 28) and that GPER1 modulates granulocyte functions through a cAMP/protein kinase A/CREB signaling pathway (27). We also observed that EDCs altered the immune response of gilthead seabream by promoting some long-lasting effects even when their estrogenic disruptive effects were not present (29, 30). Moreover, EE<sub>2</sub> bath-exposed specimens have an altered capacity to respond to an immune challenge, although the compound does not behave as an immunosuppressor (31), while the dietary intake of EE<sub>2</sub> stimulates the Ab response of vaccinated fish (29, 30). However, no information exists on the impact of EE<sub>2</sub> on fish lymphocyte proliferation and differentiation or, particularly, on natural antibodies. Natural antibodies are present in the serum of vertebrates without any apparent antigenic stimulation and are an important field of research for their relevance in autoimmunity and for their role as a bridge between innate and adaptive immunities (32, 33). Although the relevance of natural antibodies in fish is largely unknown, their existence in both cartilaginous and bony fish has been reported (32, 34–36). In addition, the opsonization and neutralizing ability of natural antibodies of rainbow trout against *Aeromonas salmonicida* has been demonstrated (37–39). We, therefore, examined whether estrogens, and in particular, GPER1

signaling, are able to modulate T- and B-lymphocyte responses to an immunological challenge and natural Ab production.

## MATERIALS AND METHODS

### Animals, *In Vivo* Treatments, and Sample Collection


Healthy specimens of gilthead seabream were from the Oceanographic Center of Murcia (Mazarrón, Spain), where they were kept in running seawater aquaria (dissolved oxygen 6 ppm, flow rate 20% aquarium vol/h) with a natural temperature and photoperiod. They were fed three times per day with a commercial pellet diet (44% protein and 22% lipids; Skretting) at a feeding rate of 1.5% of fish biomass. The environmental parameters, mortality, and food intake, as well as behavior, were recorded daily.

Two-month-old gilthead seabream specimens ( $n = 100$ /treatment), with a body weight of 26.6–63.2 (from the beginning to the end of the experiment), were exposed to EE<sub>2</sub> (Figure 1A) in 170 l aquaria. Briefly, EE<sub>2</sub> (5  $\mu$ g/g food, 98% purity; Sigma) was incorporated in the commercial food using the ethanol evaporation method (0.3 l ethanol/kg of food), as described elsewhere (40). The specimens were fed three times a day *ad libitum* with the pellet diet supplemented with EE<sub>2</sub> (treated fish) or non-supplemented (untreated fish) for 76 days (days of treatment, dt). Following this time, all the specimens were fed with the commercial food for a further 23 days (days posttreatment, dpt) (Figure 1A). In order to evaluate the effect of EE<sub>2</sub> on the immune response, specimens were intraperitoneally (i.p.) injected with keyhole limpet hemocyanin (KLH) (45  $\mu$ g/fish; Sigma-Aldrich) and Imject Alum adjuvant (4 mg/fish; Thermo Scientific) (vaccinated/immunized fish) or phosphate-buffered saline (PBS) (control/unvaccinated fish) at the end of the treatment period, 76 dt. Samples were taken 40 and 76 dt and 1, 9, and 23 days postimmunization (dpi) or dpt. Food intake was similar in all groups. Specimens ( $n = 6$  fish/treatment/time of sampling) were fasted for 24 h before sampling. They were tranquilized by 8  $\mu$ l/l of clove oil, immediately anesthetized using 40  $\mu$ l/l of clove oil, weighed and decapitated before the head kidneys and spleens were removed, and processed for gene expression and/or flow cytometry analysis, as described later. Serum samples from trunk blood were obtained by centrifugation and immediately frozen and stored at  $-80^{\circ}\text{C}$  until use. Cell suspensions from head kidney and spleen were obtained as described elsewhere (41, 42).

Another set of experiments was performed for serum Ab titer determination by ELISA: (1) 2-month-old gilthead seabream specimens were dietary treated with EE<sub>2</sub> (5  $\mu$ g/g food) or the GPER1 agonist G1 (5  $\mu$ g/g food; Sigma-Aldrich) for 110 days (30), (2) adult specimens (650 g mean weight) were treated with 0, 5, and 50 ng/l EE<sub>2</sub> for 2 months by bath immersion (31), and (3) adult specimens (225 g mean weight) were exposed to dietary G1 (0, 2, and 20  $\mu$ g/fish/day treatment) for 50 days (27).

### Determination of Serum Vitellogenin (Vtg) Levels

The serum Vtg levels were determined by the enzyme-linked immunosorbent assay using a commercial kit (Cayman



**FIGURE 1** | Experimental design: 17 $\alpha$ -ethinylestradiol (EE<sub>2</sub>) dietary exposure and immunization protocol of gilthead seabream juveniles. **(A)** Fish were exposed to a non-supplemented diet (untreated fish) or to a supplemented diet with 5 µg EE<sub>2</sub>/g food (treated fish) for 76 days (days of treatment, dt). Then, the specimens were fed with commercial food for an additional period of 23 days (days posttreatment, dpt). Fish were intraperitoneally (i.p.) injected with phosphate-buffered saline (PBS) (unvaccinated or control fish) or keyhole limpet hemocyanin (KLH) plus Imject Alum adjuvant (vaccinated fish) at 76 dt. Sampling was carried out 40 and 76 dt and 1, 9, and 23 days postinjection (dpi) (1, 9, and 23 dpt). **(B)** The levels of vitellogenin (Vtg) were determined in the serum of untreated and EE<sub>2</sub> (5 µg/g food)-treated fish 1, 9, and 23 days after ceasing the treatment (dpt) by ELISA. Levels are shown as fold increase relative to the mean of untreated fish. Each bar represents mean  $\pm$  SEM of duplicates of six independent fish. The sample size was  $n = 6$  fish/group/time of sampling. The asterisks denote statistically significant differences after a Student's  $t$ -test between the untreated and the EE<sub>2</sub>-treated group, at each time point and between the EE<sub>2</sub>-treated groups at the different time points. \* $p < 0.05$ ; \*\* $p < 0.01$ ; and \*\*\* $p < 0.001$ . **(C)** The *il1b* mRNA levels were determined in the head kidney of untreated and EE<sub>2</sub> (5 µg/g food)-treated fish (vaccinated or not) 1, 9, and 23 days postimmunization (dpi) by RT-qPCR. Gene expression levels were normalized to *rps18* mRNA levels, and data represent as the mean  $\pm$  SEM of technical triplicates of six independent fish. Levels were shown as fold increase relative to the mean of non-vaccinated fish. The sample size was  $n = 6$  fish/group/time of sampling. The asterisks denote statistically significant differences after a Student's  $t$ -test between unvaccinated and vaccinated fish at each time point. \* $p < 0.05$ ; \*\* $p < 0.01$ ; and \*\*\* $p < 0.001$ .

Chemical), following the manufacturer's instructions. In brief, an aliquot of 1:500 diluted serum from untreated and EE<sub>2</sub>-treated fish ( $n = 6$  fish/treatment/time of sampling) was added to flat-bottomed 96-well plates, followed by a commercial polyclonal Ab against gilthead seabream Vtg (1:100) and an antirabbit IgG (whole molecule)-peroxidase Ab (1:1,000) (Sigma-Aldrich). Finally, the chromogen tetramethylbenzidine (TMB) was added and the absorbance was read at 450 nm using an FLUOstart luminometer (BGM; LabTechnologies).

## Analysis of Gene Expression

Total RNA was extracted from head kidney from untreated and EE<sub>2</sub>-treated (vaccinated or not) fish ( $n = 6$  fish/treatment/time of sampling) at 1, 9, and 23 dpi/dpt with TRIzol Reagent (Thermo Fisher Scientific), following the manufacturer's instructions, and quantified with a spectrophotometer (NanoDrop, ND-1000). The RNA was then treated with amplification grade DNase I (1 U/µg RNA; Thermo Fisher Scientific) to remove genomic DNA traces that might interfere with the PCRs, and the SuperScript III RNase

H Reverse Transcriptase (Thermo Fisher Scientific) was used to synthesize first-strand cDNA with oligo-dT18 primer from 1 mg of total RNA for 50 min at 50°C. The  $\beta$ -actin (*actb*) gene was analyzed by PCR performed with an Eppendorf Mastercycler Gradient Instrument (Eppendorf) to check cDNA quality. Reaction mixtures were incubated for 2 min at 94°C, followed by 30 cycles of 45 s at 94°C, 45 s at the specific annealing temperature (55°C), 45 s at 72°C, and finally, 10 min at 72°C.

In the same samples, the expression levels of the gene coding for the proinflammatory cytokine interleukin-1 $\beta$  (*il1b*) were analyzed by real-time PCR performed with an ABI PRISM 7500 instrument (Applied Biosystems) using the SYBR Green PCR Core Reagents (Applied Biosystems). Reaction mixtures were incubated for 10 min at 95°C, followed by 40 cycles of 15 s at 95°C, 1 min at 60°C, and finally, 15 s at 95°C, 1 min at 60°C, and 15 s at 95°C. The gene expression was corrected by the ribosomal protein S18 gene (*rps18*) content in each sample using the comparative cycle threshold method, Ct method ( $2^{-\Delta\Delta Ct}$ ). The gilthead seabream-specific primers used are listed in **Table 1**. In all cases, samples were analyzed in triplicate.

## Immunofluorescence and Flow Cytometry

The percentage of Zap70-, IgM-, and Pax5-positive cells was determined by flow cytometry (Figure S1A in Supplementary Material). In brief, cytoplasmic Zap70 and Pax5, and surface and total IgM, were detected in aliquots of  $0.5 \times 10^6$  head kidney leukocytes at 1, 9, and 23 dpi/dpt and of spleen leukocytes at 23 dpi/dpt ( $n = 6$  fish/treatment/time of sampling) from untreated and EE<sub>2</sub>-treated fish (vaccinated or not). The leukocytes were washed with PBS containing 2% fetal calf serum (FCS; Thermo Fisher Scientific) and 0.05% sodium azide (FACS buffer). Cells were fixed with 4% paraformaldehyde for 15 min at room temperature. After three rinses, cells were incubated in ice-cold PBS containing 1% BSA and saponin (Thermo Fisher Scientific) at 4°C to permeabilize the plasma membrane. Cells were then stained with 0 and 2 µg/ml (0, 1:100) of the 99F2 rabbit monoclonal Ab to Zap70 (Cell Signaling) (43) or mouse monoclonal Ab specific to seabream IgM (Aquatic Diagnostics) (44), in PBS containing 2% FCS for 30 min at 4°C. After washing, cells were incubated with a 1:1,000 dilution of a phycoerythrin-conjugated antirabbit or -mouse Ig Ab, respectively, for 30 min at 4°C, and washed again twice. In other experiments, cells were also stained with 2 µg/ml (1:100) of the D19F8 rabbit monoclonal Ab to Pax5 conjugated with Alexa 488 (Cell Signaling) (43) in PBS containing 2% FCS for 1 h at room temperature. After washing, cells were analyzed in a flow cytometer (BD Biosciences).

## Proliferation Assay

To assess the proliferative activity of head kidney and spleen Zap70<sup>+</sup> and IgM<sup>+</sup> cells at 1, 9, and 23 dpi/dpt ( $n = 6$  fish/treatment/time of sampling), 5-ethynyl-2'-deoxyuridine (EdU) (Thermo Fisher Scientific) was i.p. injected 2 h before sampling. Then, aliquots of  $0.5 \times 10^6$  head kidney and spleen cells were used to determine the percentage of double positive cells, i.e., Zap70<sup>+</sup>/EdU<sup>+</sup> and IgM<sup>+</sup>/EdU<sup>+</sup>. EdU<sup>+</sup> cells were detected by fluorescent-azide coupling reaction with EdU according to the manufacturer's protocol (Click-iT; Thermo Fisher Scientific). Zap70<sup>+</sup> and IgM<sup>+</sup> were labeled as described earlier and analyzed by flow cytometry (Figure S1 in Supplementary Material). The percentage of positive cells is given on head kidney R2 region (FSC<sup>low</sup>/SSC<sup>low</sup>), which includes macrophages, lymphocytes, and hematopoietic precursor cells and excludes acidophilic granulocytes (R1: FSC<sup>high</sup>/SSC<sup>high</sup>) (44).

**TABLE 1** | Gene accession numbers and primer sequences used for expression analysis.

Gene	Accession no.	Name	Sequence (5'/30')
<i>actb</i>	X89920	F3	ATCGTGGGGCGCCCCAGGCACC
		R3	CTCCTTAATGTACGCACGATTTTC
<i>il1b</i>	AJ277166	F2	GGGCTGAACAACAGCACTCTC
		R3	TTAAGACTCTCCACCCTCCA
<i>rps18</i>	AM490061	F	AGGGTGTGGCAGACGTTAC
		R	CTTCTGCCTGTTGAGGAACC

The gene symbols followed the Zebrafish Nomenclature Guidelines ([http://zfin.org/zf\\_info/nomen.html](http://zfin.org/zf_info/nomen.html)). All primers were used for real-time PCR, except *actb* primers that were used for conventional PCR.

## Determination of Serum IgM Titer

Total IgM titers and those specific to KLH, lysozyme (unrelated antigen), *Shewanella putrefaciens* strain Pdp11 (a skin microbiota bacterium from gilthead seabream) (45), or *Vibrio anguillarum* strain R82 (a fish pathogen) were determined in serum by ELISA (Aquatic Diagnostics), following the manufacturer's instructions. In brief, serial dilutions of pooled serum samples ( $n = 6$  fish/treatment/time of sampling) were added to flat-bottomed 96-well plate precoated with KLH (1 µg/well), chicken egg white lysozyme (1 µg/well; Sigma-Aldrich), and *S. putrefaciens* or *V. anguillarum* ( $10^6$  bacteria/well), followed by a monoclonal Ab specific to gilthead seabream IgM (1:100) and then an antimouse IgG peroxidase Ab (1:1,000) (Sigma-Aldrich). Finally, the chromogen TMB was added and the absorbance was read at 450 nm using an FLUOstart luminometer (BGM; LabTechnologies). As negative controls, the serum or the primary Ab was omitted.

## Bactericidal Activity

*Vibrio anguillarum* strain R82 was grown in tryptic soy agar plates at 25°C. Fresh single colonies were diluted in 5 ml of tryptic soy broth, cultured for 16 h at 25°C on an orbital incubator at 300 rpm, and adjusted to  $10^7$  bacteria/ml. Aliquots of 50 µl containing  $10^6$  bacteria in HBSS with Ca<sup>2+</sup>/Mg<sup>2+</sup> were placed in flat-bottomed 96-well plates and incubated for 2 h with 50 µl of pool gilthead seabream serum samples diluted 1/10 in PBS. Moreover, several controls were introduced: blank, without bacteria; positive control, with 10 µg/ml gentamicin (0% growth or 100% bactericidal activity); negative control, without serum (100% growth or 0% bactericidal activity); and de complemented serum control (pre-treated at 50°C, 20 min) to assess complement-mediated killing. The bactericidal activity of serum samples was determined using a 2,3-bis(2-methoxy-4-nitro-5-sulphophenyl)-2H-tetrazolium-5-carboxanilide inner salt (XTT) assay where the yellow tetrazolium salt XTT is reduced to a colored formazan dye by dehydrogenase enzymes in metabolically active cells (46). The bactericidal activity was expressed as the percentage of bacterial growth inhibition.

## Statistical Analysis

Data were analyzed by the analysis of variance. An unpaired Student's *t*-test was applied to determine differences between two groups. The critical value for statistical significance was taken as  $p < 0.05$ . The asterisks \*, \*\*, and \*\*\* refer to  $p < 0.05$ ,  $p < 0.01$ , and  $p < 0.001$ , respectively. All statistical analyses were carried out using the GraphPad Prism 5 program.

## RESULTS


### EE<sub>2</sub> Increases Vtg Serum Levels But Hardly Affects *il1b* Expression in Head Kidney

The survival of gilthead seabream specimens was 100% during the trial (data not shown). As a control for estrogenic endocrine disruption, serum Vtg levels were analyzed. The results showed that the dietary intake of EE<sub>2</sub> significantly increased serum Vtg



levels, as we have previously described for hepatic *Vtg* transcript levels in adult and juvenile gilthead seabream fish treated with EE<sub>2</sub> (29–31, 47, 48). *Vtg* levels peaked at the end of the treatment time and then gradually decreased, although they were still

altered at 23 dpi (Figure 1B). Furthermore, when the expression of the gene coding for *il1b* was analyzed in head kidney at 1, 9, and 23 dpi to assess the activation of innate immunity in response to the immunization, increased *il1b* transcript levels were only



**FIGURE 2** | 17 $\alpha$ -ethinyloestradiol (EE<sub>2</sub>) exposure modulates the abundance and proliferation of T lymphocytes. The percentage of Zap70<sup>+</sup> cells (A,B,D) and that of proliferating Zap70<sup>+</sup> cells (i.e., EdU<sup>+</sup> cells) (C,E) were determined in head kidney leukocytes from untreated and EE<sub>2</sub> (5  $\mu$ g/g food)-treated fish (vaccinated or not) 1 (A), 9 (B,C), and 23 (D,E) days after injection and posttreatment (dpi, dpt) by flow cytometry. The sample size was *n* = 6 fish/group/time of sampling. The mean for each group of specimens is shown as a horizontal line. The percentage is given on head kidney R2 region, which contains macrophages, lymphocytes, and precursor cells, excluding acidophilic granulocytes (49). The asterisks denote statistically significant differences after a Student's *t*-test between the indicated groups. \**p* < 0.05.


evident 1 and 9 dpi as expected (Figure 1C). However, dietary EE<sub>2</sub> failed to significantly affect *il1b* mRNA levels (Figure 1C).

## EE<sub>2</sub> Affects the Abundance and Proliferation of T Lymphocytes

Both the dietary intake of EE<sub>2</sub> and immunization led to an increase in the percentage of T lymphocytes (i.e., Zap70<sup>+</sup> cells) in head kidney at 1 dpi (Figure 2A), whereas no statistically significant differences were observed at 9 and 23 dpi with any of the treatments (Figures 2B,D). Although the percentage of proliferating T lymphocytes (i.e., Zap70<sup>+</sup>/Edu<sup>+</sup> cells) in head kidney was unaffected by the treatments at 9 dpi (Figure 2C), EE<sub>2</sub> slightly increased the percentage of proliferating T lymphocytes in the head kidney at 23 dpi (Figure 2E).

## EE<sub>2</sub> Alters IgM<sup>+</sup> B-Lymphocyte Populations

Although the percentage of IgM<sup>+</sup> B lymphocytes in the head kidney was slightly higher in EE<sub>2</sub>/immunized fish than in the control at 9 dpi/dpt, it was lower in all treated group than in the control at 23 dpi/dpt (Figures 3A,C,F). Interestingly, although the percentage of IgM<sup>+</sup>/Pax5<sup>-</sup> B lymphocytes, i.e., plasma cells, was unaltered at 1 and 23 dpi/dpt, it was higher at 9 dpi/dpt in EE<sub>2</sub>/non-immunized fish but not in immunized fish treated with EE<sub>2</sub> (Figures 3B,D,G). In addition, the dietary intake of EE<sub>2</sub> and immunization resulted in a decreased proliferation of IgM<sup>+</sup> B lymphocytes at 23 dpi/dpt (Figures 3E,H). Neither immunization nor EE<sub>2</sub> exposure was seen to modulate IgM<sup>+</sup> B-lymphocyte abundance, proliferation, or differentiation in the spleen at 23 dpi/dpt (data not shown).



## EE<sub>2</sub> Induces the Production of Natural Neutralizing Antibodies


First, the KLH-specific IgM titer was analyzed by ELISA in serum from untreated and EE<sub>2</sub>-treated fish, both control and immunized, at 1, 9, and 23 dpi/dpt. Unexpectedly, the dietary intake of EE<sub>2</sub> significantly increased the KLH-specific IgM titer at 1 dpi/dpt (Figure 4A). However, immunization failed to elicit a specific IgM response at any of the times analyzed (Figures 4A–C), presumably reflecting the poor primary Ab response to KLH.

These results prompted us to analyze the IgM titer to an unrelated antigen, lysozyme, in the serum of non-immunized fish exposed to EE<sub>2</sub> by different routes (in this case dietary exposure and bath immersion). Strikingly, EE<sub>2</sub> significantly increased lysozyme-specific IgM titers during treatment (40 dt) (Figure 4D) and posttreatment (1 and 9 dpt) (Figures 4E,F), but the effect did not last until 23 dpt (Figure 4G). Similar results were obtained in fish bath-exposed to EE<sub>2</sub> for approximately 2 months (Figure 4H). The induction of natural antibodies by the administration of EE<sub>2</sub> was further confirmed by determining total serum IgM levels (Figures 4I–M). Furthermore,


EE<sub>2</sub>-induced natural IgM antibodies were not only polyreactive but also neutralizing, since they reacted against the commensal *S. putrefaciens* (Figures 5A–C) and the pathogen *V. anguillarum* (Figures 5D–F), inducing the death of the latter (Figures 5G,H). In all cases, the natural antibodies show low affinity, since high dilution of serum had to be used to detect them (Figures 4–6).

## GPER1 Signaling Promotes Natural Ab Production

As previous studies demonstrated the key role of estrogen signaling through GPER1 in the regulation of innate and adaptive immunities in gilthead seabream, we next evaluated the effect of GPER1 activation on natural Ab production using the specific agonist G1. The results showed that G1-treated fish had higher lysozyme-specific (Figure 6A) and total IgM (Figure 6B) serum titers and higher bactericidal activity against *V. anguillarum* at 30 dt than the untreated controls (Figure 6E). Moreover, the effects of EE<sub>2</sub> and G1 on natural Ab production in fish treated for 110 days were similar (Figures 6C,D).



**FIGURE 4** | 17 $\alpha$ -Ethinylestradiol (EE<sub>2</sub>) exposure induces the production of natural antibodies. Keyhole limpet hemocyanin (KLH)-specific (A–C), lysozyme-specific (D–G), and total (I–L) IgM titers were determined in the serum by ELISA 1 (A,E,I), 9 (B,F,K), and 23 (C,G,L) dpi (1, 9, and 23 dpt, respectively) and 40 dt (D,I) in untreated and EE<sub>2</sub> (5  $\mu$ g/g food)-treated fish (vaccinated or not). In addition, lysozyme-specific (H) and total (M) IgM titers were determined in the serum of specimens bath exposure for 58 days to 0, 5, and 50 ng/l EE<sub>2</sub>. The data represent the mean  $\pm$  SEM of absorbance value of the pooled sera of six individual fish. The sample size was = 6 fish/group/time of sampling. The asterisks denote statistically significant differences after Student's *t*-test between the untreated and EE<sub>2</sub>-treated fish (vaccinated or not), at each time point. \**p* < 0.05 and \*\**p* < 0.01.




## DISCUSSION

The fact that estrogens have a key as modulator role of mammals and fish immune systems is well known (4, 29–31, 47). However, the effect of estrogens on fish lymphocytes, and in particular, on the production of natural antibodies, is unknown. In this study, we report for the first time the capacity of EE<sub>2</sub> to modulate lymphocyte populations and natural Ab production in teleost

fish. We have seen that the dietary intake of EE<sub>2</sub> increased the abundance and proliferation of T lymphocytes in the head kidney of unvaccinated gilthead seabream specimens. In contrast, a reduction in the number of circulating lymphocytes was observed in fathead minnow exposed to potent estrogenic effluents (50), whereas E<sub>2</sub> bath exposure altered thymus development in European seabass (51). Unfortunately, the functional consequences of these observations were not investigated.





As regards, IgM<sup>+</sup> B lymphocytes, the dietary intake of EE<sub>2</sub> did not affect their abundance in the two main lymphomyeloid organs, namely the head kidney and the spleen, confirming previous results (29, 30). In contrast, another EDC, tamoxifen, was able to increase the number of IgM<sup>+</sup> B lymphocytes in vaccinated fish (29, 30), suggesting that each EDC may have specific effects in fish adaptive immunity. Thus, it was observed in the present study that dietary EE<sub>2</sub> exposure transiently increased the abundance of plasma cells, i.e., IgM<sup>+</sup>/Pax5<sup>-</sup> cells. This suggest that EE<sub>2</sub> is able to modulate B-cell lineage commitment, as Pax5 is (i) expressed from the pro-B cell through mature and activated B-cell stages, (ii) downregulated during terminal differentiation, and (iii) absent at the plasma cell stage in both mammals (52) and fish (53).

The most interesting observation in this study is that EE<sub>2</sub>, following dietary or bath exposure, induces the production of low affinity, polyreactive natural antibodies. Although we have previously described several long-lasting effects of dietary EE<sub>2</sub> in specific Ab production (29, 30), the modulation of natural Ab

production by dietary EE<sub>2</sub> disappears by 23 dpt. Therefore, the stimulation of natural antibodies by EE<sub>2</sub> coincides with its effect on IgM<sup>+</sup> B-lymphocyte differentiation. Moreover, it was observed that the sera from EE<sub>2</sub>-treated fish manifested an enhanced bactericidal activity, suggesting that natural antibodies are able to effectively neutralize fish pathogens. In mammals, the primary roles of natural antibodies are to confine infection locally, to enhance the IgG response, to play a protective role in autoimmunity, and to take part in homeostasis and clearing of cell debris (54). Natural antibodies are mostly IgM type and are secreted by the long-lived, self-renewing B1 subset of B cells, which are generated during fetal or neonatal development (55, 56). Teleost B cells resemble mammalian B1 cells since both of them show phagocytic and microbicidal activities (57). Although the relevance of natural antibodies in fish is largely unknown, their existence in both cartilaginous and bony fish has been reported (34–36). More recently, the existence of natural Ab in cod has also been shown even though this species is a poor Ab responder (32) and

lacks CD4 and MHCII (58). In addition, the opsonization and neutralizing ability of natural antibodies of rainbow trout against *A. salmonicida* has been demonstrated (37–39).

Although estrogens regulate Ab production, including the production of autoantibodies (19, 59, 60), their impact on natural Ab production is largely unknown. It has been reported that chronic estrogen administration (31 weeks) to mice resulted in increased IgM- and IgG-producing cells (61). However, the mechanisms involved and whether this effect results in increased natural antibodies titers are unknown. Therefore, to the best of our knowledge, this is the first study showing that estrogens regulate natural Ab production in vertebrates. In addition, this effect is mediated through GPER1 signaling, since the GPER1-specific agonist induces a similar natural Ab response to EE<sub>2</sub>. Our results pave the way for future studies aimed at shedding light on the relevance of estrogens in the protection of fish by natural antibodies, the signaling mechanism involved, including classical nuclear ERs, the impact of natural antibodies and estrogens in the crosstalk between natural antibodies, both IgM and IgT, and microbiota, and the therapeutic potential of estrogen administration to protect fish against infections in aquaculture.

## ETHICS STATEMENT

The experiments described were approved by the Consejería de Agua, Agricultura y Medio Ambiente of the Región de Murcia, Spain (approval number A13160507).

## REFERENCES

- Bouman A, Heineman MJ, Faas MM. Sex hormones and the immune response in humans. *Hum Reprod Update* (2005) 11:411–23. doi:10.1093/humupd/dmi008
- Cutolo M, Sulli A, Capellino S, Villaggio B, Montagna P, Seriolo B, et al. Sex hormones influence on the immune system: basic and clinical aspects in autoimmunity. *Lupus* (2004) 13:635–8. doi:10.1191/0961203304lu1094oa
- Karpuzoglu E, Zouali M. The multi-faceted influences of estrogen on lymphocytes: toward novel immuno-interventions strategies for autoimmunity management. *Clin Rev Allergy Immunol* (2011) 40:16–26. doi:10.1007/s12016-009-8188-0
- Straub RH. The complex role of estrogens in inflammation. *Endocr Rev* (2007) 28:521–74. doi:10.1210/er.2007-0001
- Khan D, Ansar Ahmed S. The immune system is a natural target for estrogen action: opposing effects of estrogen in two prototypical autoimmune diseases. *Front Immunol* (2015) 6:635. doi:10.3389/fimmu.2015.00635
- Bernardi AI, Andersson A, Grahne L, Nurkka-Karlsson M, Ohlsson C, Carlsten H, et al. Effects of lasofoxifene and bazedoxifene on B cell development and function. *Immun Inflamm Dis* (2014) 2:214–25. doi:10.1002/iid3.37
- Jones BG, Penkert RR, Xu B, Fan Y, Neale G, Gearhart PJ, et al. Binding of estrogen receptors to switch sites and regulatory elements in the immunoglobulin heavy chain locus of activated B cells suggests a direct influence of estrogen on antibody expression. *Mol Immunol* (2016) 77:97–102. doi:10.1016/j.molimm.2016.07.015
- Schneider AE, Karpati E, Schusztzer K, Toth EA, Kiss E, Kulcsar M, et al. A dynamic network of estrogen receptors in murine lymphocytes: fine-tuning the immune response. *J Leukoc Biol* (2014) 96:857–72. doi:10.1189/jlb.2A0214-080RR
- Deroo BJ, Korach KS. Estrogen receptors and human disease. *J Clin Invest* (2006) 116:561–70. doi:10.1172/JCI27987
- Filardo EJ, Quinn JA, Bland KI, Frackelton AR Jr. Estrogen-induced activation of Erk-1 and Erk-2 requires the G protein-coupled receptor homolog, GPR30,

## AUTHOR CONTRIBUTIONS

AG-A and VM conceived and designed the study; MR performed the research; MR, IC, NG-G, MA, JM, AG-A, and VM analyzed the data; and MR, IC, and VM wrote the manuscript with minor contribution from other authors.

## ACKNOWLEDGMENTS

We thank I. Fuentes for her excellent technical assistance and Profs. A. E. Toranzo and M. A. Moriñigo for the *V. anguillarum* and *S. putrefasciens* strains, respectively.

## FUNDING

This work was funded with support from the Spanish Ministerio de Economía y Competitividad (MINECO) (research grants BIO2014-52655-R to VM and AGL2014-53167-C3-1-R to AG-A and Ph.D. fellowship to MR), all cofunded with Fondos Europeos de Desarrollo Regional/European Regional Development Funds, and by the Fundación Séneca, CARM (19883/GERM/15).

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at <http://journal.frontiersin.org/article/10.3389/fimmu.2017.00736/full#supplementary-material>.

- and occurs via trans-activation of the epidermal growth factor receptor through release of HB-EGF. *Mol Endocrinol* (2000) 14:1649–60. doi:10.1210/mend.14.10.0532
- Thomas P, Pang Y, Filardo EJ, Dong J. Identity of an estrogen membrane receptor coupled to a G protein in human breast cancer cells. *Endocrinology* (2005) 146:624–32. doi:10.1210/en.2004-1064
  - Filardo E, Quinn J, Pang Y, Graeber C, Shaw S, Dong J, et al. Activation of the novel estrogen receptor G protein-coupled receptor 30 (GPR30) at the plasma membrane. *Endocrinology* (2007) 148:3236–45. doi:10.1210/en.2006-1605
  - Filardo EJ, Quinn JA, Frackelton AR Jr, Bland KI. Estrogen action via the G protein-coupled receptor, GPR30: stimulation of adenylyl cyclase and cAMP-mediated attenuation of the epidermal growth factor receptor-to-MAPK signaling axis. *Mol Endocrinol* (2002) 16:70–84. doi:10.1210/mend.16.1.0758
  - Funakoshi T, Yanai A, Shinoda K, Kawano MM, Mizukami Y. G protein-coupled receptor 30 is an estrogen receptor in the plasma membrane. *Biochem Biophys Res Commun* (2006) 346:904–10. doi:10.1016/j.bbrc.2006.05.191
  - Prossnitz ER, Barton M. Signaling, physiological functions and clinical relevance of the G protein-coupled estrogen receptor GPER1. *Prostaglandins Other Lipid Mediat* (2009) 89:89–97. doi:10.1016/j.prostaglandins.2009.05.001
  - Prossnitz ER, Maggiolini M. Mechanisms of estrogen signaling and gene expression via GPR30. *Mol Cell Endocrinol* (2009) 308:32–8. doi:10.1016/j.mce.2009.03.026
  - WHO/IPCS. *Global Assessment of the State of the Science of Endocrine Disruptors*. Geneva: World Health Organization/International Program on Chemical Safety (2002).
  - Chighizola C, Meroni PL. The role of environmental estrogens and autoimmunity. *Autoimmun Rev* (2012) 11:A493–501. doi:10.1016/j.autrev.2011.11.027
  - Inadera H. The immune system as a target for environmental chemicals: xenoestrogens and other compounds. *Toxicol Lett* (2006) 164:191–206. doi:10.1016/j.toxlet.2006.03.006
  - Erlandsson MC, Jonsson CA, Lindberg MK, Ohlsson C, Carlsten H. Raloxifene- and estradiol-mediated effects on uterus, bone and B

- lymphocytes in mice. *J Endocrinol* (2002) 175:319–27. doi:10.1677/joe.0.1750319
21. Prossnitz ER, Hathaway HJ. What have we learned about GPER1 function in physiology and disease from knockout mice? *J Steroid Biochem Mol Biol* (2015) 153:114–26. doi:10.1016/j.jsbmb.2015.06.014
  22. Aris AA, Shamsuddin AS, Praveena MS. Occurrence of 17 $\alpha$ -ethynylestradiol (EE<sub>2</sub>) in the environment and effect on exposed biota: a review. *Environ Int* (2014) 69:104–19. doi:10.1016/j.envint.2014.04.011
  23. Johnson AC, Williams RJ. A model to estimate influent and effluent concentrations of estradiol, estrone, and ethynylestradiol at sewage treatment works. *Environ Sci Technol* (2004) 38:3649–58. doi:10.1021/es035342u
  24. Desbrow C, Routledge EJ, Brighty G, JP S, Waldock M. Identification of estrogenic chemicals in STW effluent. Chemical fractionation and in vitro biological screening. *Environ Sci Technol* (1988) 32:1549–58. doi:10.1021/es9707973
  25. Matozzo V, Marin MG. Can 17-beta estradiol induce vitellogenin-like proteins in the clam *Tapes philippinarum*? *Environ Toxicol Pharmacol* (2008) 26:38–44. doi:10.1016/j.etap.2008.01.001
  26. Rocha MJ, Cruzeiro C, Reis M, Pardal MA, Rocha E. Pollution by oestrogenic endocrine disruptors and beta-sitosterol in a south-western European river (Mira, Portugal). *Environ Monit Assess* (2016) 188:240. doi:10.1007/s10661-016-5236-0
  27. Cabas I, Rodenas MC, Abellán E, Meseguer J, Mulero V, García-Ayala A. Estrogen signaling through the G protein-coupled estrogen receptor regulates granulocyte activation in fish. *J Immunol* (2013) 191:4628–39. doi:10.4049/jimmunol.1301613
  28. Liarte S, Chaves-Pozo E, Abellán E, Meseguer J, Mulero V, García-Ayala A. 17beta-estradiol regulates gilthead seabream professional phagocyte responses through macrophage activation. *Dev Comp Immunol* (2011) 35:19–27. doi:10.1016/j.dci.2010.07.007
  29. Rodenas MC, Cabas I, Abellán E, Meseguer J, Mulero V, García-Ayala A. Tamoxifen persistently disrupts the humoral adaptive immune response of gilthead seabream (*Sparus aurata* L.). *Dev Comp Immunol* (2015) 53:283–92. doi:10.1016/j.dci.2015.06.014
  30. Rodenas MC, Cabas I, García-Alcázar A, Meseguer J, Mulero V, García-Ayala A. Selective estrogen receptor modulators differentially alter the immune response of gilthead seabream juveniles. *Fish Shellfish Immunol* (2016) 52:189–97. doi:10.1016/j.fsi.2016.03.041
  31. Cabas I, Liarte S, García-Alcázar A, Meseguer J, Mulero V, García-Ayala A. 17alpha-ethynylestradiol alters the immune response of the teleost gilthead seabream (*Sparus aurata* L.) both in vivo and in vitro. *Dev Comp Immunol* (2012) 36:547–56. doi:10.1016/j.dci.2011.09.011
  32. Magnadottir B, Gudmundsdottir S, Gudmundsdottir BK, Helgason S. Natural antibodies of cod (*Gadus morhua* L.): specificity, activity and affinity. *Comp Biochem Physiol B Biochem Mol Biol* (2009) 154:309–16. doi:10.1016/j.cbpb.2009.07.005
  33. Panda S, Ding JL. Natural antibodies bridge innate and adaptive immunity. *J Immunol* (2015) 194:13–20. doi:10.4049/jimmunol.1400844
  34. Bisset KA. Natural antibodies in the blood serum of fresh-water fish. *J Hyg* (1948) 46:267. doi:10.1017/S0022172400036408
  35. González R, Charlemagne J, Mahana W, Avrameas S. Specificity of natural serum antibodies present in phylogenetically distinct fish species. *Immunology* (1988) 63:31–6.
  36. Vilain C, Wetzel MC, Du Pasquier L, Charlemagne J. Structural and functional analysis of spontaneous anti-nitrophenyl antibodies in three cyprinid fish species: carp (*Cyprinus carpio*), goldfish (*Carassius auratus*) and tench (*Tinca tinca*). *Dev Comp Immunol* (1984) 8:611–22. doi:10.1016/0145-305X(84)90093-4
  37. Michel C, González R, Bonjour E, Avrameas S. A concurrent increasing of natural antibodies and enhancement of resistance to furunculosis in rainbow trout. *Ann Rech Vet* (1990) 21:211–8.
  38. Nakamura A, Osonoi T, Terauchi Y. Relationship between urinary sodium excretion and pioglitazone-induced edema. *J Diabetes Investig* (2010) 1:208–11. doi:10.1111/j.2040-1124.2010.00046.x
  39. Sinyakov MS, Dror M, Zhevelev HM, Margel S, Avtalion RR. Natural antibodies and their significance in active immunization and protection against a defined pathogen in fish. *Vaccine* (2002) 20:3668–74. doi:10.1016/S0264-410X(02)00379-1
  40. Shved N, Berishvili G, D'Cotta H, Baroiller JF, Segner H, Eppler E, et al. Ethynylestradiol differentially interferes with IGF-I in liver and extrahepatic sites during development of male and female bony fish. *J Endocrinol* (2007) 195:513–23. doi:10.1677/JOE-07-0295
  41. Chaves-Pozo E, Mulero V, Meseguer J, García Ayala A. Professional phagocytic granulocytes of the bony fish gilthead seabream display functional adaptation to testicular microenvironment. *J Leukoc Biol* (2005) 78:345–51. doi:10.1189/jlb.0205120
  42. Chaves-Pozo E, Pelegrin P, Mulero V, Meseguer J, García Ayala A. A role for acidophilic granulocytes in the testis of the gilthead seabream (*Sparus aurata* L., Teleostei). *J Endocrinol* (2003) 179:165–74. doi:10.1677/joe.0.1790165
  43. Piazzon MC, Savelkoul HS, Pietretti D, Wiegertjes GF, Forlenza M. Carp Il10 has anti-inflammatory activities on phagocytes, promotes proliferation of memory T cells, and regulates B cell differentiation and antibody secretion. *J Immunol* (2015) 194:187–99. doi:10.4049/jimmunol.1402093
  44. Sepulcre MP, López-Muñoz A, Angosto D, García-Alcázar A, Meseguer J, Mulero V. TLR agonists extend the functional lifespan of professional phagocytic granulocytes in the bony fish gilthead seabream and direct precursor differentiation towards the production of granulocytes. *Mol Immunol* (2011) 48:846–59. doi:10.1016/j.molimm.2010.12.010
  45. Salinas I, Díaz-Rosales P, Cuesta A, Meseguer J, Chabrilón M, Moriño MA, et al. Effect of heat-inactivated fish and non-fish derived probiotics on the innate immune parameters of a teleost fish (*Sparus aurata* L.). *Vet Immunol Immunopathol* (2006) 111:279–86. doi:10.1016/j.vetimm.2006.01.020
  46. Sepulcre MP, López-Castejón G, Meseguer J, Mulero V. The activation of gilthead seabream professional phagocytes by different PAMPs underlines the behavioural diversity of the main innate immune cells of bony fish. *Mol Immunol* (2007) 44:2009–16. doi:10.1016/j.molimm.2006.09.022
  47. Cabas I, Chaves-Pozo E, García-Alcázar A, Meseguer J, Mulero V, García-Ayala A. The effect of 17alpha-ethynylestradiol on steroidogenesis and gonadal cytokine gene expression is related to the reproductive stage in marine hermaphrodite fish. *Mar Drugs* (2013) 11:4973–92. doi:10.3390/md11124973
  48. García-Hernández MP, Rodenas MC, Cabas I, García-Alcázar A, Chaves-Pozo E, García-Ayala A. Tamoxifen disrupts the reproductive process in gilthead seabream males and modulates the effects promoted by 17alpha-ethynylestradiol. *Comp Biochem Physiol C Toxicol Pharmacol* (2016) 179:94–106. doi:10.1016/j.cbpc.2015.09.005
  49. Esteban MA, Mulero V, Muñoz J, Meseguer J. Methodological aspects of assessing phagocytosis of *Vibrio anguillarum* by leucocytes of gilthead seabream (*Sparus aurata* L.) by flow cytometry and electron microscopy. *Cell Tissue Res* (1998) 293:133–41. doi:10.1007/s004410051105
  50. Filby AL, Neuparth T, Thorpe KL, Owen R, Galloway TS, Tyler CR. Health impacts of estrogens in the environment, considering complex mixture effects. *Environ Health Perspect* (2007) 115:1704–10. doi:10.1289/ehp.10443
  51. Seemann F, Knigge T, Oliver S, Monsinjon T. Exogenous 17 $\beta$ -oestradiol (E2) modifies thymus growth and regionalization in European sea bass *Dicentrarchus labrax*. *J Fish Biol* (2006) 86:1186–98. doi:10.1111/jfb.12626
  52. Cobaleda C, Schebesta A, Delogu A, Busslinger M. Pax5: the guardian of B cell identity and function. *Nat Immunol* (2007) 8:463–70. doi:10.1038/ni1454
  53. Zwollo P. Dissecting teleost B cell differentiation using transcription factors. *Dev Comp Immunol* (2011) 35:898–905. doi:10.1016/j.dci.2011.01.009
  54. Gronwall C, Vas J, Silverman GJ. Protective roles of natural IgM antibodies. *Front Immunol* (2012) 3:66. doi:10.3389/fimmu.2012.00066
  55. Boes M. Role of natural and immune IgM antibodies in immune responses. *Mol Immunol* (2000) 37:1141–9. doi:10.1016/S0161-5890(01)00025-6
  56. Herzenberg LA. B-1 cells: the lineage question revisited. *Immunol Rev* (2000) 175:9–22. doi:10.1111/j.1600-065X.2000.imr017520.x
  57. Li J, Barreda DR, Zhang Y, Boshra H, Gelman AE, LaPatra S, et al. B lymphocytes from early vertebrates have potent phagocytic and microbicidal abilities. *Nat Immunol* (2006) 7:1116–24. doi:10.1038/ni1389
  58. Star B, Nederbragt AJ, Jentoft S, Grimholt U, Malmstrøm M, Gregers TF, et al. The genome sequence of Atlantic cod reveals a unique immune system. *Nature* (2011) 477:207–10. doi:10.1038/nature10342

59. Murakami M, Tsubata T, Okamoto M, Shimizu A, Kumagai S, Imura H, et al. Antigen-induced apoptotic death of Ly-1 B cells responsible for autoimmune disease in transgenic mice. *Nature* (1992) 357:77–80. doi:10.1038/357077a0
60. Verthelyi D, Ahmed SA. 17 beta-estradiol, but not 5 alpha-dihydrotestosterone, augments antibodies to double-stranded deoxyribonucleic acid in nonautoimmune C57BL/6J mice. *Endocrinology* (1994) 135:2615–22. doi:10.1210/endo.135.6.7988450
61. Nilsson N, Carlsten H. Estrogen induces suppression of natural killer cell cytotoxicity and augmentation of polyclonal B cell activation. *Cell Immunol* (1995) 158:131–9. doi:10.1006/cimm.1994.1262

**Conflict of Interest Statement:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2017 Rodenas, Cabas, Gómez-González, Arizcun, Meseguer, Mulero and García-Ayala. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.