




Food and Agriculture  
Organization of the  
United Nations



# A guide to the eggs and larvae of 100 common Western Mediterranean Sea bony fish species

Cover photographs (clockwise from top left):  
Larva of *Trisopterus luscus*, © J.M. Rodríguez  
Yolk-sac-larva and egg of *Xiphias gladius*, © A. García

# **A guide to the eggs and larvae of 100 common Western Mediterranean Sea bony fish species**

by

**J.M. Rodríguez  
F. Alemany  
A. García**

The designations employed and the presentation of material in this information product do not imply the expression of any opinion whatsoever on the part of the Food and Agriculture Organization of the United Nations (FAO) concerning the legal or development status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. The mention of specific companies or products of manufacturers, whether or not these have been patented, does not imply that these have been endorsed or recommended by FAO in preference to others of a similar nature that are not mentioned.

The views expressed in this information product are those of the author(s) and do not necessarily reflect the views or policies of FAO.

ISBN 978-92-5-109895-0

© FAO, 2017

FAO encourages the use, reproduction and dissemination of material in this information product. Except where otherwise indicated, material may be copied, downloaded and printed for private study, research and teaching purposes, or for use in non-commercial products or services, provided that appropriate acknowledgement of FAO as the source and copyright holder is given and that FAO's endorsement of users' views, products or services is not implied in any way.

All requests for translation and adaptation rights, and for resale and other commercial use rights should be made via [www.fao.org/contact-us/licence-request](http://www.fao.org/contact-us/licence-request) or addressed to [copyright@fao.org](mailto:copyright@fao.org).  
FAO information products are available on the FAO website ([www.fao.org/publications](http://www.fao.org/publications)) and can be purchased through [publications-sales@fao.org](mailto:publications-sales@fao.org)

## PREPARATION OF THIS DOCUMENT

The study of ichthyoplankton is crucial in understanding the dynamics of fish populations. The knowledge of the distribution and abundance of fish eggs and larvae, obtained through plankton surveys can, among other things, help scientists improve the understanding of the spatial distribution of fisheries resources and the processes affecting the fluctuations in recruitment; two important underpinnings of fish stocks assessment and management. The FAO Regional Project “CopeMed Phase II”: Coordination to Support Fisheries Management in the Western and Central Mediterranean” has among its priorities the strengthening of research capabilities in its member countries in support of sustainable management of fishery resources. In February 2016, the Project organized a course on ichthyoplankton identification at the Spanish Institute of Oceanography, in Fuengirola (Málaga). For this course, the lecturer and co-author of this guide, J.M. Rodríguez, prepared a collection of technical sheets for several larval fish species with their corresponding description and illustrations. During the course, the scarcity of identification guides for the Mediterranean Sea was noticed. Participants asked the organizers to expand the number of sheets of the species used at the course and requested that CopeMed II publish a FAO Guide including the most common species occurring in the Mediterranean Sea. The present guide is the outcome of this process. It presents the egg and larval descriptions of 100 species of fishes belonging to 55 families, which are most likely to be present in the plankton samples collected in the continental shelf and oceanic waters of the Western Mediterranean Sea. The document is not meant to be a complete guide for ichthyoplankton of the Western Mediterranean Sea but a comprehensive tool with which to support students and researchers new to the field of ichthyoplankton identification.

**Supervision:** Pilar Hernandez (CopeMed Fishery Expert, FAO CopeMed, Malaga, Spain).

**Technical editor:** Edoardo Mostarda (FAO, Rome).

**Cover illustration:** Abla Kadiri.

**Rodriguez, J.M., Alemany, F. and Garcia A.** 2017. *A guide to the eggs and larvae of 100 common Western Mediterranean Sea bony fish species*. FAO, Rome, Italy, 256 pp.

### ABSTRACT

This guide presents the egg and larval descriptions of 100 species of fishes belonging to 55 families, which are most likely to be present in plankton samples collected in the continental shelf and oceanic waters of the Western Mediterranean Sea. The guide is structured in two parts. The first introductory part describes the different applications of ichthyoplankton studies in fisheries research, the main sampling strategies, methods and gears, the early life history of fishes and how to identify them. A brief historical account of ichthyoplankton identification studies in the Mediterranean Sea is also provided. The second part of the guide contains the identification sheets of the species. Each species sheet includes the following information: illustration of the adult fish and information on its habitat and spawning season; description of the main features useful towards identifying the egg, yolk sac and larval stage of each fish species; illustrations and (when available) photos of the different life stages.



## CONTENTS

PREPARATION OF THIS DOCUMENT .....	iii
ACKNOWLEDGEMENTS .....	x
<b>1. INTRODUCTION</b> .....	<b>1</b>
1.1 What is ichthyoplankton? .....	1
1.2 Why study ichthyoplankton? .....	2
1.3 Applications of ichthyoplankton studies in fisheries research .....	4
1.4 Ichthyoplankton sampling strategies, methods and gears .....	5
1.5 Sample representativeness .....	8
1.6 Ichthyoplankton sorting .....	9
1.7 Taxonomic identification of fish eggs and larvae .....	9
1.8 The importance of a proper taxonomic identification of ichthyoplankton .....	13
1.9 History of studies on ichthyoplankton identification in the Mediterranean Sea .....	14
1.10 Glossary .....	15
1.11 Illustrations of representative larvae of bony fish families included in this guide .....	18
<b>2. IDENTIFICATION SHEETS</b> .....	<b>23</b>
OPHICHTHIDAE .....	24
<i>Dalophis imberbis</i> .....	24
CONGRIDAE .....	26
<i>Conger conger</i> .....	26
CLUPEIDAE .....	28
<i>Sardina pilchardus</i> .....	28
<i>Sardinella aurita</i> .....	30
ENGRAULIDAE .....	32
<i>Engraulis encrasicolus</i> .....	32
ARGENTINIDAE .....	34
<i>Argentina sphyraena</i> .....	34
MICROSTOMATIDAE .....	36
<i>Microstoma microstoma</i> .....	36
<i>Nansenia oblita</i> .....	38
GONOSTOMATIDAE .....	40
<i>Cyclothone braueri</i> .....	40
STERNOPTYCHIDAE .....	42
<i>Argyropelecus hemigymnus</i> .....	42
<i>Maurolicus muelleri</i> .....	44
PHOSICHTHYIDAE .....	46
<i>Vinciguerria attenuata</i> .....	46

STOMIIDAE .....	48
<i>Stomias boa</i> .....	48
SYNODONTIDAE .....	50
<i>Synodus saurus</i> .....	50
PARALEPIDIDAE .....	52
<i>Lestidiops jayakari</i> .....	52
<i>Paralepis coregonoides</i> .....	54
EVERMANNELLIDAE .....	56
<i>Evermannella balbo</i> .....	56
MYCTOPHIDAE .....	58
<i>Benthoosema glaciale</i> .....	58
<i>Ceratoscopelus maderensis</i> .....	60
<i>Diaphus holti</i> .....	62
<i>Hygophum</i> .....	64
<i>Lampanyctus crocodilus</i> .....	66
<i>Lampanyctus pusillus</i> .....	68
<i>Lobianchia dofleini</i> .....	70
<i>Myctophum punctatum</i> .....	72
<i>Notoscopelus elongatus</i> .....	74
<i>Symbolophorus veranyi</i> .....	76
GADIDAE .....	78
<i>Gadiculus argenteus</i> .....	78
<i>Micromesistius poutassou</i> .....	80
<i>Trisopterus luscus</i> .....	82
LOTIDAE .....	84
<i>Gaidropsarus biscayensis</i> .....	84
<i>Gaidropsarus mediterraneus</i> .....	86
MERLUCCIIDAE .....	88
<i>Merluccius merluccius</i> .....	88
OPHIDIIDAE .....	90
<i>Ophidion barbatum</i> .....	90
CARAPIDAE .....	92
<i>Echiodon dentatus</i> .....	92
LOPHIIDAE .....	94
<i>Lophius piscatorius</i> .....	94
GOBIESOCIDAE .....	96
<i>Diplecogaster bimaculata</i> .....	96
BELONIDAE .....	98
<i>Belone belone</i> .....	98
ZEIDAE .....	100
<i>Zeus faber</i> .....	100



---

CENTRISCIDAE .....	102
<i>Macroramphosus scolopax</i> .....	102
SEBASTIDAE .....	104
<i>Helicolenus dactylopterus</i> .....	104
SCORPAENIDAE .....	106
<i>Scorpaena porcus</i> .....	106
TRIGLIDAE .....	108
<i>Eutrigla gurnardus</i> .....	108
<i>Lepidotrigla cavillone</i> .....	110
MORONIDAE .....	112
<i>Dicentrarchus labrax</i> .....	112
SERRANIDAE .....	114
<i>Anthias anthias</i> .....	114
<i>Serranus cabrilla</i> .....	116
<i>Serranus hepatus</i> .....	118
APOGONIDAE .....	120
<i>Apogon imberbis</i> .....	120
CARANGIDAE .....	122
<i>Trachurus mediterraneus</i> .....	122
<i>Trachurus trachurus</i> .....	124
CORYPHAENIDAE .....	126
<i>Coryphaena hippurus</i> .....	126
BRAMIDAE .....	128
<i>Brama brama</i> .....	128
SPARIDAE .....	130
<i>Boops boops</i> .....	130
<i>Diplodus sargus</i> .....	132
<i>Pagellus bogaraveo</i> .....	134
<i>Pagrus pagrus</i> .....	136
<i>Spondyliosoma cantharus</i> .....	138
CENTRACANTHIDAE .....	140
<i>Spicara smaris</i> .....	140
MULLIDAE .....	142
<i>Mullus barbatus</i> .....	142
<i>Mullus surmuletus</i> .....	144
CEPOLIDAE .....	146
<i>Cepola macrophthalma</i> .....	146
MUGILIDAE .....	148
<i>Mugil cephalus</i> .....	148
POMACENTRIDAE .....	150
<i>Chromis chromis</i> .....	150

LABRIDAE . . . . .	152
<i>Coris julis</i> . . . . .	152
<i>Ctenolabrus rupestris</i> . . . . .	154
<i>Labrus mixtus</i> . . . . .	156
<i>Symphodus melops</i> . . . . .	158
AMMODYTIDAE . . . . .	160
<i>Gymnammodytes cicereus</i> . . . . .	160
TRACHINIDAE . . . . .	162
<i>Echiichthys vipera</i> . . . . .	162
<i>Trachinus draco</i> . . . . .	164
URANOSCOPIDAE . . . . .	166
<i>Uranoscopus scaber</i> . . . . .	166
BLENNIIDAE . . . . .	168
<i>Blennius ocellaris</i> . . . . .	168
<i>Lipophrys pholis</i> . . . . .	170
<i>Parablennius pilicornis</i> . . . . .	172
<i>Parablennius gattorugine</i> . . . . .	174
CALLIONYMIDAE . . . . .	176
<i>Callionymus</i> . . . . .	176
GOBIIDAE . . . . .	178
<i>Crystallogobius linearis</i> . . . . .	178
<i>Gobius paganellus</i> . . . . .	180
<i>Lebetus guilleti</i> . . . . .	182
<i>Pomatoschistus microps</i> . . . . .	184
TRICHIURIDAE . . . . .	186
<i>Lepidopus caudatus</i> . . . . .	186
SCOMBRIDAE . . . . .	188
<i>Auxis rochei</i> . . . . .	188
<i>Sarda sarda</i> . . . . .	190
<i>Scomber colias</i> . . . . .	192
<i>Scomber scombrus</i> . . . . .	194
<i>Thunnus alalunga</i> . . . . .	196
<i>Thunnus thynnus</i> . . . . .	198
XIPHIIDAE . . . . .	200
<i>Xiphias gladius</i> . . . . .	200
CAPROIDAE . . . . .	202
<i>Capros aper</i> . . . . .	202
SCOPHTHALMIDAE . . . . .	204
<i>Lepidorhombus boscii</i> . . . . .	204
<i>Scophthalmus rhombus</i> . . . . .	206
<i>Zeugopterus regius</i> . . . . .	208

---

BOTHIDAE .....	210
<i>Arnoglossus laterna</i> .....	210
<i>Bothus podas</i> .....	212
PLEURONECTIDAE .....	214
<i>Platichthys flesus</i> .....	214
SOLEIDAE .....	216
<i>Buglossidium luteum</i> .....	216
<i>Microchirus variegatus</i> .....	218
<i>Solea solea</i> .....	220
CYNOGLOSSIDAE .....	222
<i>Symphurus nigrescens</i> .....	222
<b>3. LITERATURE CITED</b> .....	<b>225</b>
<b>4. INDEX</b> .....	<b>238</b>

## ACKNOWLEDGEMENTS

This guide was prepared in support of the training courses for identification of early life stages of fish implemented by the FAO through the Mediterranean Regional Project “CopeMed II” (Coordination to Support Fisheries Management in the Western and Central Mediterranean). CopeMed II is funded by the Government of Spain, represented by the Secretaría General de Pesca (M<sup>o</sup> de Agricultura, Pesca, Alimentación y Medio Ambiente, MAPAMA), and the European Union, represented by the European Commission (EC).

The authors would like to thank Pilar Hernández, CopeMed II Fishery Expert, who supervised the whole elaboration process and Marcelo Vasconcellos, FAO Fishery Resources Officer, who provided overall support and feedback. Many thanks to the two anonymous reviewers who, with their precious suggestions and comments, have significantly contributed to the improvement of the original manuscript.

The authors also wish to express their gratitude to the publishers and authors that gave their permission to use figures from their publication, such as:

Academic Press, Inc., New York  
American Fisheries Society  
American Society of Ichthyologists and Herpetologists  
Dr. Bill Richards  
Fishery Bulletin  
Ichthyological Bulletin of the J.L.B. Smith Institute of Ichthyology  
International Council for the Exploration of the Sea (ICES) · Conseil International pour l'Exploration de la Mer (CIEM)  
Investigación Pesquera  
Investigaciones Marinas Valparaiso  
Journal of Northwest Atlantic Fisheries Science  
Journal of Plankton Research  
Journal of the Marine Biological Association of the United Kingdom  
Scientia Marina  
South African Journal of Aquatic Science

All photos included in the introduction and species sheets were provided by the authors.

Finally, J.M. Rodríguez would like to thank his wife Rosa Vázquez for revising the manual, correcting many typographical errors and spelling mistakes, and improving the quality of several illustrations.

## 1. INTRODUCTION

### 1.1 What is ichthyoplankton?

The early life stages (eggs and larvae) of marine fishes are referred to as ichthyoplankton (Fig. 1). Some species of marine fishes produce demersal eggs that lay on the sea bottom, or are even deposited in nests. Demersal eggs of a number of species are under parental care, extended in some exceptional cases, such as in sea horses, to the larval stage. However, most marine fish species spawn pelagic eggs. Larvae hatched from pelagic and from most demersal eggs are pelagic. Due to their null (eggs) or limited (larvae) swimming abilities, they are both integrated within the plankton community as part of the meroplankton (organisms that only spend part of their life cycle within the plankton community). Ichthyoplankton is mainly found in the upper layers (from 200 m depth to the surface) of the water column, where it is subject to spatial dispersion, mainly by marine currents. The planktonic stage of fishes usually lasts from a couple of weeks to a few months (Victor, 1986; Brothers *et al.*, 1983).

During the larval stage, larvae develop specializations for the planktonic life and other important changes occur. For example, most fishes increase their weight by five orders of magnitude throughout their life and three of them occur during the planktonic stage (Werner and Gilliam, 1984; Houde, 1987; Miller *et al.*, 1988). The end of the larval stage is marked by a metamorphosis or transformation process, more or less abrupt depending on the species, during which the larva becomes a juvenile




**Figure 1.** Ichthyoplankton sample

that is morphologically similar to the adult and has the meristic characters of the species. Juveniles of pelagic species are integrated in the nekton community and those of most demersal species migrate directly to the bottom, after the metamorphosis. However, in several demersal species, where the transformation is prolonged, there is an intermediate stage (called the pre–juvenile stage) where the fish develops specializations that are distinct from both larvae and juveniles. These pelagic pre–juveniles eventually transform into demersal juveniles (Kendall *et al.*, 1984). Pelagic pre–juveniles of demersal species and the early juveniles of the pelagic ones maintain an intense relationship, mainly trophic, with the plankton community, but because of their behaviour and swimming abilities, they cannot be considered ichthyoplankton, they become part of the micronekton.

## 1.2 Why study ichthyoplankton?

There are several reasons for studying ichthyoplankton. For example, information regarding the distribution and abundance of fish eggs and larvae can provide clues to spawning locations and environmental requirements of important fish species. Moreover, the knowledge of ichthyoplankton is necessary because, as one of the components of the pelagic food web (Raymont, 1983), it can represent an important link between smaller planktonic and larger nektonic organisms. In addition, as most fishes inhabit the upper water column during their early life history and have limited avoidance capability, ichthyoplankton surveys provide a relatively low-cost efficient means to monitor marine fish populations and communities (Koslow and Wright, 2016). Finally, survival of fish larvae may directly influence future abundances of adult fish stocks. The latter has been and it still is the most important reason for studying ichthyoplankton, as most processes determining the recruitment strength and the spatial distribution of fish populations occur during the planktonic stage of fishes, resulting in important interannual fluctuations in fish stock biomasses. Such fluctuations have been known for centuries, but they started to worry scientists and fishery managers only at the end of the nineteenth century (Petersen, 1894; Garstang, 1900). Initially, it was thought that fishing pressure itself or fish migrations were responsible for such fluctuations, but at the beginning of the twentieth century, Hjort (1914, 1926), after analysing the causes of the successful herring year-class of 1904, suggested that the variability in “year-class success” was determined during the early life stages of fishes. He proposed two hypotheses to explain those fish stock fluctuations resulting from interannual recruitment variability, the *Critical Period Hypothesis* and the *Aberrant Drift Hypothesis*, or larval transport to unfavourable areas (due to interannual variability in ocean circulation), from which juveniles could not recruit to the exploited stock.

According to the *Critical Period Hypothesis*, the strength of a year-class is determined shortly after yolk-sac absorption, at the beginning of exogenous feeding, when larvae must find suitable prey and in sufficient amounts (Fig. 2). Failing to find adequate feeding conditions would lead to massive larval mortality, in a short period of time. Hjort’s hypotheses laid the groundwork for future research on recruitment variability, mainly focused on the *Critical Period Hypothesis* (Cowan and Shaw, 2002; Houde, 2008), fostering the development of “recruitment fisheries oceanography” as a multidisciplinary science to investigate the spectrum of oceanographic processes controlling recruitment (Kendall and Duker, 1998).



**Figure 2.** Illustration showing the Hjort’s Critical Period Hypothesis (1914, 1926). From Houde (2008).

Cushing (1974, 1990) merged both Hjort’s hypotheses into the *Match/Mismatch Hypothesis*. He hypothesized that a fixed time of spawning coupled with a variable time of plankton blooms generates a variability in larval fish survival and, hence, variable recruitment. In this hypothesis,


food limitation at any time of the larval period, rather than mortality being concentrated at a specific larval stage, could be a major contributor to the recruitment variability. In addition, abiotic factors that regulate water–column mixing and the timing and intensity of seasonal production cycles may be involved.

Lasker (1981), in the *Stable Ocean Hypothesis*, proposed that the occurrence and frequency of calm periods in upwelling ecosystems (named Lasker events) lead to stratification of the water column and, hence, the aggregation of fish larvae and their prey, supporting high larval feeding, survival and recruitment. A further extension of Lasker's *Stable Ocean Hypothesis* was the *Optimal Environmental Window (OEW) Hypothesis* by Cury and Roy (1989). These authors hypothesized that, in upwelling ecosystems, the relationship between annual recruitment of small pelagic fish species (sardines and anchovies) and upwelling intensity is dome shaped, confirming the existence of an optimal environmental window for recruitment, with recruitment most successful under moderate (roughly  $5\text{--}6\text{ m}^*\text{s}^{-1}$ ) wind stress that controls both ichthyoplankton advective losses and food availability for fish larvae. On the contrary, weak winds, if prolonged, disrupt the upwelling process and the renewal of nutrients in the surface layer, inducing nutrient limitation, which reduces primary productivity (Huntsman and Barber, 1977) and food availability for fish larvae. Finally, strong winds generate strong turbulence that also limits primary productivity (Huntsman and Barber, 1977) and increases the offshore transport of fish egg and larvae (Bakun and Parrish, 1982)

Based on the Hjort's *Aberrant Drift Hypothesis*, Iles and Sinclair (1982) and Sinclair (1988) proposed the *Larval Retention or Member/Vagrant Hypothesis*. In this hypothesis, larval retention, not levels of available prey, is critical in the recruitment process.

For tropical reef fish species, Sale (1978, 1991) proposed the *Lottery Hypothesis*, which states that recruitment in tropical reef fishes is strongly dependent on delivery potential to the reef of settlers and that post–settlement processes could control recruitment levels. However, nowadays, both pre– and post–settlements processes are acknowledged to influence recruitment (Jones, 1991; Doherty, 2002).

The above hypotheses consider starvation and physical processes as the main drivers of recruitment variability; however, there is strong evidence that predation is the main source of ichthyoplankton mortality and, consequently, the main factor influencing recruitment (Hunter, 1981; Bailey and Houde, 1989). Mortality from nutritional deficiencies and predation may not act independently. Slow–growing and small fish larvae remain relatively vulnerable to predation for longer periods of time. This is the foundation of the *Stage–Duration Hypothesis*, which implies that large size (“bigger is better”) and fast growth improves survival potential (Houde, 1987; Anderson, 1988; Houde, 2008).




**Figure 3.** Schematic diagram of different processes acting on the early stages of a marine fish (eggs, yolk–sac larvae, larvae and juveniles). From Houde (1987).

In summary, although Hjort focused on starvation of first-feeding larvae as the main driver of larval mortality, he already envisaged that recruitment variability was the result of complex, interacting factors, stating that “*the simultaneous investigation of meteorology, hydrography and biology seems the only way to a deeper understanding of the conditions in which the destiny of the spawned ova is being decided*” (Hjort, 1926). In this line, nowadays it is widely agreed that recruitment variability is the outcome of complex trophodynamic and physical processes, operating on different temporal and spatial scales and throughout the pre-recruit life stages of fishes (Fig. 3). That is to say, recruitment success is not determined during a particular ontogenetic stage and it depends on the species, populations and environmental conditions (Houde, 2008). In this way, the knowledge of the ecology of the early life stages of fishes is crucial to understanding the population dynamics of fish stocks, but also, the functioning of marine ecosystems.

New discoveries, technological advances and new analytical techniques, developed in the last decades, have allowed for significant advances in the knowledge of the causes of mortality of the early life fish stages. One of the major findings was the discovery by Pannella (1971) that growth rings in fish otoliths are deposited daily. Brothers *et al.* (1976) confirmed the presence of daily growth rings in otoliths of fish larvae (Fig. 4). This finding allowed for the estimation of growth and mortality rates during the larval stage, and for the cohort contributions to recruitment (Methot, 1983; Miller *et al.*, 1988; Houde, 1997). The analysis of the otolith's microstructure also allows for assessing the environmental influence on larval survival, dating, with high precision, early life events and relating them to environmental conditions.

Other indicators useful for determining the nutritional status and recent growth rate in fish larvae are the RNA/DNA ratio (Buckley, 1979, 1980, 1984) and the nitrogen (N) and carbon (C) stable



**Figure 4.** Otolith of a sardine (*Sardina pilchardus*) larva

isotope analysis (SIA). The latter is used to assess the trophic position and C flow to consumers in food webs (Minagawa and Wada, 1984, Peterson and Fry 1987, Post, 2002). Specifically,  $^{15}\text{N}$  provides an estimate of the trophic level, and  $^{13}\text{C}$  can be used to assess the sources of C for an organism when the isotopic nature of the sources is different, since C isotope ratios undergo small changes within the food web (Peterson and Fry, 1987, France and Peters, 1997). The SIA analysis has also been used recently to assess the trophic ecology of fish larvae (e.g. Laiz-Carrión *et al.*, 2013, Laiz-Carrión *et al.*, 2015). Finally, advances in telemetry have allowed synoptic sampling of environmental variables over large areas and advances in computer power the development of biophysical models, simulating larval drift and survival under real or hypothetical environmental scenarios.

### 1.3 Applications of ichthyoplankton studies in fisheries research

One of the main objectives of ichthyoplankton field studies is the direct assessment of the number or biomass of exploitable populations (Heath, 1992). This is based on the fact that for some fish



populations there may be a relationship between the number of eggs and/or larvae in a given area and the number or biomass of spawning adult fish, and this can be used to estimate the size of the adult population (Heath, 1992). There are six direct methods that use ichthyoplankton production for estimating the abundance of fish stocks, three based on eggs and three on larvae, with the latter used for demersal spawners (Stratoudakis *et al.*, 2006). The underlying principle of the ichthyoplankton-based methods is that the abundance of an early life stage of a fish species can be used to estimate the reproductive outcome of the population over a time period (Stratoudakis *et al.*, 2006).

Of the three methods based on eggs production, the *Daily Egg Production Method* (DEPM), the most widely used, was developed for small pelagic fishes (such as anchovies and sardines) that exhibit indeterminate fecundity (Lasker, 1985; Parker, 1980). Fecundity is indeterminate when the potential annual fecundity of a female is not fixed prior to the onset of spawning and unyolked oocytes continue to be matured and spawned during the spawning season (Hunter *et al.*, 1992). The advantage of the DEPM is that it only requires a single survey to estimate the daily egg production at sea and the daily specific fecundity at the peak of the spawning period (Parker, 1980). In addition, the application of this method provides information about the reproductive biology and reproductive behaviour of a population and about the distribution, mortality and development of the early life fish stages, which are of particular importance for studying the underlying mechanisms of recruitment (Murua *et al.*, 2010). The other two methods based on eggs, the *Annual Egg Production Method* (AEPM) (Lockwood *et al.*, 1981) and the *Daily Fecundity Reduction Method* (DFRM) (Lo *et al.*, 1992) were developed for species with determinate fecundity, for which the potential annual fecundity becomes fixed prior to the onset of spawning (Hunter *et al.*, 1992). Where egg production methods are impractical, a larval census has been employed to obtain a relative index of the spawning biomass (Heath, 1992).

Larval survey data can be used to obtain estimates of recruitment. The estimation or forecasting of recruitment strength, under variable environmental scenarios, is essential for the proper management of fish stocks, particularly in a context of relatively rapid climate change. Moreover, the implementation of the *Ecosystem Approach to Fisheries* (EAF), where fisheries management recognizes the full range of interactions within a marine ecosystem (Katsanevakis *et al.*, 2011), has led to an increase of studies aimed at understanding recruitment and its underlying processes. In addition, a holistic understanding of ecosystems, of the environmental influence on fish population, especially on their highly vulnerable early life stages, is required to implement the EAF. This knowledge is also essential towards implementing the *Ecosystem-Based Management of Fisheries* (EBMF), in which the order of management priorities is inverse to the customary norm, addressing the status of ecosystems, rather than that of the fishery resources (Pikitch *et al.*, 2004; Cogan *et al.*, 2009).


Furthermore, ichthyoplankton studies can provide a lot of information on the ecology and structure of fish populations, in a cheaper and simpler way than information obtained studying juvenile or adult populations. Thus, a single collection of plankton hauls can give information about most fish species spawning in a given area, both pelagic and demersal, whereas sampling of adults would require larger vessels and a variety of sampling gears and methodologies. Ichthyoplankton surveys can also be used for the prospection of new resources, determining the timing and location of spawning areas and their variations or estimating the relative abundance of different stocks and monitoring their abundance trends over time.

#### **1.4 Ichthyoplankton sampling strategies, methods and gears**

Any study on ichthyoplankton communities is based on two pillars: an adequate sampling strategy for the specific objectives to be achieved and a correct taxonomic identification of the target taxa. There is a great variety of sampling strategies, from large-scale surveys to intensive sampling of a single patch of larvae tracked by means of a Lagrangian buoy (a free-floating buoy that moves with a parcel of water). The sampling methodology to be followed and the sampling gear to be used will depend on the objectives of the study, on the target species and/or the early life stages to be sampled.

It is out of the scope of this guide to explain in detail all possible sampling strategies or describe the different types of ichthyoplankton sampling devices and sampling methods. However, it is worth illustrating those that are most frequently used and the potential sources of bias affecting sample representativeness.

As already explained, many ichthyoplankton studies are aimed at estimating the biomass of adult fish stocks, either by using eggs or larvae. When using eggs for e.g. applying the *Daily Egg Production Method* (DEPM), the gear employed is the CalVET net in vertical hauls, and the sampling strategy



**Figure 5.** Left, CalVET net and right, Bongo net

consists of sampling stations arranged in a regular grid, with a short distance between stations, covering the whole spawning area of the fish stock to be studied. When the objective is to estimate the biomass of an adult fish stock using larvae, to study the horizontal distribution of fish eggs and larvae of specific species or, in general, the ichthyoplankton community of an area, the most adequate sampling gear to be used is the Bongo net (Fig. 5). Hauls are oblique and the sampling strategy consists of sampling a regular grid of stations. Ichthyoplankton hauls must cover the depth distribution range of the target species or, if the goal is to sample the entire ichthyoplankton community, at least the upper 200 m of the water column, where most fish eggs and larvae are concentrated.

When the aim of ichthyoplankton studies is to investigate the vertical distribution of fish eggs and larvae and/or the vertical (diel and/or ontogenetic) migrations of fish larvae, a multiple net system, capable of sampling successive water layers during the same tow, is required (Fig. 6). Hauls are also oblique.



**Figure 6.** Multiple net systems. Left, MultiNet sampler and right, MOCNESS net

Neuston nets are used to sample the very most surface water layer (Fig. 7). Large micronekton nets, fitted with meshes of the size of 1 mm or larger, may also be necessary for effectively sampling larger larvae. Light traps are used for capturing advanced post-larvae in pre-settlement stages.

Ichthyoplankton samples can also be obtained with continuous sampling systems, such as the *Continuous Underway Fish Egg Sampler* (CUFES), capable of collecting plankton as the research vessel is sailing. It consists essentially of a pump that draws seawater at a given depth and sends it through a plankton collector, where plankton is retained (Fig. 8). The collector is changed and plankton collected after a fixed period of time. While CUFES is running, a data logger records the date, time, and position of each sample, as well as other environmental data from the ship's sensors.



Figure 7. Neuston net

In the last decades, new and more sophisticated sampling systems are being implemented. In situ ichthyoplankton imaging systems or plankton acoustics, despite not providing biological samples for other types of analyses, can produce impressive amounts of data on larval abundance or biomass, over large areas in relatively short periods of time (Cowen and Guigand, 2008; Garcia-Seoane *et al.*, 2016). Any sampling aimed at obtaining quantitative data on ichthyoplankton requires knowing the exact volume of water filtered by the net during each haul. This is done by using flowmeters attached to the mouth of the net. Flowmeters measure the length of the haul path, which, multiplied by the surface




Figure 8. Continuous underway fish egg sampler (CUFES) (<http://cufes.ucsd.edu/text/descr.htm>)

of the mouth of the net, gives the volume of water filtered by the net during the haul. This allows standardization of fish eggs and larvae captured in each haul to number of individuals per cubic meter, or densities. Ichthyoplankton counts can be also standardised to number of individuals beneath a unit of sea surface (square meter), by multiplying densities by the maximum depth of tow in meters (Smith and Richardson, 1977). Haul depth can be determined by using any device equipped with pressure sensors, such as a conductivity–temperature depth instrument (CTD) of small size or simple depth gauges, such as those used for scuba diving, attached to the sampling gear structure. If flowmeters or depth gauges are not available, the volume of water filtered can be estimated from the duration of the tow and the vessel's speed, which allows for estimating the distance covered by the net's path. Haul depth can be estimated by multiplying the maximum length of wire led out by the cosine of the wire angle at the moment when the maximum amount of wire has been let out. Finally, to get a reliable estimation of ichthyoplankton densities or abundances, it is necessary to homogeneously sample the water column over the whole haul depth range.

Once the haul is finished, plankton must be concentrated into the cod ends by gently washing the nets (Fig. 5). After that, plankton is transferred into jars and fixed for long-term preservation. The most commonly used preservatives are a 5% solution of buffered formalin and seawater, or ethanol at different concentrations. Formalin is more adequate for preserving samples to be used in taxonomic studies, since the egg and larval morphology is better preserved, but presents the problem that formalin is carcinogenic. Ethanol dehydrates both eggs and larvae. Eggs lose their shape, and larvae tend to curl making them difficult to measure. However, it has the advantage that specimens can be used for genetic and for larval growth studies. Other fixatives are liquid nitrogen, used to preserve larvae in biochemical and daily growth analyses, and RNAlater, when it is necessary to stabilize and protect cellular RNA. For some specific studies, larvae must be quickly sorted and identified on board, prior to fixation.

### 1.5 Sample representativeness

There are several papers dealing with problems associated to ichthyoplankton sampling (Hempel, 1974; Smith and Richardson, 1977; Lasker, 1981; Heath, 1992). Specifically, difficulties in obtaining representative samples of ichthyoplankton communities derive from the spatial and temporal heterogeneity in ichthyoplankton distribution and from the interaction of fish eggs and larvae with sampling gears (Ahlstrom *et al.*, 1973). Fish eggs and larvae are not distributed randomly, but they tend to be concentrated in patches that become part of the plankton during specific periods of the year. This is the result of the distribution of adult fishes, their spawning strategies, hydrodynamic processes, egg and larval density, and larval behaviour, which varies throughout ontogeny. The only way to deal with non-random distribution of ichthyoplankton is through an adequate sampling design.

The main sources of sampling bias, related to the direct interactions between fish eggs and larvae with the plankton nets, are escapement and avoidance. Escapement is the passage of smaller ichthyoplankton organisms through the meshes of the net. It may be active or passive and it is a function of the size, shape and behaviour of the organisms in relation to mesh size (Vannucci, 1968). Active escapement is the process by which fish larvae caught in the net may squeeze through the meshes. This involves behavioural patterns that vary with species and with the developmental stage (Vannucci, 1968). Passive escapement or extrusion is the forced passage of eggs or larvae through the net's meshes. When the organisms are larger than the meshes, extrusion is aided by the compressibility of the organisms and the flexibility of the meshes (Saville, 1958). This source of sampling bias is easily overcome by using net meshes of size adequate to retain smaller fish eggs and larvae. Considering the fact that most fish eggs have a diameter of over 0.5 mm and that most newly hatched larvae are more than 2.0 mm long, the adequate mesh size for sampling ichthyoplankton is between 300 and 500 microns. However, mesh size should not be too small for avoiding, or at least reducing, mesh clogging produced by plankton. Net clogging reduces the filtration efficiency of the net.

Avoidance is the process by which larger fish larvae avoid capture, swimming out of the path of the approaching net or migrating below the net's maximum sampling depth (Morse, 1989). This is probably the most important source of sampling bias (Clutter and Ankaru, 1968). Evidence of this phenomenon is the observed difference in larval fish abundances between night and day samples (Russell, 1976; Bridger, 1956; Ahlstrom, 1954, 1959). However, it seems that avoidance occurs both day and night (Murphy and Clutter, 1972). Larvae are not only able to see the approaching nets (visual avoidance), but they are capable of detecting the water vibrations produced by the towing wires and also the wave pressure produced by the net ahead of it (Smith and Clutter, 1965; Mahnken and Jossi, 1967; Tranter and Smith, 1968). Thus, to reduce or minimize net avoidance, it is useful to use nets without bridles in front of the net mouth, such as Bongo type nets (McGowan and Brown, 1966).

## 1.6 Ichthyoplankton sorting


For the taxonomic identification of ichthyoplankton or for other analyses, fish eggs and larvae are usually sorted from the plankton samples. The first step in ichthyoplankton sorting is to eliminate formalin from plankton samples. To do this, samples must be sieved through meshes (of smaller sizes than those used for sampling at sea), washed with seawater and placed into jars with seawater. This task must be carried out under a fume hood to avoid breathing the carcinogenic formalin vapours. Then, the sample can be placed in Petri dishes and analysed under a stereoscopic microscope, at low magnification (10x is adequate), to detect and sort fish eggs and larvae. Considering the low abundances of larvae of some species, it is recommendable, for studies on ichthyoplankton communities, to analyse the whole sample. However, for specific studies focusing on eggs or larvae of more abundant species, it is acceptable to analyse 50 or 25% of the sample or even less, depending on the abundance of the target species. Fish eggs and larvae must be handled very carefully, with soft tweezers, thin brushes or pipettes, to separate and keep them in vials with the appropriate preservative medium.

## 1.7 Taxonomic identification of fish eggs and larvae

The taxonomic identification of fish eggs and larvae is not an easy task. It is more difficult than identifying the juvenile and adult stages of fishes. This is due to several reasons. First, because of the small size of fish eggs and larvae, characters useful for their identification can only be visualized under a stereoscopic microscope. Then, in the case of fish larvae, the main problem is that they undergo continuous and, in some cases, dramatic morphological, meristic, morphometric and pigmentary changes throughout their development. This has prevented the building of dichotomous keys, such as those existing for adult fishes. In addition, other characters, such as pigmentation patterns, can present an important geographical and even individual variability. Despite these difficulties, partial dichotomous keys were developed for the Western Mediterranean Sea by Abossouan (1964) and Marinaro (1971). However, these keys only include some developmental stages of a relatively small number of the species present in the area. Therefore, they easily lead to identification dead ends or to misidentifications. To overcome this problem, the use of intelligent software, capable of managing extensive databases and integrating all the available information on larval stages of different fish species, including ecological information, such as the distribution of adults, reproduction season, etc. has been proposed (Froese and Papisassi, 1989). However, the design and implementation of such information systems is not easy, due to the heterogeneity of the available information, or even the lack of information on the early life stages of many marine fish species. Even in well-studied areas, eggs and larvae of many species are still unknown or, in many cases, early fish stages remain undescribed. More recently, molecular techniques are being used to identify fish eggs and larvae, or for validating previous descriptions of these, but they cannot be routinely applied. Because of all these difficulties, the "look-alike" method remains the method usually used to identify fish eggs and larvae. It consists of comparing the individuals with descriptions made by other authors. This elimination process ultimately results in "assigning" an individual to a species. This is also the method followed in this guide.


However, to properly use the “look-alike” method, it is first necessary to compile and analyse all the available information about the ichthyofauna of the study area. That is to say, faunistic lists and all the available information on the spawning strategies, spawning seasons and known distribution of the fish species inhabiting the study area. Another prerequisite for using this method is being able to accurately determine the developmental stage of the individual under analysis and to compare it with the corresponding developmental stage described in the literature.

The nomenclature of the different early developmental stages of fishes varies by author. In this guide, we follow the nomenclature suggested by Kendall *et al.* (1984)(Fig. 9), one of the most widely accepted. They divided the early life–history of fishes into three stages: egg, from spawning to hatching; larva, from hatching to attainment of complete fin–ray counts and beginning of squamation (juvenile); and juvenile, young fish, fundamentally like the adult in the meristic characters (excluding squamation) but smaller and reproductively inactive. Kendall *et al.* (1984) also divided the larval stage into four sub–stages: yolk–sac larva, from hatching to exhausting of yolk reserves; preflexion larva, from yolk exhausting to the beginning of upward flexion of the notochord; flexion larva, that ends when the urostyle is in its final position, at approximately 45° from the notochord axis and post–flexion larva, that ends when metamorphosis begins. Transformation, or metamorphosis is a transitional stage between larva and juvenile during which the young fish loses larval characters and acquires those of the adult (Kendall *et al.*, 1984; Moser, 1996).



**Figure 9.** Early life history stages of *Trachurus symmetricus* from Kendall *et al.* (1984)

Most marine fish eggs encountered in plankton samples are spherical and transparent, with a diameter of about 1.0 mm. Eggs are enclosed by a thin membrane or chorion. There is a space between the chorion and the yolk sac, the perivitelline space. Most fish eggs have oil globules (Fig. 10). The main anatomical, morphological and morphometric features used in fish egg identification are: egg shape (spherical or elliptic); egg size (ranges from 0.5 to more than 5.5 mm in diameter); type of surface membrane (smooth, sculptured, with a single protuberance or filaments); presence of a second internal membrane; type of yolk (homogeneous or segmented); size of the perivitelline space; absence/presence, number, position and colour (for live individuals) of oil globules. When the embryo is well developed,




**Figure 10.** Main anatomic features of a fish egg. Adapted from Munk and Nielsen (2005)

embryonic characters, such as morphological features, pigment patterns and special structures are also used to identify fish eggs (Russell, 1976; Matarese and Sandknop, 1984; Ahlstrom and Moser, 1980). Scanning electron microscopy has proved to be a good tool for the taxonomic identification of fish eggs, but cannot be used in routine analysis (Boehlert, 1984). The identification of fish eggs is usually a more difficult task than larval fish identification. This is reflected by the fact that the number of fish species with the egg stage described is much lower than those with the larval stage described. In the Mediterranean Sea, it is worth mentioning the pioneer identification guide to fish eggs developed by Marinaro (1971), which has been recently revised and extended by Crec'hriou *et al.* (2015).

In most fish species with pelagic eggs, the newly hatched larvae are in general less than 4 mm long (Russell, 1976). The size and state of development at hatching is generally related to yolk–sac diameter. Typically, the body length at hatching is 2.5 to 3.0 times the diameter of the yolk sac (Moser, 1996). In general, yolk–sac larvae hatched from eggs less than 1.5 mm in diameter have an unformed mouth, unpigmented eyes and pectoral–fin buds, while yolk–sac larvae hatched from larger eggs are comparatively well developed, with the mouth formed, eyes pigmented and pectoral fins developed (Moser, 1996). In both cases, the locomotion is aided by a prominent fin–fold that extends from the top of the head, around the caudal region, and ventrally forward to the posterior margin of the yolk sac (Fig. 11).


The identification of yolk–sac larvae is very difficult because some structures, such as fins and most of the specialized larval characters, resulting from the evolutionary adaptation to the plankton realm, are not yet well developed. Consequently, yolk–sac larvae of different species are very similar. The yolk–sac stage is characterized by the migration, coalescence and rearrangement of pigment cells or melanophores (Moser, 1996). Melanophores are amoeboid and capable of migrating from their point of origin in the neural crest to various sites in the larva, to establish the species–specific larval pigmentation pattern, at the end of the yolk–sac stage. Useful characters for the identification of yolk–sac larvae are the shape and relative size of the yolk sac, the presence and the relative position of oil globules in the yolk sac, the position of the anus in relation to the yolk sac and, in some, the species–specific pigmentation pattern.



**Figure 11.** Main anatomic features of a yolk sac larva. From Russell (1976)

The complete utilization of the yolk marks the end of the yolk–sac stage. By this time, most of the organs and the sensory system required to capture prey are functional. The mouth and gut are formed, the anus is open at the margin of the ventral fin–fold, the eyes are pigmented, and the primordial pectoral fins are present. It is now an early larva and during the larval development, the fish gradually acquires the characters of the adult, thus facilitating its identification (Moser, 1996; Russell, 1976). At first, the body is still surrounded by the primordial fin, the urostyle is straight and rudiments of the hypural elements are visible in its ventral side. As the larva grows, the urostyle bends upwards, the hypural elements become defined, caudal rays develop, and the first signs of the formation of the dorsal and anal fins appear as interspinous areas. At this stage, both the meristic characters and pigmentation pattern that are characteristic of the adult of the species, have usually appeared (Russell, 1976). The main anatomical features of a larva are shown in figure 12.

The main characters used in larval fish identification are the body form, the pigmentation pattern and meristic and morphometric characters. The body form allows for separating larvae into several major groups (Russell, 1976). For example, larvae with narrow, elongated bodies (e.g. Families Clupeidae, Engraulidae, Stomiidae, etc.); larvae with laterally compressed bodies (includes all flatfishes, e.g. Families Bothidae, Pleuronectidae, Soleidae, etc.); bodies with the typical fish shape (includes larvae of



**Figure 12.** Main anatomic features of a fish larva


most fish species, e.g. Families Gadidae, Triglidae, Gobiidae); bodies with aberrant shapes (e.g. Family Belonidae) or showing specialized larval characters for the plankton life, such as cranial armatures (e.g. Family Scorpaenidae), elongate fin rays (e.g. Families Carapidae, Lophiidae), stalked eyes (e.g. some Myctophidae species) or early developed and large fins (e.g. Family Trachinidae).

Meristic characters are countable structures appearing in series, such as the number of miomeres, vertebrae or fin rays. They have a high diagnostic value (at least the combination of several counts), because they are species–specific. However, they have the disadvantages that some of them, such as fin rays, are completely formed only in older larvae, which are scarce in plankton samples. Others, such as miomeres or vertebrae, are difficult to visualize, even using staining methods, or other techniques, such as X–rays (Pothoff, 1984; Tucker and Laroche, 1984).



Morphometric characters include the different measurements of a larva. The main measurements of a larva, shown in figure 13, are the total length (TL), or the distance from the tip of the snout to the caudal-fin end; the standard length (SL), or the distance between the tip of the snout and the urostyle end; the preanus length, or the distance between the tip of the snout and the anus; the head length, or the distance from the tip of the snout to the border of the cleitrum, and eye diameter, or the maximum diameter of the eye. When measuring the different body regions for identification purposes, the stage of development of the larvae must be taken into account, since larval growth is allometric.

Fish larvae show a variety of pigmentary cells or chromatophores. Those containing black or brown pigments are known as melanophores; those with yellow pigments, xanthophores and those with red pigments, erythrophores (Russell, 1976). However, in formalin-preserved specimens, only black pigmented cells or melanophores remain. For this reason, the latter are usually the only ones used in ichthyoplankton identification. Melanophores are situated in different parts of the body, defining a species-specific pigmentation pattern. This pattern is one of the chief diagnostic characters used for



**Figure 13.** Body regions and the most important measurements of a fish larva

the identification of the larval stage of fish species. Often, the identification of species with similar larvae is made possible thanks to the presence or absence of a single melanophore or by its position (Russell, 1976). The general appearance of preserved specimens will differ very much according to the degree of expansion or contraction of the melanophores. Moreover, in specimens preserved for a long time, the pigmentation will fade, especially if kept in the light (Russell, 1976). Besides, there may be an intraspecific and geographical variability in pigmentation patterns.

### 1.8 The importance of a proper taxonomic identification of ichthyoplankton

The first step in any ichthyoplankton study is the proper taxonomic classification of eggs and/or larvae since, as Powles and Markle (1984) pointed out, small errors in their identification can result in important misinterpretations about the biology and ecology of fish species. Moreover, there are several examples in the literature showing that wrong identifications of fish eggs or larvae have led to biased stock evaluations and, subsequently, inadequate management measures (Daniel and Graves, 1994; Armstrong *et al.*, 2001; Fox *et al.*, 2005). Unfortunately, these identification errors are probably more frequent and important than desirable. For example, a recent study focusing on the ability of researchers from five different laboratories in Taiwan to identify fish larvae, determined that the average accuracy of identification was 80.1, 41.1 and 13.5 % at the family, genus and species levels, respectively (Ko *et al.*, 2013). Recently, Puncer *et al.* (2015a) revealed that the Atlantic bluefin tuna (*Thunnus thynnus*) larvae have been misidentified in the Mediterranean Sea. These authors demonstrated, through genetic analysis, that more than half of the larvae, submitted by three Mediterranean institutions to a bluefin tuna research project, funded by the International Commission for the Conservation of Atlantic Tunas (ICCAT), were misidentified. Some of these errors are caused by the persistence in literature of wrong descriptions. However, also in species whose larvae are accurately described, the inexperience

or lack of training of the people in charge of the taxonomic identification of ichthyoplankton can lead to massive misidentifications (Puncher *et al.*, 2015b). Because of this, it is of paramount importance to produce identification guides with accurate descriptions, taking advantage of molecular genetic techniques, to validate doubtful identifications, as well as to organize courses given by experienced ichthyoplanktologists and aimed at properly training new generations of technicians and researchers in charge of the taxonomic classification of fish eggs and larvae. This book, based on a course on larval fish taxonomy, supported by the FAO COPEMED regional program, intends to be a step in this direction, for the western Mediterranean region.

### **1.9 History of studies on ichthyoplankton identification in the Mediterranean Sea**

The pioneering works on ichthyoplankton identification in the Mediterranean Sea were those of Raffaele (1888) and Marion (1894). These authors described eggs and yolk sac larvae of several fish species from the Gulf of Naples (Raffaele, 1888) and from the Gulf of Marseille (Marion, 1894). Some years later, Fage (1910) published the results of ichthyoplankton surveys carried out off the city of Nice and in the Gulf of Lion including descriptions of larvae of several fish species. Roule and Angel (1930), summarized the results of the oceanographic surveys supported by the Prince of Monaco, Albert I, carried out between 1895 and 1915. Outstanding papers on ichthyoplankton identification are the reports on the “Danish Oceanographic Expeditions in the Mediterranean and Adjacent Seas” (Dana–Reports) by e.g. Kyle (1913), Jespersen (1915), Ege (1918, 1930), Fage (1918, 1920), Schmidt (1918), Schmidt and Strubberg (1918), Täning (1918, 1923), Jespersen and Täning (1926), Guitel (1920), Ehrebaum (1924), and Schnakenbeck (1931). However, the most important publication on ichthyoplankton identification from the Mediterranean Sea is the monography “Uova, larve e stadi giovanili di Teleostei”, elaborated using material gathered by Lo Bianco and edited by Bertolini *et al.* (1931–1956). This monography compiles the description and illustrations of fish eggs and larvae of many species carried out by several authors. This monography, along with the Dana–Reports, are even nowadays the basic and essential references for any study on ichthyoplankton identification from the Mediterranean Sea. It is also worth mentioning the papers from Aboussouan (1964) and Lee (1966) that include useful keys for the identification of fish eggs and larvae and the paper by Marinaro (1971), which includes a useful key for the identification of fish eggs. In recent years, the PhD Theses of Sabatés (1988) and Alemany (1997) deserve special attention. These include the description and original illustrations of larvae of several fish species. Several illustrations of the Alemany’s PhD Thesis are included in this manual.

## 1.10 Glossary

**Anal fin:** Fin (usually single but double in some gadiforms) located on the ventral margin of the tail, behind the anus.

**Anus:** Orifice and surrounding tissue at end of the gut.

**Bathypelagic:** Pelagic zone between 1 000 and 4 000 m depth. Bathypelagic fish are those inhabiting the bathypelagic region.

**Benthic:** Referring to the sea bottom. Benthic species are those living on or in the bottom (substrate).

**Benthopelagic:** Living and feeding near the bottom as well as in midwater or even near the surface.

**Caudal fin:** Median fin situated at the posterior end of the fish.

**Caudal peduncle:** Narrow part of the tail located between the posterior end of dorsal or anal fin and the base of the caudal fin.

**Chorion/shell:** Outer membrane of a fish egg.

**Choroid tissue:** Mass of vascular tissue on the ventral side of the eye in larvae of some myctophids and other meso- and bathypelagic fishes, usually associated with elliptical eyes.

**Cleithrum:** Vertical bone in the pectoral girdle, at the junction of head and body of the larva.

**Cleithral symphysis:** Ventral junction of the cleithral bones.

**Chromatophore:** Cell containing pigment that reflects the light.

**Continental shelf:** The flattened edge of the continental land mass, between the coast and the continental slope (generally, the continental subtidal zone to a depth of about 200 m).

**Continental slope:** The sloping edge of the continental land mass, generally beginning at depth of around 200 m.

**Demersal:** Living on or near the bottom of the sea.

**Dorsal fin:** Median fin or fins located on the dorsal margin of the body.

**Early life history:** The early life stages of fishes spanning from egg to juvenile.

**Embryo:** Early stage of development, from fertilisation to hatching

**Epipelagic:** The illuminated, uppermost layer of the ocean (from 0 to 200 m depth).

**Eye stalks:** Movable peduncles of varying length bearing the eyes.

**Finfold/primordial fin:** Median fold of skin surrounding the body of young larvae, within which the dorsal, caudal and anal fins are developed.

**Flexion:** Larval stage during which the urostyle bends dorsally.

**Gut:** Alimentary tube and associated organs.

**Gut loop:** Loop, fold, or curve found along the axis of the gut.

**Head length:** Distance from the tip of the snout to the posterior margin of the cleithrum.

**Hindgut:** Posterior part of the gut.

**Homogeneous:** Uniform composition throughout; opposite to segmented in referring to egg yolk.

**Ichthyoplankton:** Zooplankton fraction including eggs and larval stages of fishes.

**Isthmus:** Ventral region of the head separating the gill openings of a fish.

**Juvenile:** A young fish, fundamentally resembling the adult in meristic characters (excluding squamation) but smaller and reproductively inactive.

**Larva:** Early life-history stage of fishes between the egg and juvenile stages.

**Leptocephalus:** The flat, transparent, and often large larvae of fishes in the orders Anguilliformes, Elopiformes and Notacanthiformes, characterized by small heads and prominent teeth

**Melanophore:** A cell containing melanin; a black or brown pigment cell.

- Meristic characters:** Countable structures occurring in series (e.g. myomeres, vertebrae, fin rays).
- Mesopelagic:** Occurring in the open ocean at middle depths, usually between 200 and 1 000 m.
- Metamorphosis:** A marked change in form or structure at the end of the larval stage involving acquisition of adult characters and loss of larval characters; synonymous of transformation.
- Myomeres:** Muscle segments occurring in series, the number is approximately equal to the number of vertebrae in adults.
- Myosepta:** Connecting tissue between adjacent myomeres.
- Nekton:** Motile, marine organisms living in the water column and capable of swimming against currents.
- Neritic:** Pelagic coastal zone extending from the low tide mark to the edge of the continental shelf.
- Neural crest:** Region of the neural ridge of the developing embryo that differentiates into many kinds of tissue and cells, including melanophores.
- Neustonic:** Occurring close to the surface of the ocean.
- Notochord:** Longitudinal flexible cartilaginous rod of cells forming the supporting axis of the body.
- Notochord length (NL):** The distance from the tip of the snout to the posterior tip of the notochord.
- Occipital crest:** A median, bony ridge, usually serrated, located on top of the head, posteriorly.
- Oceanic:** Open sea region beyond the edge of the continental shelf.
- Oil globule:** Spheres of fatty material within the yolk of some fish eggs.
- Opercular:** Relative to the operculum.
- Operculum or Opercle:** The bony plate of the gill cover.
- Ovoviviparous:** Producing eggs that develop within the maternal body.
- Pectoral fin:** Paired lateral (sometimes ventrolateral) fins located behind the head.
- Peduncle:** A narrow part or stalk that connects a structure to the body (e.g. caudal peduncle connecting the caudal fin to body).
- Pelagic:** Free living in the water column, away from the sea bottom.
- Pelvic fins:** Paired fins, usually located on the ventral edge of the body, in the abdominal region.
- Peritoneal:** Region of the body associated with the gut or the membrane of the peritoneum.
- Peritoneum:** The membrane and associated tissue lining the gut cavity.
- Perivitelline space:** Fluid-filled space between the embryo and chorion or shell of an egg.
- Photophores:** Luminous organs on some marine (mostly deep-sea) fish larvae.
- Plankton:** Small free-living organisms, passively floating or weakly swimming that drift with currents.
- Pigmentation:** Deposition of pigment in body tissues.
- Planktonic:** Passively floating, drifting, or weakly swimming with prevailing currents.
- Postanal:** posterior to the anus.
- Postflexion stage:** A stage in the development of larvae after the completion of flexion.
- Preanal/preanus:** Located anterior to the anus; preanal length (synonymous of snout-anus distance) measured from the tip of the snout to the posterior margin of the anus.
- Preflexion stage:** Larval stage before the beginning of the process of flexion.
- Preopercle:** Upper anterior bone of the gill cover.
- Preopercular:** Relative to the preopercle.
- Recruitment:** The amount of fish added to the exploitable stock each year due to growth and/or migration into the fishing area.
- Sculptured:** Egg chorion with ornamentations or surface features of different shapes and textures.

**Segmented:** Particulate or divided; opposite of homogeneous in referring to egg yolk.

**Shell:** The membrane that encloses an egg; generally, equivalent to chorion.

**Shrinkage:** The act or fact of shrinking, to contract or lessen in size.

**Swimbladder:** Sac filled with air or other gases located in the abdominal region, beneath the backbone.

**Snout:** Forward part of the head, anterior to the eye.

**Stalked eye:** Eye situated on a stalk or peduncle.

**Standard length (SL):** The distance from the tip of the snout to the tip of urostyle.

**Stellate melanophore:** Star-like pigment spot.

**Stock (Fish):** A group of individuals in a species occupying a well-defined spatial range independent of other stocks of the same species.

**Supraorbital spine:** Occurring above the eye.

**Tail:** The portion of the body posterior to the anus, the postanal region.

**Telescopic eye:** Elongate, cylindrical eye that protrudes forward or upward within an envelope of skin.

**Tentacle:** Any of various slender, flexible appendages in larvae.

**Total length (TL):** Measurement from the tip of the snout to the most posterior part of a larva, including the caudal finfold or caudal fin.

**Transformation:** The process (synonymous of metamorphosis) at the end of the larval stage, characterized by a marked change in form or structure and involving the acquisition of juvenile or adult characters and the loss of larval characters.

**Trunk:** Portion of the body between the head and the anus.

**Urostyle:** The last vertebral elements in fishes, formed by fusion or loss of several vertebrae.

**Year class:** All of the fish in a stock that were spawned in a particular year, such as all those spawned in 1990. Also referred to as a "cohort".

**Yolk:** Nutritive material of the egg forming a sac-like mass (yolk sac) below the abdominal region of a newly hatched larva.

**Yolk-sac larva:** Newly hatched larva with yolk sac.

**1.11 Illustrations of representative larvae of bony fish families included in this guide**



**Ophichthidae**



**Clupeidae**



**Congridae**



**Engraulidae**



**Microstomatidae**



**Argentinidae**



**Gonostomatidae**



**Phosichthyidae**



**Sternoptychidae**



**Stomiidae**



**Paralepididae**



**Synodontidae**




**Evermannellidae**



**Gadidae**




**Myctophidae**



**Lotidae**



**Ophidiidae**




**Merlucciidae**



**Carapidae**



**Gobiesocidae**



**Lophiidae**




**Belonidae**




**Centriscidae**




**Zeidae**



**Sebastidae**




**Triglidae**




**Scorpaenidae**




**Moronidae**




**Apogonidae**



**Serranidae**



**Carangidae**



**Bramidae**




**Coryphaenidae**




**Sparidae**




**Mullidae**



**Centracanthidae**



**Cepolidae**




**Pomacentridae**



**Mugilidae**




**Labridae**



**Trachinidae**



**Ammodytidae**



**Uranoscopidae**



**Callionymidae**




**Blenniidae**







**Gobiidae**




**Scombridae**




**Trichiuridae**




**Xiphiidae**




**Scophthalmidae**



**Caproidae**




**Bothidae**



**Soleidae**




**Pleuronectidae**



**Cynoglossidae**



## **2. IDENTIFICATION SHEETS**

**OPHICHTHIDAE***Dalophis imberbis* (Delaroche, 1809)**En:** Armless snake eel**Habitat:** Benthic, littoral, burrowing in sand or mud bottoms, between 20 and 100 m depth**Spawning season:** Between August and September**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 2.10–2.40 mm**Chorion:** smooth**Oil globules:** 1–17**Perivitelline space:** large**Yolk:** segmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 5.0 mm SL**Distinctive characters:** body elongate and slender; no teeth on jaws**Number of myomeres:** 159; 77 ahead of anus**Pigmentation:** body unpigmented**LARVAE****Figs. C–F****Distinctive characters:** leptocephalus; body elongate, transparent, laterally compressed; in young larvae, 8 long teeth on each jaw; 5 or 6 gut loops**Preanus length:** decreases with development**Number of myomeres:** 152–155; 68–76 preanal**Pigmentation:** in small larvae, 12 ventral spots, 9 over gut and 3 postanal; lower jaw and urostyle pigmented; larger larvae with up to 5 lateral melanophores located under the notochord**Length at transformation:** unknown**PHOTOS**

Not available


**Literature:** Alemany (1997), Bauchot (1986), D'Ancona (1931d), Sabatés (1988)

**OPHICHTHIDAE**


*Dalophis imberbis* (Delaroche, 1809)



A.




B. 5.0 mm




C. 5.5 mm SL

8 long teeth on each jaw




D. 9.0 mm



E. 16.5 mm SL




F. Detail of head



G. 31.0 mm

A, B, D, F: D'Ancona (1931d); C, E: Alemany (1997)

**CONGRIDAE***Conger conger* (Linnaeus, 1758)**En:** European conger – **Fr:** Congre d'Europe – **Sp:** Congrio común**Habitat:** Benthic, on rocky and sandy bottoms on the shelf, between 0 and 100 m depth**Spawning season:** Summer**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–E**

**Distinctive characters:** leptocephalus; body elongate, transparent, laterally compressed; head small; 6 long teeth on each jaw; pectoral fin apparent in small larvae; origin of dorsal fin quite ahead of anus (predorsal fin distance about 57% SL); single tubular gut; eye slightly oval

**Preanus length:** about 84% SL

**Number of myomeres:** 148–155

**Pigmentation:** small larvae with 7 melanophores over gut and 10 between anus and caudal-fin base; the number of melanophores over gut, from the 14<sup>th</sup> myomere to the anus, increases with development; older larvae show 6 melanophores in ventral side, 3 in dorsal side of postanal region and about 10 located at lateral end of body, over notochord; caudal fin pigmented

**Length at transformation:** about 150.0 mm





**PHOTOS**

Not available


**Literature:** Alemany (1997), Bauchot and Saldanha (1986), D'Ancona (1931b), Sabatés (1988)

**CONGRIDAE**


*Conger conger* (Linnaeus, 1758)



D. Detail of head  
(63.8 mm SL larva)



E. Detail of caudal region  
(63.8 mm SL larva)

**CLUPEIDAE***Sardina pilchardus* (Walbaum, 1792)**En:** European pilchard – **Fr:** Sardine commune – **Sp:** Sardina europea**Habitat:** Pelagic, coastal**Spawning season:** Between September and May**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.30–1.90 mm**Chorion:** smooth**Oil globules:** single, 0.14–0.18 mm in diameter**Perivitelline space:** large**Yolk:** segmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** 3.3–4.0 mm**Distinctive characters:** body elongate and slender, with the typical clupeid shape; oil globule situated at posterior edge of yolk sac**Preanus length:** about 80% SL**Pigmentation:** two parallel rows of small melanophores in dorsal region, extending from head to tail**LARVAE****Figs. C–H****Distinctive characters:** body elongate and slender; head length included at least 6 times in total length; gut differentiated into two sections; dorsal fin located ahead of anus**Preanus length:** about 80% SL**Pigmentation:** melanophores aligned on both sides of body, above gut; ventral rows of melanophores in posterior section of gut; caudal fin pigmented; some melanophores between anus and caudal fin (this character helps to distinguish this species from *Sardinella aurita*)**Length at flexion:** 10.0 mm SL**PHOTOS**


by J.M. Rodriguez



4.3 mm SL



5.2 mm SL



13.6 mm SL

**Literature:** Alemany (1997), Fage (1920), Russell (1976), Whitehead (1984a)




**CLUPEIDAE**

*Sardina pilchardus* (Walbaum, 1792)




A.




B. 4.0 mm



C. 7.2 mm




D. 7.6 mm SL



E. 10.3 mm SL



F. 11.1 mm




G. 16.0 mm




H. ventral view (15.0 mm)

A: D'Ancona (1931a); B, H: Russell (1976); C, F, G: Fage (1920); D, E: Alemany (1997)

**CLUPEIDAE***Sardinella aurita* Valenciennes, 1847**En:** Round sardinella – **Fr:** Allache – **Sp:** Alacha**Habitat:** Pelagic, in inner continental shelf waters**Spawning season:** Between June and September**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.20–1.40 mm**Chorion:** smooth and thin**Oil globules:** single, 0.12 mm in diameter**Perivitelline space:** large**Yolk:** segmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** 3.5 mm**Distinctive characters:** body with the typical clupeid shape; oil globule situated ventrally in yolk sac**Pigmentation:** two parallel, dorsal rows of small melanophores, extending from head to tail**LARVAE****Figs. C–F****Distinctive characters:** body elongate and slender although more robust than that of *Sardina pilchardus*; head length included less than 6 times in total length; gut differentiated into two sections; dorsal fin located ahead of anus**Preanus length:** about 90% SL**Pigmentation:** melanophores aligned on both sides of gut and over it in its posterior section; caudal fin pigmented; no melanophores between anus and caudal fin in young larvae (this character helps to distinguish this species from *S. pilchardus*)**Length at flexion:** about 10.0 mm SL**PHOTOS**

by J.M. Rodriguez



3.8 mm SL



7.3 mm SL



8.6 mm SL


**Literature:** Alemany (1997), D'Ancona (1931a), Fage (1920), Olivar and Fortuño (1991), Whitehead (1984a)

**CLUPEIDAE**


*Sardinella aurita* Valenciennes, 1847



A.




B. Not sized




C. 5.3 mm SL


No melanophores between the anus and caudal fin in young larvae



D. 6.5 mm




E. 11.0 mm




F. 15.5 mm

Preanus length about 90% SL

**ENGRAULIDAE***Engraulis encrasicolus* (Linnaeus, 1758)**En:** European anchovy – **Fr:** Anchois – **Sp:** Boquerón**Habitat:** Pelagic, coastal**Spawning season:** Summer**EGGS****Fig. A****Habitat:** pelagic**Shape:** ovoid**Size:** 1.2–1.9 x 0.5–1.2 mm**Chorion:** smooth**Oil globules:** not present**Perivitelline space:** small**Yolk:** segmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** 3.3–4.0 mm**Distinctive characters:** body shape similar to yolk-sac larvae of clupeid species; yolk sac very elongate, stretching nearly to anus (this feature along with the absence of an oil globule allows for distinguishing yolk-sac larvae of this species from those of clupeid species, *Sardina pilchardus* and *Sardinella aurita*)**Pigmentation:** unpigmented**LARVAE****Figs. C–G****Distinctive characters:** body elongate and slender; gut relatively shorter than that of *S. pilchardus* and *S. aurita*; dorsal fin located over anus (allows for distinguishing this species from *S. aurita* and *S. pilchardus*)**Preanus length:** about 75% SL**Pigmentation:** similar to *S. pilchardus*; melanophores aligned on both sides of body, above gut; row of melanophores ventrally, in posterior section of gut; caudal fin pigmented; some melanophores between anus and caudal fin**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



3.2 mm SL



5.5 mm SL



10.8 mm SL


**Literature:** Alemany (1997), Fage (1920), Russell (1976), Whitehead (1984b)

**ENGRAULIDAE**

*Engraulis encrasicolus* (Linnaeus, 1758)



A.



Body unpigmented

B. ca. 2.0 mm

Yolk sac very elongate



C. 2.4 mm SL



Dorsal fin over anus

D. 5.2 mm SL

Preanus length about 75% SL




E. 6.5 mm SL


Melanophores between anus and caudal fin present at all sizes



F. 11.0 mm



G. 19.0 mm

**ARGENTINIDAE***Argentina sphyraena* Linnaeus, 1758**En:** Argentine – **Fr:** Petite argentine – **Sp:** Pez plata**Habitat:** Pelagic, coastal**Spawning seasons:** Winter and spring**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.70–1.85 mm**Chorion:** smooth**Oil globules:** single, 0.37–0.47 mm in diameter**Perivitelline space:** small**Yolk:** segmented**YOLK-SAC LARVAE****Figs. B, C****Hatch size:** may be between 7.0 and 7.5 mm**Distinctive characters:** body elongate**Pigmentation:** large stellate melanophores on yolk sac and groups of melanophores along dorsal side of gut; caudal–dorsal and ventral groups of melanophores**LARVAE****Figs. D–F****Distinctive characters:** body elongate and slender**Preanus length:** about 76% SL**Pigmentation:** 6 groups of melanophores, approximately equidistant, situated along ventral region of trunk (above gut) and tail; two opposing melanophores near caudal end; tips of upper and lower jaw pigmented; caudal fin pigmented**Length at flexion:** flexion begins at about 13.0 mm**PHOTOS**

by J.M. Rodriguez



6.0 mm SL




11.8 mm SL


**Literature:** Cohen (1984), Russell (1976), Sanzo (1931a), Schmidt (1918)

**ARGENTININIDAE**


*Argentina sphyraena* Linnaeus, 1758



A.




B. 7.6 mm



C. 7.3 mm

Body elongate and slender




D. 13.0 mm

Six spots along ventral margin of trunk and tail




E. 17.0 mm

Tips of upper and lower jaw pigmented



F. 23.3 mm

**MICROSTOMATIDAE***Microstoma microstoma* (Risso, 1810)**En:** Slender argentine**Habitat:** Mesopelagic**Spawning season:** Throughout the year, mainly in winter**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.60–1.72 mm**Chorion:** with 'pustules' on inner surface**Oil globules:** single, 0.48–0.52 mm in diameter**Perivitelline space:** small**Yolk:** segmented**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. B–F****Distinctive characters:** body moderately elongate and laterally compressed in early larvae, becomes rounded in older larvae; gut locally swollen and detached from body at its end**Preanus length:** about 75% TL**Pigmentation:** some melanophores on snout; continuous row of melanophores running along ventral side of body, from posterior margin of eye to caudal fin; row of ventral melanophores from pectoral fins to caudal peduncle; short series of spots under developing dorsal fin, becomes longer in older larvae**Length at flexion:** flexion occurs between 7.0 and 11.0 mm**PHOTOS**


Not available

**Literature:** Alemany (1997), Cohen (1984), Sanzo (1931c), Schmidt (1918)




**MICROSTOMATIDAE**

*Microstoma microstoma* (Risso, 1810)




A.



B. 2.0 mm SL




C. 3.4 mm SL




Gut swollen and detached from body at its end

D. 7.0 mm




E. 9.5 mm



F. 11.0 mm

A: Sanzo (1931c); B, C: Alemany (1997); D-F: Schmidt (1918)

**MICROSTOMATIDAE***Nansenia oblita* (Facciola, 1887)**Habitat:** Mesopelagic**Spawning season:** Winter**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.48–1.50 mm**Chorion:** with 'pustules' on inner surface**Oil globules:** single, 0.40–0.43 mm in diameter**Perivitelline space:** small**Yolk:** segmented**YOLK-SAC LARVAE****Hatch size:** about 4.0 mm SL**Pigmentation:** not pigmented**LARVAE****Figs. B–F****Distinctive characters:** body moderately elongate and laterally compressed, becomes rounded in larger larvae; end of gut detached from body in early larvae**Preanus length:** about 75% TL**Pigmentation:** a row of melanophores along both the dorsal and ventral margin of the notochord; scattered melanophores on head appear as a bar through the eye; rows of small melanophores along ventral side of trunk and tail, from pectoral-fin base to level of anus; row of melanophores along lower jaw; pigment spreads over most of body with development, except on caudal peduncle**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



3.4 mm SL



3.5 mm SL

**Literature:** Alemany (1997), Cohen (1984), Sabatés (1988), Sanzo (1931c), Schmidt (1918)

**MICROSTOMATIDAE**

*Nansenia oblita* (Facciolà, 1887)



A.




B. 5.3 mm SL




End of gut detached  
from body


C. 6.0 mm SL



D. 7.8 mm




E. 9.0 mm



F. 13.0 mm

A: Sanzo (1931c); B, C: Alemany (1997); D-F: Schmidt (1918)

**GONOSTOMATIDAE***Cylothone braueri* Jespersen & Täning, 1926**En:** Garrick**Habitat:** Oceanic, mesopelagic, between 250 and 900 m depth**Spawning season:** Mainly between April and October*Cylothone* sp.**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–F****Distinctive characters:** body quite elongate with a large swimbladder; eye round and small; dorsal and anal fins located at same level, just after anus**Preanus length:** about 50% SL**Pigmentation:** prominent spot on ventral side of caudal peduncle; 3 pairs of melanophores along lateral sides of gut: one close to pectoral-fin base, another at mid-gut and a third one over hindgut; dorsum of swimbladder pigmented; a series of 9 to 12 equidistant melanophores along the postanal ventral region; later larvae with a regular ventral postanal row of internal melanophores in correspondence with the external postanal row; row of 5 or 6 internal melanophores on ventrolateral anterior region of body and about 3 internal melanophores on upper part of caudal peduncle**Length at flexion:** about 4.8 mm**PHOTOS**

by J.M. Rodriguez



3.6 mm SL



5.0 mm SL



10.5 mm SL

**Literature:** Alemany (1997), Badcock (1984a), Jespersen and Täning (1926), Sabatés (1988)

**GONOSTOMATIDAE**

*Cyclothone braueri* Jespersen & Täning, 1926



A. 2.3 mm SL

Eye round and small  
9 to 12 equidistant melanophores along ventral tail region




B. 2.9 mm SL




C. 4.0 mm SL

Body elongate with prominent swimbladder




D. 6.1 mm SL

Dorsum of swimbladder pigmented




E. 10.2 mm SL


Preanus length ca. 50% SL



F. 13.7 mm

**STERNOPTYCHIDAE***Argyropelecus hemigymnus* Cocco, 1829**En:** Half-naked hatchetfish – **Sp:** Hacha de plata**Habitat:** Oceanic, mesopelagic, between 200 and 800 m during the day and 100 to 600 m at night, making marked vertical migrations**Spawning season:** Throughout the year, peaking in winter and early summer*Argyropelecus* sp.**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 0.92–1.04 mm**Chorion:** smooth, with a secondary membrane inside chorion**Oil globules:** single, 0.26–0.28 mm in diameter**Perivitelline space:** small**Yolk:** segmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** 2.5 mm SL**Distinctive characters:** body elongate and slender; yolk sac very large; oil globule situated at ventral side of yolk sac**Pigmentation:** unpigmented**LARVAE****Figs. C–F****Distinctive characters:** body moderately elongate in young larvae, shortens at transformation; eye oval and narrow in early stages, becomes slightly telescopic in older larvae; gut detached from body**Preanus length:** 44–59% SL in early larvae**Pigmentation:** body unpigmented in preflexion stage; photophores present in older larvae**Length at flexion:** 10.0–11.0 mm**Length at transformation:** 7.8–12.0 mm; initially, during transformation, larvae shrink (mainly the anterior part of body), gut shortens, head deepens, and eyes become telescopic**PHOTOS**

by J.M. Rodriguez



4.9 mm SL



6.7 mm SL


**Literature:** Alemany (1997), Badcock (1984c), Sanzo (1931b), Täning (1918)

**STERNOPTYCHIDAE**

*Argyropelecus hemigymnus* Cocco, 1829




A.



B. 2.5 mm

Body unpigmented in  
preflexion stage




C. 6.9 mm SL




D. 9.0 mm

Gut detached from body




E. 5.6 mm SL




F. 6.2 mm SL

A, B: Sanzo (1931b); C, E, F: Alemany (1997); D: Täning (1918)

**STERNOPTYCHIDAE***Maurolicus muelleri* (Gmelin, 1789)**En:** Silvery lightfish**Habitat:** Mesopelagic, between 40 and 400 m depth**Spawning season:** Over the whole year*Maurolicus* sp.**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.32–1.58 mm**Chorion:** sculptured with hexagonal structures**Oil globules:** single, 0.26–0.28 mm in diameter**Perivitelline space:** narrow**Yolk:** segmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** around 3.0 mm**Distinctive characters:** body relatively elongate and slender, transparent; yolk sac very large, oval and projected beyond snout; oil globule situated at ventral side of yolk sac; anus reaches the border of finfold**Pigmentation:** unpigmented**LARVAE****Figs. C–F****Distinctive characters:** body relatively elongate with a prominent swimbladder; eye vertically elliptical, becoming round in late larvae; at about 6 mm SL, a photophore under eye and two in ventral region, under swimbladder**Preanus length:** about 50% SL, increasing with development**Pigmentation:** young larvae unpigmented; pigment first appears on dorsum of swimbladder; in later larvae, pigmentation limited to dorsum of swimbladder and photophores**Length at flexion:** flexion completed at 5.8 mm SL**PHOTOS**

by J.M. Rodriguez



5.4 mm SL



7.6 mm SL




10.5 mm SL

**Literature:** Alemany (1997), Badcock (1984c), Jespersen and Täning (1926), Sanzo (1935)




**STERNOPTYCHIDAE**


*Maurolicus muelleri* (Gmelin, 1789)




A.




B. 2.5 mm



C. 2.9 mm SL




D. 4.2 mm SL




E. 6.4 mm

At about 6 mm SL, a photophore under eye and two in ventral region, at level of swimbladder



F. 10.0 mm

**PHOSICHTHYIDAE***Vinciguerria attenuata* (Cocco, 1838)**En:** Silvery lightfish**Habitat:** Mesopelagic, between 250 and 600 m depth**Spawning season:** Spring–summer

Phosichthyidae

**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–F****Distinctive characters:** body elongate and slender with a prominent swimbladder (not present in young larvae); dorsal and anal fins located at level of anus**Preanus length:** about 75% SL**Pigmentation:** larvae less than 6 mm SL with 6–8 stellate melanophores on both sides of body; in larger larvae, lateral rows disappear and a prominent median caudal spot develops; dorsum of swimbladder pigmented**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



7.2 mm SL



10.2 mm SL



10.7 mm SL

**Literature:** Alemany (1997), Badcock (1984b), Jespersen and Täning (1926)

**PHOSICHTHYIDAE**

*Vinciguerria attenuata* (Cocco, 1838)



A. 4.0 mm SL

Larvae less than 6 mm SL with 6–8 stellate melanophores on each side of body




Body elongate and slender

B. 7.0 mm SL



Large median caudal spot

C. 9.7 mm




Dorsum of swimbladder pigmented

D. 13.5 mm SL



E. 18.3 mm



F. 18.5 mm

A, B, D: Alemany (1997); C, E, F: Jespersen and Täning (1926)

**STOMIIDAE***Stomias boa* (Risso, 1810)**En:** Boa dragonfish – **Sp:** Estomias**Habitat:** Mesopelagic, in deep oceanic waters to more than 1 000 m depth**Spawning season:** Spring–summer**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–E****Distinctive characters:** body very elongate with a long head and large jaws; eye oval; small dorsal and anal fins located at same level, at end of body**Preanus length:** about 88% SL**Pigmentation:** in young larvae, row of small melanophores over gut, between pectoral and caudal-fin base; some melanophores appear in dorsal–caudal region in larvae of about 6 mm SL, extending forward with growth; dorsal and ventral series of melanophores disappear in older larvae**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



6.0 mm SL








16.6 mm SL

**Literature:** Alemany (1997), Fahay (1983), Gibbs (1984), Sabatés (1988)

**STOMIIDAE**

*Stomias boa* (Risso, 1810)



**SYNODONTIDAE***Synodus saurus* (Linnaeus, 1758)**En:** Atlantic lizardfish – **Sp:** Pez lagarto**Habitat:** Benthic, between 20 and 400 m depth, mostly in depths less than 50 m**Spawning season:** Unknown**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.10–1.35 mm**Chorion:** ornamented, covered with a hexagonal reticulum**Oil globules:** none**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Hatch size:** 4.0–4.5 mm SL**Pigmentation:** 5 large ventral spots approximately equidistant, 4 over gut, between pectoral fin and anus and another in caudal region; melanophores arranged radially over caudal end of primordial fin**LARVAE****Figs. B–F****Distinctive characters:** body elongate and slender**Preanus length:** about 80% SL**Pigmentation:** similar to yolk-sac larvae, difference lies in the number of ventral patches that may be 6; caudal fin pigmented**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez




3.4 mm SL


**Literature:** Alemany (1997), Sulak (1984), Täning (1918), Tortonese (1956b)

**SYNODONTIDAE**


*Synodus saurus* (Linnaeus, 1758)



A.



B. 4.1 mm SL




C. 4.5 mm SL


5 (6 in larger larvae) large spots  
along ventral side of trunk and tail



D. 4.6 mm



E. 5.0 mm



F. 12.0 mm

A, D: Tortonese (1956b); B, C: Alemany (1997); E, F: Täning (1918)

## PARALEPIDIDAE

*Lestidiops jayakari* (Boulenger, 1889)

**Habitat:** Mesopelagic, between 200 and 2000 m depth

**Spawning season:** Probably throughout all year



## EGGS

Undescribed

## YOLK-SAC LARVAE

Undescribed

## LARVAE

Figs. A–G

**Distinctive characters:** body elongate and slender; head short and deep in small larvae, becomes larger with development; eye large and round

**Preanus length:** about 20% SL in young larvae, increases with development

**Pigmentation:** two large stellate postanal, ventral melanophores (congener species, *L. sphyrenoides*, has 3 postanal ventral patches) and two series of small melanophores, one above and another below urostyle; tip of upper and lower jaw pigmented; up to 12 peritoneal patches of pigment form during development

**Length at flexion:** 12.0–16.0 mm

## PHOTOS

by J.M. Rodriguez



5.5 mm SL



10.5 mm SL

**Literature:** Alemany (1997), Ege (1930), Fahay (2007), Post (1984), Richards (2006)




**PARALEPIDIDAE**


*Lestidiops jayakari* (Boulenger, 1889)



A. 2.8 mm SL




Two large postanal, ventral melanophores (3 in *L. sphyrenoides*) B. 6.6 mm SL



C. 8.0 mm

Series of melanophores above and below urostyle (also in *L. sphyrenoides*)




D. 12.0 mm



E. 15.0 mm




F. 16.5 mm



Up to 12 peritoneal patches

G. 20.0 mm

A, B: Alemany (1997); C-G: Ege (1930)

**PARALEPIDIDAE***Paralepis coregonoides* Risso, 1820**Habitat:** Meso- to bathypelagic**Spawning season:** Spring–summer**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–G****Distinctive characters:** body elongate, snout relatively long; eye big and ovoid at first stages of development, becomes round in older larvae**Preanus length:** about 30% SL in early larvae, increases with development**Pigmentation:** postanal region unpigmented up to about 5 mm when a deep spot appears above notochord, close to its end; long series of internal melanophores above notochord and a short one, below it; the number of peritoneal patches increases from 1 in larvae of 6 mm SL to 3 in older larvae**Length at flexion:** about 10.0–15.0 mm**PHOTOS**


by J.M. Rodriguez



8.8 mm SL



10.2 mm SL



15.0 mm SL

**Literature:** Alemany (1997), Ege (1930), Fahay (2007), Post (1984), Richards (2006)

**PARALEPIDIDAE**

*Paralepis coregonoides* Risso, 1820



A. 3.3 mm SL

Postanal region unpigmented until about 5 mm




B. 5.0 mm




C. 7.5 mm



D. 11.0 mm




E. 15.0 mm




Long series of internal melanophores, above, and short series below notochord in large larvae

Up to 3 peritoneal patches

F. 18.5 mm



G. 21.0 mm

**EVERMANNELLIDAE***Evermannella balbo* (Risso, 1820)**En:** Balbo sabretooth**Habitat:** Mesopelagic, between 100 and 800 m depth**Spawning season:** No information**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–E****Distinctive characters:** body fairly elongate; snout long and pointed; eye elliptical; large teeth in both jaws**Preanus length:** less than 50% SL**Pigmentation:** 3 large peritoneal pigment patches; in young larvae, melanophores scattered on lateral preanal region and some melanophores on caudal-fin base; in later larvae, melanophores arranged in transversal bars, following myosepta; melanophores on head, snout and neck; caudal-fin base pigmented**Length at flexion:** flexion completed at 6.8 mm SL**PHOTOS**

by J.M. Rodriguez



5.2 mm SL



6.5 mm SL




7.4 mm SL

**Literature:** Alemany (1997), Fahay (2007), Johnson (1984), Richards (2006), Schmidt (1918)


**EVERMANNELLIDAE**

*Evermannella balbo* (Risso, 1820)



A. 3.0 mm SL

3 large peritoneal pigment patches




B. 5.75 mm

Melanophores arranged in transversal bars




C. 8.0 mm

Large teeth in both jaws




D. 11.5 mm

Eye elliptical



E. 14.0 mm

**MYCTOPHIDAE***Benthoosema glaciale* (Reinhardt, 1837)**En:** Glacier lantern fish – **Fr:** Lanterne glaciale – **Sp:** Linterna oscura**Habitat:** Mesopelagic, between 275 and 850 m depth during daytime; from surface down to 225 m depth at night**Spawning season:** Spawning peaks in late spring, summer**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–E****Distinctive characters:** body moderately elongate; eye slightly elliptical with choroid tissue ventrally; gap between anus and anal fin present in young larvae, closes at about 8.0 mm**Preanus length:** about 33% of SL in young larvae, increases with development**Pigmentation:** melanophores at posterior edge of opercle, at tip of snout and lower jaw; 3 ventral melanophores on cleithral symphysis, lateral sides of gut and hindgut; several melanophores on postanal ventral region, decreasing in number to a single one over mid of anal fin in older larvae; pectoral-fin rays pigmented; premature preopercular photophore**Length at flexion:** 5.0–7.0 mm**PHOTOS**

by J.M. Rodriguez



4.1 mm SL



6.2 mm SL



8.2 mm SL


**Literature:** Alemany (1997), Fahay (1983), Hulley (1984), Täning (1918)

**MYCTOPHIDAE**


*Benthosema glaciale* (Reinhardt, 1837)



A. 2.3 mm SL




B. 4.8 mm SL




Gap between anus and  
anal fin

C. 5.7 mm SL




Eye elliptical with  
choroid tissue ventrally

D. 7.6 mm SL



E. 11.0 mm

**MYCTOPHIDAE***Ceratoscopelus maderensis* (Lowe, 1839)**En:** Madeira lantern fish – **Fr:** Lanterne de Madère – **Sp:** Carpintero**Habitat:** Mesopelagic, between 330 and 600 m depth during daytime, 0 to 175 m at night**Spawning season:** Spawning peaks in spring, summer**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–F****Distinctive characters:** body elongate, moderately slender; eye spherical and relatively large**Preanus length:** more than 50% SL in early larvae, increases to more than 60% SL in older larvae**Pigmentation:** a single melanophore on each side of gut and on hindgut; a continuous line of melanophores between anus and caudal fin decreasing in number with development; 3 or 4 melanophores on dorsal side of caudal peduncle (not present in young larvae); no melanophores at cleithral symphysis**Length at flexion:** about 6 mm SL**PHOTOS**

by J.M. Rodriguez



4.3 mm SL



6.0 mm SL



7.9 mm SL

**Literature:** Alemany (1997), Fahay (1983), Hulley (1984), Täning (1918)



**MYCTOPHIDAE**

*Ceratoscopelus maderensis* (Lowe, 1839)




A. 2.0 mm SL



No melanophores at cleithral symphysis

Continuous ventral line of melanophores between anus and caudal fin

B. 2.8 mm SL




3 dorsal melanophores on caudal peduncle, not present in youngest larvae

C. 5.0 mm




D. 7.0 mm




E. 11.5 mm

No melanophores at ventral caudal-fin base



F. 16.5 mm

**MYCTOPHIDAE***Diaphus holti* Täning, 1918**En:** Small lantern fish – **Fr:** Lanterne courte – **Sp:** Rafino corto**Habitat:** Mesopelagic, between 80 and 675 m depth**Spawning season:** Spring and summer**EGGS**


Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–E****Distinctive characters:** body moderately slender; eye round and relatively large**Preanus length:** less than 50% SL**Pigmentation:** melanophores on cleithral symphysis; a single melanophore on lateral sides of gut and over hindgut; regular row of ventral melanophores between anus and caudal region; single, large melanophore on lower half of caudal-fin base**Length at flexion:** less than 5 mm**PHOTOS**

by J.M. Rodriguez



3.9 mm SL



4.6 mm SL



5.5 mm SL


**Literature:** Alemany (1997), Fahay (1983), Hulley (1984), Täning (1918)

**MYCTOPHIDAE**

*Diaphus holti* Täning, 1918




A. 3.0 mm SL



Melanophores at  
cleithral symphysis


B. 5.0 mm




Prominent spot at  
caudal-fin base

Continuous line of  
melanophores between  
anus and caudal fin

C. 6.5 mm



D. 10.0 mm




E. 11.5 mm

**MYCTOPHIDAE***Hygophum* Bolin, 1939

Larvae of the two species of this genus inhabiting the Mediterranean Sea, *H. benoiti* and *H. hygomii*, are very similar. They are almost only differentiated by their pigmentation patterns

**Habitat:** Mesopelagic, between 100 and 1000 m depth during daytime, 12 to 400 m and 700 to 1000 m at night (*H. benoiti*); 600 to 750 m during the day, 0 to 225 m at night (*H. hygomii*)

**Spawning season:** Spawning peaks in spring–summer (*H. benoiti*); end of summer–autumn (*H. hygomii*)

**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–F**

**Distinctive characters:** body moderately slender; body depth increases with development; eye unstalked, moderately elliptical with a prominent conical mass of choroid tissue ventrally; anus situated at anterior margin of anal fin; gut thick with a fold–end section; dorsal anterior part of finfold globose

**Preanus length:** 50% SL in young larvae, increases with development

**Pigmentation:** ventral series of melanophores on isthmus, continuing to cleithrum; a series of usually 3 melanophores on lateral sides of gut (disappearing with development in *H. hygomii*); a melanophore on hindgut; melanophores on caudal–fin base in some individuals


**Length at flexion:** 5.0–5.5 mm SL

**PHOTOS**

by J.M. Rodriguez



8.1 mm SL




10.3 mm SL

**Literature:** Badcock and Merrett (1976), Fahay (2007), Hulley (1984), Olivar and Palomera (1994), Täning (1918)

**MYCTOPHIDAE**


*Hygophum* Bolin, 1939

*Hygophum benoiti*




A. 5.5 mm SL

Dorsal anterior part of finfold globose




B. 7.8 mm SL




Prominent mass of choroid tissue

C. 9.2 mm SL


*Hygophum hygomii*



D. 5.3 mm SL




E. 7.0 mm SL



F. 9.9 mm SL

All illustrations reproduced from *Journal of Plankton Research* (1994) 16(8): 977–991 with permission

A–F: Olivar and Palomera (1994)

**MYCTOPHIDAE***Lampanyctus crocodilus* (Risso, 1810)**En:** Jewel lanternfish – **Fr:** Lanterne crocodile – **Sp:** Madre de la anchoa**Habitat:** Mesopelagic between 100 and 200 m depth and 700 to 1000 m during daytime, 45 to 150 and 400 to 1000 m at night**Spawning season:** Spring–summer**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–F****Distinctive characters:** body initially elongate, soon deepens, especially at level of pectoral region; eye moderately large and round; well-developed, protruding teeth in the upper jaw from young larvae on**Preanus length:** increases from about 25% SL, in young larvae, to about 60% SL, in older larvae**Pigmentation:** prominent, single melanophore on top of head; a single melanophore on body dorsum, between dorsal and adipose fins, from the flexion stage on; melanophores at tip of lower jaw and over hindgut; peritoneal pigment develops in older larvae; spots on pectoral-fin base; late stages add pigment to anterior myosepta; in some individuals, melanophores at top of upper jaw and at middle of forehead**Length ant flexion:** about 5 mm**PHOTOS**

by J.M. Rodriguez



4.5 mm SL




6.3 mm SL


**Literature:** Alemany (1997), Fahay (1983), Hulley (1984), Richards (2006), Täning (1918)

**MYCTOPHIDAE**

*Lampanyctus crocodilus* (Risso, 1810)



A-C: Alemany (1997); D-F: Täning (1918)

**MYCTOPHIDAE***Lampanyctus pusillus* (Johnson, 1890)**En:** Pigmy lanternfish**Habitat:** Mesopelagic, between 500 and 1000 m depth during the day, 50 to 150 m at night**Spawning season:** Spawning peaks at the end of summer–autumn**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–F****Distinctive characters:** body initially elongate, soon becomes very deep and stout; head large; eye large and round; gut thick; well-developed teeth in upper jaw from young larvae on**Preanus length:** increases from about 30% SL, in young larvae, to about 65% SL, in late larvae**Pigmentation:** 1 to 3 melanophores from snout to top of head, these melanophores become internal, located at bases of brain and forebrain, in older larvae; spots at tip of lower jaw, over opercle, pectoral-fin base, anterior gut and over hindgut; internal pigment on swimbladder; ventral spots on abdominal region; a series of paired melanophores along dorsum and another along lateral midline in older larvae**Length at flexion:** about 4.0–5.0 mm**PHOTOS**

by J.M. Rodriguez



3.4 mm SL



4.3 mm SL

**Literature:** Alemany (1997), Fahay (1983), Hulley (1984), Täning (1918)




**MYCTOPHIDAE**


*Lampanyctus pusillus* (Johnson, 1890)



A. 1.9 mm SL




B. 3.0 mm SL




C. 3.3 mm SL


Body shorter and thicker than that of *L. crocodilus*




D. 5.0 mm



E. 6.5 mm



F. 8.5 mm

**MYCTOPHIDAE***Lobianchia dofleini* (Zugmayer, 1911)**En:** Dofleini's lantern fish – **Fr:** Lanterne de Dofleini – **Sp:** Japonés**Habitat:** Mesopelagic, between 375 and 600 m depth during daytime, 25 to 400 m at night**Spawning season:** Throughout the year, peaking in February–June**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–E**

**Distinctive characters:** body relatively elongate in young larvae, soon becoming deep and stout, especially its anterior half; eye small and round in early stages, becoming slightly elliptical, with a mass of choroid tissue ventrally; pectoral fin well developed, with elongate (almost reaching anus) upper rays, these rays shortening with development

**Preanus length:** 50–60% SL in late larvae

**Pigmentation:** swimbladder pigmented; spots on each side of hindgut; melanophores on gut, on ventral midline, anterior to cleithral symphysis, over the latter and along base of anal fin; base and rays of pectoral fin pigmented; in late stages, spots appear on dorsum of body

**Length at flexion:** 5–6 mm

**PHOTOS**

by J.M. Rodriguez



5.2 mm SL



6.7 mm SL


**Literature:** Alemany (1997), Fahay (1983), Hulley (1984), Täning (1918)

**MYCTOPHIDAE**

*Lobianchia dofleini* (Zugmayer, 1911)




A. 2.2 mm SL




B. 3.0 mm SL




C. Detail of pectoral fin




C. 4.5 mm SL



D. 5.5 mm



E. 8.0 mm

**MYCTOPHIDAE***Myctophum punctatum* Rafinesque, 1810**En:** Spotted lanternfish**Habitat:** Mesopelagic, between 100 to 150 and 700 to 1000 m depth during daytime, less than 190 and 700 to 800 m at night**Spawning season:** Late winter–summer**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–E****Distinctive characters:** body elongate in young larvae, becoming slightly deeper with development; head large and flat; eye elliptical, stalked in young larvae, with tapered choroid mass, ventrally**Preanus length:** about 50–60% SL**Pigmentation:** ventral series of spots from anus to head; spots on edges of both jaws and on upper part of opercle; several ventral postanal melanophores; two single, well developed melanophores, in an opposite position over and under urostyle; pigment may be present on posterior rays of dorsal, anal and adipose fins; rays and base of pectoral fin pigmented; spots on edges of both jaws and on upper part of opercle; dorsal melanophore on caudal region**Length at flexion:** 7 mm SL**PHOTOS**

by J.M. Rodriguez



5.3 mm SL



7.0 mm SL




9.1 mm SL

**Literature:** Alemany (1997), Fahay (1983), Hulley (1984), Täning (1918)

**MYCTOPHIDAE**

*Myctophum punctatum* Rafinesque, 1810




A-C: Alemany (1997); D, E: Täning (1918)

**MYCTOPHIDAE***Notoscopelus elongatus* (Costa, 1844)

**Habitat:** Mesopelagic, between 375 to 1 000 m depth during daytime, 45 to 150 m at night

**Spawning season:** Winter–spring

**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–E**

**Distinctive characters:** body elongate in young larvae, becomes deep and compressed with development; eye large and round; large swimbladder

**Preanus length:** increases from about 25 to 50–60% SL with development

**Pigmentation:** increases with development; initially, melanophores at jaw tips, and two on head (increasing to 4 in older larvae); peritoneum and dorsum of swimbladder, pigmented; two parallel dorsal rows of melanophores in older larvae

**Length at flexion:** almost completed at 6.0 mm SL

**PHOTOS**

by J.M. Rodriguez



3.7 mm SL



4.2 mm SL



6.8 mm SL


**Literature:** Alemany (1997), Hulley (1984), Palomera (1983)

**MYCTOPHIDAE**


*Notoscopelus elongatus* (Costa, 1844)




A. 3.0 mm SL




B. 4.6 mm SL



C. 5.6 mm SL




D. 6.5 mm SL




Two parallel, dorsal rows of melanophores in late larvae

E. Dorsal view



E. 11.3 mm SL

**MYCTOPHIDAE***Symbolophorus veranyi* (Moreau, 1888)**En:** Large scale lantern fish – **Fr:** Lanterne à grandes écailles – **Sp:** Linterna**Habitat:** Oceanic, mesopelagic, between 100 and 700–800 m during daytime, surface to 150 m depth at night**Spawning season:** No information**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–F****Distinctive characters:** body moderately elongate and slender, similar to that of *Myctophum punctatum*; narrow eyes on short stalks; small ventral cone of choroid tissue; snout pointed, flat in young larvae; large pectoral fins with elongated bases, extending beyond anus**Preanus length:** about 50% SL**Pigmentation:** few preanal ventral spots; spots on tips of snout and lower jaw; large spot on posterior edge of opercle; row of ventral–postanal melanophores; row of spots along gut; pigment on pectoral–fin rays (heavier at ray base); pigment decreases towards the end of the larval period**Length at flexion:** flexion almost completed at 7.5 mm SL**PHOTOS**

by J.M. Rodriguez



7.0 mm SL




9.1 mm SL

**Literature:** Alemany (1997), Fahay (1983), Hulley (1984), Täning (1918)




**MYCTOPHIDAE**

*Symbolophorus veranyi* (Moreau, 1888)



A. 3.8 mm SL




Narrow eyes on short stalks

B. 4.8 mm SL




C. 5.1 mm SL




Large pectoral fins with elongated bases

D. 7.5 mm SL




Ventral cone of choroid tissue

E. 8.3 mm



Large spot on posterior edge of opercle, in older larvae

F. 17.0 mm

**GADIDAE*****Gadiculus argenteus*** Guichenot, 1850**En:** Silvery pout – **Fr:** Merlan argenté – **Sp:** Faneca plateada**Habitat:** Bathypelagic, forming shoals in midwater over the continental-shelf edge, between 200 and 1 000 m depth**Spawning season:** December and January**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–D****Distinctive characters:** body shape typical of a gadid species, relatively short, with a relatively large head and deep abdominal region; pelvic fins, short, develop in later larvae**Preanus length:** less than 50% SL**Pigmentation:** two opposing pigment patches (one dorsal and another ventral) about midway along tail, growing on lateral sides of body to form a continuous bar in later larvae; some melanophores on head, and upper and lower jaw tips; peritoneal region pigmented; fins, except the caudal, unpigmented**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



2.1 mm SL




2.7 mm SL




**Literature:** D'Ancona (1933a), Russell (1976), Sabatés (1988), Svetovidov (1986a)

**GADIDAE**


*Gadiculus argenteus* Guichenot, 1850



Postanal pigment reduced to 2 opposing pigment patches at about mid-tail



Fins, except the caudal, unpigmented

**GADIDAE***Micromesistius poutassou* (Risso, 1827)**En:** Blue whiting(=Poutassou) – **Fr:** Merlan bleu – **Sp:** Bacaladilla**Habitat:** Mesopelagic, between 30 to 400 m depth, over depths of 160 to 3 000 m**Spawning season:** Between January and March**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.0–1.14 mm**Chorion:** smooth**Oil globules:** no oil globule**Perivitelline space:** narrow**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** 2.0–2.2 mm**Pigmentation:** melanophores scattered over body, except caudal region; posteriorly, dorsal and ventral rows of paired melanophores over trunk and a cap of melanophores on head; yolk sac unpigmented**LARVAE****Figs. C–G****Distinctive characters:** body shape typical of a gadid species, relatively short with a relatively large head and deep abdominal region**Preanus length:** less than 50% SL**Pigmentation:** dorsal (from head) and ventral (from anus) rows of paired melanophores to about mid-caudal region, with dorsal rows longer than ventral ones; melanophores on head; peritoneal region pigmented; no melanophores on lateral sides of trunk in larvae less than 6 mm long; dorsal and ventral rows of pigment extend backwards with development**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



2.4 mm SL




6.0 mm SL


**Literature:** D'Ancona (1933a), Sabatés (1988), Seaton and Bailey (1971), Schmidt (1905), Svetovidov (1986a)

**GADIDAE**


*Micromesistius poutassou* (Risso, 1827)



A.



B. 2.1 mm



C. 3.4 mm


Dorsal rows of paired melanophores  
longer than ventral ones




D. 6.5 mm



G. 3.29 mm  
(dorsal view)




E. 9.0 mm



F. 15.0 mm


A-C, G: Seaton and Bailey (1971); D, E: Schmidt (1905); F: D'Ancona (1933a)

**GADIDAE*****Trisopterus luscus*** (Linnaeus, 1758)**En:** Pouting(=Bib) – **Fr:** Tacaud commun – **Sp:** Faneca**Habitat:** Benthic, between 30 and 100 m depth**Spawning season:** March and April**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.0–1.23 mm**Chorion:** smooth**Oil globules:** no oil globule**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** less than 3.0 mm**Pigmentation:** recently hatched larvae, very similar to those of *Micromesistius poutassou*; posteriorly, well-marked dorsal (from head) and a ventral (from anus) paired rows of melanophores, extending to about two-thirds of postanal region**LARVAE****Figs. C–F****Distinctive characters:** body shape typical of a gadid species, relatively short with a relatively large head and deep abdominal region**Preanus length:** about 50% of SL**Pigmentation:** dorsal (from head) and ventral (from anus) rows of paired melanophores, ending at same level, at about mid-tail region**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



3.3 mm SL




12.0 mm SL


**Literature:** Russell (1976), Svetovidov (1986a)

**GADIDAE**


*Trisopterus luscus* (Linnaeus, 1758)




A.




B. 3.0 mm



C. 4.5 mm




D. 7.3 mm




Dorsal and ventral rows of paired melanophores end at same level

E. 11.0 mm



F. 18.0 mm

**LOTIDAE*****Gaidropsarus biscayensis* (Collett, 1890)****En:** Mediterranean bigeye rockling – **Fr:** Motelle – **Sp:** Barbada**Habitat:** Benthic, possibly bathypelagic over soft bottoms, between 140 and 600 m depth**Spawning season:** Winter**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–E**

**Distinctive characters:** body shape typical of a gadid species, relatively short with a relatively large head and deep abdominal region; body depth decreases gradually with development starting from a size of 3.5–4.0 mm; anus does not reach finfold edge in early larvae; 2 prominent cephalic spines on each side of head; large pelvic fins with 4 rays, located on sides of gut

**Preanus length:** less than 60% SL in early larvae, decreases slightly with development

**Pigmentation:** dorsal side of peritoneum strongly pigmented; melanophores on top of head and on lateral surface of trunk, these latter increase in number and widen joining those on head and forming a continuous bar; upper and lower jaw tips pigmented; pelvic fins heavily pigmented; postanal region unpigmented, in young larvae

**Length at flexion:** unknown

**PHOTOS**

by J.M. Rodriguez



2.6 mm SL



3.9 mm SL




5.2 mm SL

**Literature:** Alemany (1997), Demir (1982), Svetovidov (1986a)




**LOTIDAE**

*Gaidropsarus biscayensis* (Collett, 1890)




A. 2.3 mm SL




Two prominent cephalic spines  
on each side of head

B. 3.2 mm SL




Postanal region  
unpigmented

C. 3.3 mm SL




C. Dorsal  
view




Pelvic fins located on  
sides of gut

D. 5.9 mm SL



E. 10.4 mm SL

**LOTIDAE*****Gaidropsarus mediterraneus*** (Linnaeus, 1758)**En:** Shore rockling – **Fr:** Motelle de Méditerranée – **Sp:** Bertorella**Habitat:** Benthic, littoral, from about 20 to 120 m depth**Spawning season:** Winter to early spring**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 0.66–0.72 mm**Chorion:** smooth**Oil globules:** single, 0.15–0.19 mm in diameter**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** 1.8–1.9 mm**Pigmentation:** similar to that of the larva, with the only difference of a pigment patch located at end of gut that extends to the dorsal region of the body**LARVAE****Figs. C–F****Distinctive characters:** body shape typical of a gadid species, relatively short with a relatively large head and deep abdominal region; cephalic spines on each side of head; pelvic fins large with 4 rays**Preanus length:** less than 50% SL**Pigmentation:** two opposing pigment patches (one dorsal and another ventral) about midway along tail; pigment patch on ventral part of caudal fin; melanophores on neck, head, lower jaw and peritoneal region; pelvic fins heavily pigmented**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



2.4 mm SL



3.5 mm SL


**Literature:** Aboussouan (1964), Alemany (1997), Cipria (1939), D'Ancona (1933a), Demir (1982)

**LOTIDAE**


*Gaidropsarus mediterraneus* (Linnaeus, 1758)




A.




B. 3.0 mm




C. 2.3 mm SL



D. 3.2 mm SL




E. 6.1 mm SL



F. 11.2 mm SL


A, B: Cipria (1939); C-F: Demir (1982)

**MERLUCCIIDAE***Merluccius merluccius* (Linnaeus, 1758)**En:** European hake – **Fr:** Merlu européen – **Sp:** Merluza europea**Habitat:** Midwater or close to the bottom, mainly between 70 and 700 m depth**Spawning season:** Throughout the whole year, mostly in winter and spring**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 0.94–1.03 mm**Chorion:** smooth**Oil globules:** single, 0.25–0.28 mm in diameter**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 3 mm SL**Pigmentation:** 3 postanal stellate melanophores**LARVAE****Figs. C–F****Distinctive characters:** body relatively short with a large and deep head and abdominal region; pelvic fins appear early, at 4–6 mm in length**Preanus length:** about 45% SL**Pigmentation:** three stellate melanophores in lateral–postanal region; peritoneal region and pelvic fins pigmented; melanophores on snout, lower jaw and on neck; no melanophores on the postanal ventral side of body**Length at flexion:** no information; for other *Merluccius* species, it has been established at about 9.0–10.0 mm**PHOTOS**

by J.M. Rodriguez



3.5 mm SL




7.0 mm SL


**Literature:** D'Ancona (1933a), Palomera *et al.* (2005), Russell (1976), Svetovidov (1986b)

**MERLUCCIIDAE**


*Merluccius merluccius* (Linnaeus, 1758)




A.



B. 2.5 mm SL




C. 3.8 mm SL




D. 6.5 mm SL

3 melanophores on lateral side of tail



E. 9.1 mm SL

No melanophores on the postanal ventral side of body




F. 12.5 mm

**OPHIDIIDAE***Ophidion barbatum* Linnaeus, 1758

Larvae of *Ophidion barbatum* and *Parophidion vassali* are very similar and many times their identification has been confusing. Even the systematics and the taxonomic validity of these two genera and their corresponding species have been questioned

**Habitat:** Benthic, over sand and mud bottoms, from shallow down to 150 m depth

**Spawning season:** Summer

**EGGS****Fig. A**

**Habitat:** pelagic

**Shape:** spherical

**Diameter:** 0.76 mm

**Chorion:** smooth

**Oil globules:** none

**Perivitelline space:** small

**Yolk:** unsegmented

**YOLK-SAC LARVAE****Fig. B**

**Hatch size:** may be about 2.5 mm SL

**Distinctive characters:** body slender

**Pigmentation:** dotted melanophores under gut; a series of dotted melanophores extending from pectoral fin to urostyle end; some melanophores on dorsal posterior half of tail

**LARVAE****Figs. C-E**

**Distinctive characters:** body elongate and slender

**Preanus length:** less than 50% SL

**Pigmentation:** in early stages, similar to that of yolk-sac larvae; in late larvae, dorsal melanophores decrease in number and melanophores on lower jaw and posterior edge of operculum appear

**Length at flexion:** unknown

**PHOTOS**

by J.M. Rodriguez



3.0 mm SL




6.8 mm SL


**Literature:** Alemany (1997), Nielsen (1986b), Padoa (1956j), Sabatés (1988), Tortonese (1975)

**OPHIDIIDAE**

*Ophidion barbatum* Linnaeus, 1758




A.




B. 2.5 mm SL

Series of dotted melanophores from pectoral fin to urostyle end




C. 6.2 mm SL



D. 11.1 mm SL

Some melanophores on dorsal, posterior half of tail




E. 15.5 mm SL

**CARAPIDAE***Echiodon dentatus* (Cuvier, 1829)

**Habitat:** Benthic; young swimming in the water column

**Spawning season:** Winter

**EGGS****Fig. A**

**Habitat:** pelagic, in slimy masses

**Shape:** ellipsoidal

**Size:** 1.43 x 0.89 mm

**Chorion:** smooth

**Oil globules:** single, about 0.23 mm in diameter

**Perivitelline space:** small

**Yolk:** unsegmented

**YOLK-SAC LARVAE****Fig. B**

**Hatch size:** about 3.84 mm

**Distinctive characters:** incipient dorsal appendage situated above posterior end of yolk sac; oil globule situated at anterior end of yolk sac

**Pigmentation:** pigment patch over trunk, at base of dorsal appendage; some melanophores on dorsal surface of yolk sac; 2 melanophores in ventral tail region; opposing dorsal and ventral melanophores at caudal-end region

**LARVAE****Figs. C-E**

**Distinctive characters:** body very elongate with a dorsal appendage situated a little posterior to anus

**Preanus length:** about 20% SL

**Pigmentation:** melanophores on snout, on lower jaw and over gut; 3 relatively large and another small postanal ventral melanophores; opposing dorsal and ventral melanophores at caudal-end region; dorsal appendage pigmented

**Length at flexion:** unknown

**PHOTOS**

by J.M. Rodriguez



Not sized



Not sized

**Literature:** Alemany (1997), Padoa (1956f), Russell (1976)




**CARAPIDAE**

*Echiodon dentatus* (Cuvier, 1829)




A.

Incipient dorsal appendage  
in yolk-sac larvae




B. 3.8 mm



4 ventral tail melanophores


C. 4.3 mm SL

Body very elongate with a large and  
pigmented dorsal appendage




Opposing melanophores  
at caudal end region

D. 6.4 mm



E. 16.2 mm SL


**LOPHIIDAE***Lophius piscatorius* Linnaeus, 1758**En:** Angler(=Monk) – **Fr:** Baudroie commune – **Sp:** Rape**Habitat:** Benthic, from shallow, inshore waters down to 500 m depth**Spawning season:** Between February and July**EGGS****Habitat:** pelagic**Shape:** slightly oval, embedded in a gelatinous ribbon**Size:** 2.3–3.1 mm wide, 0.62–0.70 mm high**Chorion:** smooth**Oil globules:** 1–9 oil globules, the largest 0.51–0.88 mm in diameter**Yolk:** unsegmented**YOLK-SAC LARVAE****Figs. A, B****Hatch size:** may be about 4.5 mm**Distinctive characters:** yolk sac very large and globular; rudiments of pelvic fins apparent; at end of yolk sac stage, dorsal fin shows 3 rays and pelvic fins, formed by a single ray, are strongly developed**Pigmentation:** large and branched melanophores on head and shoulder region, on gut, yolk sac and oil globules; pelvic fins heavily pigmented; postanal region unpigmented**LARVAE****Figs. C–F****Distinctive characters:** body head and trunk globose; dorsal fin formed by 5 rays and pelvic fin formed by 4 rays, strongly developed**Preanus length:** about 50% SL**Pigmentation:** early larvae similar to yolk-sac larvae; larvae of more than 11.0–12.0 mm in length develop two postanal bars of melanophores**Length at flexion:** unknown**PHOTOS**

Not available


**Literature:** Caruso (1986), Lebour (1925), Padoa (1956i), Russell (1976), Täning (1923)

**LOPHIIDAE**

*Lophius piscatorius* Linnaeus, 1758




A. 4.5 mm




B. 5.6 mm

Rudiments of pectoral and pelvic fins apparent




C. 6.5 mm




D. 7.0 mm

Pelvic fin formed by 4, strongly developed rays




E. 11.5 mm

Dorsal fin formed by 5, strongly developed rays




F. 16.5 mm

**GOBIESOCIDAE***Diplecogaster bimaculata* (Bonnaterre, 1788)**En:** Two-spotted clingfish – **Sp:** Chafarrocas**Habitat:** Benthic, attached to substratum with a thoracic sucker**Spawning season:** No information

Gobiesocidae

**EGGS****Fig. A****Habitat:** demersal**Shape:** flattened, oval in shape**Size:** 1.37–1.54 mm long, 1.08–1.24 mm wide, 0.62–0.70 mm high, attached to empty shells**Chorion:** smooth**Oil globules:** single, 0.24–0.28 mm in diameter**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** may be about 4.26 mm**Pigmentation:** rounded stellate melanophores arranged in fairly regular lines along sides of body; no melanophores on head; caudal region free of pigment**LARVAE****Figs. C–E****Distinctive characters:** body relatively elongate and slender; sucker is developed in larvae of about 6.0 mm in length.**Preanus length:** more than 70% SL**Pigmentation:** similar to yolk-sac larvae, melanophores extend backwards on caudal region**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



3.6 mm SL




4.9 mm SL

**Literature:** Briggs (1986), Russell (1976)


**GOBIESOCIDAE**

*Diplecogaster bimaculata* (Bonnaterre, 1788)




A.

Body covered with rounded stellate melanophores




B. 3.0 mm



C. 4.5 mm


Body relatively elongate and slender

No melanophores on head




D. 6.5 mm

Preanus length more than 70% SL



E. 8.0 mm

**BELONIDAE***Belone belone* (Linnaeus, 1761)**En:** Garfish – **Fr:** Orphie – **Sp:** Aguja**Habitat:** Epipelagic (neustonic), neritic**Spawning season:** February to May (Algeria)**EGGS****Fig. A****Habitat:** demersal**Shape:** spherical**Diameter:** 3.0–3.5 mm**Chorion:** filamentous**Oil globules:** no oil globule**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 9.0 mm, in an advanced stage of development**Distinctive characters:** lower jaw projects slightly beyond upper jaw**Pigmentation:** body covered with melanophores; yolk sac barely visible due to the abundant pigmentation**LARVAE****Figs. C–E****Distinctive characters:** body relatively elongate and slender, very similar to that of yolk-sac larvae; the only apparent change that occurs during development is the lengthening of the lower jaw**Preanus length:** about 45% SL**Pigmentation:** strongly pigmented; melanophores arranged along myomeres in dorsal half of body**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



13.0 mm SL




Not sized


**Literature:** Alemany (1997), Collette and Parin (1986), D'Ancona (1931e), Russell (1976)

**BELONIDAE**

*Belone belone* (Linnaeus, 1761)



A.




B. 9.0 mm




C. 13.5 mm

Lower jaw projects  
beyond upper jaw




D. 14.0 mm


Body strongly  
pigmented



E. 18.3 mm

**ZEIDAE***Zeus faber* Linnaeus, 1758**En:** John dory – **Fr:** Saint Pierre – **Sp:** Pez de San Pedro**Habitat:** Near the bottom or in midwater, from close to shore down to 400 m depth, mainly between 50 and 150 m depth**Spawning season:** Spring, in the South Mediterranean**Fig. A****EGGS****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.96–2.0 mm**Chorion:** smooth**Oil globules:** single, 0.35–0.40 mm in diameter**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** 4.3 mm**Pigmentation:** body, except caudal peduncle, covered with stellate melanophores; dorsal and ventral group of melanophores on finfold in postanal region**LARVAE****Figs. C–E****Distinctive characters:** body rhomboid-shaped; pelvic fins, well developed from young larvae on, have 6 rays; no spines on body**Preanus length:** less than 50% SL in young larvae, increases with development**Pigmentation:** similar to yolk-sac larvae; pelvic fins pigmented**Length at flexion:** 6 mm**PHOTOS**

by J.M. Rodriguez



4.3 mm SL

**Literature:** Alemany (1997), Quero (1986b), Sanzo (1956a)




**ZEIDAE**

*Zeus faber* Linnaeus, 1758




A.




B. 4.3 mm

Pigmentation extends into the finfold in postanal region




C. 2.5 mm SL

Pelvic fins well developed from young larvae on




D. 4.5 mm

Body rhomboid-shaped from young larvae on




E. 7.8 mm

No spines on body

**CENTRISCIDAE***Macroramphosus scolopax* (Linnaeus, 1758)**En:** Longspine snipefish – **Fr:** Bécasse de mer – **Sp:** Trompetero**Habitat:** Juveniles to 10 cm are epipelagic, oceanic; adults are benthic, between 50 and 150 m depth, occasionally down to 500 m**Spawning season:** Between October and March (in the subtropical northeastern Atlantic)**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.0 mm**Chorion:** smooth**Oil globules:** single, 0.2 mm in diameter**Perivitelline space:** small**YOLK-SAC LARVAE****Fig. B****Hatch size:** 3.0 mm TL**Pigmentation:** continuous line of melanophores along ventral surface of trunk and tail, from eye to caudal region; group of dorsal melanophores in postanal region; spots on top of head and snout**LARVAE****Figs. C–F****Distinctive characters:** body moderately elongate; supraorbital and occipital crests develop in larvae of 4.0–6.0 mm in length; series of spines over preopercular edge; spinous scales develop along lateral line in larvae of about 4.2 mm in length and cover entire body by 6.2 mm**Preanus length:** about 50% SL**Pigmentation:** early larvae similar to yolk-sac larvae; later larvae show a lateral line of melanophores in postanal region; pigmentation increases with development spreading onto flanks, covering most of the body, except caudal peduncle**Length at flexion:** about 4.0 mm TL**PHOTOS**

by J.M. Rodriguez



3.7 mm SL



Not sized



5.7 mm SL


**Literature:** Alemany (1997), D'Ancona (1933b), Ehrich (1986), Sparta (1936)

**CENTRISCIDAE**

*Macroramphosus scolopax* (Linnaeus, 1758)



A.




B. 3.0 mm

Melanophores on top of head and snout




C. 1.8 mm SL

Mid-lateral line of melanophores




D. 2.5 mm SL

Continuous line of melanophores, from eye to caudal region, along ventral side of trunk




E. 2.8 mm SL

Supraorbital and occipital crests




Spinous scales along lateral line


F. 4.8 mm

**SEBASTIDAE***Helicolenus dactylopterus* (Delaroche, 1809)**En:** Blackbelly rosefish – **Fr:** Sébaste chèvre – **Sp:** Gallineta**Habitat:** Benthic, on the continental slope between 200 and 1 000 m depth**Spawning season:** February and March**Fig. A****EGGS****Habitat:** pelagic**Shape:** ovoid**Size:** 0.92–0.98 × 0.88–0.93 mm**Chorion:** smooth**Oil globules:** single, about 0.2 mm in diameter**Perivitelline space:** small**Yolk:** segmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** 1.9–2.6 mm**Distinctive characters:** body moderately elongate**Pigmentation:** oil globule, dorsum of body and finfold pigmented**LARVAE****Figs. C–F****Distinctive characters:** body moderately large with a large head; 3 or 4 stout preopercular spines; few small spines on lateral ridge; supraocular ridge with small, simple spines; parietal spine well developed with secondary serrations; in young larvae, primordial fin prolonged to snout**Preanus length:** increases from about 45% to less than 60% SL with development**Pigmentation:** peritoneum, ventral side of gut and tip of lower jaw pigmented; large ventral patch of pigment close to caudal region; some dotted melanophores on head and on pectoral-fin border**Length at flexion:** 6.0–8.0 mm**PHOTOS**

by J.M. Rodriguez



Not sized



3.0 mm SL


**Literature:** Alemany (1997), Brownell (1979), Fahay (2007), Hureau and Litvinenko (1986), Täning (1961)

**SEBASTIDAE**

*Helicolenus dactylopterus* (Delaroche, 1809)




A.



B. 2.2 mm


Finfold extending to snout (in young larvae)



Body moderately elongate

Peritoneum pigmented


C. 2.3 mm SL



Tip of lower jaw pigmented


Large patch of pigment (located ventrally, close to caudal region)

D. 3.6 mm




E. 5.6 mm

Prominent head armature in larger larvae




F. 10.0 mm

**SCORPAENIDAE***Scorpaena porcus* Linnaeus, 1758**En:** Black scorpionfish – **Fr:** Rascasse brune – **Sp:** Rascacio**Habitat:** Benthic, littoral, common among rocks and algae**Spawning season:** Between June and August**Fig. A****EGGS****Habitat:** pelagic**Shape:** elliptical**Size:** 0.92 mm x 0.84 mm**Chorion:** smooth**Oil globules:** single**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 1.7 mm**Distinctive characters:** anus does not reach finfold border**Pigmentation:** unpigmented**LARVAE****Figs. C–H****Distinctive characters:** body short, increases in height with development; head relatively large and armed with spines and crests; pectoral fins strongly developed**Preanus length:** about 50% SL**Pigmentation:** peritoneum strongly pigmented; a row of postanal melanophores in young larvae, decreasing in number (to 2–4) with development; pectoral-fin edges pigmented**Length at flexion:** completed at 3.4 mm SL**PHOTOS**

by J.M. Rodriguez



3.5 mm SL



Not sized


**Literature:** Alemany (1997), Froese and Pauly (2016), Hureau and Litvinenko (1986), Sparta (1956b)

**SCORPAENIDAE**


*Scorpaena porcus* Linnaeus, 1758




A.



B. 1.4 mm




C. 1.8 mm SL



G. Dorsal view of head  
(3.4 mm SL individual)

Head armed with  
spines and crests




D. 2.9 mm SL


Pectoral fin strongly developed  
with pigmented edges




H. Detail of pigment  
pattern (4.4 mm  
individual)



E. 4.0 mm




F. 6.5 mm

**TRIGLIDAE***Eutrigla gurnardus* (Linnaeus, 1758)**En:** Grey gurnard – **Fr:** Grondin gris – **Sp:** Borracho**Habitat:** Benthic, littoral, over muddy, sandy and rocky bottoms**Spawning season:** Between January and June**EGGS****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.27–1.55 mm**Chorion:** smooth**Oil globules:** single, 0.25–0.33 mm in diameter**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. A****Hatch size:** 3.0–4.0 mm**Pigmentation:** body, yolk and oil globule covered with chromatophores; during development, pigmentation decreases and black pigment predominates; postanal ventral row of melanophores**LARVAE****Figs. B–E****Distinctive characters:** body relatively large compared to other Triglidae species; head depressed and armed with a supraorbital and a double occipital crest, and three opercular spines; pectoral fins large**Preanus length:** increases with development, from about 38% to about 50% TL**Pigmentation:** peritoneum strongly pigmented; postanal ventral row of melanophores; several melanophores on head, upper jaw and ventral abdominal region; border of pectoral-fin rays pigmented**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



5.5 mm SL




Not sized

**Literature:** Alemany (1997), Hureau (1986b), Padoa (1956p), Russell (1976), Sabatés (1988)



**TRIGLIDAE**

*Eutrigla gurnardus* (Linnaeus, 1758)




A. 3.2 mm

Body, yolk and oil globule covered with chromatophores



B. 4.7 mm




C. 8.3 mm

Lateral sides of body unpigmented


Head depressed and armed with spines and crests

Pectoral fin large with pigmented borders




D. 10.0 mm

Postanal ventral row of melanophores



E. 17.0 mm

**TRIGLIDAE***Lepidotrigla cavillone* (Lacepède, 1801)**En:** Large-scaled gurnard**Habitat:** Benthic, over mud, sand and gravel, between 30 and 450 m depth**Spawning season:** Between May and July**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.16 mm**Chorion:** smooth**Oil globules:** single, 0.21–0.22 mm in diameter**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** unknown**Distinctive characters:** yolk sac ovoid; oil globule situated at posterior edge of yolk sac; anus reaches finfold border and opens a little bit behind yolk sac**Pigmentation:** melanophores uniformly distributed over body, yolk sac and finfold**LARVAE****Figs. C–F****Distinctive characters:** body relatively high; pectoral fins large; occipital and supraorbital crests and opercular spines are present**Preanus length:** about 40% SL, increases with development**Pigmentation:** peritoneum pigmented; dorsal and ventral melanophores at about mid-postanal region; dorsal melanophore disappears when larvae reach about 4.0 mm in length**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



2.9 mm SL



5.0 mm SL


**Literature:** Alemany (1997), Fage (1918), Hureau (1986b), Padoa (1956p)

**TRIGLIDAE**


*Lepidotrigla cavillone* (Lacepède, 1801)




A.



B. Not sized




C. 2.1 mm NL




Larva scarcely pigmented


D. 3.3 mm SL



E. 6.0 mm



F. 10.0 mm

**MORONIDAE***Dicentrarchus labrax* (Linnaeus, 1758)**En:** European seabass – **Fr:** Bar européen – **Sp:** Lubina**Habitat:** Benthic, over the continental shelf**Spawning season:** Between January and March**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.20–1.51 mm**Chorion:** smooth**Oil globules:** single, 0.36–0.46 mm in diameter**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** 3.61–4.05 mm**Distinctive characters:** body elongate**Preanus length:** more than 50% SL**Pigmentation:** strongly pigmented with melanophores forming bands along body; yolk sac and oil globule pigmented**LARVAE****Figs. C–G****Distinctive characters:** body elongate and narrow**Preanus length:** more than 50% SL**Pigmentation:** continuous line of melanophores spreading from snout to caudal-fin base; dorsal melanophores restricted with development to tail end; tip of snout and lower jaw pigmented**Length at flexion:** about 6 mm SL**PHOTOS**

by J.M. Rodriguez



5.6 mm SL



6.7 mm SL


**Literature:** Bertolini (1933b), Russell (1976), Sabatés (1988), Tortonese (1986d)

**MORONIDAE**

*Dicentrarchus labrax* (Linnaeus, 1758)



A.




B. 4.5 mm



C. 5.0 mm

Tips of upper  
and lower jaw  
pigmented



Continuous line of melanophores  
spreading from snout to caudal-fin base


D. 6.0 mm



E. 9.0 mm




F. 11.0 mm



G. 17.0 mm

A, G: Bertolini (1933b); B, C: Kennedy & Fitzmaurice (1968); D-F: Russell (1976)

**SERRANIDAE***Anthias anthias* (Linnaeus, 1758)**En:** Swallowtail seaperch – **Sp:** Tres colas**Habitat:** Benthic, on the continental shelf and upper slope, down to 200 m depth**Spawning season:** Between March and August**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–E**

**Distinctive characters:** body, head and trunk height increases with development; a supraorbital crest, two series (inner and outer) of preopercular and a series of opercular spines are present; a spine of both the outer preopercular and opercular series is quite large; second ray of dorsal fin and first and second rays of pelvic fins are quite large


**Preanus length:** more than 50% SL

**Pigmentation:** some peritoneal melanophores; a relatively large spot over hindgut and another one opposite it in dorsal region; a series of melanophores between anus and caudal fin that decrease in number to 3 in late larval stages; a melanophore on caudal-fin base

**Length at flexion:** 3.3–6.0 mm SL

**PHOTOS**

by J.M. Rodriguez



1.6 mm SL



3.6 mm SL

**Literature:** Alemany (1997), Bertolini (1933b), Froese and Pauly (2016), Sabatés (1988), Tortonese (1986d)


**SERRANIDAE**

*Anthias anthias* (Linnaeus, 1758)




A. 1.9 mm SL

Dorsal melanophore at level of anus




B. 2.5 mm SL

Second dorsal-fin ray, elongate




C. 4.0 mm




D. 6.0 mm

First two pelvic-fin rays, elongate



E. 16.0 mm

**SERRANIDAE***Serranus cabrilla* (Linnaeus, 1758)**En:** Comber – **Fr:** Serran-chèvre – **Sp:** Cabrilla**Habitat:** Benthic, on the continental shelf and upper slope, down to 200 m depth**Spawning season:** Between April and July**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 0.90–0.97 mm**Chorion:** smooth**Oil globules:** single, 0.14–0.15 mm in diameter**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** 1.84–2.3 mm SL**Pigmentation:** row of dorsal melanophores extending from head to level of anus; a large melanophore on hindgut and another opposite it in dorsal region; a dorsal and ventral melanophore situated at about midpoint of the postanal region**LARVAE****Figs. C–F****Distinctive characters:** body height increases with development; three series of preopercular spines, one of which goes beyond the edge of the operculum in older larvae; an opercular spine in upper part of operculum; pelvic fins and third spiny ray of dorsal fin very elongate**Preanus length:** more than 50% SL**Pigmentation:** ventral abdominal region pigmented in young larvae and several postanal ventral melanophores may be present; pair of opposing melanophores, one dorsal and another ventral, located at midway along postanal region; caudal-fin base pigmented; spot on hindgut and another opposite it in dorsal region in young larvae; pelvic fins pigmented**Length at flexion:** flexion begins at about 4.8 mm SL and is completed at 9.7 mm SL**PHOTOS**

by J.M. Rodriguez



3.0 mm SL



6.2 mm SL

**Literature:** Bertolini (1933b), Fage (1918), Russell (1976), Sabatés (1988), Tortonese (1986d)




**SERRANIDAE**


*Serranus cabrilla* (Linnaeus, 1758)




A.



B. ca. 2.3 mm




C. 5.0 mm



D. 6.5 mm


Third spiny ray  
of dorsal fin very  
elongate

Pelvic fins very  
elongate




E. 8.0 mm

Pair of opposing  
melanophores at about  
mid-tail region



F. 16.0 mm

**SERRANIDAE***Serranus hepatus* (Linnaeus, 1758)**En:** Brown comber – **Fr:** Serran tambour – **Sp:** Merillo**Habitat:** Benthic, between 5 and 320 m depth over grasslands or sandy and muddy bottoms**Spawning season:** Between March and August**EGGS****Habitat:** pelagic**Shape:** spherical**Diameter:** 0.78 mm**Chorion:** smooth**Oil globules:** single, 0.14 mm in diameter**Yolk:** unsegmented**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–E****Distinctive characters:** body very similar to that of *Serranus cabrilla*; differences lie on weakly developed pelvic fins in *S. hepatus* and on the pigmentation pattern**Preanus length:** more than 50% SL**Pigmentation:** early larvae with two dorsal and a single ventral melanophore over finfold; three regularly spaced postanal, ventral melanophores; a melanophore over hindgut and another on cleithral symphysis; peritoneum pigmented; some melanophores on ventral surface of gut; no dorsal melanophores**Length at flexion:** almost completed at 5.7 mm SL**PHOTOS**

by J.M. Rodriguez



2.1 mm SL




6.5 mm SL

**Literature:** Alemany (1997), Bertolini (1933b), Sabatés (1988), Tortonese (1986d)


**SERRANIDAE**

*Serranus hepatus* (Linnaeus, 1758)




A. 1.5 mm SL

Two dorsal and a single ventral melanophore on finfold




B. 3.1 mm SL

3 evenly spaced postanal ventral melanophores




C. 4.1 mm SL




D. 5.5 mm SL

No dorsal melanophores

Pelvic fins weakly developed



E. 12.4 mm SL

**APOGONIDAE***Apogon imberbis* (Linnaeus, 1758)**En:** Cardinal fish – **Fr:** Coq – **Sp:** Salmonete real**Habitat:** Benthic, down to 320 m depth**Spawning season:** Between June and September**EGGS****Habitat:** pelagic**Shape:** quasi spherical**Diameter:** 0.77 mm**Chorion:** with filaments**Oil globules:** single, 0.23 mm in diameter**Yolk:** segmented**YOLK-SAC LARVAE****Hatch size:** may be about 2.5 mm SL**Pigmentation:** unpigmented**LARVAE****Figs. A-F****Distinctive characters:** body relatively short and high; a series of preopercular spines in larvae of more than 2.4 mm SL; at 6.5 mm, larvae have a long preopercular and 3 shorter opercular spines; small supraorbital crest present; swimbladder apparent**Preanus length:** 50% SL**Pigmentation:** ventral side of trunk and tail, from pectoral-fin base to mid-tail region, pigmented; group of dorsal melanophores in caudal region; two melanophores over head and some melanophores under gut; young larvae with postanal, dorsal group of melanophores; pigmentation diminishes with development**Length at flexion:** flexion begins at 3.0 mm SL**PHOTOS**

by J.M. Rodriguez



3.3 mm SL



3.5 mm SL

**Literature:** Alemany (1997), Bertolini (1933a), Tortonese (1986a)

**APOGONIDAE**

*Apogon imberbis* (Linnaeus, 1758)




A. 1.5 mm SL

Ventral side of trunk and tail pigmented




B. 2.2 mm SL




C. 2.4 mm SL

Postanal, dorsal group of melanophores in young larvae




D. 4.0 mm SL




E. 6.5 mm

Pigmentation diminishes with development



F. 14.0 mm

**CARANGIDAE***Trachurus mediterraneus* (Steindachner, 1868)**En:** Mediterranean horse mackerel – **Fr:** Chinchard à queue jaune**Sp:** Jurel mediterráneo**Habitat:** Benthopelagic, between 40 and 500 m depth**Spawning season:** No information**EGGS****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.0–1.04 mm**Chorion:** smooth**Oil globules:** single, about 0.24 mm in diameter**Yolk:** segmented**YOLK-SAC LARVAE****Fig. A****Hatch size:** may be about 1.5 mm; oil globule situated at anterior region of yolk sac**Pigmentation:** 3 or 4 dorsal melanophores in trunk region; melanophores over gut and hindgut; oil globule pigmented**LARVAE****Figs. B–F****Distinctive characters:** body height increases with development; an occipital crest and a series of preopercular spines are present; body very similar to that of *Trachurus trachurus* (differences between the two species mainly lie on the pigmentation pattern)**Preanus length:** about 50% SL**Pigmentation:** young larvae slightly pigmented, although melanophores are large; 3 or 4 dorsal preanal melanophores; melanophores over gut and hindgut; some postanal melanophores; melanophores over lateral body walls, over head, ventral tail region, under gut and lower jaw increases with development**Length at flexion:** almost completed at 4.8 mm SL**PHOTOS**

by J.M. Rodriguez



2.0 mm SL




3.1 mm SL


**Literature:** Alemany (1997), Padoa (1956e), Sabatés (1988), Smith–Vaniz (1986)

**CARANGIDAE**

*Trachurus mediterraneus* (Steindachner, 1868)




A. 1.7 mm SL




3 or 4 dorsal melanophores

B. 1.9 mm SL




Body faintly pigmented

C. 3.1 mm SL




D. 4.6 mm SL




Occipital crest

E. 6.1 mm SL



F. 8.3 mm SL

**CARANGIDAE***Trachurus trachurus* (Linnaeus, 1758)**En:** Atlantic horse mackerel – **Fr:** Chinchard d'Europe – **Sp:** Jurel**Habitat:** Benthic, over sandy bottoms between 100 and 200 m depth, occasionally down to 500 m**Spawning season:** Between October and July (Greece)**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 0.81–1.04 mm**Chorion:** smooth**Oil globules:** single, 0.19–0.28 mm in diameter**Perivitelline space:** small**Yolk:** segmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 2.5 mm**Distinctive characters:** oil globule situated at anterior edge of yolk sac (similar to *T. mediterraneus*)**Pigmentation:** melanophores are irregularly spread over body, except in caudal region; oil globule pigmented**LARVAE****Figs. C–G****Distinctive characters:** body very similar to that of *Trachurus mediterraneus*, has preopercular spines and an occipital crest (differences between the two *Trachurus* species mainly lie on the pigmentation pattern)**Preanus length:** about 50% SL**Pigmentation:** dorsal and ventral body contours of melanophores; dorsal row of about 10 melanophores, ends at about mid-way between anus and caudal fin; ventral row consists of 4 or 5 large postanal melanophores, followed by small melanophores extending nearly to posterior end of notochord; numerous melanophores on lower jaw, along ventral surface of abdomen, upper surface of gut and on head**Length at flexion:** may begin about 5.0 mm SL**PHOTOS**

by J.M. Rodriguez



2.7 mm SL



4.8 mm SL

**Literature:** Demir (1961), Froese and Pauly (2016), Padoa (1956e), Sabatés (1988), Smith–Vaniz (1986)




**CARANGIDAE**

*Trachurus trachurus* (Linnaeus, 1758)




A.




B. 2.7 mm


Body more pigmented than that of *T. mediterraneus*




C. 3.2 mm




D. 4.6 mm




E. 5.8 mm



F. 7.0 mm




G. 10.5 mm

**CORYPHAENIDAE***Coryphaena hippurus* Linnaeus, 1758**En:** Common dolphinfish – **Fr:** Coryphène commune – **Sp:** Lampuga**Habitat:** Pelagic, oceanic but it also approaches the coast**Spawning season:** Between May and September (in waters of Gran Canary Island)**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.3 mm**Chorion:** smooth**Oil globules:** single, 0.3–0.4 mm in diameter**Perivitelline space:** small**Yolk:** segmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** 3.0 mm**Distinctive characters:** body elongate; anus located at about mid-body, reaches finfold border; oil globule located at posterior end of yolk-sac**Pigmentation:** body, yolk-sac and oil globule strongly pigmented**LARVAE****Figs. C–G****Distinctive characters:** body elongate with a moderately blunt head**Preanus length:** about 60% SL**Pigmentation:** body heavily pigmented, from head to about urostyle; melanophores arranged in bars in older larvae; stellate melanophores on preorbital region and on lower jaw; melanophores under gut**Length at flexion:** 7.5–9.0 mm SL**PHOTOS**

by J.M. Rodriguez



Not sized



Not sized



Not sized


**Literature:** Collette (1986a), Fahay (2007), Froese and Pauly (2016), Mito (1960), Richards (2006)

**CORYPHAENIDAE**

*Coryphaena hippurus* Linnaeus, 1758




A.



B. 4.0 mm TL

Body, yolk-sac and oil globule strongly pigmented



C. 5.7 mm SL



D. 5.0 mm SL

Larvae strongly pigmented




E. 7.1 mm SL




F. 9.5 mm SL

Melanophores arranged in bars in older larvae



G. 11.0 mm SL

**BRAMIDAE***Brama brama* (Bonnaterre, 1788)**En:** Atlantic pomfret – **Fr:** Grande castagnole – **Sp:** Japuta**Habitat:** Pelagic, offshore**Spawning season:** August and September**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.50–1.60 mm**Chorion:** smooth**Oil globules:** single, oval, 0.40 x 0.32 mm**Perivitelline space:** reduced**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 4.8 mm**Distinctive characters:** relatively large head, rounded gut and long tail**Pigmentation:** head, yolk sac, pectoral and caudal fins pigmented; two prominent pigment patches in tail region, one dorsal and another ventral that extend on finfold**LARVAE****Figs. C–G****Distinctive characters:** body in early stages similar to that of yolk-sac larvae, although laterally compressed; during development, head and trunk, globose, increase in size and tail decreases in length; pectoral fins relatively large and rounded**Preanus length:** about 30% SL in early larvae, increases with development**Pigmentation:** head and gut heavily pigmented; tail region free of pigment; pectoral and caudal fins pigmented**Length at flexion:** flexion completed at 4.9 mm**PHOTOS**

by J.M. Rodriguez



3.3 mm SL



3.9 mm SL


**Literature:** Alemany (1997), Fahay (2007), Haedrich (1986), Padoa (1956d), Sabatés (1988), Schmidt (1918)

**BRAMIDAE**

*Brama brama* (Bonnaterre, 1788)




A.




B. 4.8 mm



C. 3.0 mm SL



D. 3.0 mm SL




Trunk and gut globose and heavily pigmented

Tail region free of pigment


E. 3.3 mm SL



G. 5.5 mm (dorsal view)



F. 4.9 mm SL

**SPARIDAE*****Boops boops*** (Linnaeus, 1758)**En:** Bogue – **Fr:** Bogue – **Sp:** Boga**Habitat:** Benthic, down to 200 m, occasionally to 350 m depth**Spawning season:** April and May**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 0.89 mm**Chorion:** smooth**Oil globules:** single, 0.2 mm in diameter**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** less than 2 mm**Distinctive characters:** body elongate, anus does not reach finfold edge; oil globule pigmented**Pigmentation:** opposing dorsal and ventral bars of pigment at about mid-tail region; a series of postanal, ventral melanophores and one or two under urostyle**LARVAE****Figs. C-G****Distinctive characters:** body elongate with a prominent swimbladder; two series of preopercular spines in older larvae**Preanus length:** about 40% SL**Pigmentation:** young larvae, same as yolk-sac larvae; in older larvae, postanal ventral bar of melanophores disappears; peritoneum pigmented; a single melanophore under hindgut; melanophores under gut and on head**Length at flexion:** completed at 7.6 mm SL**PHOTOS**

by J.M. Rodriguez



3.1 mm SL



6.1 mm SL



8.1 mm SL


**Literature:** Alemany (1997), Bauchot and Hureau (1986), De Gaetani (1937), Ranzi (1933), Sabatés (1988)

**SPARIDAE**


*Boops boops* (Linnaeus, 1758)




A.




B. 3.2 mm




C. 5.2 mm




D. 6.0 mm



E. 7.1 mm



F. 8.2 mm



G. 11.9 mm


**SPARIDAE***Diplodus sargus* (Linnaeus, 1758)

The genus *Diplodus* includes several species whose larvae are very similar, mainly during the first stages of development. The only species for which these stages are described is *D. sargus*.

**En:** White seabream – **Fr:** Sar commun – **Sp:** Sargo

**Habitat:** Benthic, down to 50 m depth

**Spawning season:** Between March and June

**Fig. A****EGGS**

**Habitat:** pelagic

**Shape:** spherical

**Diameter:** about 0.90 mm

**Chorion:** smooth

**Oil globules:** single, about 0.2 mm in diameter

**Perivitelline space:** small

**Yolk:** unsegmented

**YOLK-SAC LARVAE****Figs. B, C**

**Hatch size:** about 2.5 mm

**Distinctive characters:** body elongate, anus reaches finfold edge

**Pigmentation:** scattered melanophores over dorsum, no ventral melanophores; with development, dorsal melanophores disappear and ventral melanophores appear; oil globule pigmented

**LARVAE****Figs. D–G**

**Distinctive characters:** body elongate (the most among the species belonging to this genus); older larvae with two series of preopercular spines

**Preanus length:** about 30% SL in young larvae, increases with development

**Pigmentation:** postanal ventral row of melanophores; peritoneum, ventral region of gut, occipital and shoulder regions, pigmented

**Length at flexion:** about 5.6 mm

**PHOTOS**

by J.M. Rodriguez



*Diplodus* sp.  
4.5 mm SL




*Diplodus* sp.  
5.4 mm SL

**Literature:** Alemany (1997), Bauchot and Hureau (1986), Brownell (1979), Sabatés (1988)




**SPARIDAE**

*Diplodus sargus* (Linnaeus, 1758)



A.




B. 2.4 mm NL




C. 3.4 mm SL

Body elongate



Postanal ventral row of melanophores

D. 3.3 mm SL




Ventral region of gut pigmented

E. 4.5 mm SL




E. Dorsal view




Two series of preopercular spines

Pigmentation increases with development

F. 8.0 mm SL



G. 14.4 mm SL

**SPARIDAE***Pagellus bogaraveo* (Brünnich, 1768)**En:** Blackspot(=red) seabream – **Fr:** Dorade rose**Sp:** Besugo**Habitat:** Benthic, down to 400 m depth**Spawning season:** All year round, with a peak between January and May**Fig. A****EGGS****Habitat:** pelagic**Shape:** spherical**Diameter:** about 1.2 mm**Chorion:** smooth with a double capsule**Oil globules:** single, about 0.28 mm in diameter**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 3.8 mm**Distinctive characters:** body elongate; anus reaches finfold edge; oil globule pigmented, situated at posterior edge of yolk sac**Pigmentation:** dorsal melanophores between head and mid-trunk; two opposing groups of melanophores at posterior tail region**LARVAE****Figs. C–G****Distinctive characters:** body relatively elongate**Preanus length:** about one third of SL, increases with development**Pigmentation:** some occipital melanophores; peritoneum pigmented; at least two dorsal melanophores on tail; pre- and postanal melanophores on ventral side of body**Length at flexion:** about 6.0 mm**PHOTOS**


by J.M. Rodriguez



2.0 mm SL



3.5 mm SL



6.2 mm SL


**Literature:** Alemany (1997), Bauchot and Hureau (1986), De Gaetani (1934)

**SPARIDAE**

*Pagellus bogaraveo* (Brünnich, 1768)



A.




B. 3.8 mm




C. 3.5 mm

At least two postanal dorsal spots




D. 6.0 mm




E. 6.5 mm


Pre- and postanal ventral melanophores



F. 7.7 mm



G. 13.0 mm

**SPARIDAE***Pagrus pagrus* (Linnaeus, 1758)**En:** Red porgy – **Fr:** Pagre rouge – **Sp:** Pargo**Habitat:** Benthic, over sandy and rocky bottoms, down to 200 m depth**Spawning season:** April to June**Fig. A****EGGS****Habitat:** pelagic**Shape:** spherical**Diameter:** 0.89–0.93 mm**Chorion:** smooth**Oil globules:** single, 0.18–0.20 mm in diameter**Perivitelline space:** narrow**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** between 1.92 and 3.04 mm NL**Distinctive characters:** body relatively elongate; anus does not reach finfold border; oil globule situated at posterior end of yolk sac**Pigmentation:** melanophores along dorsal and ventral midlines of body; oil globule pigmented**LARVAE****Figs. C–F****Distinctive characters:** body, head and trunk quite high; an occipital crest from young larvae on (specific character of this Sparidae species); preopercular spines well developed; older larvae show a supraorbital spinous arch**Preanus length:** less than 50% SL**Pigmentation:** scantily pigmented; peritoneum pigmented; about 6 postanal, ventral melanophores; a melanophore on head and another on hindgut**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



2.9 mm SL




7.0 mm SL


**Literature:** Alemany (1997), Bauchot and Hureau (1986), Machinandiarena *et al.* (2003), Ranzi (1933)

**SPARIDAE**


*Pagrus pagrus* (Linnaeus, 1758)




A.



B. 2.4 mm SL



C. 2.7 mm SL




D. 3.4 mm SL


Large preopercular spines

Body scantily pigmented


Supraorbital spinous arch  
The only Sparidae species with an occipital crest



E. 9.0 mm



F. 15.5 mm

**SPARIDAE***Spondyliosoma cantharus* (Linnaeus, 1758)**En:** Black seabream – **Fr:** Dorade grise – **Sp:** Chopa**Habitat:** Benthic, on rocky or sandy bottoms and *Posidonia* beds, down to 50 m (young) and 300 m (adults) depth**Spawning season:** Between February and May**EGGS****Habitat:** demersal**Shape:** spherical**Diameter:** 1.0–1.2 mm**Chorion:** smooth**Oil globules:** single, 0.20–0.25 mm in diameter**Perivitelline space:** moderate**Yolk:** unsegmented**YOLK-SAC LARVAE****Figs. A, B****Hatch size:** about 2.2 mm**Distinctive characters:** oil globule situated at posterior edge of yolk sac**Pigmentation:** yolk sac strongly pigmented; oil globule pigmented; row of ventral melanophores from anus to about mid-postanal region in early yolk-sac larvae, reaches caudal region later; melanophores on head**LARVAE****Figs. C–E****Distinctive characters:** body relatively elongate and slender, becoming more robust with development; small preopercular spines appearing at about 5.0 mm SL**Preanus length:** about 50% SL until after flexion**Pigmentation:** early stages similar to yolk-sac larvae; melanophores extend to lateral sides in early larvae and posteriorly to dorsal sides of body; gut pigmented**Length at flexion:** about 6.0 mm**PHOTOS**

by J.M. Rodriguez



3.7 mm SL



4.8 mm SL








5.8 mm SL

**Literature:** Bauchot and Hureau (1986), Camus and Besseau (1986), Fage (1918), Ranzi (1933), Russell (1976)

**SPARIDAE**

*Spondyliosoma cantharus* (Linnaeus, 1758)



**CENTRACANTHIDAE***Spicara smaris* (Linnaeus, 1758)**En:** Picarel – **Fr:** Picarel – **Sp:** Caramel**Habitat:** Benthic, on muddy and sandy bottoms, down to 130 m depth**Spawning season:** Between February and May**EGGS****Fig. A****Habitat:** demersal, laid in nests excavated in detritic or sandy bottoms**Shape:** assumes the form of a spherical cap**Size:** about 0.89 mm in diameter by about 0.72 mm in height**Chorion:** thick and resistant**Oil globules:** single, 0.19–0.21 mm in diameter**Perivitelline space:** no information**Yolk:** transparent**YOLK-SAC LARVAE****Fig. B****Hatch size:** not known**Distinctive characters:** body elongate; short preanus length; oil globule situated at center of yolk sac**Pigmentation:** unpigmented**LARVAE****Figs. C–G****Distinctive characters:** body elongate and slender becoming more robust with development; small preopercular and opercular spines appear with development**Preanus length:** increases with development from about 30% to more than 40% SL**Pigmentation:** dotted occipital melanophore; two peritoneal pigment patches; about 15 postanal ventral melanophores; 1–3 dotted melanophores under urostyle that move to caudal-fin base in later larvae; two melanophores between dorsal and caudal fins in later larvae**Length at flexion:** flexion almost completed at 6.5 mm SL**PHOTOS**

by J.M. Rodriguez



3.4 mm SL



6.8 mm SL

**Literature:** Alemany (1997), Froese and Pauly (2016), Montalenti (1937b), Tortonese (1986b)



**CENTRACANTHIDAE**

*Spicara smaris* (Linnaeus, 1758)




A.



Oil globule situated at the center of yolk sac


Short preanus length

B. Not sized




Larvae scantily pigmented

C. 3.3 mm SL




About 15 postanal ventral melanophores

D. 3.9 mm SL




Two peritoneal patches of pigment


E. 5.3 mm SL



F. 6.5 mm SL



G. 8.3 mm SL

**MULLIDAE***Mullus barbatus* Linnaeus, 1758**En:** Red mullet – **Fr:** Rouget de vase – **Sp:** Salmonete de fango**Habitat:** Benthic, on muddy bottoms of the continental shelf, between 10 and 500 m depth**Spawning season:** Between April and August**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–E****Distinctive characters:** body relatively elongate**Preanus length:** about 37% SL**Pigmentation:** increases with development; early larvae show an occipital melanophore, peritoneum strongly pigmented, a row of postanal ventral melanophores, from anus to about two-thirds of tail region, a short series of mid-lateral melanophores and a single melanophore under urostyle; later larvae with several melanophores over head and on dorsal region of trunk: entire surface of gut, except its ventral side, pigmented**Length at flexion:** flexion occurs between about 3.5 and 5.2 mm SL**PHOTOS**

by J.M. Rodriguez



3.3 mm SL



5.5 mm SL




8.3 mm SL

**Literature:** Alemany (1997), Hureau (1986a), Montalenti (1937a), Sabatés (1988)

**MULLIDAE**


*Mullus barbatus* Linnaeus, 1758

No dorsal melanophores  
in tail region




A. 2.6 mm SL

Short series of mid-lateral  
melanophores




B. 2.9 mm SL




C. 3.7 mm SL


Postanal row of ventral  
melanophores



D. 4.7 mm SL



E. 5.2 mm SL

**MULLIDAE***Mullus surmuletus* Linnaeus, 1758**En:** Surmullet – **Fr:** Rouget de roche – **Sp:** Salmonete de roca**Habitat:** Benthic, over rocky bottoms, down to 400 m depth**Spawning season:** Between May and July**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 0.81–0.91 mm**Chorion:** smooth**Oil globules:** single, 0.23–0.25 mm in diameter**Perivitelline space:** small**Yolk:** segmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 2.83 mm**Distinctive characters:** yolk sac projected beyond snout; oil globule located at anterior edge of yolk sac; anus situated close behind to posterior edge of yolk sac**Pigmentation:** melanophores scattered over body; oil globule strongly pigmented**LARVAE****Figs. C–F****Distinctive characters:** body relatively elongate**Preanus length:** about 30% SL**Pigmentation:** at about 4.0 mm SL shows a ventral row of 10–12 postanal melanophores extending from anus to about two-thirds of tail region; 2 or 3 melanophores over and below urostyle; mid-lateral line of melanophores; peritoneum pigmented; later larvae develop dorsal and dorsolateral rows of melanophores**Length at flexion:** it occurs between less than 4.0 and 7.6 mm SL**PHOTOS**

by J.M. Rodriguez



2.8 mm SL



5.4 mm SL


**Literature:** Alemany (1997), Hureau (1986a), Montalenti (1937a), Russell (1976)

**MULLIDAE**

*Mullus surmuletus* Linnaeus, 1758



A.




B. Not sized




C. 2.3 mm SL

Mid-lateral line of melanophores,  
longer than that of *M. barbatus*




D. 3.1 mm SL




E. 3.6 mm SL

Dorsal side of body  
strongly pigmented  
(older larvae)



F. 7.6 mm SL

**CEPOLIDAE***Cepola macrophthalmma* (Linnaeus, 1758)**En:** Cardinal fish – **Fr:** Coq – **Sp:** Salmonete real**Habitat:** Benthic, down to 320 m depth**Spawning season:** Spring to autumn**EGGS****Habitat:** pelagic**Shape:** quasi spherical**Diameter:** 0.77 mm**Chorion:** with filaments**Oil globules:** single, 0.23 mm in diameter**Yolk:** segmented**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–D****Distinctive characters:** body relatively short and high; a series of preopercular spines in larvae more than 2.4 mm SL; at 6.5 mm, larvae have a long preopercular spine, 3 shorter opercular spines and a small supraorbital crest; swimbladder apparent**Preanus length:** 50% SL**Pigmentation:** ventral side of trunk pigmented, from pectoral-fin base to mid-tail region; group of dorsal melanophores in caudal region; two melanophores over head and some melanophores under gut; pigmentation diminishes with development**Length at flexion:** flexion begins at 3.0 mm SL**PHOTOS**

by J.M. Rodriguez



2.0 mm SL




3.5 mm SL


**Literature:** Alemany (1997), Russell (1976), Tortonese (1986c)

**CEPOLIDAE**


*Cepola macrophthalmal* (Linnaeus, 1758)



A. 1.7 mm SL




B. 3.0 mm



C. 3.9 mm



D. 7.2 mm SL

**MUGILIDAE***Mugil cephalus* Linnaeus, 1758**En:** Flathead grey mullet – **Fr:** Mulet à grosse tête – **Sp:** Pardete**Habitat:** Pelagic, inshore, entering lagoons and estuaries**Spawning season:** Between July and October**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 0.72 mm**Chorion:** smooth, fine raised striations**Oil globules:** single, 0.28 mm in diameter**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 2.5 mm**Pigmentation:** heavily pigmented, except caudal region; a dorsal and another ventral row of about 4 dotted melanophores each at caudal region; oil globule strongly pigmented**LARVAE****Fig. C–F****Distinctive characters:** body relatively stubby with a large, bulky gut, moderate head and small mouth; no swimbladder**Preanus length:** up to 70% SL**Pigmentation:** strongly pigmented, except lateral sides of head and caudal region where dotted melanophores persist; pigment is heaviest on dorsum, dorsal surface of gut and on postanal ventral region; apparent mid-lateral row of melanophores**Length at flexion:** 4.0–5.0 mm SL**PHOTOS**

by J.M. Rodriguez



2.4 mm SL



2.5 mm SL



7.5 mm SL

**Literature:** Alemany (1997), Ben-Tuvia (1986), Fahay (1983), Froese and Pauly (2016), Sanzo (1936)




**MUGILIDAE**

*Mugil cephalus* Linnaeus, 1758



A.




B. ca. 2.5 mm



C. 2.1 mm SL

Body, except caudal region,  
strongly pigmented



D. 2.3 mm SL


Dorsal and ventral  
dotted melanophores on  
caudal region

Apparent mid lateral row  
of melanophores




E. 3.4 mm SL

No swimbladder



F. 5.1 mm

**POMACENTRIDAE***Chromis chromis* (Linnaeus, 1758)**En:** Damselfish**Habitat:** Littoral, mainly in rocky areas, between 3 and 35 m depth**Spawning season:** Between May and August**Fig. A****EGGS****Habitat:** demersal**Shape:** ovoid, attached to substrate with filaments**Size:** 0.85–0.90 x 0.70–0.72 mm**Oil globules:** single, 0.20 mm in diameter**Perivitelline space:** small**Yolk:** segmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** 2.6 mm SL**Distinctive characters:** body relatively slender; oil globule situated ventrally in yolk sac**Preanus length:** about 33% SL**Pigmentation:** yolk sac pigmented; row of postanal ventral melanophores; occipital melanophores; melanophores dorsal and ventral to urostyle**LARVAE****Fig. C–F****Distinctive characters:** body of early larvae similar to that of yolk-sac larvae, becoming short and robust with development; gut triangular-shaped**Preanus length:** slightly less than 50% SL**Pigmentation:** similar to yolk-sac larvae in earlier larvae; with development, postanal pigmentation reduces to a transversal band located around mid-tail region; posteriorly, melanophores appear on mid-lateral line and on dorsum, at same level of ventral group; peritoneum pigmented; melanophores on head**Length at flexion:** begins at 3.60 mm and is completed at 4.75 mm SL**PHOTOS**

by J.M. Rodriguez



1.9 mm SL









7.1 mm SL

**Literature:** Alemany (1997), Fage (1918), Froese and Pauly (2016), Padoa (1956k), Quignard and Pras (1986b)


**POMACENTRIDAE**

*Chromis chromis* (Linnaeus, 1758)




**LABRIDAE***Coris julis* (Linnaeus, 1758)**En:** Rainbow wrasse – **Fr:** Girelle – **Sp:** Julia**Habitat:** Benthic, littoral, over rocky bottoms or eelgrass beds, down to 120 m depth**Spawning season:** Between April and August**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 0.60–0.67 mm**Chorion:** smooth**Oil globules:** single, 0.12–0.16 mm in diameter**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 2.2 mm**Distinctive characters:** oil globule situated at anterior edge of yolk sac**Pigmentation:** several dorsal melanophores decrease in number to 2 and migrate to finfold with development; row of ventral melanophores along ventral side of trunk and tail, ending in a large melanophore**LARVAE****Figs. C–G****Distinctive characters:** body elongate; gut with a prominent loop**Preanus length:** about 50% SL**Pigmentation:** in young larvae, 2 dorsal melanophores on finfold which, in older larvae, remain on dorsal fin; two large ventral melanophores, one over gut, close to gut loop and another in caudal region**Length at flexion:** completed at 9.8 mm**PHOTOS**


by J.M. Rodriguez



2.2 mm SL



2.7 mm SL




5.6 mm SL


**Literature:** Alemany (1997), Quignard and Pras (1986a), Sabatés (1988)

**LABRIDAE**


*Coris julis* (Linnaeus, 1758)



A.




B. 2.2 mm




C. 2.5 mm SL

Two melanophores on dorsal finfold and later on dorsal fin




D. 3.1 mm SL

Two large ventral melanophores




E. 5.0 mm SL

Prominent gut loop



F. 6.4 mm SL



G. 9.8 mm SL

**LABRIDAE***Ctenolabrus rupestris* (Linnaeus, 1758)**En:** Goldsinny-wrasse – **Fr:** Rouquié – **Sp:** Tabernero**Habitat:** Benthic, on rocky, weed-covered shores, from 1 to 50 m depth**Spawning season:** Between January and July**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 0.72–1.01 mm**Chorion:** smooth**Oil globules:** none**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** 1.95–2.19 mm SL**Distinctive characters:** anus located some distance behind yolk sac**Pigmentation:** in newly hatched larvae, melanophores scattered over most of the body; during yolk-sac larval development, all melanophores from dorsal side of body disappear, while on ventral side, only 4 melanophores remain: over gut, hindgut, mid-tail and caudal region**LARVAE****Figs. C–H****Distinctive characters:** body relatively elongate and slender, becoming more robust with development**Preanus length:** about 50% SL**Pigmentation:** similar to yolk-sac larvae in earlier larvae; later, two melanophores appear under gut increasing in number with development; peritoneal pigment also develops**Length at flexion:** at less than 4.82 mm SL**PHOTOS**

by J.M. Rodriguez



2.6 mm SL



4.4 mm SL




7.1 mm SL


**Literature:** Fives (1976), Quignard and Pras (1986a), Russell (1976), Sabatés (1988), Sparta (1956a)

**LABRIDAE**

*Ctenolabrus rupestris* (Linnaeus, 1758)



A.




B. 2.2 mm



C. 3.0 mm TL


Melanophores over gut, hindgut,  
mid-tail and ventral caudal fin




D. 5.0 mm TL




H. Dorsal view of  
head (7.5 mm TL  
larva)



E. 6.9 mm TL




F. 9.0 mm TL



Melanophores on  
ventral side of gut

G. 10.0 mm TL

**LABRIDAE***Labrus mixtus* Linnaeus, 1758**En:** Cuckoo wrasse – **Fr:** Vieille coquette – **Sp:** Gallano**Habitat:** Benthic, on algal zones of rocky, weed-covered shores, from 2 to 200 m depth**Spawning season:** Between March and June**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–E****Distinctive characters:** body relatively elongate and slender, becoming more robust with development**Preanus length:** about 50% SL**Pigmentation:** five evenly spaced melanophores along dorsal contour of body, three along ventral postanal region, one on caudal fin, several on lower jaw and along dorsal and ventral contours of abdomen; a single melanophore on head**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



3.6 mm SL



4.3 mm SL








8.8 mm SL


**Literature:** Froese and Pauly (2016), Fives (1976), Quignard and Pras (1986a), Russell (1976)



**LABRIDAE**

*Labrus mixtus* Linnaeus, 1758



**LABRIDAE***Symphodus melops* (Linnaeus, 1758)**En:** Corkwing wrasse – **Fr:** Crénilabre mélops – **Sp:** Porredana**Habitat:** Benthic, littoral, near rocks and eelgrass beds, from 1 to 50 m depth**Spawning season:** Between March and May (possibly June)**EGGS****Habitat:** demersal, attached to weed in nests**Shape:** spherical**Diameter:** 0.75–0.80 mm**Oil globules:** none**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. A****Hatch size:** 2.4–2.85 mm SL**Distinctive characters:** body relatively elongate**Preanus length:** about 50% SL**Pigmentation:** two rows of dorsal melanophores from behind eye to about midway along tail region; three postanal ventral melanophores**LARVAE****Figs. B–D****Distinctive characters:** body relatively elongate and slender, becoming more robust with development**Preanus length:** about 50% SL**Pigmentation:** a small number of melanophores on head; two parallel rows of dorsal melanophores; lower jaw pigmented; lateral sides of body, except caudal region, and anal fin pigmented; 3 or 4 melanophores between caudal and anal fins; melanophore on caudal-fin base**Length at flexion:** probably begins at about 4.0 mm SL**PHOTOS**

by J.M. Rodriguez



2.2 mm SL




4.6 mm SL


**Literature:** Fives (1976), Quignard (1967), Quignard and Pras (1986a), Russell (1976)

**LABRIDAE**

*Symphodus melops* (Linnaeus, 1758)



A. 2.9 mm SL



A. Dorsal view


Two parallel rows of dorsal melanophores




B. 3.2 mm SL



C. Dorsal view of head



C. 4.5 mm TL



Anal fin pigmented

D. 6.2 mm TL

**AMMODYTIDAE***Gymnammodytes cicereus* (Rafinesque, 1810)**En:** Mediterranean sand eel – **Fr:** Cicerelle de Méditerranée**Sp:** Barrinaire**Habitat:** Littoral, benthic, on sandy bottoms**Spawning season:** Between November and January**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed


**LARVAE****Figs. A–F****Distinctive characters:** body elongate and slender in all larval stages**Preanus length:** more than 60% SL**Pigmentation:** 8 dorsal melanophores evenly distributed along body; several melanophores on dorsal side of gut; 2 or 3 postanal ventral melanophores; pigmentation diminishes with development**Length at flexion:** almost completed at 11.0 mm SL**PHOTOS**

Not available

**Literature:** Aboussouan (1964), Alemany (1997), Froese and Pauly (2016), Reay (1986), Sabatés *et al.* (1990)

**AMMODYTIDAE**

*Gymnamodytes cicereus* (Rafinesque, 1810)



A. 2.5 mm SL

8 dorsal melanophores  
evenly distributed along body




2 or 3 postanal ventral  
melanophores

B. 3.1 mm SL



C. 4.1 mm SL




D. 7.7 mm SL



E. 11.0 mm SL



F. 12.9 mm SL

**TRACHINIDAE***Echiichthys vipera* (Cuvier, 1829)**En:** Lesser weever**Habitat:** Littoral and benthic, on sandy, muddy or gravelly bottoms, from a few meters to about 150 m depth**Spawning season:** No information**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.0–1.37 mm**Chorion:** smooth**Oil globules:** multiple, 6–30**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 3.3 mm**Distinctive characters:** body relatively elongate; anus situated close behind yolk sac; oil globules situated ventrally in yolk sac; pelvic fins well developed**Pigmentation:** dorsal and ventral rows of melanophores extending posteriorly a little beyond mid-tail region in newly hatched larvae; pigmentation soon becomes characteristic with melanophores on snout, head, peritoneum and two bars of pigment, one in the anal region and another midway along postanal region**LARVAE****Figs. C–F****Distinctive characters:** body relatively elongate and slender, becoming stout and deep with development; a series of preopercular and opercular spines; pelvic fins well developed from young larvae on**Preanus length:** about 45% SL**Pigmentation:** anal bars of pigment disappear in larvae of about 4.5 mm SL, with only a group of melanophores remaining; melanophores on head and on shoulder; pelvic fins and peritoneum strongly pigmented**Length at flexion:** flexion begins at about 6 mm SL**PHOTOS**

by J.M. Rodriguez



2.7 mm SL



7.7 mm SL


**Literature:** Alemany (1997), Froese and Pauly (2016), Padoa (1956a), Russell (1976)

**TRACHINIDAE**

*Echiichthys vipera* (Cuvier, 1829)




A.




B. 3.3 mm

Oil globules situated ventrally in yolk sac



C. 3.5 mm



D. 5.2 mm


Pelvic fins well developed

Peritoneum strongly pigmented




E. 9.2 mm

Only a group of melanophores in tail region in older larvae



F. 13.5 mm

**TRACHINIDAE***Trachinus draco* Linnaeus, 1758**En:** Greater weever – **Fr:** Grande vive – **Sp:** Escorpión**Habitat:** Benthic, littoral, over muddy, sandy or gravel bottoms, down to 150 m depth**Spawning season:** No information**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 0.96–1.11 mm**Chorion:** smooth**Oil globules:** single, 0.19–0.23 mm in diameter**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 3.2 mm**Distinctive characters:** oil globule and anus situated at anterior and posterior edges of yolk sac, respectively; primordial fin globose reaching the snout**Pigmentation:** melanophores on snout, two behind eye, on dorsal anterior part of body, and near anus; row of melanophores on postanal ventral region with a postanal bar at midway between anus and tail end**LARVAE****Figs. C–F****Distinctive characters:** body relatively elongate and slender, becoming more robust with development; pelvic fins appear at about 4 mm SL; young larvae, with primordial fin globose, reaching the snout; older larvae have 5 preopercular spines**Preanus length:** about 40% SL**Pigmentation:** early stages similar to yolk-sac larvae; with development, the origin of ventral postanal row of melanophores moves backwards and melanophores decrease in number; postanal bar of melanophores disappears and peritoneal pigment increases; pelvic fins strongly pigmented**Length at flexion:** almost completed at 5.1 mm**PHOTOS**

by J.M. Rodriguez



3.3 mm SL



7.0 mm SL

**Literature:** Alemany (1997), Munk and Nielsen (2005), Padoa (1956n), Russell (1976), Sabatés (1988)



**TRACHINIDAE**

*Trachinus draco* Linnaeus, 1758




A.




B. 3.0 mm

Primordial fin globose, reaching the snout

Row of postanal ventral melanophores




C. 3.5 mm




Pelvic fins well developed and strongly pigmented

D. 5.3 mm SL


Body scarcely pigmented




E. 5.7 mm SL




F. 12.0 mm

**URANOSCOPIDAE***Uranoscopus scaber* Linnaeus, 1758**En:** Stargazer – **Fr:** Uranoscope – **Sp:** Rata**Habitat:** Benthic on sandy or muddy bottoms, burrowing in sediment, on the continental shelf and upper slope, between 15 and 400 m depth**Spawning season:** Between April and August**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.6–2.0 mm**Chorion:** sculptured with hexagonal reticules**Oil globules:** none**Perivitelline space:** small**Yolk:** segmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 4.0 mm**Pigmentation:** trunk heavily pigmented, only caudal peduncle unpigmented**LARVAE****Figs. C–E****Distinctive characters:** body short with an extremely high preanus region**Preanus length:** more than 50% SL**Pigmentation:** body, except caudal peduncle, covered with small melanophores**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



2.4 mm SL




4.8 mm SL


**Literature:** Fage (1918), Hureau (1986c), Padoa (1956q), Sabatés (1988)

**URANOSCOPIDAE**

*Uranoscopus scaber* Linnaeus, 1758




A.




B. ca. 4.0 mm

Body strongly pigmented




C. 3.2 mm SL


Body short with extremely high preanus region




D. 3.3 mm SL



E. 8.0 mm



E. Dorsal view

**BLENNIIDAE***Blennius ocellaris* Linnaeus, 1758**En:** Butterfly blenny – **Sp:** Torillo**Habitat:** Benthic, on hard bottoms, between 30 and 100 m depth**Spawning season:** April (Marseille, France)**Fig. A****EGGS****Habitat:** demersal, attached to shells and hollow objects**Shape:** sub-spherical**Diameter:** 1.12–1.20 mm**Oil globules:** some**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Hatch size:** about 4.5 mm**Distinctive characters:** pectoral fin apparent**Pigmentation:** melanophores on head, snout and peritoneum; pectoral fin strongly pigmented with melanophores arranged in longitudinal rows between incipient rays; row of melanophores on ventral tail end region**LARVAE****Figs. B–F****Distinctive characters:** body relatively elongate and slender becomes more robust with development; pectoral fin large (reaching beyond anus) and rounded in shape, with 12 rays**Preanus length:** about 33% SL**Pigmentation:** pectoral fin heavily pigmented with melanophores located between fin rays; some melanophores on head; 5 or 6 postanal ventral melanophores at posterior half of postanal region; peritoneum heavily pigmented**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



3.6 mm SL



3.9 mm SL


**Literature:** Fives (1986), Ford (1922), Padoa (1956b), Russell (1976), Zander (1986)

**BLENNIIDAE**


*Blennius ocellaris* Linnaeus, 1758



A.




B. 4.6 mm




C. 3.6 mm SL

Peritoneum heavily pigmented




D. 5.0 mm




5 or 6 postanal ventral melanophores at posterior half of postanal region

E. 8.5 mm

Large pectoral fin with 12 rays



F. 13.3 mm

**BLENNIIDAE***Lipophrys pholis* (Linnaeus, 1758)**En:** Shanny**Habitat:** Benthic, in shallow waters of rocky coasts, often in rock-pools, sometimes among algae**Spawning season:** Between April and August**EGGS****Habitat:** demersal**Shape:** hemispherical**Diameter:** 1.18–1.6 mm**Chorion:** smooth**Oil globules:** several**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. A****Hatch size:** about 4.4 mm**Pigmentation:** pectoral fins heavily pigmented; melanophores on head and snout; peritoneum pigmented; postanal region unpigmented**LARVAE****Figs. B–E****Distinctive characters:** body relatively elongate; pectoral fins very large with 13 rays; apparent teeth in older larvae**Preanus length:** less than 50% SL**Pigmentation:** pectoral fins heavily pigmented; melanophores on head and close to operculum; peritoneum and base of caudal fin pigmented; row of melanophores on ventral tail end region**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



4.6 mm SL




9.0 mm SL

**Literature:** Alemany (1997), Fives (1986), Ford (1922), Padoa (1956b), Russell (1976), Zander (1986)


**BLENNIIDAE**

*Lipophrys pholis* (Linnaeus, 1758)




A. 4.4 mm

Postanal region unpigmented  
in yolk-sac larvae




B. 5.0 mm

Row of melanophores on  
ventral tail-end region




C. 5.5 mm

Pectoral fin very  
large with 13 rays




D. 9.0 mm

Teeth apparent in  
larger larvae



E. 17.5 mm TL

**BLENNIIDAE***Parablennius pilicornis* (Cuvier, 1829)**En:** Ringneck blenny**Habitat:** Benthic, littoral**Spawning season:** No information**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–F****Distinctive characters:** body elongate; pectoral fins fairly large; preopercular spines appear at 4.0 mm SL**Preanus length:** 36% of SL in young larvae, diminishes to 32% in older larvae**Pigmentation:** young larvae with only one stellate melanophore on head (many in older larvae); dorsal surface of gut heavily pigmented, pigmentation of abdominal region increases with development; a single melanophore ventrally over hindgut; ventral row of melanophores from anus to notochord tip; older larvae show a melanophore on caudal-fin base; dorsal pigmentation in tail region appears in larvae larger than 13 mm SL**Length at flexion:** between 5 and 6 mm SL**PHOTOS**

by J.M. Rodriguez



2.8 mm SL



3.3 mm SL




8.4 mm SL

**Literature:** Olivar (1986), Olivar and Fortuño (1991), Zander (1986)




**BLENNIIDAE**

*Parablennius pilicornis* (Cuvier, 1829)



A. 2.4 mm SL

Ventral row of melanophores  
from anus to notochord tip




B. 4.1 mm SL

A single melanophore  
on hindgut




C. 5.6 mm SL


Preopercular spines in  
larvae larger than 4 mm



D. 8.6 mm SL




E. 11.1 mm SL



F. 13.4 mm SL

All illustrations reproduced from the *African Journal of Marine Science* (1986) 4: 193-201 with permission – © NISC (Pty) Ltd

A-F: Olivar (1986)

**BLENNIIDAE***Parablennius gattorugine* (Linnaeus, 1758)**En:** Tompot blenny – **Sp:** Cabruza**Habitat:** Benthic, littoral**Spawning season:** No information**EGGS****Habitat:** demersal**Shape:** hemispherical**Diameter:** 1.6 mm**Oil globules:** none**Yolk:** unsegmented**YOLK-SAC LARVAE****Hatch size:** 4.9 mm**Pigmentation:** dorsal side of gut pigmented, from behind eye to anus; a few melanophores on snout and top of head; apparently, no postanal ventral row of melanophores; pectoral fin unpigmented**LARVAE****Figs. A–D****Distinctive characters:** body elongate; pectoral fin fairly large; preanus length short**Preanus length:** about one third of SL**Pigmentation:** no melanophore on hindgut; black peritoneal pigment covering dorsal side of gut, to almost anus; postanal ventral row of 19–21 regularly spaced melanophores begins some distance behind anus and extends to caudal peduncle; melanophores on head and snout and some caudal melanophores; apparent paired dorsal bands of melanophores across dorsal fin in larvae larger than 18 mm; lateral row of melanophores above notochord**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



3.8 mm SL



4.5 mm SL




9.4 mm SL

**Literature:** Ford (1922), Padoa (1956b), Russell (1976), Zander (1986)

**BLENNIIDAE**


*Parablennius gattorugine* (Linnaeus, 1758)



No melanophore on hindgut

A. 6.0 mm

Melanophores on head and snout




B. 10.0 mm


Ventral row of melanophores begins some distance behind anus

B. Detail of pectoral fin

Lateral row of melanophores above notochord



C. 13.0 mm



D. 18.5 mm

## CALLIONYMIDAE

*Callionymus* Linnaeus, 1758

In the Western Mediterranean Sea there are 6 species of the genus *Callionymus*. Younger larvae (< 3 mm SL) of this genus are indistinguishable. The identification at the species level is only possible in larvae larger than 3 mm, when preopercular spines develop, and pelvic fins and pigmentation patterns become species-specific

**Habitat:** Benthic, over sandy bottoms down to 650 m depth

**Spawning season:** Between January and August (varying by species)




Fig. A

### EGGS

**Habitat:** pelagic

**Shape:** spherical

**Diameter:** 0.55–0.97 mm

**Chorion:** in some species it shows hexagonal sculptures

**Oil globules:** none

**Yolk:** segmented

### YOLK-SAC LARVAE

Fig. B

**Hatch size:** about 2 mm (*Callionymus lyra*); yolk appears fully absorbed at 2.3 mm SL

### LARVAE

Figs. C–H

**Distinctive characters:** body short with thick head and abdomen; it shows a bifurcated preopercular spine; urostyle strongly developed and curved up at its end


**Preanus length:** about 50% SL

**Pigmentation:** strongly pigmented except for urostyle region

**Length at flexion:** no information

### PHOTOS

by J.M. Rodriguez



*Callionymus* sp.  
2.5 mm SL



*Callionymus* sp.  
5.9 mm SL


**Literature:** Demir (1972), Fage (1918), Fricke (1986), Padoa (1956s), Russell (1976), Sabatés (1988)

**CALLIONYMIDAE**


*Callionymus* Linnaeus, 1758



A.




B. *C. lyra* 2.2 mm




Body short; thick  
head and abdomen

C. *C. lyra* 5.8 mm




D. *C. lyra* 7.5 mm


Body strongly  
pigmented




E. *C. lyra* 8.3 mm




F. *C. reticulatus* 6.6 mm TL



G. *C. lyra* 6.5 mm TL



H. *C. maculatus* 6.6 mm TL

**GOBIIDAE***Crystallogobius linearis* (Düben, 1845)**En:** Crystal goby**Habitat:** Pelagic, in coastal to offshore waters, down to over 400 m depth. Males live on the bottom during the breeding season**Spawning season:** Between May and August in the NE Atlantic

Gobiidae

**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–E****Distinctive characters:** body elongate and slender; first dorsal fin absent in females and showing two rays in males**Preanus length:** less than 50% SL**Pigmentation:** larvae up to 5 mm length, strongly pigmented with melanophores on head, in front of eye, at lower jaw tip and oocyst region; 5 large evenly spaced melanophores along dorsal contour of body; dorsal and ventral sides of gut and swimbladder pigmented; postanal ventral row of melanophores; pigmentation diminishes with development, so that larvae of ca. 11.0 mm SL have a row of postanal ventral melanophores and a series of melanophores on anterior ventral side of gut; swimbladder pigmented**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez




10.4 mm SL

**Literature:** Alemany (1997), Miller (1986), Padoa (1956h), Sabatés (1988)


**GOBIIDAE**

*Crystallogobius linearis* (Düben, 1845)



A. 4.9 mm

Young larvae,  
strongly pigmented




B. 9.0 mm

In older larvae, pigment reduced to  
ventral side of body and swimbladder




C. 11.2 mm


Melanophores on  
ventral side of gut



D. 14.5 mm SL



E. 15.2 mm SL

**GOBIIDAE***Gobius paganellus* Linnaeus, 1758**En:** Rock goby – **Fr:** Gobie paganel – **Sp:** Bobí**Habitat:** Benthic, in inshore and intertidal waters over rocky bottoms with weed cover**Spawning season:** Between January and June**EGGS****Fig. A****Habitat:** demersal**Shape:** fusiform, with a rounded, pointed apex**Size:** 2.24 x 0.84 mm**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 4.76 mm with mouth open and swimbladder apparent**Pigmentation:** branched melanophores on ventral profile of abdomen; large, opposing dorsal and ventral melanophores at about mid-postanal region; swimbladder pigmented; a melanophore on hindgut; no melanophores on head**LARVAE****Figs. C–F****Distinctive characters:** body relatively elongate and slender**Preanus length:** about 50% SL**Pigmentation:** early larvae similar to yolk-sac larvae; larvae larger than 7.0 mm with melanophores on head and a row of postanal ventral melanophores, while postanal dorsal melanophore disappears in postflexion larvae**Length at flexion:** unknown**PHOTOS**


Not available

**Literature:** Alemany (1997), Lebour (1919), Miller (1986), Padoa (1956h)




**GOBIIDAE**

*Gobius paganellus* Linnaeus, 1758



A.



B. 4.0 mm


Large, opposing dorsal and ventral melanophores at about mid-tail region



C. 2.1 mm SL




D. 2.6 mm SL




E. 4.8 mm

Dorsal, mid-tail melanophore, disappears in late larvae



F. 11.5 mm

**GOBIIDAE***Lebetus guilleti* (Le Danois, 1913)**En:** Guillet's goby**Habitat:** Benthic, littoral, over coarse grounds**Spawning season:** Larvae in plankton from February to December in the Balearic Sea

Gobiidae

**EGGS**


Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–E****Distinctive characters:** body short and robust; small to moderate head; no cephalic spines; eyes rounded and large; prominent swimbladder; pelvic fins develop from young larvae on**Preanus length:** about 67% SL**Pigmentation:** body covered with melanophores except for caudal region; well-marked lateral row of melanophores and another ventral row, from anus to base of caudal fin**Length at flexion:** 5.0–7.0 mm SL**PHOTOS**

by J.M. Rodriguez



2.2 mm SL



2.8 mm SL




4.5 mm SL

**Literature:** Alemany (1997), Miller (1986), Russell (1976)


**GOBIIDAE**

*Lebetus guilleti* (Le Danois, 1913)




A. 1.9 mm SL

Body short and robust



B. 2.5 mm SL

Body, except caudal region, covered with melanophores




C. 3.1 mm SL

Well-marked lateral row of melanophores




D. 3.2 mm SL

Large pelvic fins



E. 3.5 mm SL

**GOBIIDAE***Pomatoschistus microps* (Krøyer, 1838)**En:** Common goby – **Fr:** Gobie commun**Habitat:** Benthic, coastal, frequently in estuaries and intertidal pools**Spawning season:** Spring and summer

Gobiidae

**EGGS****Fig. A****Habitat:** demersal**Shape:** pear-shaped, with a rounded apex**Size:** 1.0–0.7 x 0.65–0.8 mm**Oil globules:** several, maximum size 0.08 mm in diameter**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 3.0 mm**Pigmentation:** row of melanophores along ventral side of trunk and tail; yolk sac and ventral side of gut strongly pigmented; large branched melanophore on dorsal mid-tail region**LARVAE****Figs. C–F****Distinctive characters:** body relatively elongate and slender**Preanus length:** about 50% SL**Pigmentation:** melanophores on head and lower jaw; an almost continuous row of ventral melanophores from throat, along trunk and tail region, one of which is highly ramified; highly ramified melanophore on mid dorsal tail region, opposite to ventral one**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



3.2 mm SL



5.9 mm SL

**Literature:** Alemany (1997), Miller (1986), Padoa (1956h), Russell (1976), Sabatés (1988)

**GOBIIDAE**

*Pomatoschistus microps* (Krøyer, 1838)




A.




B. 3.0 mm

A dorsal melanophore strongly ramifies at mid-tail region



C. 3.6 mm



Continuous row of melanophores along ventral side of body

D. 8.0 mm




E. 11.0 mm

Large larvae strongly pigmented



F. 14.0 mm

**TRICHIURIDAE***Lepidopus caudatus* (Euphrasen, 1788)**En:** Silver scabbardfish – **Fr:** Sabre argenté – **Sp:** Pez cinto**Habitat:** Benthopelagic, on the continental shelf and along its edge, down to 400 m depth; usually over sandy and muddy bottoms between 100 and 250 m depth**Spawning season:** Between June and November in the Adriatic Sea**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.6–1.7 mm**Chorion:** smooth**Oil globules:** single, 0.40 mm in diameter**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 6.0 mm**Distinctive characters:** oil globule and anus situated at posterior edge of yolk sac; high number of myomeres**Preanus length:** about 25% SL**Pigmentation:** two dorsal and two ventral melanophores on tail region, one at level of caudal peduncle; some melanophores on head**LARVAE****Figs. C–F****Distinctive characters:** body very elongate and slender; long cephalic appendix (first ray of dorsal fin) in younger larvae, shortens with development**Preanus length:** about 25% SL in young larvae increases to about 50% SL in later larvae**Pigmentation:** early stages, similar to yolk-sac larvae: melanophores on head, snout, and post-occipital region, tail patches move into finfold and later into dorsal fin; peritoneum and lateral sides of gut pigmented; pigment over caudal peduncle reduces to its ventral side with development; anterior dorsal pigment patch disappears and others decrease in size; melanophores on dorsal-fin base extends backwards**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



7.3 mm SL


**Literature:** Froese and Pauly (2016), Padoa (1956o), Parin (1986), Sabatés (1988), Schmidt (1918)

**TRICHIURIDAE**


*Lepidopus caudatus* (Euphrasen, 1788)



A.




B. ca. 6.0 mm




Long cephalic appendix  
in young larvae

C. 9.0 mm




Typical pigment  
patches in tail region

D. 10.2 mm




Body slender and  
very elongate


E. 12.6 mm



F. 12.9 mm

**SCOMBRIDAE***Auxis rochei* (Risso, 1810)**En:** Bullet tuna – **Fr:** Bonitou – **Sp:** Melva (=Melvera)**Habitat:** Epipelagic, in inshore waters and near islands**Spawning season:** Between May and November**EGGS****Habitat:** pelagic**Shape:** spherical**Diameter:** 0.82–1.10 mm**Chorion:** smooth**Oil globules:** single, 0.24–0.29 mm in diameter**Yolk:** unsegmented**YOLK-SAC LARVAE****Hatch size:** 2.14 mm SL**Pigmentation:** melanophores on head, and a row of melanophores on postanal ventral region; gut pigmented**LARVAE****Figs. A–F****Distinctive characters:** body moderately elongate becoming deeper with development; gut compact and triangular; jaws relatively short, compared to other tuna-like species; preopercular spines well-developed**Preanus length:** increases from about 37% to 50% SL with development**Pigmentation:** early stages, similar to yolk-sac larvae; in late larvae, the origin of ventral postanal row of melanophores moves backwards and melanophores decrease in number; pigmentation of peritoneal region increases; some melanophores appear in the occipital region together with one or two dorsal melanophores, a little ahead of caudal peduncle; these latter ones extend forward with development**Length at flexion:** 4.5–6.0 mm**PHOTOS**

by J.M. Rodriguez



2.4 mm SL




7.0 mm SL

**Literature:** Alemany (1997), Collette (1986b), Fahay (2007), Padoa (1956l), Richards (2006), Sabatés (1988)




**SCOMBRIDAE**

*Auxis rochei* (Risso, 1810)



A-C: Alemany (1997); D-F: Padoa (1956l)

**SCOMBRIDAE***Sarda sarda* (Bloch, 1793)**En:** Atlantic bonito – **Fr:** Bonite à dos rayé**Sp:** Bonito del Atlántico**Habitat:** Epipelagic, migratory**Spawning season:** Between May and July**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.2–1.32 mm**Chorion:** smooth**Oil globules:** 1–5; 0.28–0.32 mm in diameter, when single; 0.02–0.24 mm in diameter, when multiple**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 4.0 mm**Distinctive characters:** yolk sac voluminous; oil globules situated at posterior edge of yolk sac**Pigmentation:** finfold pigmented in its anterior dorsal part; small melanophores on head and along dorsal region; some isolated melanophores on lateral and ventral sides of body**LARVAE****Figs. C–F****Distinctive characters:** body moderately elongate, deepest at level of pectoral region; head large with pointed snout and jaws; prominent teeth from early larvae on; preopercular spines well developed; gut compact and triangular**Preanus length:** increases from about 50% SL to 60% SL with development**Pigmentation:** series of large spots along ventral tail region, move up internally between myomeres with development; top of head and peritoneal region well-pigmented; melanophores at cleithral symphysis, at lower jaw tip and caudal-fin base; patch of pigment in lateral caudal region in older larvae**Length at flexion:** 6.0–7.0 mm SL**PHOTOS**

by J.M. Rodriguez



3.5 mm SL



4.0 mm SL


**Literature:** Collette (1986b), Fahay (2007), Padoa (1956l), Richards (2006)

**SCOMBRIDAE**

*Sarda sarda* (Bloch, 1793)




A.



B. 4.3 mm


Oil globules situated at posterior edge of yolk sac



C. 4.2 mm

Head large and snout and jaws pointed


Teeth prominent



D. 4.7 mm SL


Gut compact and triangular

Series of large spots along ventral tail region




E. 6.4 mm

Preopercular spines well developed



F. 9.8 mm SL

**SCOMBRIDAE***Scomber colias* Gmelin, 1789**En:** Atlantic chub mackerel – **Fr:** Maquereau espagnol**Sp:** Estornino**Habitat:** Epipelagic or meso–demersal, down to 250–300 m depth**Spawning season:** Summer**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.04–1.14 mm**Chorion:** smooth**Oil globules:** single, 0.26–0.27 mm in diameter**Perivitelline space:** small**Yolk:** unsegmented**YOLK–SAC LARVAE****Fig. B****Hatch size:** about 3.0 mm SL**Distinctive characters:** body relatively elongate; oil globule located at posterior edge of the yolk sac**Pigmentation:** melanophores irregularly distributed along dorsal and ventral contours of body; melanophores on snout and behind eye; yolk sac and oil globule pigmented**LARVAE****Figs. C–F****Distinctive characters:** body moderately elongate, becoming stubby with development, deeper than that of larvae of comparable sizes of *Scomber scombrus*; head moderate; gut compact and triangular; jaws relatively short; no head spines; teeth prominent in larvae larger than 4.0 mm SL; the number of myomeres allows for distinguishing this species (32) from *Diplodus* species (24)**Preanus length:** increases from about 50% SL to more than 60% SL with development**Pigmentation:** postanal ventral rows of melanophores starting some distance from anus; larvae less than 7 mm long may only show two dorsal melanophores, situated on caudal peduncle; peritoneum pigmented; melanophores at caudal–fin base; melanophores on head; no melanophores on body sides in preflexion stages; no melanophores at cleithral symphysis and on isthmus**Length at flexion:** about 6 mm**PHOTOS**

by J.M. Rodriguez



4.4 mm SL




7.0 mm SL


**Literature:** Berrien (1978), Collette (1986b), Kramer (1969), Sabatés (1988)

**SCOMBRIDAE**


*Scomber colias* Gmelin, 1789



A.




B. 3.3 mm



C. 3.8 mm SL

No melanophores at isthmus and cleithral symphysis




Only two dorsal melanophores in larvae less than 7 mm


D. 5.3 mm SL

Preamus length greater and body deeper than that of *S. scombrus*


Myomere count, 32 in *Scomber*, 24 in Sparidae species



E. 7.3 mm SL



F. 11.3 mm SL

**SCOMBRIDAE***Scomber scombrus* Linnaeus, 1758**En:** Atlantic mackerel – **Fr:** Maquereau commun**Sp:** Caballa del Atlántico**Habitat:** Epipelagic or meso–demersal, down to 200–250 m depth**Spawning season:** March and April**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.0–1.38 mm**Chorion:** smooth**Oil globules:** single, 0.28–0.35 mm in diameter**Perivitelline space:** small**Yolk:** unsegmented**YOLK–SAC LARVAE****Fig. B****Hatch size:** 3.3 mm SL**Distinctive characters:** body relatively elongate; oil globule located at posterior edge of yolk sac**Pigmentation:** melanophores on head; double rows of irregularly distributed melanophores along dorsal and ventral contours of body; peritoneum pigmented; melanophores on snout and behind eye; oil globule pigmented**LARVAE****Figs. C–G****Distinctive characters:** body moderately elongate, becoming stubbier with development; head moderate; gut compact and triangular; jaws relatively short; no head spines; teeth prominent in larvae larger than 4.0 mm SL**Preanus length:** increases from about 40% SL to more than 60% SL with development**Pigmentation:** postanal rows of dorsal and ventral melanophores; in earliest larval stages, there may be fewer melanophores in the dorsal row, equaling their number (14–15) with development; peritoneum pigmented; some melanophores on urostyle and along caudal–fin base; melanophores on head and sometimes on snout and lower jaw; no melanophores on body sides in preflexion larvae; melanophores at cleithral symphysis and on itsmus**Length at flexion:** 5.0–7.0 mm**PHOTOS**

by J.M. Rodriguez



3.7 mm SL



7.0 mm SL


**Literature:** Berrien (1978), Collette (1986b), Froese and Pauly (2016), Padoa (1956), Sabatés (1988)

**SCOMBRIDAE**

*Scomber scombrus* Linnaeus, 1758




A.



B. 2.4 mm




C. 2.9 mm SL



D. 3.8 mm SL


Melanophores at isthmus and cleithral symphysis

Postanal row of dorsal melanophores




Prianus length shorter and body more slender than that of *S. colias*


E. 4.8 mm SL



F. 7.2 mm SL



G. 11.7 mm SL

**SCOMBRIDAE***Thunnus alalunga* (Bonnaterre, 1788)**En:** Albacore – **Fr:** Germon – **Sp:** Atún blanco**Habitat:** Epi–mesopelagic, down to 100 m depth**Spawning season:** Between June and August**EGGS****Habitat:** pelagic**Shape:** spherical**Diameter:** 0.84–0.94 mm**Chorion:** smooth**Oil globules:** single, 0.24 mm in diameter**Yolk:** unsegmented**YOLK–SAC LARVAE****Fig. A****Hatch size:** about 2.6 mm SL**Pigmentation:** melanophores on yolk and on gut; dotted melanophores dorsal and ventral to urostyle**LARVAE****Figs. B–F****Distinctive characters:** body moderately stocky, deepest at level of pectoral region, tapering to a narrow caudal peduncle; large head with pointed snout and jaws; preopercular spines present; gut compact and triangular**Preanus length:** increases from about 40% SL to about 55% SL with development**Pigmentation:** few dotted dorsal melanophores over urostyle disappear with development; peritoneal region strongly pigmented; occipital region pigmented; melanophore at snout tip in older larvae**Length at flexion:** 5.0–7.0 mm SL**PHOTOS**

by J.M. Rodriguez



3.4 mm SL



7.0 mm SL

**Literature:** Alemany (1997), Fahay (2007), Froese and Pauly (2016), Richards (2006)




**SCOMBRIDAE**

*Thunnus alalunga* (Bonnaterre, 1788)


Melanophores dorsal and ventral to urostyle



A. 2.5 mm SL




B. 2.6 mm SL




C. 3.1 mm SL

Preopercular spines




D. 4.3 mm

Tail region unpigmented




E. 5.1 mm SL

First dorsal fin unpigmented



F. 6.4 mm SL

**SCOMBRIDAE***Thunnus thynnus* (Linnaeus, 1758)**En:** Atlantic bluefin tuna – **Fr:** Thon rouge de l'Atlantique**Sp:** Atún rojo del Atlántico**Habitat:** Epi-mesopelagic, carrying out trans-oceanic migrations**Spawning season:** Between May and July**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.0–1.12 mm**Chorion:** smooth**Oil globules:** single, 0.25–0.28 mm in diameter**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Hatch size:** 3.0 mm TL**Pigmentation:** large mid-trunk dorsal melanophore extending to finfold**LARVAE****Figs. B–F****Distinctive characters:** body stout, deepest at level of pectoral region, tapering to a narrow caudal peduncle; large head with pointed snout and jaws; gut compact and triangular; preopercular spines present**Preanus length:** increases from about 40% SL to about 55% SL with development**Pigmentation:** about 4 dotted postanal-ventral melanophores; a single melanophore on dorsal side of mid-tail, migrating towards dorsal fin with development; peritoneal region strongly pigmented**Length at flexion:** 5.0–7.0 mm SL**PHOTOS**

by J.M. Rodriguez



3.8 mm SL



6.7 mm SL


**Literature:** Alemany (1997), Collette (1986b), Fahay (2007), Richards (2006)

**SCOMBRIDAE**

*Thunnus thynnus* (Linnaeus, 1758)




A.



B. 4.0 mm SL


About 4 dotted postanal-ventral melanophores



C. 5.1 mm SL


Gut compact and triangular

A single melanophore on dorsal side of mid-tail, migrating towards dorsal fin with development




D. 6.8 mm

Preopercular spines well developed




E. 7.3 mm



F. 9.4 mm

Large jaws with apparent teeth

**XIPHIIDAE***Xiphias gladius* Linnaeus, 1758**En:** Swordfish – **Fr:** Espadon – **Sp:** Pez espada**Habitat:** Pelagic, oceanic down to 800 m depth, strongly migratory**Spawning season:** June to September**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.6–1.8 mm**Chorion:** smooth**Oil globules:** single, 0.40 mm in diameter**Perivitelline space:** small**Yolk:** segmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 4.2 mm**Distinctive characters:** yolk sac large; oil globule situated at posterior edge of yolk sac**Pigmentation:** body and finfold covered with small melanophores, grouped at myosepta**LARVAE****Figs. C–F****Distinctive characters:** body relatively stubby, with bulky gut, becoming elongate with development; head shallow, with elongate jaws; mouth very large, extending posteriorly beyond eye; teeth well formed by 6.0 mm SL; spinous scales in larger larvae**Preanus length:** between 70 and 80% SL, increasing with development**Pigmentation:** body, except caudal peduncle, covered with small branched melanophores**Length at flexion:** 8.0–12.0 mm SL**PHOTOS**

by J.M. Rodriguez



3.7 mm SL



Not sized


**Literature:** Alemany (1997), Arata Jr. (1954), Nakamura (1986), Padoa (1956r), Richards (2006)

**XIPHIIDAE**


*Xiphias gladius* Linnaeus, 1758



A.




B. 4.2 mm SL



Body, except caudal peduncle, covered with small branched melanophores


C. 4.1 mm SL




Gut bulky

D. 6.1 mm SL

Mouth large with elongate jaws and prominent teeth




E. 7.8 mm SL



Spinous scales in larger larvae


F. 9.0 mm SL

**CAPROIDAE***Capros aper* (Linnaeus, 1758)**En:** Boarfish – **Fr:** Sanglier – **Sp:** Ochavo**Habitat:** Benthic, between 25 and 600 m depth**Spawning season:** Between March and August**Fig. A****EGGS****Habitat:** pelagic**Shape:** spherical**Diameter:** 0.90–1.01 mm**Chorion:** smooth**Oil globules:** single, 0.15–0.17 mm in diameter**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** 2.02–2.46 mm**Pigmentation:** body, except caudal region, covered with large stellate melanophores**LARVAE****Figs. C–F****Distinctive characters:** body thin during first stages of development, acquires a rhomboid shape at about 3 mm SL; opercular spines and a spiny crest over head appear with development; first dorsal fin spinous; body covered with small spines in later larvae**Preanus length:** about 50% SL**Pigmentation:** early larvae with a row of ventral melanophores above gut to about midway, along postanal region; dorsal row of six large melanophores extending from mid-gut to level of ventral row; row of melanophores along ventral abdominal surface; some melanophores on head and on mid-lateral side of body; upper and lower jaw tips pigmented; only caudal peduncle remains free of pigment in later larvae**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



2.0 mm SL



3.0 mm SL


**Literature:** Alemany (1997), Froese and Pauly (2016), Quero (1986a), Russell (1976)

**CAPROIDAE**

*Capros aper* (Linnaeus, 1758)



A.




B. 2.9 mm




C. 2.9 mm

Body rhomboid-shaped  
in late larvae



D. 2.4 mm SL




Spinous crest

First dorsal fin  
spinous

Opercular  
spines


E. 3.2 mm SL

Body covered with small  
spines in older larvae



Caudal peduncle  
free of pigment

F. 8.7 mm SL

**SCOPHTHALMIDAE***Lepidorhombus boscii* (Risso, 1810)**En:** Four-spot megrim – **Fr:** Cardine à quatre taches**Sp:** Gallo de cuatro manchas**Habitat:** Benthic, down to 800 m depth**Spawning season:** Between May and October**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A–E**

**Distinctive characters:** body relatively slender at early stages; trunk and abdominal regions increase considerably in depth with development; larvae more than 5.0 mm long show preopercular spines and a spine on each side of upper posterior side of head

**Preanus length:** less than 50% SL

**Pigmentation:** 4 melanophores evenly arranged along dorsal side of body and 3 postanal ventral melanophores; gut, head and frontal region pigmented

**Length at flexion:** unknown

**Remarks:** asymmetry starts when the larva is about 9.7 mm long

**PHOTOS**

by J.M. Rodriguez



3.2 mm SL



6.9 mm SL


**Literature:** Alemany (1997), Nielsen (1986d), Padoa (1956c), Russell (1976), Sabatés (1991)



**SCOPHTHALMIDAE**


*Lepidorhombus boscii* (Risso, 1810)

4 evenly spaced dorsal melanophores




A. 3.9 mm SL

3 postanal ventral melanophores




B. 5.1 mm SL

Swimbladder apparent




C. 6.8 mm SL


Gut curved



D. 8.4 mm




E. 10.0 mm

**SCOPHTHALMIDAE***Scophthalmus rhombus* (Linnaeus, 1758)**En:** Brill – **Fr:** Barbue – **Sp:** Rémol**Habitat:** Benthic, between 5 and 70 m depth**Spawning season:** Between February and April (probably)**Fig. A****EGGS****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.24–1.50 mm**Chorion:** smooth**Oil globules:** 0.16–0.25 mm in diameter**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** 3.8–4.0 mm**Pigmentation:** body (except caudal region) and yolk sac pigmented; melanophores extend onto finfold to form a wide postanal bar and a dorsal patch above gut**LARVAE****Figs. C–F****Distinctive characters:** body relatively elongate and slender becomes laterally compressed and ovoid with development; a spiny ridge above eye and opercular spines**Preanus length:** about 50% SL**Pigmentation:** body covered with melanophores (caudal region free of pigment in early larvae); body pigment extends onto dorsal and anal fins forming almost regular bands in older larvae**Length at flexion:** unknown**PHOTOS**

by J.M. Rodriguez



2.8 mm SL



3.2 mm SL


**Literature:** Jones (1972), Padoa (1956c), Russell (1976), Sabatés (1988)

**SCOPHTHALMIDAE**


*Scophthalmus rhombus* (Linnaeus, 1758)



A.




B. 3.8 mm TL




C. 5.8 mm TL

Body covered with melanophores (caudal region free of pigment in early larvae)




D. 6.9 mm TL

Spiny ridge above eye and opercular spines




E. 8.0 mm TL

In old larvae, body pigment extends onto dorsal and caudal fins forming almost regular bands




F. 10.1 mm TL

**SCOPHTHALMIDAE***Zeugopterus regius* (Bonnaterre, 1788)**En:** Eckström's topknot – **Sp:** Pelaya miserés**Habitat:** Benthic, from a few to 300 m depth**Spawning season:** Between May and August in the British Isles**EGGS****Habitat:** pelagic**Shape:** spherical**Diameter:** 0.90–0.99 mm**Chorion:** smooth**Oil globules:** 0.16–0.18 mm in diameter**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. A****Hatch size:** about 2.4 mm**Pigmentation:** body, yolk sac and finfold covered with small melanophores**LARVAE****Figs. B–E****Distinctive characters:** body relatively short and relatively deep at level of pectoral region; two large otocystic spines on each side of head**Preanus length:** about 50% SL**Pigmentation:** body and fins covered with small melanophores**Length at flexion:** completed at about 6.7 mm**PHOTOS**

by J.M. Rodriguez



4.8 mm SL




9.0 mm SL


**Literature:** Padoa (1956c), Russell (1976), Sabatés (1988), Tortonese (1970)

**SCOPHTHALMIDAE**


*Zeugopterus regius* (Bonnaterre, 1788)



A. 2.4 mm




B. 4.7 mm




C. 6.5 mm SL

2 large otocystic spines on each side of head



D. 6.7 mm

Body and fins covered with small melanophores




E. 8.0 mm

**BOTHIDAE***Arnoglossus laterna* (Walbaum, 1792)**En:** Mediterranean scaldfish – **Fr:** Arnoglosse de Méditerranée – **Sp:** Serrandell

Five *Arnoglossus* species are known to occur in the Mediterranean Sea. The identification of their larvae is difficult due to their similar morphology and pigmentation pattern. *A. laterna* is the only species of this genus for which all early life stages have been described and is here presented.

**Habitat:** Benthic, over muddy and sandy bottoms down to 200 m depth, occasionally to 400 m

**Spawning season:** Between April and August

**EGGS****Fig. A**

**Habitat:** pelagic

**Shape:** spherical

**Diameter:** 0.60–0.76 mm

**Chorion:** smooth

**Oil globules:** single, 0.11–0.15 mm in diameter

**Perivitelline space:** small

**Yolk:** unsegmented

**YOLK-SAC LARVAE****Fig. B**

**Hatch size:** about 2.5 mm

**Pigmentation:** free of black pigment

**LARVAE****Figs. C–F**

**Distinctive characters:** body relatively elongate, increases in height and becomes laterally compressed with development; gut becomes curved; elongate tentacle over head, disappears in older larvae; swimbladder prominent; minute spines on different regions of body in larger larvae

**Preanus length:** about 40% SL

**Pigmentation:** two rows of evenly spaced ventral melanophores, extending from anus to caudal end; dorsal bar of melanophores close to caudal region; melanophores along abdominal ventral contour, on lower jaw and over hindgut; swimbladder pigmented


**Length at flexion:** unknown

**PHOTOS**

by J.M. Rodriguez



3.9 mm SL



6.2 mm SL


**Literature:** Alemany (1997), Nielsen (1986a), Padoa (1956c), Russell (1976), Sabatés (1988)

**BOTHIDAE**


*Arnoglossus laterna* (Walbaum, 1792)



A.




B. 2.6 mm



C. 2.8 mm

Elongate tentacle  
over head


Body laterally compressed



D. 4.7 mm


Gut curved

2 rows of evenly spaced ventral  
melanophores, from anus to caudal end




E. 8.5 mm

Prominent swimbladder



F. 12.5 mm

**BOTHIDAE***Bothus podas* (Delaroche, 1809)**En:** Wide-eyed flounder**Habitat:** Benthic, over sand and mud of the continental shelf and slope, from a few to 400 m depth**Spawning season:** Between May and August**EGGS**


Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A-D****Distinctive characters:** body very similar to that of *Arnoglossus* larvae; relatively elongate, increases in height and becomes laterally compressed with development; gut curved; elongate tentacle over head; swimbladder prominent**Pigmentation:** large melanophores on dorsal and ventral side of urostyle; rest of body, unpigmented**Length at flexion:** flexion begins at about 7.0 mm**PHOTOS**

by J.M. Rodriguez



3.9 mm SL


**Literature:** Alemany (1997), Padoa (1956c), Sabatés (1988), Tortonese (1975)




**BOTHIDAE**

*Bothus podas* (Delaroche, 1809)

Elongated  
tentacle over head  
→




A. 3.6 mm SL




B. 6.5 mm SL


Body  
unpigmented



C. 7.2 mm SL




D. 13.5 mm SL

**PLEURONECTIDAE***Platichthys flesus* (Linnaeus, 1758)**En:** European flounder – **Fr:** Flet d'Europe – **Sp:** Platija europea**Habitat:** Benthic, at shallow depths on soft bottoms**Spawning season:** Between December and March (Black Sea)**Fig. A****EGGS****Habitat:** pelagic**Shape:** spherical**Diameter:** 0.8–1.13 mm**Chorion:** smooth**Oil globules:** none**Perivitelline space:** small**Yolk:** unsegmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 2.3 mm**Pigmentation:** no pigmentation on primordial fin in earlier larvae; as development proceeds, a group of melanophores develops about mid-way along postanal region, spreading out dorsally and ventrally onto the primordial fin; other melanophores scattered on body; postanal end region unpigmented**LARVAE****Figs. C–F****Distinctive characters:** body elongate and slender, becoming more robust and laterally compressed with development**Preanus length:** less than 50% SL**Pigmentation:** many scattered melanophores along ventral sides of body, spreading out over anal and caudal fin where melanophores tend to be aligned with fin rays**Length at flexion:** unknown**Remarks:** asymmetry starts at about 7 mm and is completed at about 10 mm length**PHOTOS**

by J.M. Rodriguez



2.7 mm SL



7.0 mm SL


**Literature:** Froese and Pauly (2016), Nielsen (1986c), Russell (1976)

**PLEURONECTIDAE**

*Platichthys flesus* (Linnaeus, 1758)



A.




B. 2.8 mm




C. 4.0 mm

Body elongate and slender in young larvae




D. 7.2 mm




E. 6.0 mm


Scattered melanophores along ventral sides of body spreading out over anal and caudal fins




F. 8.0 mm

**SOLEIDAE***Buglossidium luteum* (Risso, 1810)**En:** Solenette**Habitat:** Benthic, over sandy bottoms, between 5 and 450 m depth**Spawning season:** February**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 0.64–0.94 mm**Chorion:** smooth**Oil globules:** 12–15 small oil globules**Perivitelline space:** small**Yolk:** segmented**YOLK-SAC LARVAE****Figs. B, C****Hatch size:** about 2.0 mm**Pigmentation:** melanophores on head; 4 spots on dorsal and 1 or 2 on ventral margins of primordial fin; some melanophores on body margins**LARVAE****Figs. D–F****Distinctive characters:** body short with large head and deep abdominal region; body becomes laterally compressed with development; swimbladder prominent**Preanus length:** less than 50% SL**Pigmentation:** evenly spaced melanophores along dorsal (9–13) and ventral (8–11) body contours; swimbladder and pectoral fin pigmented**Length at flexion:** unknown**Remarks:** asymmetry begins at about 6.0 mm SL, almost completed at 8.0 mm SL**PHOTOS**

by J.M. Rodriguez



3.5 mm SL



6.5 mm SL


**Literature:** Munk and Nielsen (2005), Padoa (1956m), Quero *et al.* (1986b), Russell (1976)

**SOLEIDAE**


*Buglossidium luteum* (Risso, 1810)




A.




B. 2.1 mm



C. 2.4 mm  
Lateral sides of body slightly pigmented




D. 3.7 mm



Swimbladder prominent


Evenly spaced melanophores along dorsal (9-13) and ventral (8-11) body contours

E. 6.3 mm




Body short, large head and deep abdominal region


F. 7.5 mm

**SOLEIDAE***Microchirus variegatus* (Donovan, 1808)**En:** Thickback sole – **Sp:** Golleta**Habitat:** Benthic, over sand and mud on the continental shelf and slope, at depths between 80 and 400 m**Spawning season:** Spring**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.28–1.42 mm**Chorion:** smooth**Oil globules:** 50 or more scattered oil globules**Perivitelline space:** small**Yolk:** segmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** about 2.5 mm**Pigmentation:** body and primordial fins covered with small stellate melanophores; apparent rows of stellate melanophores along the body contours**LARVAE****Figs. C–F****Distinctive characters:** body short with a prominent head and deep abdominal region; no swimbladder**Preanus length:** about 50% SL**Pigmentation:** body and fins covered with small stellate melanophores; dorsal and ventral rows of melanophores (dorsal of about 70 and ventral of about 50) along body contours, these melanophores are larger and tend to merge and form continuous rows**Length at flexion:** unknown**Remarks:** asymmetry completed at 12.0 mm SL**PHOTOS**

by J.M. Rodriguez



3.0 mm SL



5.1 mm SL


**Literature:** Alemany (1997), Padoa (1956m), Quero *et al.* (1986b), Russell (1976)

**SOLEIDAE**


*Microchirus variegatus* (Donovan, 1808)



A.




B. ca. 3.7 mm



C. 4.0 mm


Body short, prominent head and deep abdominal region



D. 5.8 mm


Body and fins covered with small melanophores

No swimbladder




E. 8.0 mm


Melanophores of dorsal and ventral contours of body form continuous rows




F. 9.8 mm

**SOLEIDAE***Solea solea* (Linnaeus, 1758)**En:** Common sole – **Fr:** Sole commune – **Sp:** Lengüado común**Habitat:** Benthic, over sand and mud of continental shelf from a few to 150–200 m depth**Spawning season:** Between January and May**EGGS****Fig. A****Habitat:** pelagic**Shape:** spherical**Diameter:** 1.0–1.6 mm**Chorion:** smooth**Oil globules:** many small oil globules aggregated in clusters**Perivitelline space:** small**Yolk:** peripherally segmented**YOLK-SAC LARVAE****Fig. B****Hatch size:** 3.8 mm**Pigmentation:** body (except caudal end), primordial fin and yolk sac covered with small stellate melanophores; rows of melanophores along dorsal (9–14 melanophores) ventral (4–12) contours of body, number of melanophores in these rows increases with age**LARVAE****Figs. C–F****Distinctive characters:** body short with a prominent head and deep abdominal region; swimbladder small, apparent at 6–7 mm long**Preanus length:** about 50% SL**Pigmentation:** similar to that of yolk-sac larvae in earlier stages; larger larvae, body covered with large stellate melanophores; dorsal and ventral rows of melanophores almost forming an unbroken line of pigment**Length at flexion:** unknown**Remarks:** asymmetry begins at 7–8 mm, almost completed at 9.5 mm**PHOTOS**

by J.M. Rodriguez



3.6 mm SL



7.3 mm SL

**Literature:** Aboussouan (1964), Padoa (1956m), Quero *et al.* (1986b), Russell (1976), Sabatés (1988)




**SOLEIDAE**

*Solea solea* (Linnaeus, 1758)




A.



B. 4.0 mm




C. 4.2 mm




D. 6.8 mm

Swimbladder apparent  
in larvae more than  
6-7 mm long

Body covered with large  
stellate melanophores



E. 7.5 mm



F. 13.0 mm

**CYNOGLOSSIDAE***Symphurus nigrescens* Rafinesque, 1810**En:** Tonguesole – **Fr:** Plagusie sombre**Habitat:** Benthic, over sand and mud of the continental shelf and slope, from 20 to 1 140 m depth**Spawning season:** Between May and October**EGGS**

Undescribed

**YOLK-SAC LARVAE**

Undescribed

**LARVAE****Figs. A-C**

**Distinctive characters:** body relatively short with a large head and abdominal region, laterally compressed since early stages of development; finger-shaped prolongation of gut at posterior ventral side of gut loop; 4 first anterior rays of dorsal fin considerably enlarged (these rays enlarge in a sequential order, from anterior to posterior)

**Preanus length:** less than 50% SL

**Pigmentation:** melanophores on ventral side of gut and on gut prolongation; melanophore on hindgut; group of ventral melanophores on anterior region of tail; group of dorsal and another ventral melanophore at about mid-tail in later larvae

**Length at flexion:** unknown

**PHOTOS**

by J.M. Rodriguez




6.0 mm SL

**Literature:** Alemany (1997), Padoa (1956g), Quero *et al.* (1986a), Sabatés (1988)


**CYNOGLOSSIDAE**

*Symphurus nigrescens* Rafinesque, 1810




A. 3.1 mm SL

Finger-shaped  
prolongation of gut



B. 4.6 mm SL



C. 7.5 mm SL



### 3. LITERATURE CITED

- Aboussouan, A.** 1964. Contribution à l'étude des oeufs et larves pelagiques des poissons téléostéens dans le Golfe de Marseille. *Revue Travaux de la Station Marine d'Endoume*, 32: 87–171
- Ahlstrom, E.H.** 1954. Distribution and abundance of egg and larval populations of the Pacific sardine. *Fishery Bulletin*, 93: 83–140
- Ahlstrom, E.H.** 1959. Vertical distribution of pelagic fish eggs and larvae off California and Baja California. *Fishery Bulletin*, 161: 107–146
- Ahlstrom, E.H. & Moser, H.G.** 1980. Characters useful in identification of pelagic marine fish eggs. *California Cooperative Oceanic Fisheries Investigations Reports*, 21: 121–131
- Ahlstrom, E.H., Sherman, K. & Smith, P.E.** 1973. Seagoing Operations in ichthyoplankton. In G. Hempel, ed. *Fish Egg and Larval Surveys (Contributions to a Manual)*, FAO Fisheries Technical Paper No. 122: 14–26
- Aleman, F.** 1997. *Ictioplancton del Mar Balear*. Univ. Illes Balears, Palma de Mallorca. (Ph.D. Thesis)
- Anderson, J.T.** 1988. A review of size dependent survival during pre-recruit stages of fishes in relation to recruitment. *Journal of Northwest Atlantic Fisheries Science*, 8: 55–56
- Armstrong, M.J., Connolly, P., Nash, R.D.M., Pawson, M.G., Alesworth, E., Coullahan, P.J., Dickey-Collas, M., Milligan, S.P., O'Neill, M.F., Witthames, P.R. & Woolner, L.** 2001. An application of the annual egg production method to estimate the spawning biomass of cod (*Gadus morhua* L.), plaice (*Pleuronectes platessa* L.) and sole (*Solea solea* L.) in the Irish Sea. *ICES Journal of Marine Science*, 58: 183–203
- Arata Jr., G.F.** 1954. A contribution to the life history of the swordfish, *Xiphias gladius*, Linnaeus, from the south Atlantic coast of the United States and the Gulf of Mexico. *Bulletin of Marine Science of the Gulf and Caribbean*, 4: 183–243
- Badcock, J.** 1984a. Gonostomatidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume I. pp. 284–301. UNESCO, Paris
- Badcock, J.** 1984b. Photichthyidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume I. pp. 318–324. UNESCO, Paris
- Badcock, J.** 1984c. Sternoptychidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume I. pp. 302–317. UNESCO, Paris
- Badcock, J. & Merret, N.R.** 1976. Midwater fishes in the eastern North Atlantic – I. Vertical distribution and associated biology in 30°N, 23°W, with developmental notes on certain myctophids. *Progress in Oceanography*, 7: 3–58
- Bailey, K.M. & Houde, E.D.** 1989. Predation on eggs and larvae of marine fishes and the recruitment problem. *Advances in Marine Biology*, 25: 1–83
- Bakun, A. & Parrish, R.H.** 1982. Turbulence, transport, and pelagic fish in the California and Peru Current systems. *California Cooperative Oceanic Fisheries Investigations Reports*, 23: 99–112
- Bauchot, M.–L.** 1986. Ophichthidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume II. pp. 577–585. UNESCO, Paris
- Bauchot, M.–L. & Hureau, J.–C.** 1986. Sparidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume II. pp. 883–907. UNESCO, Paris

- Bauchot, M.-L. & Saldanha, L.** 1986. Congridae. In P.J.P. Whitehead, M.-L. Bauchot, J.-C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume II. pp. 567–574. UNESCO, Paris
- Ben-Tuvia, A.** 1986. Mugilidae. In P.J.P. Whitehead, M.-L. Bauchot, J.-C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume III. pp. 1197–1204. UNESCO, Paris
- Berrien, P.L.** 1978. *Scomber scombrus* and *Scomber japonicus* in continental shelf waters between Massachusetts and Florida. *Fishery Bulletin*, 78: 95–115
- Bertolini, F.** 1933a. Famiglia Apogonidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 306–310
- Bertolini, F.** 1933b. Famiglia Serranidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 310–331
- Boehlert, G.W.** 1984. Scanning electron microscopy. In G. Moser, W.J. Richards, D.M. Cohen, M.P. Fahay, A.W. Kendall Jr. & S.L. Richardson, eds. *Ontogeny and systematics of fishes*. American Society of Ichthyologists and Herpetologists. Special Publication, No. 1, pp. 43–48
- Bridger, J.P.** 1956. On Day and Night Variation in Catches of Fish Larvae. *Journal du Conseil / Conseil Permanent International pour l'Exploration de la Mer*, 22: 42–57
- Briggs, J.C.** 1986. Gobiesocidae. In P.J.P. Whitehead, M.-L. Bauchot, J.-C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume III. pp. 1351–1359. UNESCO, Paris
- Brothers, E.B., Matthews, C.P. & Lasker, R.** 1976. Daily growth increments in otoliths from larval and adult fishes. *Fishery Bulletin*, 74: 1–8
- Brothers, E.B., Williams, D.M. & Sale, P.F.** 1983. Length of larval life in twelve families of fishes at “One Tree Lagoon”, Great Barrier Reef, Australia. *Marine Biology*, 76: 319–324
- Brownell, C.L.** 1979. Stages in the early development of 40 marine fish species with pelagic eggs from the Cape of Good Hope. *Ichthyological Bulletin of the J.L.B. Smith Institute of Ichthyology*, 40: 1–84
- Buckley, L.J.** 1979. Relationships between RNA/DNA ratio, prey density, and growth rate in Atlantic Cod (*Gadus morhua*) larvae. *Journal of the Fisheries Research Board of Canada*, 36: 1497–1502
- Buckley, L.J.** 1980. Changes in ribonucleic acid, deoxyribonucleic acid, and protein content during ontogenesis in wintert flounder, *Pseudopleuronectes americanus*, and effect of starvation. *Fishery Bulletin*, 77: 703–708
- Buckley, L.J.** 1984. RNA–DNA ratio: An index of larval fish growth in the sea. *Marine Biology*, 80: 291–298
- Camus, P. & Besseau, L.** 1986. Sparidae, *Spondyliosoma cantharus*. In G.A. Robinson, ed. *Fiches d'Identification du Zooplancton*, No. 177: 4 pp.
- Caruso, J.H.** 1986. Lophiidae. In P.J.P. Whitehead, M.-L. Bauchot, J.-C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume III. pp. 1362–1363. UNESCO, Paris
- Cipria, G.** 1939. Contributo alla conoscenza delle uova dello sviluppo embrionale e post-embrionale in varie specie di Gadidi. *Memorie del Regio Comitato Talassografico Italiano*, 263
- Clutter, R.I. & Anraku, M.** 1968. Avoidance of samplers in zooplankton sampling. In D.J. Tranter, ed. *UNESCO Monographs on Oceanographic Methodology, 2: Zooplankton sampling*. pp. 57–76. UNESCO Press, Paris
- Cogan, C.B., Todd, B.J., Lawton, P. & Noji, T.T.** 2009. The role of marine habitat mapping in ecosystem-based management. *ICES Journal of Marine Science*, 66: 2033–2042

- Cohen, D.M.** 1984. Argentinidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume I. pp. 386–391, UNESCO, Paris
- Collette, B.B.** 1986a. Coryphaenidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume II. pp. 845–846. UNESCO, Paris
- Collette, B.B.** 1986b. Scombridae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume II. pp. 981–997. UNESCO, Paris
- Collette, B.B. & Parin, N.V.** 1986. Belonidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume II. pp. 604–609. UNESCO, Paris
- Collette, B.B., Potthoff, T., Richards, W.J., Ueyanagi, S., Russo, J.L. & Nishikawa, Y.** 1984. Scombroidei: development and relationships. Development and Relationships. In G. Moser, W.J. Richards, D.M. Cohen, M.P. Fahay, A.W. Kendall Jr. & S.L. Richardson, eds. *Ontogeny and systematics of fishes*. American Society of Ichthyologists and Herpetologists, Special Publication, No. 1, pp. 591–620
- Conand, F. & Fagetti, E.** 1971. Description et distribution saisonnière des larves de sardinelles des côtes du Sénégal et de la Gambie en 1968 et 1969. *Cahiers ORSTOM, Series Oceanography*, IX: 293–318
- Cowan, J.H.J. & Shaw, R.F.** 2002. Recruitment. In L.A. Fuiman & R.G. Werner, eds. *Fishery science: the unique contributions of early life stages*. pp. 88–111. Blackwell Publishing, Oxford
- Cowen, R.K. & Guigand, C.M.** 2008. In situ ichthyoplankton imaging system (ISIIS): system design and preliminary results. *Limnology and Oceanography: Methods*, 6: 126–132
- Crec'hriou, R., Marinaro, J.Y. & Planes, S.** 2015. Advance in identification of pelagic eggs of Mediterranean teleostean fish: development of a new identification key. *Vie et Milieu*, 65: 47–61
- Cury, P. & Roy, C.** 1989. Optimal environmental window and pelagic fish recruitment success in upwelling areas. *Canadian Journal of Fisheries and Aquatic Sciences*, 46: 670–680
- Cushing, D.H.** 1974. The natural regulation of populations. In H.F.R. Jones, ed. *Sea fisheries research*. pp. 399–412. Elek Science, London
- Cushing, D.H.** 1990. Plankton production and year class strength in fish populations: an update of the match/mismatch hypothesis. *Advances in Marine Biology*, 26: 249–293
- D'Ancona, U.** 1931a. Famiglia Clupeidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 1–1
- D'Ancona, U.** 1931b. Famiglia Congridae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 102–114
- D'Ancona, U.** 1931c. Famiglia Engraulidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 1–16
- D'Ancona, U.** 1931d. Famiglia Ophichthyidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 128–139
- D'Ancona, U.** 1931e. Famiglia Scomberesocidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 157–164
- D'Ancona, U.** 1933a. Famiglia Gadidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 178–255
- D'Ancona, U.** 1933b. Famiglia Macrorhamphosidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 299–306

- Daniel, L.B. & Graves, J.E.** 1994. Morphometric and genetic identification of eggs of spring-spawning sciaenids in lower Chesapeake Bay. *Fishery Bulletin*, 92: 254–261
- De Gaetani, D.** 1934. Uova, sviluppo embrionale e stadi post-embrionali negli Sparidi. 2. – *Pagellus centrodontus* De La Roche. *Memorie del Regio Comitato Talassografico Italiano*, 209: 18 pp.
- De Gaetani, D.** 1937. Uova, sviluppo embrionale e stadi post-embrionali negli Sparidi. 5. – *Box boops* L. *Memorie del Regio Comitato Talassografico Italiano*, 241: 14 pp.
- Demir, M.** 1961. On the eggs and larvae of the *Trachurus trachurus* (L.) and *Trachurus mediterraneus* (Stdhnr.) from the Sea of Marmara and the Black Sea. *Rapports et Procès-Verbaux des Reunions, Conseil International pour L'exploration de la Mer*, 16: 317–320
- Demir, N.** 1972. The abundance and distribution of the eggs and larvae of some teleost fishes off Plymouth in 1969 and 1970. II. The postlarvae of *Callionymus*. *Journal of the Marine Biological Association of the United Kingdom*, 52 (4): 997–1010
- Demir, N.** 1982. On postlarvae and pelagic juveniles of rocklings *Gaidropsaurus mediterraneus* and *G. biscayensis*. *Journal of Marine Biological Association of the United Kingdom*, 62: 647–665
- Ditty, J.G., Shaw, R.F., Grimes, C.B. & Cope, J.S.** 1994. Larval development, distribution, and abundance of the common dolphin, *Coryphaena hippurus*, and pompano dolphin, *C. equiselis* (Family: Coryphaenidae), in the northern Gulf of Mexico. *Fishery Bulletin*, 92: 275–291
- Doherty, P.J.** 2002. Postrecruitment processes in the ecology of coral reef fish populations: a multifactorial perspective. In P.F. Sale, ed. *Coral reef fishes: dynamics and diversity in a complex ecosystem*. pp. 327–355. Academic Press, San Diego, California, USA
- Ehrenbaum, E.** 1924. Scombriformes. In *Report on the Danish Oceanographical Expeditions 1908–10 to the Mediterranean and Adjacent Seas*, Volume II, Biology (A11): 42 pp.
- Ege, V.** 1918. Stomiidae (*Stomias*). In *Report on the Danish Oceanographical Expeditions 1908–10 to the Mediterranean and Adjacent Seas*, Volume II, Biology (A4): 28 pp.
- Ege, V.** 1930. Sudidae (*Paralepis*). In *Report on the Danish Oceanographical Expeditions 1908–10 to the Mediterranean and Adjacent Seas*, Volume II, Biology (A13): 173 pp.
- Ehrich, S.** 1986. Macroramphosidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume II. pp. 627–627. UNESCO, Paris
- Fage, L.** 1910. Recherches sur les stades pélagiques de quelques Téléostéens de la mer de Nice (parages de Mônaco) et du golfe du Lion. *Annales de l'Institut Océanographique*, 1(7), 53 pp.
- Fage, L.** 1918. – Shore-fishes: Macroramphosidae, Ammodytidae, Atherinidae, Serranidae, Chlopteridae, Cepolidae, Scaridae, Mullidae, Pomacentridae, Labroidae, Caproidae, Gobiidae, Scorpaenidae, Triglidae, Cyclopteridae, Trachinidae, Uranoscopidae, Callionymidae, Blenniidae, Ophidiidae. In *Report on the Danish Oceanographical Expeditions 1908–10 to the Mediterranean and Adjacent Seas*, Volume II, Biology (A3): 154 pp.
- Fage, L.** 1920. Engraulidae, Clupeidae. In *Report on the Danish Oceanographical Expeditions 1908–10 to the Mediterranean and Adjacent Seas*, Volume II, Biology (A9): 140 pp.
- Fahay, M.P.** 1983. Guide to the Early Stages of Marine Fishes Occurring in the Western North Atlantic Ocean, Cape Hatteras to the Southern Scotian Shelf. *Journal of Northwest Atlantic Fishery Science*, 423 pp.
- Fahay, M.P.** 2007. Early stages of fishes in the western North Atlantic Ocean (Davis Strait, Southern Greenland and Flemish Cap to Cape Hatteras), Vol I and II. Northwest Atlantic Fisheries Organization. P.O. Box 638, Dartmouth, Nova Scotia, Canada
- Fives, J.M.** 1976. Labridae of the eastern North Atlantic. *ICES Fiches d'Identification du Zooplancton*, 7 pp.



- Fives, J.M.** 1986. Blenniidae of the North Atlantic (revised). *ICES Fiches d'Identification du Zooplancton*, 172: 6
- Ford, E.** 1922. On the young stages of *Blennius ocellaris* L., *Blennius pholis* L., and *Blennius gattorugine* L. *Journal of Marine Biological Association of the United Kingdom*, 12: 688–692
- Fox, C.J., Taylor, M.I., Pereyra, R., Villasana, M.I. & Rico, C.** 2005. TaqMan DNA technology confirms likely overestimation of cod (*Gadus morhua* L.) egg abundance in the Irish Sea: implications for the assessment of the cod stock and mapping of spawning areas using egg-based methods. *Molecular Ecology*, 14: 879–984
- France, R.L. & Peters, R.H.** 1997. Ecosystem differences in the trophic enrichment of  $^{13}\text{C}$  in aquatic food webs. *Canadian Journal of Fisheries and Aquatic Sciences*, 54: 1255–1258
- Fricke, R.** 1986. Callionymidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume III. pp. 1086–1093. UNESCO, Paris
- Froese, R. & Papisassi, C.** 1989. The use of modern relational databases for identification of fish larvae. *ICES CM/L*: 12, 10 pp.
- Froese, R. & Pauly, D.** 2016. FishBase. World Wide Web electronic publication. [www.fishbase.org](http://www.fishbase.org), (01/2016)
- García-Seoane, E., Álvarez-Colombo, G., Miquel, J., Rodríguez, J.M., Guevara-Fletcher, C., Álvarez, P. & Saborido-Rey, F.** 2016. Acoustic detection of larval fish aggregations in Galician waters (NW Spain). *Marine Ecology Progress Series*, 551: 31–44
- Garstang, W.** 1900. The impoverishment of the sea. *Journal of the Marine Biological Association of the United Kingdom*, 6: 1–69
- Gibbs, R.H.** 1984. Stomiidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume I. pp. 338–340. UNESCO, Paris
- Guittel, F.** 1920. *Lepadogaster*. In *Report on the Danish Oceanographical Expeditions 1908–10 to the Mediterranean and Adjacent Seas*, Volume II, Biology (A8): 9 pp.
- Haedrich, R.L.** 1986. Bramidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume III. pp. 847–853. UNESCO, Paris
- Heath, M.R.** 1992. Field investigations of the early life stages of marine fish. *Advances in Marine Biology*, 28: 2–174
- Hempel, G., ed.** 1974. Fish Egg and Larval Surveys (Contributions to a Manual), FAO Fisheries Technical Paper No. 122, 87 pp.
- Hjort, J.** 1914. Fluctuations in the great fisheries of Northern Europe viewed in the light of biological research. *Rapports et Procès – Verbaux des Réunions du Conseil International pour l'Exploration de la Mer*, 20: 228 pp.
- Hjort, J.** 1926. Fluctuations in the year classes of important food fishes. *ICES Journal of Marine Science*, 1: 5–38
- Houde, E.D.** 1987. Fish early life dynamics and recruitment variability. *American Fisheries Society Symposium*, 2: 17–29
- Houde, E.D.** 1997. Patterns and trends in larval-stage growth and mortality in teleost fish. *Journal of Fish Biology*, 51: 52–83
- Houde, E.D.** 2008. Emerging from Hjort's Shadow. *Journal of Northwest Atlantic Fisheries Science*, 41: 53–70

- Hulley, P.A.** 1984. Myctophidae. In P.J.P. Whitehead, M.-L. Bauchot, J.-C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume I. pp. 429–483. UNESCO, Paris
- Hunter, J.R.** 1981. Feeding ecology and predation of marine fish larvae. In R. Lasker, ed. *Marine fish larvae. Morphology, ecology and relation to fisheries*. pp. 34–77. University of Washington Press, Seattle
- Hunter, J.R., Macewicz, B.J., Lo, N.C. & Kimbrell, C.A.** 1992. Fecundity, spawning, and maturity of female Dover sole, *Microstomus pacificus*, with an evaluation of assumptions and precision. *Fishery Bulletin*, 90: 101–128
- Huntsman, S.A. & Barber, R.T.** 1977. Primary production off northwest Africa: the relationship to wind and nutrient conditions. *Deep Sea Research*, 24: 25–33
- Hureau, J.-C.** 1986a. Mullidae. In P.J.P. Whitehead, M.-L. Bauchot, J.-C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume II. pp. 877–882. UNESCO, Paris
- Hureau, J.-C.** 1986b. Triglidae. In P.J.P. Whitehead, M.-L. Bauchot, J.-C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume III. pp. 1230–1238. UNESCO, Paris
- Hureau, J.-C.** 1986c. Uranoscopidae. In P.J.P. Whitehead, M.-L. Bauchot, J.-C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume II. pp. 955–956. UNESCO, Paris
- Hureau, J.-C. & Litvinenko, N.I.** 1986. Scorpaenidae. In P.J.P. Whitehead, M.-L. Bauchot, J.-C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume III. pp. 1211–1229. UNESCO, Paris
- Iles, T.D. & Sinclair, M.** 1982. Atlantic Herring: Stock Discreteness and Abundance. *Science*, 215: 627–633
- Jespersen, P.** 1915. Sternoptychidae (*Argyropelecus* and *Sternoptyx*). In *Report on the Danish Oceanographical Expeditions 1908–10 to the Mediterranean and Adjacent Seas*, Volume II, Biology (A2): 49 pp.
- Jespersen, P. & Täning, A.V.** 1926. Mediterranean Sternoptychidae. In *Report on the Danish Oceanographical Expeditions 1908–10 to the Mediterranean and Adjacent Seas*, Volume II, Biology (A12): 59 pp.
- Johnson, R.K.** 1984. Evermannellidae. In P.J.P. Whitehead, M.-L. Bauchot, J.-C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume I. pp. 489–493. UNESCO, Paris
- Jones, A.** 1972. Studies on egg development and larval rearing of turbot, *Scophthalmus maximus* L. and brill *Scophthalmus rhombus* L., in the laboratory. *Journal of Marine Biological Association of the United Kingdom*, 52: 965–986
- Jones, G.P.** 1991. Postrecruitment processes in the ecology of coral reef fish populations: a multifactorial perspective. In P.F. Sale, ed. *The ecology of fishes on coral reefs*. pp. 294–328. Academic Press, San Diego, California, USA
- Katsanevakis, S., Stelzenmüller, V., South, A., Sørensen, T.K., Jones, P.J.S., Kerr, S., Badalamenti, F., Anagnostou, C., Breen, P., Chust, G., D’Anna, G., Duijn, M., Filatova, T., Fiorentino, F., Hulsman, H., Johnson, K., Karageorgis, A.P., Kröncke, I., Mirto, S., Pipitone, C., Portelli, S., Qiu, W., Reiss, H., Sakellariou, D., Salomidi, M., van Hoof, L., Vassilopoulou, V., Vega-Fernández, T., Vöge, S., Weber, A., Zenetos, A. & Hofstede, R.T.** 2011. Ecosystem-based marine spatial management: Review of concepts, policies, tools, and critical issues. *Ocean & Coastal Management*, 54: 807–820
- Kendall, A.W. & Duker, G.J.** 1998. The development of recruitment fisheries oceanography in the United States. *Fisheries Oceanography*, 7: 69–88

- Kendall, J., A.W., Ahlstrom, E.H. & Moser, G.** 1984. Early life history of fishes and their characters. In G. Moser, Richards, W.J., Cohen, D.M., Fahay, M.P., Kendall, J., A.W., Richardson, S.L., eds. *Ontogeny and systematics of fishes*. pp. 11–22. American Society of Ichthyologists and Herpetologists, La Jolla, California
- Kennedy, M. & Fitzmaurice, P.** 1968. Occurrence of Eggs of Bass *Dicentrarchus labrax* on the Southern Coasts of Ireland. *Journal of the Marine Biological Association of the United Kingdom*, 43: 585–592
- Ko, H.-L., Wang, Y.-T., Chiu, T.-S., Lee, M.-A., Leu, M.-Y., Chang, K.-Z., Chen, W.-Y. & Shao, K.-T.** 2013. Evaluating the Accuracy of Morphological Identification of Larval Fishes by Applying DNA Barcoding. *PLoS ONE*, 8: e53451
- Koslow, J.A. & Wright, M.** 2016. Ichthyoplankton sampling design to monitor marine fish populations and communities. *Marine Policy*, 68: 55–64
- Kramer, D.** 1969. Synopsis of the biological data on the Pacific mackerel. *FAO Fisheries Synopsis*, No. 40
- Kyle, H.M.** 1913. Flat-fishes (Heterostomata). In *Report on the Danish Oceanographical Expeditions 1908–10 to the Mediterranean and Adjacent Seas*, Volume II, Biology (A1), 150 pp.
- Laiz-Carrión, R., Quintanilla, J.M., Torres, A.P., Alemany, F. & García, A.** 2013. Hydrographic patterns conditioning variable trophic pathways and early life dynamics of bullet tuna *Auxis rochei* larvae in the Balearic Sea. *Marine Ecology Progress Series*, 475: 203–212
- Laiz-Carrión, R., Gerard, T., Uriarte, A., Malca, E., Quintanilla, J.M., Muhling, B.A., Alemany, F., Privoznik, S.L., Shiroza, A., Lamkin, J.T. & García, A.** 2015. Trophic ecology of Atlantic bluefin tuna (*Thunnus thynnus*) [corrected] larvae from the Gulf of Mexico and NW Mediterranean spawning grounds: A comparative stable isotope study. *PLoS ONE*, 10: e0138638
- Lasker, R.** 1981. Factors contributing to variable recruitment of the northern anchovy (*Engraulis mordax*) in the California Current: contrasting years, 1975 through 1978. *Rapports et Procès – Verbaux des Réunions du Conseil International pour l'Exploration de la Mer*, 178: 375–388
- Lasker, R.** 1985. Introduction: An Egg Production Method for Anchovy Biomass Assessment. In R. Lasker, ed. *An Egg Production Method for Estimating Spawning Biomass of Pelagic Fish: Application to the Northern Anchovy, Engraulis mordax*, Volume 36. pp. 1–4. NOAA Technical Report NMFS, Springfield, USA
- Lebour, M.V.** 1919. The young of the Gobiidæ from the neighbourhood of Plymouth. *Journal of the Marine Biological Association of the United Kingdom*, 12: 48–80
- Lebour, M. V.** 1925. Young anglers in captivity and some of their enemies. A study in a plunger jar. *Journal of the Marine Biological Association of the United Kingdom*, 13 (3): 721–734
- Lee, J.Y.** 1966. Oeufs et larves planctoniques de Poissons. *Revue des Travaux de l'Institut des Pêches Maritimes*, 30: 171–208
- Lo, N.C.-h., Hunter, J.R., Moser, H.G., Smith, P.E. & Methot, R.D.** 1992. The daily fecundity reduction method: a new procedure for estimating adult fish biomass. *ICES Journal of Marine Science*, 49: 209–215
- Lo Bianco, S., Bertolini, F., D'Ancona, U., Montalenti, G., Padoa, E., Ranzi, Sanzo, L., Sparta, A., Trtonese, E. & Vialli, M.** 1931–1956. Uova e stadi giovanili di Teleostei. Fauna e Flora del Golfo di Napoli, Monografia 38: 1–1164
- Lockwood, S.J., Nichols, J.H. & Dawson, W.A.** 1981. The estimation of a mackerel (*Scomber scombrus* L.) spawning stock size by plankton survey. *Journal of Plankton Research*, 3: 217–233
- Machinandarena, L., Mülle, M. & López, A.** 2003. Early life stages of development of the red porgy *Pagrus pagrus* (Pisces, Sparidae) in captivity, Argentina. *Investigaciones Marinas Valparaíso*, 31: 5–13
- Mahnken, C.V.W. & Jossi, J.W.** 1967. Flume experiments on the hydrodynamics of plankton nets. *ICES Journal of Marine Science*, 31: 38–45
- Marinaro, J.Y.** 1971. Contribution à l'étude des oeufs et larves pélagiques des poissons Méditerranéens. V. Oeufs pélagiques de la Baie d'Alger. *Pelagos*, 3: 1–118

- Marion, A.F.** 1894. Oeufs flottants et alevins observés dans le Golfe de Marseille durant l'année 1890. *Annales du Musée d'Histoire Naturelle de Marseille*, 4 (XI): 112–121, pls. I–II
- Matarese, A.C. & Sandknop E.M.** 1984. Identification of fish eggs. In G. Moser, Richards, W.J., Cohen, D.M., Fahay, M.P., Kendall, J., A.W., Richardson, S.L., eds. *Ontogeny and systematics of fishes*. pp. 27–31. American Society of Ichthyologists and Herpetologist, La Jolla, California
- McGowan, J.A. & Brown, D. M.** 1966. A new opening–closed paired zooplankton net. *University of California, Scripps Institution of Oceanography*. No. 66–23, pp. 1–56
- Methot, R.D.** 1983. Seasonal variation in survival of larval northern anchovy, *Engraulix mordax*, estimated from the age distribution of juveniles. *Fishery Bulletin*, 81: 741–750
- Miller, P.J.** 1986. Gobiidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume III. pp. 1019–1085. UNESCO, Paris
- Miller, T.J., Crowder, L.B., Rice, J.A. & Marschall, E.A.** 1988. Larval size and recruitment mechanisms in fishes: Toward a conceptual framework. *Canadian Journal of Fisheries and Aquatic Sciences*, 45: 1657–1670
- Minagawa, M. & Wada, E.** 1984. Stepwise enrichment of  $\delta^{15}\text{N}$  along food chains: Further evidence and the relation between  $\delta^{15}\text{N}$  and animal age. *Geochimica and Cosmochimica Acta*, 48: 1113–1140
- Mito, S.** 1960. Egg development and hatched larvae of the common dolphin–fish, *Coryphaena hippurus* Linne. *Bulletin of the Japanese Society of Scientific Fisheries*, 26: 223–226
- Montalenti, G.** 1937a. Famiglia Mullidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 391–398
- Montalenti, G.** 1937b. Famiglia Sciaenidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 399–406
- Morse, W.W.** 1989. Catchability, growth, and mortality of larval fishes. *Fishery Bulletin*, 87: 417–446
- Moser, H.G.** 1996. Introduction. In H.G. Moser, ed. *The early stages of fishes in the California Current region*. pp. 1–72. California Cooperative Oceanic Fisheries Investigations. Atlas, 33
- Munk, P. & Nielsen, J.G.** 2005. Eggs and larvae of North Sea fishes. *Biofilia*, Frederiksberg, Demark
- Murphy, G.I. & Clutter, R.I.** 1972. Sampling anchovy larvae with a plankton purse seine. *Fishery Bulletin*, 70: 769–798
- Murua, H., Ibaibarriaga, L., Álvarez, P., Santos, M., Korta, M., Santurtun, M. & Motos, L.** 2010. The daily egg production method: A valid tool for application to European hake in the Bay of Biscay? *Fisheries Research*, 104: 100–110
- Nakamura, I.** 1986. Xiphiidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume II. pp. 1006–1007. UNESCO, Paris
- Nichols, J.H.** 1976. Soleidae. *ICES Fiches d'Identification du Zooplancton*, 150/151: 1–10
- Nielsen, J.G.** 1986a. Bothidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume III. pp. 1294–1298. UNESCO, Paris
- Nielsen, J.G.** 1986b. Ophidiidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume III. pp. 1158–1166. UNESCO, Paris
- Nielsen, J.G.** 1986c. Pleuronectidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume III. pp. 1299–1307. UNESCO, Paris

- Nielsen, J.G.** 1986d. Scophthalmidae. In P.J.P. Whitehead, M.-L. Bauchot, J.-C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume III. pp. 1287–1293. UNESCO, Paris
- Olivar, M.P.** 1986. Development and distribution of *Parablennius pilicornis* (Cuvier) larvae (teleostei: Blenniidae) off Namibia. *South African Journal of Aquatic Science*, 4: 193–201
- Olivar, M.P. & Fortuño, J.M.** 1991. Guide to Ichthyoplankton of the Southeast Atlantic (Benguela Current Region). *Scientia Marina*, 55: 1–363
- Olivar, M.P. & Palomera, I.** 1994. Ontogeny and distribution of *Hygophum benoiti* (Pisces, Myctophidae) of the north western Mediterranean. *Journal of Plankton Research*, 16: 977–991
- Padoa, E.** 1956a. Famiglia Ammodytidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 709–714
- Padoa, E.** 1956b. Famiglia Blenniidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 720–745
- Padoa, E.** 1956c. Famiglia Bothidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 783–831
- Padoa, E.** 1956d. Famiglia Bramidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 522–526
- Padoa, E.** 1956e. Famiglia Carangidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 548–570
- Padoa, E.** 1956f. Famiglia Carapidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 761–774
- Padoa, E.** 1956g. Famiglia Cynoglossidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 870–877
- Padoa, E.** 1956h. Famiglia Gobiidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 649–678
- Padoa, E.** 1956i. Famiglia Lophiidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 878–888
- Padoa, E.** 1956j. Famiglia Ophidiidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 755–761
- Padoa, E.** 1956k. Famiglia Pomacentridae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 573–576
- Padoa, E.** 1956l. Famiglia Scombridae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 471–478
- Padoa, E.** 1956m. Famiglia Soleidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 838–869
- Padoa, E.** 1956n. Famiglia Trachinidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 687–697
- Padoa, E.** 1956o. Famiglia Trichiuridae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 508–513
- Padoa, E.** 1956p. Famiglia Triglididae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 627–640
- Padoa, E.** 1956q. Famiglia Uranoscopidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 714–720
- Padoa, E.** 1956r. Famiglia Xiphiidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 516–521

- Padoa, E.** 1956s. Famiglia Callionymidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 697–708
- Padoa, E.** 1956t. Famiglia Pleuronectidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 831–838
- Palomera, I.** 1983. Desarrollo larvario de *Notoscopelus elongatus elongatus* (Costa, 1844) y *Notoscopelus bolini* (Nafpaktitis, 1975). *Investigación Pesquera*, 47: 263–276
- Palomera, I., Olivar, M.P. & Morales-Nin, B.** 2005. Larval development and growth of the European hake *Merluccius merluccius* in the northwestern Mediterranean. *Scientia Marina*, 69: 251–258
- Pannella, G.** 1971. Fish otoliths: daily growth layers and periodical patterns. *Science*, 173: 1124–1127
- Parin, N.V.** 1986. Trichiuridae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume II. pp. 976–980. UNESCO, Paris
- Parker, K.** 1980. A direct method for estimating northern anchovy, *Engraulis mordax*, spawning biomass. *Fishery Bulletin*, 78: 541–547
- Petersen, C.J.G.** 1894. On the biology of our flatfishes and on the decrease of our flatfisheries. *Report of the Danish Biological Station*, 4: 147 pp.
- Peterson, B.J. & Fry, B.** 1987. Stable Isotopes in Ecosystem Studies. *Annual Review of Ecology and Systematics*, 18: 293–320
- Pikitch, E.K., Santora, C., Babcock, E.A., Bakun, A., Bonfil, R., Conover, D.O., Dayton, P., Doukakis, P., Fluharty, D., Heneman, B., Houde, E.D., Link, J., Livingston, P.A., Mangel, M., McAllister, M.K. Pope, J. & Sainsbury, K.J.** 2004. Ecosystem–Based Fishery Management. *Science*, 305: 346–347
- Post, A.** 1984. Paralepididae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume I. pp. 498–508. UNESCO, Paris
- Post, D.M.** 2002. Using stable isotopes to estimate trophic position: models, methods, and assumptions. *Ecology*, 83: 703–718
- Pothoff, T.** 1984. Clearing and Staining Techniques. In H.G. Moser, W.J. Richards, D.M. Cohen, M.P. Fahay, A.W. Kendall & S.L. Richardson, eds. *Ontogeny and systematics of fishes*. pp. 35–37. American Society of Ichthyologists and Herpetologists, La Jolla, California
- Powles, H. & Markle, D.F.** 1984. Identification of larvae. In H.G. Moser, W.J. Richards, D.M. Cohen, M.P. Fahay, A.W. Kendall & S.L. Richardson, eds. *Ontogeny and Systematics of Fishes*. pp. 31–33. American Society of Ichthyologists and Herpetologists.
- Puncher, G.N., Arrizabalaga, H., Alemany, F., Cariani, A., Oray, I.K., Karakulak, F.S., Basilone, G., Cuttitta, A., Mazzola, S. & Tinti, F.W.** 2015a. Molecular Identification of Atlantic Bluefin Tuna (*Thunnus thynnus*, Scombridae) larvae and development of a DNA character–based identification key for Mediterranean Scombrids. *PLoS ONE*, 10: e0130407
- Puncher, G.N., Alemany, F., Arrizabalaga, H., Cariani, A. & Tinti, F.W.** 2015b. Misidentification of bluefin tuna larvae: a call for caution and taxonomic reform. *Reviews in Fish Biology and Fisheries*, 25: 485–502
- Quero, J.–C.** 1986a. Caproidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume II. pp. 777–779. UNESCO, Paris
- Quero, J.–C.** 1986b. Zeidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume II. pp. 769–772. UNESCO, Paris
- Quero, J.–C., Desoutter, M. & Lagardère, F.** 1986a. Cynoglossidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume III. pp. 1325–1328. UNESCO, Paris

- Quero, J.-C., Desoutter, M. & Lagardère, F.** 1986b. Soleidae. In P.J.P. Whitehead, M.-L. Bauchot, J.-C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume III. pp. 1308–1324. UNESCO, Paris
- Quignard, J.-P.** 1967. Lœuf et la larve du Labridé *Symphodus (Crenilabrus) melops* (Linné, 1758). Influence de différents facteurs physicochimique sur la durée du développement embryonnaire. *Revue des travaux de l'Institut des pêches maritimes*, 31: 355–358
- Quignard, J.-P. & Pras, A.** 1986a. Labridae. In P.J.P. Whitehead, M.-L. Bauchot, J.-C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume II. pp. 919–942. UNESCO, Paris
- Quignard, J.-P. & Pras, A.** 1986b. Pomacentridae. In P.J.P. Whitehead, M.-L. Bauchot, J.-C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume II. pp. 916–918. UNESCO, Paris
- Raffaele, F.** 1888. Le uova galleggianti e le larve dei Teleostei nel golfo di Napoli. *Mittheilungen aus der Zoologischen Station zu Neapel*, 8: 1–85, pls. I–V
- Ranzi, S.** 1933. Famiglia Sparidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 332–382
- Raymont, J.E.G.** 1983. *Plankton and Productivity in the Oceans. Volume 2 – Zooplankton*. 2nd edition. Pergamon Press
- Reay, P.J.** 1986. Ammodytidae. In P.J.P. Whitehead, M.-L. Bauchot, J.-C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North-eastern Atlantic and the Mediterranean*, Volume II. pp. 945–950. UNESCO, Paris
- Richards, W.J.** 2006. *Early stages of Atlantic fishes: an identification guide for the western central North Atlantic, Volumes I and II*. Taylor and Francis, New York
- Roule, L. & Angel, F.** 1930. Larves et Alevins de Poissons provenant des Croissières du Prince Albert I de Monaco. *Résultats Des Campagnes Scientifiques Accomplies Sur Son Yacht Par Albert Ier Prince Souverain De Monaco*, Fascicule LXXIX: 148 pp, pls. I–VI
- Russell, F.S.** 1976. *The eggs and planktonic stages of British marine fishes*. Academic Press, London
- Sabatés, A.** 1988. *Sistemática y distribución espacio-temporal del ictioplancton en la costa catalana*. Universidad de Barcelona, Barcelona (Tesis Doctoral)
- Sabatés, A.** 1991. Larval development of *Lepidorhombus boscii* (Risso, 1810) (Pleuronectiformes) in the Northwestern Mediterranean. *Scientia Marina*, 55: 543–546
- Sabatés, A., Demestre, M. & Sanchez, P.** 1990. Revision of the family Ammodytidae (Perciformes) in the Mediterranean with the first record of *Gymnammodytes semisquamatus*. *Journal of the Marine Biological Association of the United Kingdom*, 70: 493–504
- Sale, P.F.** 1978. Coexistence of coral reef fishes — a lottery for living space. *Environmental Biology of Fishes*, 3: 85–102
- Sale, P.F.** 1991. Reef fish communities: open nonequilibrium systems. In P.F. Sale, ed. *The ecology of fishes on coral reefs*. pp. 564–598. Academic Press, San Diego
- Sanzo, L.** 1931a. Famiglia Argentinidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 22–31
- Sanzo, L.** 1931b. Famiglia Sternoptichidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 42–81
- Sanzo, L.** 1931c. Famiglia Microstomidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: pp. 32–42
- Sanzo, L.** 1931d. Uova e larve di *Zeus faber* L. *Archivio Zoologico Italiano* 15: 475–482
- Sanzo, L.** 1935. Uova, sviluppo embrionale, stadi larvali, postalarvali e giovanili di Sternoptichidae e Stomiidae. III *Maurolicus pennanti* (Walb.). *Memorie del Regio Comitato Talassografico Italiano*, 2: 122–180

- Sanzo, L.** 1936. Contributi alla conoscenza dello sviluppo embrionario e postembrionario nei Mugilidae. I Uova e larve di *Mugil cephalus* Cuv. ottenute per fecondazione artificiale. II. Uova e larve di *Mugil chelo* Cuv. *Memorie del Regio Comitato Talassografico Italiano*, 230: 1–15
- Sanzo, L.** 1956a. Famiglia Zeidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 461–466
- Sanzo, L.** 1956b. Famiglia Caproidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 466–471
- Saville, A.** 1958. Mesh Selection in Plankton Nets. *Journal du Conseil / Conseil Permanent International pour l'Exploration de la Mer*, 23: 192–201
- Schmidt, J.** 1905. The pelagic postlarval stages of the Atlantic species of *Gadus*. *Medd Komm Havunders, Ser Fisk.*, 1: 1–74
- Schmidt, J.** 1918. Argentinidae, Microstomidae, Opisthoproctidae, Mediterranean Odontostomidae. In *Report on the Danish oceanographical expeditions 1908–10 to the Mediterranean and adjacent seas*, Volume II: Biology (A5): 40 pp.
- Schmidt, J. & Strubberg, A.** 1918. Mediterranean Bramidae and Trichiuridae. In *Report on the Danish oceanographical expeditions 1908–10 to the Mediterranean and adjacent seas*, Volume II: Biology (A6): 15 pp.
- Schnakenbeck, W.** 1931. Carangidae. *Report on the Danish Oceanographical Expeditions 1908–10 to the Mediterranean and Adjacent Seas*, Volume II: Biology (A14): 20 pp.
- Seaton, D.D. & Bailey, R.S.** 1971. The identification and development of the eggs and larvae of the blue whiting *Micromesistius poutassou* (Risso). *Journal du Conseil / Conseil Permanent International pour l'Exploration de la Mer*, 34: 76–83
- Sinclair, M.** 1988. *Marine populations: an essay on population regulation and speciation*. University of Washington Press, Seattle
- Smith, P.E. & Clutter, R.I.** 1965. Hydrodynamics of flow and collection in plankton nets. In *Ocean Science and Ocean engineering*, Volume 1. pp. 515. American Society of Limnology and Oceanography, Marine Technology Society, Washington D.C.
- Smith, P.E. & Richardson, S.L.** 1977. *Standard techniques for pelagic fish egg and larva surveys*. FAO Technical Paper No. 175
- Smith–Vaniz, W.F.** 1986. Carangidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume II. pp. 815–844. UNESCO, Paris
- Sparta, A.** 1936. Contributo alla conoscenza di uova, stadi embrionali e post–embrionali in *Macrorhamphosus scolopax*. *Memorie del Regio Comitato Talassografico Italiano*, 225: 14 pp.
- Sparta, A.** 1956a. Famiglia Labridae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 576–593
- Sparta, A.** 1956b. Famiglia Scorpaenidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 599–626
- Sulak, K.J.** 1984. Synodontidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume I. pp. 405–411 UNESCO, Paris
- Svetovidov, A.N.** 1986a. Gadidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume II. pp. 680–710. UNESCO, Paris
- Svetovidov, A.N.** 1986b. Merlucciidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume II. pp. 677–679. UNESCO, Paris



- Stratoudakis, Y., Bernal, M., Ganiats, K. & Uriarte, A.** 2006. The daily egg production method: recent advances, current applications and future challenges. *Fish and Fisheries*, 7: 35–57
- Täning, A.V.** 1918. Mediterranean Scopelidae (*Saurus*, *Aulopus*, *Chlorophthalmus* and *Myctophum*). In *Report on the Danish Oceanographical Expeditions 1908–10 to the Mediterranean and Adjacent Seas*, Volume II, Biology (A7): 154 pp.
- Täning, A. V.** 1923. *Lophius*. In *Report on the Danish Oceanographical Expeditions 1908–10 to the Mediterranean and Adjacent Seas*, Volume II, Biology (A10): 30 pp.
- Täning, A.V.** 1961. Larval and postlarval stages of *Sebastes* species and *Helicolenus dactylopterus*. *Rapports et procès-verbaux des réunions / Conseil permanent international pour l'exploration de la mer*, 1501: 234–240
- Tortonese, E.** 1956a. Famiglia Paralepididae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 899–913
- Tortonese, E.** 1956b. Famiglia Synodidae. In *Uova, larve e stadi giovanili di Teleostei*. Fauna e flora del Golfo di Napoli, Monografia 38: 890–894
- Tortonese, E.**, 1970. Osteichthyes (Pesci ossei), Parte Prima, Fauna italiana, Calderini. Bologna, Vol. 10
- Tortonese, E.** 1975. Osteichthyes (Pesci ossei), Parte Seconda. Fauna italiana, Calderini, Bologna, Vol. 11
- Tortonese, E.** 1986a. Apogonidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume II. pp. 803–809. UNESCO, Paris
- Tortonese, E.** 1986b. Centranchidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume II. pp. 908–911. UNESCO, Paris
- Tortonese, E.** 1986c. Cepolidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume II. pp. 810–811. UNESCO, Paris
- Tortonese, E.** 1986d. Serranidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume II. pp. 780–792. UNESCO, Paris
- Tranter, D.J. & Smith, P.E.** 1968. Filtration performance. In D.J. Tranter, ed. *Zooplankton sampling*. pp. 27–56. UNESCO Press, Paris
- Tucker, J.W. & Laroche, J.L.** 1984. Radiographic Techniques in Studies of young Fishes. In G. Moser, Richards, W.J., Cohen, D.M., Fahay, M.P., Kendall, J., A.W., Richardson, S.L., ed. *Ontogeny and Systematics of Fishes*. pp. 37–40. American Society of Ichthyologists and Herpetologists. Special Publication Number 1
- Vannucci, M.** 1968. Loss of organisms through the meshes. In D.J. Tranter, ed. *Zooplankton Sampling*. pp. 77–86. UNESCO Press, Paris
- Victor, B.C.** 1986. Duration of the planktonic larval stage of one hundred species of Pacific and Atlantic wrasses (family Labridae). *Marine Biology*, 90: 317–326
- Werner, E.E. & Gilliam, J.F.** 1984. The ontogenetic niche and species interactions in size–structured populations. *Annual Review of Ecology and Systematics*, 15: 393–425
- Whitehead, P.J.P.** 1984a. Clupeidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume II. pp. 268–281. UNESCO, Paris
- Whitehead, P.J.P.** 1984b. Engraulidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume II. pp. 282–283. UNESCO, Paris
- Zander, C.D.** 1986. Blenniidae. In P.J.P. Whitehead, M.–L. Bauchot, J.–C. Hureau, J. Nielsen & E. Tortonese, eds. *Fishes of the North–eastern Atlantic and the Mediterranean*, Volume III. pp. 1096–1112. UNESCO, Paris

## 4. INDEX

### A

Aguja	98
Alacha	30
<i>alalunga</i> , <i>Thunnus</i>	196, 197
Albacore	196
Allache	30
AMMODYTIDAE	20, 160, 161
Anchois	32
Angler(=Monk)	94
<i>anthias</i> , <i>Anthias</i>	114, 115
<i>Anthias anthias</i>	114, 115
<i>aper</i> , <i>Capros</i>	202, 203
APOGONIDAE	19, 120, 121
<i>Apogon imberbis</i>	120, 121
<i>argenteus</i> , <i>Gadiculus</i>	78, 79
Argentina sphyraena	34, 35
Argentine	34
ARGENTINIDAE	18, 34, 35
<i>Argyropelecus hemigymnus</i>	42, 43
Armless snake eel	24
Arnoglosse de Méditerranée	210
<i>Arnoglossus</i>	210, 212
<i>Arnoglossus laterna</i>	210, 211
Atlantic bluefin tuna	13, 198
Atlantic bonito	190
Atlantic chub mackerel	192
Atlantic horse mackerel	124
Atlantic lizardfish	50
Atlantic mackerel	194
Atlantic pomfret	128
<i>attenuata</i> , <i>Vinciguerria</i>	46, 47
Atún blanco	196
Atún rojo del Atlántico	198
<i>aurita</i> , <i>Sardinella</i>	30, 31
<i>Auxis rochei</i>	188, 189

### B

Bacaladilla	80
<i>balbo</i> , <i>Evermannella</i>	56, 57
Balbo sabretooth	56
Barbada	84
<i>barbatum</i> , <i>Ophidion</i>	90, 91
<i>barbatus</i> , <i>Mullus</i>	142, 143, 145
Barbue	206
Bar européen	112
Barrinaire	160
Baudroie commune	94
Bécasse de mer	102
<i>belone</i> , <i>Belone</i>	98, 99
<i>Belone belone</i>	98, 99
BELONIDAE	12, 19, 98, 99
<i>benoiti</i> , <i>Hygophum</i>	65
<i>Benthosema glaciale</i>	58, 59
Bertorella	86
Besugo	134
<i>bimaculata</i> , <i>Diplecogaster</i>	96, 97
<i>biscayensis</i> , <i>Gaidropsarus</i>	84, 85

Blackbelly rosefish	104
Black scorpionfish	106
Black seabream	138
Blackspot(=red) seabream	134
BLENNIIDAE	20, 168–175
<i>Blennius ocellaris</i>	168, 169
Blue whiting(=Poutassou)	80
Boa dragonfish	48
Boarfish	202
<i>boa</i> , <i>Stomias</i>	48, 49
Bobí	180
Boga	130
<i>bogaraveo</i> , <i>Pagellus</i>	134, 135
Bogue	130
Bonite à dos rayé	190
Bonito del Atlántico	190
Bonitou	188
<i>boops</i> , <i>Boops</i>	130, 131
<i>Boops boops</i>	130, 131
Boquerón	32
Borracho	108
<i>boscii</i> , <i>Lepidorhombus</i>	204, 205
BOTHIDAE	12, 21, 210–213
<i>Bothus podas</i>	212, 213
<i>brama</i> , <i>Brama</i>	128, 129
<i>Brama brama</i>	128, 129
BRAMIDAE	20, 128, 129
<i>braueri</i> , <i>Cyclothone</i>	40, 41
Brill	206
Brown comber	118
<i>Buglossidium luteum</i>	216, 217
Bullet tuna	188
Butterfly blenny	168

### C

Caballa del Atlántico	194
Cabrilla	116
<i>cabrilla</i> , <i>Serranus</i>	116, 117
Cabruza	174
CALLIONYMIDAE	20, 176, 177
<i>Callionymus</i>	176, 177
<i>Callionymus lyra</i>	176, 177
<i>Callionymus maculatus</i>	177
<i>Callionymus reticulatus</i>	177
<i>cantharus</i> , <i>Spondyliosoma</i>	138, 139
CAPROIDAE	21, 202, 203
<i>Capros aper</i>	202, 203
Caramel	140
CARANGIDAE	20, 122–125
CARAPIDAE	12, 19, 92, 93
Cardinal fish	120, 146
Cardine à quatre taches	204
Carpintero	60
<i>caudatus</i> , <i>Lepidopus</i>	186, 187
<i>cavillone</i> , <i>Lepidotrigla</i>	110, 111
CENTRACANTHIDAE	20, 140, 141
CENTRISCIDAE	19, 102, 103
<i>cephalus</i> , <i>Mugil</i>	148, 149
<i>Cepola macrophthalma</i>	146, 147
CEPOLIDAE	20, 146, 147

*Ceratoscopelus maderensis* . . . . . 60, 61  
 Chafarrocas . . . . . 96  
 Chinchard à queue jaune . . . . . 122  
 Chinchard d'Europe . . . . . 124  
 Chopa . . . . . 138  
*chromis*, *Chromis* . . . . . 150, 151  
*Chromis chromis* . . . . . 150, 151  
 Cicerelle de Méditerranée . . . . . 160  
*cicerelus*, *Gymnammodytes* . . . . . 160, 161  
 CLUPEIDAE . . . . . 12, 18, 28–31  
*colias*, *Scomber* . . . . . 192, 193  
 Comber . . . . . 116  
 Common dolphinfish . . . . . 126  
 Common goby . . . . . 184  
 Common sole . . . . . 220  
*conger*, *Conger* . . . . . 26, 27  
*Conger conger* . . . . . 26, 27  
 Congre d'Europe . . . . . 26  
 CONGRIDAE . . . . . 18, 26, 27  
 Congrio común . . . . . 26  
 Coq . . . . . 120, 146  
*coregonoides*, *Paralepis* . . . . . 54, 55  
*Coris julis* . . . . . 152, 153  
 Corkwing wrasse . . . . . 158  
*Coryphaena hippurus* . . . . . 126, 127  
 CORYPHAENIDAE . . . . . 20, 126, 127  
 Coryphène commune . . . . . 126  
 Crénilabre mélops . . . . . 158  
*crocodilus*, *Lampanyctus* . . . . . 66, 67, 69  
 Crystal goby . . . . . 178  
*Crystallogobius linearis* . . . . . 178, 179  
*Ctenolabrus rupestris* . . . . . 154, 155  
 Cuckoo wrasse . . . . . 156  
*Cylothone braueri* . . . . . 40, 41  
 CYNOGLOSSIDAE . . . . . 21, 222, 223

## D

*dactylopterus*, *Helicolenus* . . . . . 104, 105  
*Dalophis imberbis* . . . . . 24, 25  
 Damsel fish . . . . . 150  
*dentatus*, *Echiodon* . . . . . 92, 93  
*Diaphus holti* . . . . . 62, 63  
*Dicentrarchus labrax* . . . . . 112, 113  
*Diplecogaster bimaculata* . . . . . 96, 97  
*Diplodus* . . . . . 132  
*Diplodus sargus* . . . . . 132, 133  
*dofleini*, *Lobianchia* . . . . . 70, 71  
 Dofleini's lantern fish . . . . . 70  
 Dorade grise . . . . . 138  
 Dorade rose . . . . . 134  
*draco*, *Trachinus* . . . . . 164, 165

## E

*Echiichthys vipera* . . . . . 162, 163  
*Echiodon dentatus* . . . . . 92, 93  
 Eckström's topknot . . . . . 208  
*elongatus*, *Notoscopelus* . . . . . 74, 75  
*encrasicolus*, *Engraulis* . . . . . 32, 33  
 ENGRAULIDAE . . . . . 12, 18, 32, 33  
*Engraulis encrasicolus* . . . . . 32, 33

Escorpión . . . . . 164  
 Espadon . . . . . 200  
 Estomias . . . . . 48  
 Estornino . . . . . 192  
 European anchovy . . . . . 32  
 European conger . . . . . 26  
 European flounder . . . . . 214  
 European hake . . . . . 88  
 European pilchard . . . . . 28  
 European seabass . . . . . 112  
*Eutrigla gurnardus* . . . . . 108, 109  
*Evermannella balbo* . . . . . 56, 57  
 EVERMANNELLIDAE . . . . . 18, 56, 57

## F

*faber*, *Zeus* . . . . . 100, 101  
 Faneca . . . . . 82  
 Faneca plateada . . . . . 78  
 Flathead grey mullet . . . . . 148  
*flesus*, *Platichthys* . . . . . 214, 215  
 Flet d'Europe . . . . . 214  
 Four-spot megrim . . . . . 204

## G

*Gadiculus argenteus* . . . . . 78, 79  
 GADIDAE . . . . . 12, 18, 78–83  
*Gaidropsarus biscayensis* . . . . . 84, 85  
*Gaidropsarus mediterraneus* . . . . . 86, 87  
 Gallano . . . . . 156  
 Gallineta . . . . . 104  
 Gallo de cuatro manchas . . . . . 204  
 Garfish . . . . . 98  
 Garrick . . . . . 40  
*gattorugine*, *Parablennius* . . . . . 174, 175  
 Germon . . . . . 196  
 Girelle . . . . . 152  
*glaciale*, *Benthoosema* . . . . . 58, 59  
 Glacier lantern fish . . . . . 58  
*gladius*, *Xiphias* . . . . . 200, 201  
 Gobie commun . . . . . 184  
 Gobie paganel . . . . . 180  
 GOBIESOCIDAE . . . . . 19, 96, 97  
 GOBIIDAE . . . . . 12, 21, 178–185  
*Gobius paganellus* . . . . . 180, 181  
 Goldsinny-wrasse . . . . . 154  
 Golleta . . . . . 218  
 GONOSTOMATIDAE . . . . . 18, 40, 41  
 Grande castagnole . . . . . 128  
 Grande vive . . . . . 164  
 Greater weever . . . . . 164  
 Grey gurnard . . . . . 108  
 Grondin gris . . . . . 108  
*guilleti*, *Lebetus* . . . . . 182, 183  
 Guillet's goby . . . . . 182  
*gurnardus*, *Eutrigla* . . . . . 108, 109  
*Gymnammodytes cicerelus* . . . . . 160, 161

## H

Hacha de plata . . . . . 42

Half-naked hatchetfish	42
<i>Helicolenus dactylopterus</i>	104, 105
<i>hemigymnus</i> , <i>Argyropelecus</i>	42, 43
<i>hepatus</i> , <i>Serranus</i>	118, 119
<i>hippurus</i> , <i>Coryphaena</i>	126, 127
<i>holti</i> , <i>Diaphus</i>	62, 63
<i>hygommii</i> , <i>Hygophum</i>	65
<i>Hygophum benoiti</i>	65
<i>Hygophum hygommii</i>	65
<i>Hygophum</i>	64, 65

**I**

<i>imberbis</i> , <i>Apogon</i>	120, 121
<i>imberbis</i> , <i>Dalophis</i>	24, 25

**J**

Japonés	70
Japuta	128
<i>jayakari</i> , <i>Lestidiops</i>	52, 53
John dory	100
Julia	152
<i>julis</i> , <i>Coris</i>	152, 153
Jurel	124
Jurel méditerranéen	122

**L**

<i>labrax</i> , <i>Dicentrarchus</i>	112, 113
LABRIDAE	20, 152–159
<i>Labrus mixtus</i>	156, 157
<i>Lampanyctus crocodilus</i>	66, 67, 69
<i>Lampanyctus pusillus</i>	68, 69
Lampuga	126
Lanterne à grandes écailles	76
Lanterne courte	62
Lanterne de Dofleini	70
Lanterne de Madère	60
Lanterne glaciaire	58
Large-scaled gurnard	110
Large scale lantern fish	76
<i>laterna</i> , <i>Arnoglossus</i>	210, 211
<i>Lebetus guillesti</i>	182, 183
Lenguado común	220
<i>Lepidopus caudatus</i>	186, 187
<i>Lepidorhombus boschii</i>	204, 205
<i>Lepidotrigla cavillone</i>	110, 111
Lesser weever	162
<i>Lestidiops jayakari</i>	52, 53
<i>Lestidiops sphyrenoides</i>	52, 53
<i>linearis</i> , <i>Crystalllogobius</i>	178, 179
Linterna	76
Linterna oscura	58
<i>Lipophrys pholis</i>	170, 171
<i>lyra</i> , <i>Callionymus</i>	176, 177
<i>Lobianchia dofleini</i>	70, 71
Longspine snipefish	102
LOPHIIDAE	12, 19, 94, 95
<i>Lophius piscatorius</i>	94, 95
LOTIDAE	19, 84–87
Lubina	112

<i>luscus</i> , <i>Trisopterus</i>	82, 83
<i>luteum</i> , <i>Buglossidium</i>	216, 217
<i>lyra</i> , <i>Callionymus</i>	176, 177

**M**

<i>macrophthalmia</i> , <i>Cepola</i>	146, 147
<i>Macroramphosus scolopax</i>	102, 103
<i>maculatus</i> , <i>Callionymus</i>	177
Madeira lantern fish	60
<i>maderensis</i> , <i>Ceratoscopelus</i>	60, 61
Maquereau commun	194
Maquereau espagnol	192
Maurolucus muelleri	44, 45
Mediterranean bigeye rockling	84
Mediterranean horse mackerel	122
Mediterranean sand eel	160
Mediterranean scaldfish	210
<i>mediterraneus</i> , <i>Gaidropsarus</i>	86, 87
<i>mediterraneus</i> , <i>Trachurus</i>	122–124
<i>melops</i> , <i>Symphodus</i>	158, 159
Melva(=Melvera)	188
Merillo	118
Merlan argenté	78
Merlan bleu	80
MERLUCCIIDAE	19, 88, 89
<i>merluccius</i> , <i>Merluccius</i>	88, 89
<i>Merluccius merluccius</i>	88, 89
Merlu européen	88
Merluza europea	88
<i>Microchirus variegatus</i>	218, 219
<i>Micromesistius poutassou</i>	80, 81
<i>microps</i> , <i>Pomatoschistus</i>	184, 185
<i>microstoma</i> , <i>Microstoma</i>	36, 37
<i>Microstoma microstoma</i>	36, 37
MICROSTOMATIDAE	18, 36–39
<i>mixtus</i> , <i>Labrus</i>	156, 157
MORONIDAE	19, 112, 113
Motelle	84
Motelle de Méditerranée	86
<i>muelleri</i> , <i>Maurolucus</i>	44, 45
<i>Mugil cephalus</i>	148, 149
MUGILIDAE	20, 148, 149
Mulet à grosse tête	148
MULLIDAE	20, 142–145
<i>Mullus barbatus</i>	142, 143, 145
<i>Mullus surmuletus</i>	144, 145
MYCTOPHIDAE	12, 18, 58–77
<i>Myctophum punctatum</i>	72, 73, 76

**N**

<i>Nansenia oblita</i>	38, 39
<i>nigrescens</i> , <i>Symphurus</i>	222, 223
<i>Notoscopelus elongatus</i>	74, 75

**O**

<i>oblita</i> , <i>Nansenia</i>	38, 39
<i>ocellaris</i> , <i>Blennius</i>	168, 169
Ochavo	202
OPHICHTHIDAE	18, 24, 25

OPHIDIIDAE . . . . . 19, 90, 91  
*Ophidion barbatum* . . . . . 90, 91  
 Orphie . . . . . 98

**P**

*paganellus*, *Gobius* . . . . . 180, 181  
*Pagellus bogaraveo* . . . . . 134, 135  
 Pagre rouge . . . . . 136  
*pagrus*, *Pagrus* . . . . . 136, 137  
*Pagrus pagrus* . . . . . 136, 137  
*Parablennius gattorugine* . . . . . 174, 175  
*Parablennius pilicornis* . . . . . 172, 173  
 PARALEPIDIDAE . . . . . 18, 52–55  
*Paralepis coregonoides* . . . . . 54, 55  
 Pardete . . . . . 148  
 Pargo . . . . . 136  
*Parophidion vassali* . . . . . 90  
 Pelaya misereres . . . . . 208  
 Petite argentine . . . . . 34  
 Pez cinto . . . . . 186  
 Pez de San Pedro . . . . . 100  
 Pez espada . . . . . 200  
 Pez lagarto . . . . . 50  
 Pez plata . . . . . 34  
*pholis*, *Lipophrys* . . . . . 170, 171  
 PHOSICHTHYIDAE . . . . . 18, 46, 47  
 Picarel . . . . . 140  
 Pigmy lanternfish . . . . . 68  
*pilchardus*, *Sardina* . . . . . 28, 29  
*pilicornis*, *Parablennius* . . . . . 172, 173  
*piscatorius*, *Lophius* . . . . . 94, 95  
 Plagusie sombre . . . . . 222  
*Platichthys flesus* . . . . . 214, 215  
 Platija europea . . . . . 214  
 PLEURONECTIDAE . . . . . 12, 21, 214, 215  
*podas*, *Bothus* . . . . . 212, 213  
 POMACENTRIDAE . . . . . 20, 150, 151  
*Pomatoschistus microps* . . . . . 184, 185  
*porcus*, *Scorpaena* . . . . . 106, 107  
 Porredana . . . . . 158  
*poutassou*, *Micromesistius* . . . . . 80, 81  
 Pouting(=Bib) . . . . . 82  
*punctatum*, *Myctophum* . . . . . 72, 73, 76  
*pusillus*, *Lampanyctus* . . . . . 68, 69

**R**

Rafino corto . . . . . 62  
 Rainbow wrasse . . . . . 152  
 Rape . . . . . 94  
 Rascacio . . . . . 106  
 Rascasse brune . . . . . 106  
 Rata . . . . . 166  
 Red mullet . . . . . 142  
 Red porgy . . . . . 136  
*regius*, *Zeugopterus* . . . . . 208, 209  
 Rémol . . . . . 206  
*reticulatus*, *Callionymus* . . . . . 177  
*rhombus*, *Scophthalmus* . . . . . 206, 207  
 Ringneck blenny . . . . . 172  
*rochei*, *Auxis* . . . . . 188, 189

Rock goby . . . . . 180  
 Rouget de roche . . . . . 144  
 Rouget de vase . . . . . 142  
 Round sardinella . . . . . 30  
 Rouquié . . . . . 154  
*rupestris*, *Ctenolabrus* . . . . . 154, 155

**S**

Sabre argenté . . . . . 186  
 Saint Pierre . . . . . 100  
 Salmonete de fango . . . . . 142  
 Salmonete de roca . . . . . 144  
 Salmonete real . . . . . 120, 146  
 Sanglier . . . . . 202  
*sarda*, *Sarda* . . . . . 190, 191  
*Sarda sarda* . . . . . 190, 191  
*Sardina europea* . . . . . 28  
*Sardina pilchardus* . . . . . 28, 29  
 Sardine commune . . . . . 28  
*Sardinella aurita* . . . . . 30, 31  
*sargus*, *Diplodus* . . . . . 132, 133  
*saurus*, *Synodus* . . . . . 50, 51  
*scaber*, *Uranoscopus* . . . . . 166, 167  
*scolopax*, *Macroramphosus* . . . . . 102, 103  
*Scomber colias* . . . . . 192, 193  
*Scomber scombrus* . . . . . 192, 194, 195  
 SCOMBRIDAE . . . . . 21, 188–199  
*scombrus*, *Scomber* . . . . . 192, 194, 195  
 SCOPHTHALMIDAE . . . . . 21, 204–209  
*Scophthalmus rhombus* . . . . . 206, 207  
*Scorpaena porcus* . . . . . 106, 107  
 SCORPAENIDAE . . . . . 12, 19, 106, 107  
 Sébaste chèvre . . . . . 104  
 SEBASTIDAE . . . . . 19, 104, 105  
 Serran-chèvre . . . . . 116  
 Serrandell . . . . . 210  
 SERRANIDAE . . . . . 19, 114–119  
 Serran tambour . . . . . 118  
*Serranus cabrilla* . . . . . 116, 117  
*Serranus hepatus* . . . . . 118, 119  
 Shanny . . . . . 170  
 Shore rockling . . . . . 86  
 Silver scabbardfish . . . . . 186  
 Silvery lightfish . . . . . 44, 46  
 Silvery pout . . . . . 78  
 Slender argentine . . . . . 36  
 Small lantern fish . . . . . 62  
*smaris*, *Spicara* . . . . . 140, 141  
*solea*, *Solea* . . . . . 220, 221  
*Solea solea* . . . . . 220, 221  
 Sole commune . . . . . 220  
 SOLEIDAE . . . . . 12, 21, 216–221  
 Solenette . . . . . 216  
 SPARIDAE . . . . . 20, 130–139  
*sphyraena*, *Argentina* . . . . . 34, 35  
*sphyrenoides*, *Lestidiops* . . . . . 52, 53  
*Spicara smaris* . . . . . 140, 141  
*Spondyliosoma cantharus* . . . . . 138, 139  
 Spotted lanternfish . . . . . 72  
 Stargazer . . . . . 166

- STERNOPTYCHIDAE . . . . . 18, 42–45  
*Stomias boa* . . . . . 48, 49  
 STOMIIDAE . . . . . 12, 18, 48, 49  
*surmuletus*, *Mullus* . . . . . 144, 145  
 Surmullet . . . . . 144  
 Swallowtail seaperch . . . . . 114  
 Swordfish . . . . . 200  
*Symbolophorus veranyi* . . . . . 76, 77  
*Symphodus melops* . . . . . 158, 159  
*Symphurus nigrescens* . . . . . 222, 223  
 SYNODONTIDAE . . . . . 18, 50, 51  
*Synodus saurus* . . . . . 50, 51
- T**
- Tabernero . . . . . 154  
 Tacaud commun . . . . . 82  
 Thickback sole . . . . . 218  
 Thon rouge de l'Atlantique . . . . . 198  
*Thunnus alalunga* . . . . . 196, 197  
*Thunnus thynnus* . . . . . 13, 198, 199  
*thynnus*, *Thunnus* . . . . . 13, 198, 199  
 Tompot blenny . . . . . 174  
 Tonguesole . . . . . 222  
 Torillo . . . . . 168  
 TRACHINIDAE . . . . . 12, 20, 162–165  
*Trachinus draco* . . . . . 164, 165  
*Trachurus* . . . . . 124  
*Trachurus mediterraneus* . . . . . 122–124  
*trachurus*, *Trachurus* . . . . . 122, 124, 125  
*Trachurus trachurus* . . . . . 122, 124, 125  
 Tres colas . . . . . 114  
 TRICHIURIDAE . . . . . 21, 186, 187
- TRIGLIDAE . . . . . 12, 19, 108–111  
*Trisopterus luscus* . . . . . 82, 83  
 Trompetero . . . . . 102  
 Two-spotted clingfish . . . . . 96
- U**
- Uranoscope . . . . . 166  
 URANOSCOPIDAE . . . . . 20, 166, 167  
*Uranoscopus scaber* . . . . . 166, 167
- V**
- variegatus*, *Microchirus* . . . . . 218, 219  
*vassali*, *Parophidion* . . . . . 90  
*veranyi*, *Symbolophorus* . . . . . 76, 77  
 Vieille coquette . . . . . 156  
*Vinciguerria attenuata* . . . . . 46, 47  
*vipera*, *Echiichthys* . . . . . 162, 163
- W**
- Wide-eyed flounder . . . . . 212
- X**
- Xiphias gladius* . . . . . 200, 201  
 XIPHIIDAE . . . . . 21, 200, 201
- Z**
- ZEIDAE . . . . . 19, 100, 101  
*Zeugopterus regius* . . . . . 208, 209  
*Zeus faber* . . . . . 100, 101



This guide presents the egg and larval descriptions of 100 species of fishes belonging to 55 families, which are most likely to be present in plankton samples collected in the continental shelf and oceanic waters of the Western Mediterranean Sea. The guide is structured in two parts. The first introductory part describes the different applications of ichthyoplankton studies in fisheries research, the main sampling strategies, methods and gears, the early life history of fishes and how to identify them. A brief historical account of ichthyoplankton identification studies in the Mediterranean Sea is also provided. The second part of the guide contains the identification sheets of the species. Each species sheet includes the following information: illustration of the adult fish and information on its habitat and spawning season; description of the main features useful towards identifying the egg, yolk sac and larval stage of each fish species; illustrations and (when available) photos of the different life stages.

ISBN 978-92-5-109895-0



9 7 8 9 2 5 1 0 9 8 9 5 0

I7708EN/1/08.17