

A trans-Atlantic
assessment and
deep-water
ecosystem based
spatial management
plan for Europe

ATLAS 2ND General Assembly

Hotel Blau Colonia Sant Jordi Mallorca, Spain 24 – 28 April 2017

Conference Programme and Abstracts



This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 678760 (ATLAS). This output reflects only the author's views, and the European Union cannot be held responsible for any use that may be made of the information contained therein.

Biodiversity and benthic megafaunal communities inhabiting the Formigas Bank (NE Azores)

Marina Carreiro-Silva¹, Lea-Anne Henry², Meri Bilan¹, Iris Sampaio^{1,3}, J Rivera⁴, Maria Rakka¹, Gerald H. Taranto¹, M Ramos⁵, José L. Rueda⁶, A Mateo-Ramirez⁷, Javier Urra^{6,7}, Marina Gallardo⁶, Sophie Arnaud-Haond⁸, Juancho Movilla ^{9,10}, Miriam Hermida¹¹, Telmo Morato¹, F Tempera¹² and Cova Orejas⁹

The Formigas Bank is an offshore seamount located in the easternmost part of the Azores archipelago (northeast Atlantic). It rises from abyssal depths to the surface, including a small set of islets. The bank holds multiple nature conservation designations, including a Natura 2000 Special Area of Conservation, an OSPAR Marine Protected Area, a RAMSAR site and a Nature Reserve declared under the Azores network of protected areas. The protection is based on the presence of sublittoral biotopes of high conservation interest, and importance as feeding grounds, spawning and nursery areas for many marine species, including fish, cetaceans and turtles. Although some information exists on the sublittoral communities occurring on the seamount summit (e.g., infralittoral Cystoseira and Laminaria beds, circalittoral hydrarian and sponge gardens, rich pelagic fauna), virtually no information was available on the deep-sea communities inhabiting the seamount flanks. Therefore, during the MEDWAVES cruise, the flanks of the Formigas bank have been surveyed using multibeam sonar, an ROV and oceanographic profiles, with the objective to characterise deep-sea biodiversity and megafaunal communities as well as the environment where they occur. This communication will present results from the video annotations of the ten dives made on the seamount slopes between ~500m and ~1,500 m depth. Diverse communities of sedentary suspension-feeding organisms were observed, with more than 20 cold-water coral species (mainly octocorals) being recorded, as well as many different sponge morphotypes. Dense coral garden habitats and sponge grounds were identified on several occasions, confirming the presence of vulnerable marine

¹IMAR – Institute of Marine Research and MARE – Marine and Environmental Sciences Centre, Universidade dos Acores

² School of Geosciences, University of Edinburgh

³ Senckenberg am Meer, Abteilung Meeresforschung

⁴Instituto Español de Oceanografía, Madrid

⁵ Marine and Environmental Sciences Centre (MARE), MARE-Azores, Faculdade de Ciências da Universidade de Lisboa (FCUL)

⁶ Instituto Español de Oceanografía, Málaga

⁷ Universidad de Málaga

⁸ Ifremer, Station de Sète

⁹ Instituto Español de Oceanografía, Palma de Mallorca

¹⁰ Instituto de Ciencias del Mar (ICM-CSIC), Passeig Maritim de la Barceloneta

¹¹ University of Alcalá

¹² Ifremer, Issy-les-Moulineaux