

Monroig O, Guinot D, Hontoria F, Tocher DR & Navarro JC
(2012) Biosynthesis of essential fatty acids in *Octopus vulgaris*
(Cuvier, 1797): Molecular cloning, functional characterisation
and tissue distribution of a fatty acyl elongase, *Aquaculture*,
360-361, pp. 45-53.

This is the peer reviewed version of this article

NOTICE: this is the author's version of a work that was accepted for publication in Aquaculture. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in Aquaculture, [VOL 360-361 (2012)] DOI: <http://dx.doi.org/10.1016/j.aquaculture.2012.07.016>

1 **Title**

2 Biosynthesis of essential fatty acids in *Octopus vulgaris* (Cuvier, 1797): Molecular
3 cloning, functional characterisation and tissue distribution of a fatty acid elongase

4

5 **Authors**

6 Óscar Monroig^{1*}, Diana Guinot¹, Francisco Hontoria¹, Douglas R Tocher², Juan C
7 Navarro¹

8

9 **Addresses**

10 ¹ Instituto de Acuicultura Torre de la Sal (IATS-CSIC), 12595 Ribera de Cabanes,
11 Castellón, Spain

12 ² Institute of Aquaculture, School of Natural Sciences, University of Stirling, Stirling
13 FK9 4LA, Scotland, UK

14

15

16 ***Corresponding author**

17 Óscar Monroig

18 Instituto de Acuicultura Torre de la Sal (IATS-CSIC), 12595 Ribera de Cabanes,
19 Castellón, Spain.

20 Tel: +34 964319500; Fax: +34 964319500; E-mail: oscar@iats.csic.es

21

22 **Keywords**

23 Elongases; essential fatty acids; non-methylene interrupted fatty acids; *Octopus*
24 *vulgaris*; polyunsaturated fatty acids.

25

25 **Summary**

26 Polyunsaturated fatty acids (PUFA) have been identified as key nutrients for the
27 common octopus (*Octopus vulgaris*), particularly for its early life-cycle stages
28 (paralarvae). Our overarching aim is to establish the essential fatty acid (FA)
29 requirements for octopus paralarvae through determination of the enzymes of
30 endogenous PUFA biosynthetic pathways. We here report on the molecular cloning and
31 functional characterisation of a cDNA encoding a putative elongase of very long-chain
32 fatty acids (Elovl), a critical enzyme that mediate the elongation of FA including PUFA.
33 Our results suggested that the octopus Elovl is phylogenetically related to Elovl5 and
34 Elovl2, two elongases with demonstrated roles in PUFA biosynthesis in vertebrates.
35 Further evidence supporting a role of the octopus Elovl in PUFA biosynthesis was
36 provided through functional characterisation of its activity in yeast. It was confirmed
37 that expression of the octopus Elovl conferred on yeast the ability to elongate some C18
38 and C20 PUFA, while C22 PUFA substrates remained unmodified. The substrate
39 specificities exhibited by the octopus elongase were consistent with those of vertebrate
40 Elovl5. Interestingly, the octopus Elovl elongated n-6 PUFA substrates more efficiently
41 than their analogous n-3 substrates, suggesting that n-6 PUFA may have particular
42 biological significance in *O. vulgaris*. Finally, we investigated the potential role of the
43 newly cloned Elovl in the biosynthesis of non-methylene-interrupted FA, compounds
44 typically found in marine invertebrates and confirmed to be also present in the common
45 octopus.

46

46 **Introduction**

47 Cephalopods have emerged as prime candidates for diversifying aquaculture. Among
48 the species studied, the common octopus (*Octopus vulgaris*, Cuvier, 1797) has received
49 special attention and relevant aspects for its culture such as husbandry (Iglesias et al.,
50 2006; Estefanell et al., 2012), behaviour (Di Cristo et al., 2005; Valverde and García,
51 2005), reproduction (Otero et al., 2007; Wodinsky, 2008; Estefanell et al., 2010),
52 pathologies (Castellanos-Martínez and Gestal, 2011) and nutrition (Villanueva, 1994,
53 Navarro and Villanueva, 2000, 2003; Villanueva et al., 2004, 2009; Quintana, 2006;
54 Villanueva and Bustamante, 2006; Seixas et al., 2010; Estefanell et al., 2011; Fuentes et
55 al., 2011; Viciano et al., 2011) have been studied. Despite considerable effort, the
56 production of the common octopus in captivity is limited to on-growing wild-captured
57 specimens in floating cages (Iglesias et al., 2007), as the octopus life cycle has not yet
58 been closed. While limited success in the production of juvenile octopuses has been
59 achieved (Villanueva, 1995; Iglesias et al., 2002, 2004; Lenzi et al., 2009), the massive
60 mortalities occurring during early life-cycle stages (paralarvae) have become an, as yet,
61 unresolved zootechnical issue that requires further investigation.

62 Polyunsaturated fatty acids (PUFA) have been previously suggested to be critical
63 dietary components for octopus paralarvae (Navarro and Villanueva, 2003). We have
64 recently initiated a series of studies to establish the essential fatty acid (FA)
65 requirements for octopus paralarvae, so that balanced diets matching their endogenous
66 biosynthetic capability can be formulated and thus promote paralarval survival and
67 development. Due to the obvious difficulties in conducting feeding trials with octopus
68 paralarvae, our approach aims to characterise the gene products encoding enzymes
69 involved in the PUFA biosynthetic pathway, which themselves dictate the ability of

70 species to endogenously produce long-chain PUFA (LC-PUFA) (Bell and Tocher,
71 2009).

72 Previously, we reported the molecular cloning and functional characterisation of a
73 fatty acyl desaturase (Fad) from *O. vulgaris* (Monroig et al., 2012a). The substrate
74 specificity of the octopus Fad revealed that this enzyme was a $\Delta 5$ -like Fad and thus we
75 provided for first time molecular evidence of such an enzymatic activity in molluscs
76 (Monroig et al., 2012a). Interestingly, the $\Delta 5$ Fad enables the common octopus to
77 endogenously convert 20:4n-3 and 20:3n-6 into eicosapentaenoic acid (EPA, 20:5n-3)
78 and arachidonic acid (ARA, 20:4n-6), respectively, the latter regarded as critical LC-
79 PUFA in a variety of physiological processes ensuring normal cellular function (Funk,
80 2001). Rather than a role in the biosynthesis of EPA, we hypothesised that $\Delta 5$ Fad
81 activity may actually be retained in the octopus for the endogenous biosynthesis of
82 ARA, as high contents of ARA encountered in adult octopus tissues were unlikely to be
83 exclusively from dietary origin. In addition to the potential participation of the octopus
84 $\Delta 5$ Fad in ARA biosynthesis, the common octopus $\Delta 5$ Fad might also have a role in the
85 biosynthesis of non-methylene interrupted (NMI) FA, compounds with unusual
86 unsaturation features that have been found in a variety of marine invertebrates
87 (Barnathan, 2009; Kornprobst and Barnathan, 2010).

88 The biosynthesis of PUFA including NMI FA in marine molluscs has been
89 investigated previously (De Moreno et al., 1976; Waldock and Holland, 1984; Zhukova,
90 1986, 1991, 2007). Although the biosynthetic ability of molluscs was species-specific
91 (Waldock and Holland, 1984), it has been shown that some molluscs have active PUFA
92 biosynthetic pathways and, in addition to the above mentioned $\Delta 5$ desaturase, active FA
93 elongation systems also appear to be present. Using radioactive FA, it was reported that
94 the clam *Mesoderma mactroides* could elongate both 18:3n-3 and 18:2n-6 (De Moreno

95 et al., 1976). Later, Waldock and Holland (1984) demonstrated that the Pacific oyster
96 *Crassostrea gigas* had the ability to produce 20:5n-3 and 22:6n-3. Investigations in
97 other bivalves like *Scapharca broughtoni*, *Callista brevisiphonata* and *Mytilus edulis*
98 (Zhukova, 1986, 1991) demonstrated that the biosynthesis of the NMI dienes $\Delta 7,13$
99 22:2 and $\Delta 7,15$ 22:2 was achieved by elongation from $\Delta 5,11$ 20:2 and $\Delta 5,13$ 20:2,
100 respectively. In addition to biochemical assays with radiotracers, indirect evidence of
101 FA elongase activity in molluscs was provided analytically (Joseph, 1982). For
102 instance, the unusual NMI FA $\Delta 5,9,15$ 24:3 and $\Delta 5,9,17$ 24:3 found in the limpets
103 *Cellana grata* and *Collisella dorsuosa* were suggested to derive from the typical NMI
104 dienes $\Delta 7,13$ 22:2 and $\Delta 7,15$ 22:2, respectively, by chain elongation and subsequent $\Delta 5$
105 desaturation (Kawashima, 2005).

106 The elongases of very long-chain fatty acids (Elovl), a protein family with seven
107 distinct members (Elovl 1-7) in vertebrates, account for the condensation of 2C into
108 activated preexisting fatty acyl chains (Jakobsson et al., 2006). Investigation of FA
109 biosynthetic pathways has allowed the molecular and functional characterisation of a
110 number of genes encoding Elovl enzymes from vertebrates (see reviews by Jakobsson et
111 al., 2006; Guillou et al., 2010; Monroig et al., 2011a). In contrast, studies of elongase-
112 encoding genes from non-vertebrate organisms are scarce, with only few examples such
113 as elongases from the nematode *Caenorhabditis elegans* (Beaudoin et al., 2000) and the
114 marine protist *Thraustochytrium* sp. (Heinz et al., 2001; Jiang et al., 2008), and no
115 elongases from molluscs have been reported.

116 In order to expand our knowledge of EFA requirements of common octopus, the
117 present study reports the molecular cloning, functional characterisation and tissue
118 distribution of transcripts (mRNA) of a cDNA encoding a putative elongase involved in
119 PUFA biosynthesis. In order to understand a potential role of the newly cloned elongase

120 in the NMI FA biosynthesis in the common octopus, we also analysed the double bond
121 features of NMI FA found in specific tissues of octopus adult specimens.

122 **Materials and methods**

123 *Tissue samples*

124 Tissue samples from common octopus were obtained from the dissection of two
125 (male and female) adult individuals (~1.5 kg) captured through artisanal fisheries along
126 the Mediterranean East coast of Spain. The octopuses were cold anaesthetised and
127 sacrificed by direct brain puncture and tissues including nerve, nephridium,
128 hepatopancreas, brain, caecum, gill, muscle, heart and gonad were sampled and
129 immediately frozen at -80 °C until further analysis.

130 *Elongase cDNA cloning*

131 Total RNA was extracted from octopus tissues using TRIzol® (Gibco BRL, Grand
132 Island, NY, USA) reagent following manufacturer's instructions. Subsequently, first
133 strand cDNA was synthesised using a Verso™ cDNA kit (ABgene, Rockford, IL, USA)
134 primed with random hexamers. In order to amplify the first fragment of the elongase
135 cDNA, the amino acid (aa) sequences of Elov15 proteins from *Homo sapiens*
136 (NP_068586.1), *Rattus norvegicus* (NP_599209.1), *Bos taurus* (NP_001040062.1),
137 *Danio rerio* (NP_956747.1) and *Pagrus major* (ADQ27303.1) were aligned using
138 BioEdit v5.0.6 (Tom Hall, Department of Microbiology, North Carolina State
139 University, USA). Conserved regions were used for *in silico* searches of mollusc
140 expressed sequence tags (EST) using NCBI tblastn tool (<http://www.ncbi.nlm.nih.gov/>).
141 Several EST displaying high homology with Elov1 encoding genes were identified from
142 the molluscs *Mytilus galloprovincialis* (gb|FL495089.1| and gb|FL499406.1|),
143 *Euprymna scolopes* (gb|DW256301.1|), and *Lymnaea stagnalis* (gb|FC701557.1|,
144 gb|FC773093.1|, gb|FC770692.1| and gb|FC696214.1|). Additionally, a search of the

145 owl limpet *Lottia gigantea* genome was performed using the zebrafish Elov15
146 (NP_956747.1) sequence with the tblastn tool at [http://genome.jgi-](http://genome.jgi-psf.org/Lotgi1/Lotgi1.home.html)
147 [psf.org/Lotgi1/Lotgi1.home.html](http://genome.jgi-psf.org/Lotgi1/Lotgi1.home.html). After processing, the mollusc Elov1-like sequences
148 were aligned (Bioedit) for the design of the primers UNIEloF (5'-
149 TTGTGGTGGTACTTCTC-3') and UNIEloR (5'-
150 GTAATATACTTTTCCACCA-3') that were used for polymerase chain reaction
151 (PCR) using GoTaq® Colorless Master Mix (Promega, Southampton, UK), and using a
152 mixture of cDNA from gonads, brain, nerve and caecum as template. The PCR
153 consisted of an initial denaturing step at 95 °C for 2 min, followed by 35 cycles of
154 denaturation at 95 °C for 30 s, annealing at 50 °C for 30 s, extension at 72 °C for 1 min,
155 followed by a final extension at 72 °C for 5 min. The PCR fragment was sequenced at
156 the DNA Sequencing Service of the IBMCP-UPV (Valencia, Spain) and gene-specific
157 primers were designed for 5' and 3' rapid amplification of cDNA ends (RACE) PCR
158 (FirstChoice® RLM-RACE kit, Ambion, Applied Biosystems, Warrington, UK) to
159 produce a full-length cDNA. Details of all primers used for RACE PCR are given in
160 Table 1.

161 For 5'RACE PCR, a positive fragment was obtained by two-round PCR. The first
162 round PCR was performed using the adapter-specific 5'RACE OUTER primer and the
163 gene-specific forward primer OVEloR1, with an initial denaturing step at 95 °C for 2
164 min, followed by 32 cycles of denaturation at 95 °C for 30 s, annealing at 58 °C for 30 s,
165 extension at 72 °C for 75 s, followed by a final extension at 72 °C for 5 min (GoTaq®
166 Colorless Master Mix, Promega). First round PCR products were used as template for
167 nested PCR with primers 5'RACE INNER and OVEloR2 in a 32-cycle reaction under
168 the same thermal conditions as above. For 3'RACE PCR, a similar two-round approach
169 was followed with first round PCR performed with primers OVEloF1 and 3'RACE

170 OUTER, with an initial denaturing step at 95 °C for 1 min, followed by 32 cycles of
171 denaturation at 95 °C for 30 s, annealing at 56 °C for 30 s, extension at 72 °C for 2 min,
172 followed by a final extension at 72 °C for 5 min (GoTaq[®] Colorless Master Mix,
173 Promega). First round PCR product was then used as template for nested PCR with
174 primers 5'RACE INNER and OVEloF2, with thermal conditions as above. RACE PCR
175 products were cloned into pGEM-T Easy Vector (Promega) and sequenced as above.

176 *Sequence and phylogenetic analyses*

177 Using ClustalW (Bioedit), the deduced aa sequence of the newly cloned *O. vulgaris*
178 elongase cDNA was aligned with those of a predicted elongase found in the gastropod
179 owl limpet (termed 'L. gigantea Elov1 transcript 1', jgi|Lotgi1|224291|), as well as those
180 of protein homologues including the human ELOVL5 (gb|NP_068586|) and ELOVL2
181 (gb|NP_060240|), and the zebrafish Elov15 (gb|NP_956747|) and Elov12
182 (gb|NP_001035452|). The aa sequence identity between Elov1-like proteins was
183 compared using the EMBOSS Needle Pairwise Sequence Alignment tool
184 (http://www.ebi.ac.uk/Tools/psa/emboss_needle/). Phylogenetic analysis of the aa
185 sequences deduced from the Elov1-like cDNA from common octopus and those from
186 other organisms including several marine invertebrates was performed by constructing a
187 tree using the Neighbour Joining method (Saitou and Nei 1987), with confidence in the
188 resulting tree branch topology measured by bootstrapping through 10000 iterations.

189 *Functional characterisation of the octopus elongase by heterologous expression in* 190 *Saccharomyces cerevisiae*

191 PCR fragments corresponding to the open reading frame (ORF) of the putative
192 elongase were amplified from a mixture of cDNA synthesised from gonads, brain, nerve
193 and caecum RNA extracts, and using the high fidelity *Pfu* DNA Polymerase (Promega).
194 PCR conditions consisted of an initial denaturing step at 95 °C for 2 min, followed by

195 35 cycles of denaturation at 95°C for 30 s, annealing at 57 °C for 30 s, extension at 72
196 °C for 2 min 15 s, followed by a final extension at 72 °C for 5 min. The primers
197 containing restriction sites (underlined in Table 1) OVEloVF (*Hind*III) and OVEloVR
198 (*Sac*I) were used for PCR, and the DNA fragments produced were subsequently
199 purified, digested with the corresponding restriction enzymes (Promega), and ligated
200 into a similarly restricted pYES2 yeast expression vector (Invitrogen, Paisley, UK). The
201 purified plasmids (GenElute™ Plasmid Miniprep Kit, Sigma) containing the octopus
202 elongase ORF were then used to transform *Saccharomyces cerevisiae* competent cells
203 (S.c. EasyComp Transformation Kit, Invitrogen). Transformation and selection of yeast
204 with recombinant pYES2-OVElo plasmids, and yeast culture were performed as
205 described in detail previously (Agaba et al., 2004).

206 In order to test the ability of octopus Elov1 cDNA ORF to elongate either saturated or
207 monounsaturated FA, yeast transformed with pYES2 vector containing the octopus
208 elongase as an insert (pYES2-OVElo) and no insert (control) were grown in *S.*
209 *cerevisiae* minimal medium^{-uracil} with no exogenously added FA substrates.
210 Additionally, the ability of *O. vulgaris* Fad to desaturate PUFA substrates was tested by
211 growing pYES2-OVElo transgenic yeast in medium supplemented with one of the
212 following substrates: 18:3n-3, 18:2n-6, 18:4n-3, 18:3n-6, 20:5n-3, 20:4n-6, 22:5n-3 and
213 22:4n-6. The FA were added to the yeast cultures at final concentrations of 0.5 (C18),
214 0.75 (C20) and 1.0 (C22) mM as uptake efficiency decreases with increasing chain
215 length (Zheng et al., 2009). Yeast transformed with empty pYES2 were also grown in
216 presence of PUFA substrates as control treatments. After 2-day culture at 30 °C, yeast
217 were harvested, washed, and lipid extracted by homogenisation in chloroform/methanol
218 (2:1, v/v) containing 0.01% butylated hydroxy toluene (BHT) as antioxidant. All fatty
219 acid substrates, except stearidonic acid (18:4n-3), were purchased from Nu-Chek Prep,

220 Inc (Elysian, MN, USA). Stearidonic acid and chemicals used to prepare the *S.*
221 *cerevisiae* minimal medium-^{uracil} were from Sigma Chemical Co. Ltd. (Dorset, UK),
222 except for the bacteriological agar obtained from Oxoid Ltd. (Hants, UK).

223 *Tissue distribution of elongase transcripts*

224 Expression of the octopus elongase was examined in adult tissues by RT-PCR. Total
225 RNA from a series of tissues including nerve, nephridium, hepatopancreas, brain,
226 caecum, gill, muscle, heart, and female and male gonads was extracted as described
227 above, and 1 µg of total RNA was reverse transcribed into cDNA (M-MLV reverse
228 transcriptase, Promega). In order to determine the mRNA distribution of the octopus
229 elongase, the tissue cDNAs were used as templates in PCR consisting of a denaturing
230 step at 95 °C for 1 min, followed by 35 cycles of denaturation at 95 °C for 30 s,
231 annealing at 58 °C for 30 s, extension at 72 °C for 30 s, followed by a final extension at
232 72 °C for 5 min (GoTaq[®] Green Master Mix, Promega). Additionally, the expression of
233 the housekeeping β-actin was determined to check the cDNA integrity. Primers used for
234 RT-PCR are shown in Table 1.

235 *Fatty acid analysis by GC-MS*

236 FA from the transgenic yeast were analysed by preparing methyl esters (FAME) as
237 previously described (Hastings et al. 2001). Briefly, FAME were identified and
238 quantified using an Agilent 6850 Gas Chromatograph system coupled to a 5975 series
239 MSD (Agilent Technologies, Santa Clara, CA, USA). The elongation efficiency from
240 potential substrates including the yeast endogenous FA and the exogenously added
241 PUFA substrates (18:3n-3, 18:2n-6, 18:4n-3, 18:3n-6, 20:5n-3, 20:4n-6, 22:5n-3 and
242 22:4n-6) were calculated by the proportion of substrate FA converted to elongated FA
243 product as [product area/(product area + substrate area)] x 100. When further
244 confirmation of double bond positions was required, picolinyl esters were prepared

245 from FAME according to the methodology described by Destailats and Angers (2002)
246 and modified according to Li et al. (2010).

247 In order to investigate the potential participation of the octopus elongase in the
248 biosynthesis of NMI FA, the FA compositions of specific tissues in which we had
249 previously detected NMI FA (Monroig et al., 2012a) were determined through
250 preparation of both methyl and picolinyl ester derivatives from polar lipid (PL) fractions
251 prepared as follows. Lipid extracts (2 mg) from nephrydium, male gonad, eye and
252 caecum were applied to 20x20 silica gel plates (Merck, Darmstadt, Germany) and
253 eluted with a solvent mixture of n-hexane / diethyl ether / glacial acetic acid (85:15:1.5,
254 v/v/v). PL fractions, identified by comparison with known standards, were scraped from
255 the plates and FAME prepared (Monroig et al., 2012a) and analysed as described above.
256 FAME samples were subsequently derivatised to FA picolinyl prepared for
257 identification of the double bond patterns in NMI FA.

258

259 **Results**

260 *Octopus elongase sequence and phylogenetics*

261 The ORF of the newly cloned Elov1 from *O. vulgaris* consisted of 885 bp encoding a
262 putative protein of 294 aa. Its sequence was deposited in the GenBank database with
263 accession number JX020803. The deduced aa sequence from the octopus elongase
264 showed identity scores ranging from 39.3 and 43.2 % with several Elov1 proteins
265 (Elov12, Elov14 and Elov15) from vertebrates including *H. sapiens*, *X. tropicalis* and *D.*
266 *rerio*. When compared with the two full-length elongases found in the genome of the
267 gastropod *L. gigantea*, the octopus Elov1 was 58.1 % identical to the so-called ‘*L.*
268 *gigantea* Elov1 transcript 1’ and 39.5 % identical to the ‘Elov1 transcript 2’. When the
269 octopus Elov1 aa sequence was compared with incomplete elongase sequences from *E.*

270 *scolopes*, *L. stagnalis*, *M. galloprovincialis* and *A. californica* the identity scores were
271 relatively low, ranging from 31.9 to 43.9 %.

272 Similar to vertebrate Elovl-like proteins, the deduced aa sequence of the octopus
273 elongase contained the diagnostic histidine box (HXXHH) conserved in all members of
274 the Elovl protein family (Fig. 1). It also possessed two lysine (K) residues at the
275 carboxyl terminus (KKXX), regarded as putative ER retrieval signals. Additionally, five
276 putative transmembrane-spanning regions containing hydrophobic aa stretches were
277 predicted in residues 32-50, 65-83, 117-137, 158-192 and 239-259 by InterProScan
278 (version 4.2) (Fig. 1).

279 A phylogenetic tree was constructed on the basis of aa sequence comparisons of the
280 octopus Elovl and other predicted elongases from molluscs, as well as several Elovl
281 types (Elovl 1-7) from a variety of vertebrates (Fig. 2). Our results showed that the
282 octopus Elovl protein clustered with other Elovl-like from molluscs including the
283 cephalopod *E. scolopes* and the gastropods *L. stagnalis* and *L. gigantea* ('transcript 1'),
284 altogether forming a group close to Elovl2 and Elovl5 proteins from vertebrates. More
285 distantly, three main clusters could be distinguished including Elovl3/Elovl6,
286 Elovl1/Elovl7 and Elovl4 representatives. Interestingly, the Elovl4 cluster included the
287 well-studied proteins from vertebrates, but also other mollusc Elovl-like proteins from
288 *L. gigantea* ('transcript 2'), *A. californica* and *M. galloprovincialis*.

289 *Functional characterisation in yeast*

290 The octopus Elovl-like encoding cDNA was functionally characterised by expressing
291 the ORF in yeast *S. cerevisiae*. The FA composition of wild yeast consists basically of
292 the main endogenous FA of *S. cerevisiae*, namely 16:0, 16:1 isomers (16:1n-9 and
293 16:1n-7), 18:0, 18:1n-9 and 18:1n-7 (Monroig et al., 2010a). Total lipid of yeast
294 transformed with the empty pYES2 vector (control) contained these FA together with

295 whichever exogenous FA (if any) was added as substrate (data not shown), indicating
296 that no elongase activity towards any of the exogenously added PUFA substrates
297 assayed. This result is in agreement with the well-know inability of *S. cerevisiae*
298 elongases to operate towards PUFA substrates (Hastings et al., 2001; Agaba et al.,
299 2004).

300 In order to test the ability of the octopus Elovl to elongate saturated and
301 monounsaturated FA, yeast transformed with pYES2-OVElovl were grown in absence
302 of exogenously added substrates. Our results showed that none of the yeast endogenous
303 FA, whether saturated or monounsaturated, were elongated. Conversely, yeast
304 transformed with pYES2-OVElovl showed activity towards PUFA substrates producing
305 the corresponding 2-carbon elongation product. As shown in Fig. 3, the exogenously
306 added C18 (18:3n-3, 18:2n-6, 18:4n-3 and 18:3n-6) and C20 (20:5n-3 and 20:4n-6)
307 substrates were elongated to C20 (20:3n-3, 20:2n-6, 20:4n-3 and 20:3n-6) and C22
308 (22:5n-3 and 22:4n-6) products, respectively. Conversion rates derived from the yeast
309 assays suggested that the octopus Elovl generally elongated n-6 PUFA substrates more
310 efficiently than n-3 substrates for each pair of homologous substrates considered. Thus,
311 the substrates 18:2n-6, 18:3n-6, 20:4n-6 were consistently elongated at higher rates than
312 the corresponding n-3 PUFA substrates 18:3n-3, 18:4n-3 and 20:5n-3, respectively.
313 Interestingly, no activity towards C22 (22:5n-3 and 22:4n-6) PUFA substrates was
314 detected.

315 *Tissue distribution of octopus elongase transcripts*

316 Tissue expression of the common octopus Elovl was studied by RT-PCR on cDNA
317 samples obtained from a range of tissues (Fig. 4). Except for nephridium, transcripts of
318 the octopus Elovl gene were detected in all tissues analysed. Although RT-PCR

319 analyses should not be regarded as strictly quantitative data, our results indicate that
320 both the male and female gonads showed higher expression signals.

321 *Fatty acid composition from polar lipids of adult octopus tissues*

322 FA from PL were analysed in several tissues of adult octopus individuals (Table 3).
323 DHA appeared the most abundant FA for each tissue considered, with up to 27.0 % of
324 total FA in eye PL. Other PUFA relatively abundant in the tissues studied were ARA
325 (with up to 16.4 % in male gonad PL) and EPA (up to 13.7 % in caecum PL).
326 Interestingly, 20:3n-3 content in eye was 13.5 % of total FA in the PL fraction.

327 GC-MS analysis of picolinyl esters enabled us to identify four different NMI FA in
328 the octopus tissues, namely $\Delta 5,11$ 20:2, $\Delta 7,13$ 20:2, $\Delta 5,11,14$ 20:3 and $\Delta 7,13$ 22:2
329 (Table 3). Although we specifically analysed the PL fractions, where NMI FA are
330 believed to accumulate (Klingensmith, 1982; Pirini et al., 2007), the amounts of all the
331 NMI FA identified were generally low, and only relatively higher contents were
332 detected for $\Delta 5,11$ 20:2 in nephrydium (1.8 %) and its corresponding elongation product
333 $\Delta 7,13$ 22:2 in male gonad (2.2 %).

334 **Discussion**

335 The FA biosynthesis pathways have been investigated in both terrestrial (van der
336 Horst 1973, 1974; Weinert et al. 1993; Zhu et al. 1994) and aquatic mollusc species
337 (Chu and Greaves 1991; de Moreno et al. 1976; Waldock and Holland 1984; Zhukova
338 1986, 1991, 2007; Delaporte et al., 2005). It was shown that some molluscs have active
339 FA elongation systems (Waldock and Holland, 1984; Zhukova, 1986; Delaporte et al.,
340 2005). In the present study we provide compelling evidence of the existence of an Elovl
341 cDNA that encodes an enzyme potentially involved in the biosynthesis of PUFA in the
342 cephalopod *O. vulgaris*.

343 The deduced aa sequence of the Elovl-like cDNA from *O. vulgaris* contains all the
344 features of the vertebrate Elovl protein family members, including five membrane-
345 spanning regions, an ER retrieval signal at the C terminus containing lysine residues
346 (KKXX) and a diagnostic histidine box (HXXHH) (Leonard et al., 2004; Jakobsson et
347 al., 2006). Moreover, the histidine (H) box and its N-terminal side (QVTFLHVFHH)
348 show a typical aa pattern of the PUFA elongase subfamily of eukaryotic elongases, with
349 a glutamine (Q) at position -5 and a leucine (L) at position -1 from the first H
350 (Hashimoto et al., 2008). Further evidence supporting a potential role of this octopus
351 Elovl cDNA in the PUFA biosynthetic pathways was provided by phylogenetic
352 analysis. Thus, the octopus Elovl aa sequence, as well as those of other mollusc
353 elongases, obtained by *in silico* searches, including the cephalopod *Euprymna scolopes*
354 and the gastropods *Lymnaea stagnalis* and *Lottia gigantea* (transcript 1), showed great
355 similarity to the sequences of Elovl2 and Elovl5 proteins, critical enzymes participating
356 in the biosynthesis of LC-PUFA in vertebrates (Leonard et al., 2004; Jakobsson et al.,
357 2006). More distantly, the other elongase identified in the *L. gigantea* genome
358 (transcript 2) and also other Elovl-like proteins from *A. californica* and *M.*
359 *galloprovincialis* grouped together with vertebrate Elovl4 elongases, another type of
360 elongase involved in the biosynthesis of very long-chain FA (C>24) including both
361 saturates and polyenes (Agbaga et al., 2008; Monroig et al., 2010b, 2011b, 2012b).
362 While these results suggest that another elongase with similarity to Elovl4 might also be
363 present in the common octopus, the functional characterisation of the present Elovl
364 cDNA confirmed, not only its participation in the PUFA elongation pathway, but also
365 that it has substrate specificities more similar to Elovl5 than Elovl2.

366 Clearly, transgenic yeast expressing the octopus Elovl efficiently converted C18 and
367 C20 PUFA substrates to their corresponding 2-carbon elongated products, but no

368 activity towards C22 PUFA was detected. Generally, this pattern of substrate specificity
369 of the octopus elongase is consistent with that of vertebrate Elovl5 proteins (Jakobsson
370 et al., 2006). For instance, the human ELOVL5 (also termed HELO1) and the rat
371 ELOVL5 (also termed rELO1) were shown to efficiently elongate C18 and C20 PUFA,
372 whereas C22 PUFA did not appear to be substrates for these enzymes (Leonard et al.,
373 2000; Inagaki et al., 2002). Similarly, fish Elovl5 demonstrated high activity for the
374 elongation of C18 and C20 PUFA substrates, whereas C22 substrates were only
375 elongated to a lesser extent (Agaba et al., 2004; Morais et al., 2009; Mohd-Yusof et al.,
376 2010; Monroig et al., 2012b). Importantly, elongation of C22 PUFA including 22:5n-3
377 and 22:4n-6 in vertebrates is basically mediated by Elovl2, whose substrate chain-length
378 specificity also includes C20, but not C18, PUFA substrates, the latter being only
379 marginally or not elongated (Tvrdik et al., 2000; Leonard et al., 2002; Monroig et al.,
380 2009; Morais et al., 2009). Overall it can be concluded that the *O. vulgaris* elongase
381 cloned here is phenotypically an Elovl5-like elongase, but its sequence similarity to
382 vertebrate Elovl2 suggests an interesting evolutionary scenario that is worth exploring
383 in future investigations.

384 The functional characterisation of the octopus Elovl revealed, however, that the gene
385 product might have conserved/acquired a different PUFA family specificity compared
386 to vertebrate Elovl5 proteins during evolution. Unlike mammalian (Leonard et al., 2000;
387 Inagaki et al., 2002) and fish Elovl5 (Agaba et al., 2005; Mohd-Yusof et al., 2010;
388 Morais et al., 2011; Monroig et al., 2012b), which are generally more efficient in
389 elongating n-3 rather than n-6 FA substrates, the octopus Elovl exhibited higher
390 elongation rates towards n-6 compared to n-3 substrates for each homologous pair
391 considered. Thus 18:2n-6, 18:3n-6, 20:4n-6 were all elongated at higher rates than the
392 corresponding n-3 FA, namely 18:3n-3, 18:4n-3 and 20:5n-3, respectively. These results

393 emphasise that n-6 FA in general, and especially ARA (20:4n-6), might play
394 particularly important physiological roles in the common octopus. Consistent with this,
395 several studies have reported unexpectedly high levels of ARA in tissues of common
396 that were unlikely to derive purely from dietary origin and, thus, an active biosynthesis
397 of ARA in the common octopus was postulated (Milou et al., 2006; García-Garrido et
398 al., 2010; Monroig et al., 2012a). In the present study, the efficiency shown by the
399 octopus Elovl to elongate certain PUFA substrates indicates that this enzyme could
400 contribute to the endogenous biosynthesis of ARA in this species.

401 In vertebrates, ARA is biosynthesised from the dietary essential C18 PUFA 18:2n-6
402 through two alternative pathways, the ‘classical’ $\Delta 6$ -pathway ($\Delta 6$ desaturation \rightarrow
403 elongation $\rightarrow \Delta 5$ desaturation), or alternatively through the so-called ‘ $\Delta 8$ -pathway’
404 (elongation $\rightarrow \Delta 8$ desaturation $\rightarrow \Delta 5$ desaturation) (Monroig et al., 2011c). In addition
405 to the ability of the formerly characterised Fad cDNA to mediate the $\Delta 5$ -desaturation
406 steps of these pathways (Monroig et al., 2012a), we here demonstrate that the newly
407 cloned octopus Elovl can efficiently catalyse the elongation reactions required for ARA
408 biosynthesis from the dietary essential 18:2n-6, namely 18:3n-6 \rightarrow 20:3n-6 for the $\Delta 6$ -
409 pathway and 18:2n-6 \rightarrow 20:3n-6 for the $\Delta 8$ -pathway. Although genes responsible for
410 elongation and $\Delta 5$ desaturation steps of these pathways have now been identified in
411 octopus, no Fad cDNA with $\Delta 6$ or $\Delta 8$ -desaturase activity has yet been identified and,
412 consequently, it remains unclear whether the common octopus can biosynthesise ARA
413 from the dietary essential 18:2n-6. This appears to be the case for some abalone species
414 (Dunstan et al., 1996; Durazo-Beltrán et al., 2003) but other species like *C. gigas*
415 (Waldock and Holland, 1984) and *Mytilus edulis* (Zhukova, 1991) appear unable to
416 biosynthesise ARA from 18:2n-6.

417 In addition to the biosynthesis of conventional PUFA, the octopus Elovl can also
418 have a role in the production of non-methylene-interrupted (NMI) FA. Thus, the
419 biosynthesis of $\Delta 7,13$ 22:2 encountered in male gonad, eye and caecum may be
420 accounted for by the elongation of $\Delta 5,11$ 20:2, as described for other marine
421 invertebrates (Kornprobst and Barnathan, 2010). Although we cannot directly conclude
422 that the octopus Elovl has the ability to elongate $\Delta 5,11$ 20:2 as this substrate was not
423 available, some of our results suggest a role for the elongase in the production of $\Delta 7,13$
424 22:2 from $\Delta 5,11$ 20:2. First, the increased expression signal of Elovl in the male gonad
425 is consistent with this tissue containing the highest amount of $\Delta 7,13$ 20:2. Second, it is
426 reasonable to assume that, similar to the elongation rates exhibited towards other C20
427 PUFA like 20:4n-3 and 20:3n-6, the octopus Elovl might also efficiently operate
428 towards another C20 PUFA like $\Delta 5,11$ 20:2. Whereas these circumstantial data suggest
429 that the octopus Elovl may contribute to the endogenous biosynthesis of NMI FA in this
430 cephalopod, the extent to which this biosynthetic pathway is operative in the common
431 octopus is difficult to predict. On one hand, the ability of the octopus $\Delta 5$ Fad to convert
432 20:3n-3 ($\Delta 11,14,17$ 20:3) and 20:2n-6 ($\Delta 11,14$ 20:2) to the NMI FA $\Delta 5,11,14,17$ 20:4
433 and $\Delta 5,11,14$ 20:3, respectively (Monroig et al., 2012a), supports the hypothesis of a
434 notable production of NMI FA by *O. vulgaris* itself. On the other, the endogenous
435 biosynthesis of NMI FA in the common octopus appears to be limited as, despite the
436 likely intake of preformed NMI FA through the diet, they still present relative low
437 levels compared to those found in some bivalves (Klingensmith, 1982) or nudibranchs
438 (Zhukova, 2007).

439 In summary, the present study demonstrates that the common octopus possesses an
440 Elovl-like cDNA with high homology to vertebrate Elovl5 and Elovl2 enzymes. The
441 functions of the octopus Elovl, while generally consistent those of vertebrate Elovl5,

442 have some novel particularities. Thus, the octopus Elovl showed higher elongation
443 efficiency towards n-6 than n-3 PUFA suggesting that these compounds, and especially
444 ARA, might play particularly pivotal physiological roles in the common octopus.
445 Moreover, the Elovl might be involved in the biosynthesis of NMI FA, although the
446 quantitative significance of this biosynthetic pathways in *O. vulgaris* requires further
447 investigation.

448 **Acknowledgements**

449 This research and OM were supported by a Marie Curie Reintegration Grant within the
450 7th European Community Framework Programme (PERG08-GA-2010-276916,
451 LONGFA), with additional support from “Ministerio de Ciencia e Innovación” through
452 the OCTOPHYS Project (AGL-2010-22120-C03-02) and a Juan de la Cierva
453 postdoctoral contract for OM, and the Generalitat Valenciana through a PROMETEO
454 Project (2010/006). The authors would also like to thank Mr. Miguel Ángel Montolio
455 for assistance in lipid analyses.

456

457 **References**

- 458 Agaba M, Tocher DR, Dickson C, Dick JR, Teale AJ (2004) Zebrafish cDNA encoding
459 multifunctional fatty acid elongase involved in production of eicosapentaenoic
460 (20:5n-3) and docosahexaenoic (22:6n-3) acids. *Mar Biotechnol* 6:251-261.
- 461 Agaba, M.K., Tocher, D.R., Dickson, C.A., Zheng, X., Dick, J.R., Teale, A.J., 2005.
462 Cloning and functional characterisation of polyunsaturated fatty acid elongases from
463 marine and freshwater teleost fish. *Comp. Biochem. Physiol.* 142B, 342–352.
- 464 Agbaga, M.P., Brush, R.S., Mandal, M.N.A., Henry, K., Elliott, M.H., Anderson, R.E.,
465 2008. Role of Stargardt-3 macular dystrophy protein (ELOVL4) in the biosynthesis
466 of very long chain fatty acids, *Proc. Natl. Acad. Sci. USA.* 105, 12843-12848.

467 Barnathan G. (2009) Non-methylene-interrupted fatty acids from marine invertebrates:
468 Occurrence, characterization and biological properties. *Biochimie* 91:671-678.

469 Beaudoin F, Michaelson LV, Lewis MJ, Shewry PR, Sayanova O, Napier JA (2000)
470 Production of C20 polyunsaturated fatty acids (PUFAs) by pathway engineering:
471 identification of a PUFA elongase component from *Caenorhabditis elegans*.
472 *Biochem Soc Trans* 28:661-663.

473 Bell MV, Tocher DR (2009) In: Arts MT, Brett M, Kainz M. (eds) *Lipids in Aquatic*
474 *Ecosystems*. Springer-Verlag, New York.

475 Castellanos-Martinez S., Gestal C. 2011. Immune response of *Octopus vulgaris* against
476 the infection by the gastrointestinal parasite *Aggregata octopiana*. *Journal of*
477 *Shellfish* 30: 997-998.

478 Chu FE, Greaves J. (1991) Metabolism of palmitic, linoleic, and linolenic acid in adult
479 oyster, *Crassostrea virginica*. *Mar Biol* 110:229–236.

480 Delaporte, M., Soudant, P., Moal, J., Kraffe, E., Marty, Y., Samain, J.F., 2005.
481 Incorporation and modification of dietary fatty acids in gill polar lipids by two
482 bivalves species *Crassostrea gigas* and *Ruditapes philippinarum*. *Comp. Biochem.*
483 *Physiol.* 140 A, 460–470.

484 De Moreno JEA, Moreno VJ, Brenner RR (1976) Lipid metabolism of the yellow clam,
485 *Mesodesma mactroides*: 2-polyunsaturated fatty acid metabolism. *Lipids* 11:561-
486 566.

487 Destailats F, Angers P (2002) One-step methodology for the synthesis of FA picolinyl
488 esters from intact lipids. *J Am Oil Chem Soc* 79:253–256.

489 Di Cristo C; Van Minnen J; Di Cosmo A., 2005. The presence of APGWamide in
490 *Octopus vulgaris*: a possible role in the reproductive behavior. *Peptides* 26, 53-62.

491 Dunstan GA, Baillie HJ, Barrett SM, Volkman JK (1996) Effect of diet on the lipid
492 composition of wild and cultured abalone. *Aquacult Abalone Cult* 140:115–127.

493 Durazo-Beltrán E, D’Abramo LR, Toro-Vazquez JF, Vasquez-Peláez C, Viana MT
494 (2003) Effect of triacylglycerols in formulated diets on growth and fatty acid
495 composition in tissue of green abalone (*Haliotis fulgens*). *Aquaculture* 224:257–
496 270.

497 Estefanell J, Socorro J, Roo FJ, Fernández-Palacios H, Izquierdo M (2010) Gonad
498 maturation in *Octopus vulgaris* during on-growing, under different conditions of sex
499 ratio. *ICES J Mar Sci* 67:1487–1493.

500 Estefanell, J., Socorro, J., Tuya, F., Izquierdo, M., Roo, J. 2011. Growth, protein
501 retention and biochemical composition in *Octopus vulgaris* fed on different diets
502 based on crustaceans and aquaculture by-products. *Aquaculture* 322, 91-98.

503 Estefanell J.; Roo J.; Fernandez-Palacios H.; Izquierdo, M., Socorro, J., Guirao, R. 2012
504 Comparison between individual and group rearing systems in *Octopus vulgaris*
505 (Cuvier, 1797). *Journal of the World Aquaculture Society* 43, 63-72.

506 Fuentes, L., Sanchez F.J., Lago M.J., Iglesias J., Pazos G., Linares, F. 2011. Growth and
507 survival of *Octopus vulgaris* (Cuvier 1797) paralarvae fed on three Artemia-based
508 diets complemented with frozen fish flakes, crushed zooplankton and marine
509 microalgae. *Scientia Marina* 75, 771-777.

510 C.D. Funk, Prostaglandins and leukotrienes: Advances in eicosanoid biology, *Science*
511 294 (2001) 1871–1875.

512 García-Garrido, S., Hachero-Cruzado, I., Garrido, D. Rosas, C., Domingues, P. (2010).
513 Lipid composition of the mantle and digestive gland of *Octopus vulgaris* juveniles
514 (Cuvier, 1797) exposed to prolonged starvation. *Aquacult Int* 18:1223–1241

515 Guillou H, Zadavec D, Martin PGP, Jacobsson A (2010). The key roles of elongases
516 and desaturases in mammalian fatty acid metabolism: Insights from transgenic mice.
517 Prog Lipid Res 49:186-199.

518 Hashimoto K, Yoshizawa AC, Okuda S, Kuma K, Goto S, Kanehisa M. (2008) The
519 repertoire of desaturases and elongases reveals fatty acid variations in 56 eukaryotic
520 genomes. J Lipid Res 49:183-191.

521 Hastings N, Agaba M, Tocher DR, Leaver MJ, Dick JR, Sargent JR, Teale AJ (2001) A
522 vertebrate fatty acid desaturase with $\Delta 5$ and $\Delta 6$ activities. Proc Natl Acad Sci USA
523 98:14304-14309.

524 Heinz E, Zank T, Zaehring U, Lerchl J, Renz A. Patent: WO 0159128-A, 16 August
525 2001.

526 Iglesias, J., Otero, J.J., Moxica, C., Fuentes, L., Sánchez, F.J., 2002. Paralarvae culture
527 of octopus (*Octopus vulgaris* Cuvier) using *Artemia* and crab zoeas and first data
528 on juvenile growth up to eight months of age. Aquac. Eur., vol. 32. European
529 Aquaculture Society, pp. 268–269. Special Publication.

530 Iglesias, J., Otero, J.J., Moxica, C., Fuentes, L., Sánchez, F.J., 2004. The completed life
531 cycle of the octopus (*Octopus vulgaris*, Cuvier) under culture conditions: paralarval
532 rearing using *Artemia* and zoeae, and first data on juvenile growth up to 8 months
533 of age. Aquac. Int. 12, 481–487.

534 Iglesias J, Fuentes L, Sánchez J, Otero JJ, Moxica C, Lago MJ (2006) First feeding of
535 *Octopus vulgaris* Cuvier, 1797 paralarvae using *Artemia*: Effect of prey size, prey
536 density and feeding frequency. Aquaculture 261:817–822.

537 Iglesias J, Sánchez FJ, Bersano JGF, Carrasco JF, Dhont J, Fuentes L, Linares F, Muñoz
538 JL, Okumura S, Roo J, van der Meeren T, Vidal EAG, Villanueva R (2007)

539 Rearing of *Octopus vulgaris* paralarvae: Present status, bottlenecks and trends.
540 Aquaculture 266:1–15.

541 Inagaki K, Aki T, Fukuda Y, Kawamoto S, Shigeta S, Ono K, Suzuki O. 2002.
542 Identification and expression of a rat fatty acid elongase involved in the biosynthesis
543 of C18 fatty acids. Biosci Biotechnol Biochem 66:613–21.

544 Jakobsson A, Westerberg R, Jacobsson A (2006) Fatty acid elongases in mammals:
545 Their regulation and roles in metabolism. Prog Lipid Res 45:237-249.

546 Jiang X, Qin L, Tian B, Shu Z, Huang J., 2008. Cloning and expression of two elongase
547 genes involved in the biosynthesis of docosahexaenoic acid in *Thraustochytrium* sp.
548 FJN-10. Acta microbiologica Sinica 48, 176-83.

549 Joseph JD (1982) Lipid composition of marine and estuarine invertebrates. Part II:
550 Mollusca. Prog Lipid Res 21:109-153.

551 Kawashima H. (2005) Unusual minor nonmethylene-interrupted di-, tri-, and tetraenoic
552 fatty acids in limpet gonads. Lipids 40:627-630.

553 Klingensmith J (1982) Distribution of methylene and nonmethylene-interrupted dienoic
554 fatty acids in polar lipids and triacylglycerols of selected tissues of the hardshell clam
555 (*Mercenaria mercenaria*). Lipids 17:976–981

556 Kornprobst JM, Barnathan G (2010) Demospongiac acids revisited. Mar Drugs 8:2569-
557 2577.

558 Leonard, A.E., Kelder, B., Bobik, E.G., Kroeger, P.E., Chuang, L.-T., Thurmond, J.M.,
559 Parker-Barnes, J.M., Kopchick, J.J., Huang, Y.-S., Murkerji, P., 2000. cDNA cloning
560 and characterisation of human $\Delta 5$ desaturase involved in the synthesis of arachidonic
561 acid. Biochem. J. 347, 719–724.

562 Leonard, A.E., Kelder, B., Bobik, E.G., Chuang, L., Lewis, C.J., Kopchick, J.J.,
563 Mukerji, P., Huang, Y., 2002. Identification and expression of mammalian long-
564 chain PUFA elongation enzymes. *Lipids* 37, 733–740.

565 Leonard, A.E., Pereira, S.L., Sprecher, H., Huang, Y.-S., 2004. Elongation of long-
566 chain fatty acids. *Prog. Lipid Res.* 43, 36–54.

567 Li Y, Monroig Ó, Zhang L, Wang S, Zheng X, Dick JR, You C, Tocher DR (2010)
568 Vertebrate fatty acyl desaturase with $\Delta 4$ activity. *Proc Natl Acad Sci USA*
569 107:16840-16845.

570 Milou H, Fintikaki M, Tzitzinakis M, Kountouris T, Virriopoulos G (2006) Fatty acid
571 composition of the common octopus, *Octopus vulgaris*, in relation to rearing
572 temperature and body weight. *Aquaculture* 256:311-322.

573 Mohd-Yusof, N.Y., Monroig, Ó., Mohd-Adnan, A., Wan, K.-L., Tocher, D.R., 2010.
574 Investigation of highly unsaturated fatty acid metabolism in the Asian sea bass,
575 *Lates calcarifer*. *Fish Physiol. Biochem.* 3, 827–843.

576 Monroig Ó, Rotllant J, Sánchez E, Cerdá-Reverter JM, Tocher DR (2009) Expression of
577 long-chain polyunsaturated fatty acid (LC-PUFA) biosynthesis genes during
578 zebrafish *Danio rerio* early embryogenesis. *Biochim Biophys Acta* 1791:1093–
579 1101.

580 Monroig Ó, Zheng X, Morais S, Leaver MJ, Taggart JB, Tocher DR (2010a) Multiple
581 genes for functional $\Delta 6$ fatty acyl desaturases (Fad) in Atlantic salmon (*Salmo salar*
582 L.): Gene and cDNA characterization, functional expression, tissue distribution and
583 nutritional regulation. *Biochim Biophys Acta* 1801:1072–1081.

584 Monroig, Ó., Rotllant, J., Cerdá-Reverter, J.M., Dick, J.R., Figueras, A., Tocher, D.R.,
585 2010b. Expression and role of Elovl4 elongases in biosynthesis of very long-chain
586 fatty acids during zebrafish *Danio rerio* early embryonic development. *Biochim.*

587 Biophys. Acta 1801, 1145-1154.

588 Monroig, Ó., Navarro, J.C., Tocher, D.R., 2011a. Long-chain polyunsaturated fatty
589 acids in fish: recent advances on desaturases and elongases involved in their
590 biosynthesis. In: Cruz-Suarez, L.E., Ricque-Marie, D., Tapia-Salazar, M., Nieto-
591 López, M.G., Villarreal-Cavazos, D. A., Gamboa-Delgado, J., Hernández-
592 Hernández, L.H. (Eds), Proceedings of the XI International Symposium on
593 Aquaculture Nutrition, 23-25 November 2011, San Nicolás de los Garza, N.L.
594 Universidad Autónoma de Nuevo León, Monterrey, Nuevo León, México. ISBN
595 978-607-433-775-4. Universidad Autónoma de Nuevo León, Monterrey, México.
596 pp. 257-282.

597 Monroig, Ó., Webb, K., Ibarra-Castro, L., Holt, G.J., Tocher, D.R. 2011b. Biosynthesis
598 of long-chain polyunsaturated fatty acids in marine fish: Characterization of an
599 Elovl4-like elongase from cobia *Rachycentron canadum* and activation of the
600 pathway during early life stages. Aquaculture 312, 145–153.

601 Monroig Ó, Li Y, Tocher DR, 2011c. Delta-8 desaturation activity varies among fatty
602 acyl desaturases of teleost fish: high activity in delta-6 desaturases of marine
603 species. Comp Biochem Physiol 159B, 206-213.

604 Monroig, Ó., Navarro, J.C., Dick, J.R., Alemany, F., Tocher, D.R. (2012a).
605 Identification of a Δ 5-like fatty acyl desaturase from the cephalopod *Octopus*
606 *vulgaris* (Cuvier 1797) involved in the biosynthesis of essential fatty acids. Marine
607 Biotechnology 14, 411-422.

608 Monroig, Ó., Wang S., Zhang L., You, C., Tocher D.R., Li Y. (2012b). Elongation of
609 long-chain fatty acids in rabbitfish *Siganus canaliculatus*: Cloning, functional
610 characterisation and tissue distribution of Elovl5- and Elovl4-like elongases.
611 Aquaculture 350-353, 63-70.

612 Morais, S., Monroig, Ó., Zheng, X., Leaver, M.J., Tocher, D.R., 2009. Highly
613 unsaturated fatty acid synthesis in Atlantic salmon: characterization of Elovl5- and
614 Elovl2-like elongases. *Mar. Biotechnol.* 11, 627–639.

615 Morais, S., Mourente, G., Ortega, A., Tocher, J.A., Tocher, D.R., 2011. Expression of
616 fatty acyl desaturase and elongase genes, and evolution of DHA:EPA ratio during
617 development of Atlantic bluefin tuna (*Thunnus thynnus* L.). *Aquaculture* 313, 129-
618 139.

619 Navarro JC, Villanueva R (2000) Lipid and fatty acid composition of early stages of
620 cephalopods: an approach to their lipid requirements. *Aquaculture* 183:161–177.

621 Navarro JC, Villanueva R (2003) The fatty acid composition of *Octopus vulgaris*
622 paralarvae reared with live and inert food: deviation from their natural fatty acid
623 profile. *Aquaculture* 219:613–631.

624 Otero J, González ÁF, Sieiro MP, Guerra Á (2007) Reproductive cycle and energy
625 allocation of *Octopus vulgaris* in Galician waters, NE Atlantic. *Fish Res* 85:122-
626 129.

627 Pirini M, Manuzzi MP, Pagliarani A, Trombetti F, Borgatti AR, Ventrella V (2007)
628 Changes in fatty acid composition of *Mytilus galloprovincialis* (Lmk) fed on
629 microalgal and wheat germ diets. *Comp Biochem Physiol* 147B:616–626.

630 Quintana D (2009) Valoración de los requerimientos nutricionales de reproductores de
631 pulpo común (*Octopus vulgaris*). PhD Thesis. Universidad de La Laguna.

632 Saitou N, Nei M (1987) The neighbor-joining method. A new method for reconstructing
633 phylogenetic trees. *Mol Biol Evol* 4:406-425.

634 Seixas P, Otero A, Valente LMP, Dias J, Rey-Méndez M (2010) Growth and fatty acid
635 composition of *Octopus vulgaris* paralarvae fed with enriched *Artemia* or co-fed
636 with an inert diet. *Aquacult Int* 18:1121-1135.

637 Tvrdik P, Westerberg R, Silve S, Asadi A, Jakobsson A, Cannon B, Loison G,
638 Jacobsson A. 2000. Role of a new mammalian gene family in the biosynthesis of
639 very long chain fatty acids and sphingolipids. *J Cell Biol* 149:707-18.

640 Valverde JC, Garcia BG, 2005. Suitable dissolved oxygen levels for common octopus
641 (*Octopus vulgaris* Cuvier, 1797) at different weights and temperatures: analysis of
642 respiratory behaviour. *Aquaculture* 244, 303-314.

643 Van der Horst DJ (1973) Biosynthesis of saturated and unsaturated fatty acids in the
644 pulmonate land snail *Cepaea nemoralis* (L.). *Comp Biochem Physiol* 46:551–560.

645 Van der Horst DJ (1974) In vivo biosynthesis of fatty acids in the pulmonate land snail
646 *Cepaea nemoralis* (L.) under anoxic conditions. *Comp Biochem Physiol* 47B:181-
647 187.

648 Viciano E, Iglesias J, Lago MJ, Sánchez FJ, Otero JJ, Navarro JC (2011) Fatty acid
649 composition of polar and neutral lipid fractions of *Octopus vulgaris* Cuvier, 1797
650 paralarvae reared with enriched on-grown *Artemia*. *Aquacult Res* 42:704-709.

651 Villanueva R (1994) Decapod crab zoeae as food for rearing cephalopod paralarvae.
652 *Aquaculture* 128:143-152.

653 Villanueva R (1995) Experimental rearing and growth of planktonic *Octopus vulgaris*
654 from hatching to settlement. *Can J Fish Aquat Sci* 52:2639–2650.

655 Villanueva R, Riba J, Ruíz-Capillas C, González AV, Baeta M (2004) Amino acid
656 composition of early stages of cephalopods and effects of amino acid dietary
657 treatments on *Octopus vulgaris* paralarvae. *Aquaculture* 242:455–478.

658 Villanueva R, Bustamante P (2006) Composition in essential and non-essential elements
659 of early stages of cephalopods and dietary effects on the elemental profiles of
660 *Octopus vulgaris* paralarvae. *Aquaculture* 261:225-240.

661 Villanueva R, Escudero JM, Deulofeu R, Bozzano A, Casoliva C (2009) Vitamin A and
662 E content in early stages of cephalopods and their dietary effects in *Octopus*
663 *vulgaris paralarvae*. *Aquaculture* 286:277-282.

664 Waldock MJ, Holland DL (1984) Fatty acid metabolism in young oysters, *Crassostrea*
665 *gigas*: Polyunsaturated fatty acids. *Lipids* 19:332-336.

666 Weinert J, Blomquist GJ, Borgeson CE (1993) De novo biosynthesis of linoleic acid in
667 two non-insect invertebrates: the land slug and the garden snail. *Experientia*
668 49:919-921.

669 Wodinsky J (2008) Reversal and transfer of spermatophores by *Octopus vulgaris* and *O.*
670 *hummelincki*. *Mar Biol* 155:91-103.

671 Zheng X, Ding Z, Xu Y, Monroig O, Morais S, Tocher DR (2009) Physiological roles
672 of fatty acyl desaturase and elongase in marine fish: Characterisation of cDNAs of
673 fatty acyl $\Delta 6$ desaturase and Elovl5 elongase of cobia (*Rachycentron canadum*).
674 *Aquaculture* 290:122-131.

675 Zhu N, Dai X, Lin DS, Connor WE (1994) The lipids of slugs and snails: Evolution,
676 diet and biosynthesis. *Lipids* 29:869-875.

677 Zhukova NV (1986) Biosynthesis of non-methylene-interrupted dienoic fatty acids from
678 ^{14}C -acetate in molluscs. *Biochim Biophys Acta* 878:131-133.

679 Zhukova NV (1991) The pathway of the biosynthesis of non-methylene-inter- rupted
680 dienoic fatty acids in Molluscs. *Comp Biochem Physiol* 100B:801–804.

681 Zhukova NV (2007) Lipid classes and fatty acid composition of the tropical nudibranch
682 mollusks *Chromodoris* sp. and *Phyllidia coelestis*. *Lipids* 42:1169–1175.

683

684

684 **Tables**

685 Table 1. Sequences of the primer pairs used and accession numbers of the sequences
 686 used as references for primer design in the cloning of the octopus elongase of very long-
 687 chain fatty acids (Elovl) ORF and for RT-PCR analysis of gene expression in octopus
 688 tissues.

Aim	Transcript	Primer	Primer sequence	Ac
<i>RACE PCR</i>	Elovl	OVEloF1	5'-GACTTGGTTCGGTGCTTGTT-3'	
		OVEloF2	5'-ATGGCCTGTCTGCTATACCAT-3'	
		OVEloR1	5'-ATGGTATAGCAGACAGGCCAT-3'	
		OVEloR2	5'-ATGATGGAAGACATGCAGGAA-3'	
<i>ORF cloning</i>	Elovl	OVEloVF	5'-CCCAAGCTTAAAATGGCGGACGTTGTG-3'	
		OVEloVR	5'-CCGGAGCTCCTATTGAGCTTTCTTCACC-3'	
<i>RT-PCR</i>	Elovl	OVEloF1	5'-GACTTGGTTCGGTGCTTGTT-3'	
		OVEloR3	5'-GTCTGCCTTTGATGTAAGCCTG-3'	
	β-actin	OVACTF	5'-CTTGACTCCGGAGATGGTGT-3'	
		OVACTR	5'-CGCATTTCATGATGGAGTTG-3'	

689 ^a GenBank (<http://www.ncbi.nlm.nih.gov/>)

690

691 Table 2. Functional characterisation of the octopus elongase in *Saccharomyces*
 692 *cerevisiae*. Results are expressed as a percentage of total fatty acid (FA) substrate
 693 converted to elongated products.

FA Substrate	Product	% Conversion	Activity
18:3n-3	20:3n-3	13.4	C18→20
18:2n-6	20:2n-6	40.8	C18→20
18:4n-3	20:4n-3	36.9	C18→20
18:3n-6	20:3n-6	52.3	C18→20
20:5n-3	22:5n-3	2.4	C20→22
20:4n-6	22:4n-6	15.9	C20→22
22:5n-3	24:5n-3	0.0	C22→24
22:4n-6	24:4n-6	0.0	C22→24

694


695 Table 3. Fatty acid (FA) composition (% of totals) from the polar lipids of tissues
 696 collected from *Octopus vulgaris* adult individuals. FA are designated using the ‘n-’
 697 nomenclature, except for non-methylene-interrupted FA where the ‘Δ’ nomenclature
 698 was used.

	Nephrydium	Male gonad	Eye	Caecum
14:0	0.7	0.5	0.7	1.3
15:0	0.3	0.3	0.3	0.3
16:0	14.2	14.7	18.9	14.8
16:1n-9	nd	0.5	0.2	0.1
16:1n-7	0.5	0.4	0.4	1.9
16:0 iso	0.2	0.1	0.2	0.2
16:0 anteiso	0.2	nd	0.1	nd
17:0	2.7	1.4	1.2	1.9
17:1	nd	0.1	0.2	nd
17:0 iso	0.3	nd	0.2	0.2
18:0	13.6	8.4	7.4	14.9
18:1n-13	0.6	0.9	1.4	0.3
18:1n-9	2.0	2.1	1.1	3.1
18:1n-7	2.0	1.3	1.4	2.6
18:1n-5	0.3	0.1	nd	nd
18:2n-6	0.1	nd	0.8	0.3
18:3n-3	nd	nd	0.1	0.2
18:4n-3	nd	nd	nd	0.1
20:0	0.1	0.1	0.1	0.2
20:1n-11	0.5	0.5	0.5	1.0
20:1n-9	9.2	10.5	2.4	2.8
20:1n-7	0.2	0.2	0.1	0.2
Δ5,11 20:2	1.8	nd	nd	nd
Δ7,13 20:2	nd	0.2	0.1	0.1
20:2n-6	0.3	0.1	0.8	0.4
Δ5,11,14 20:3	0.8	nd	nd	nd
20:3n-6	0.1	nd	0.2	0.1
20:4n-6	11.9	16.4	5.1	13.6
20:3n-3	0.1	nd	13.5	0.1
20:5n-3	10.0	7.3	11.4	13.7
22:0	0.2	0.1	0.2	0.3
22:1n-11	2.0	2.3	0.4	1.8
22:1n-9	0.1	0.1	nd	0.3
Δ7,13 22:2	nd	2.2	0.4	0.8
21:5n-3	0.1	nd	0.1	nd
22:2n-6	nd	nd	nd	0.3
22:4n-6	1.1	7.7	0.3	1.1
22:5n-6	1.0	0.8	0.2	0.8
22:5n-3	1.3	2.0	1.0	1.5
22:6n-3	20.3	17.4	27.0	15.5
24:1n-9	0.1	0.1	0.2	0.4

699 nd, no detected.

700


700 **Legends to Figures**



701


702 Fig. 1. Alignment of the deduced amino acid (aa) sequence of the elongase from
 703 *Octopus vulgaris* (Ov). The aa sequence of the octopus Elov1-like protein was aligned
 704 with the *Homo sapiens* (Hs) ELOVL2 (gb|NP_060240|) and ELOVL5 (gb|NP_068586|),
 705 the *Danio rerio* (Ds) Elov12 (gb|NP_001035452|) and Elov15 (gb|NP_956747|) and the
 706 so-called Elov1-like transcript 1 (jgi|Lotg1|224291|) from *Lottia gigantea* (Lg) using
 707 ClustalW (Bioedit). Identical residues are shaded black and similar residues are shaded
 708 grey. Identity/similarity shading was based on the BLOSUM62 matrix, and the cut-off
 709 for shading was 70%. The histidine box (HXXHH) conserved among Elov1 family
 710 members is highlighted with a grey square. Five (I-V) transmembrane-regions predicted
 711 by InterProScan (<http://www.ebi.ac.uk/Tools/pfa/iprscan/>) are dot-underlined.

712




713

714 Fig. 2. Phylogenetic tree comparing the deduced amino acid (aa) sequence of the
715 *Octopus vulgaris* elongase of very long-chain fatty acids (Elovl)-like with a series of
716 protein sequences including representatives of the seven (1-7) Elovl subtypes and other
717 Elovl-like sequences from invertebrate organisms. All accession numbers are from
718 GenBank database, except for *Lottia gigantea* elongases where JGI protein ID are given
719 (<http://genome.jgi-psf.org/Lotgi1/Lotgi1.home.html>). Asterisks indicate the aa
720 sequences deduced from searches and subsequent assembly of expressed sequence tags
721 (EST) using NCBI tblastn tool (<http://www.ncbi.nlm.nih.gov/>) as described in Materials
722 and Methods. The tree was constructed using the Neighbour Joining method (Saitou and
723 Nei 1987) with MEGA4. The horizontal branch length is proportional to aa substitution
724 rate per site. The numbers represent the frequencies (%) with which the tree topology
725 presented was replicated after 10000 iterations.



727

728 Fig. 3. Functional characterisation of the *Octopus vulgaris* elongase of very long-chain
 729 fatty acids (Elov1) in yeast (*Saccharomyces cerevisiae*). The fatty acid (FA) profiles of
 730 yeast transformed with pYES2 containing the ORF of the putative Elov1 cDNA as an
 731 insert, were determined after the yeast was grown in the presence of one of the
 732 exogenously added substrates 18:2n-6 (A), 18:3n-6 (B) and 20:3n-6 (C). Peaks 1-5 in all
 733 panels are the main endogenous FA of *S. cerevisiae*, namely 16:0 (1), 16:1 isomers (2),
 734 18:0 (3), 18:1n-9 (4) and 18:1n-7 (5). Additionally peaks derived from exogenously
 735 added substrates (“*”) or elongation products are indicated accordingly in panels A-C.
 736 Vertical axis, FID response; horizontal axis, retention time.



737

738 Fig. 4. RT-PCR analyses showing the tissue distribution of octopus elongase of very
 739 long-chain fatty acids (Elov1) transcripts. Expression of the housekeeping gene β -actin
 740 is also shown. NTC, no template control.