

JOHNSON O. OLALERU* AND HUDSON AKEWE

THE CONVERGENCE OF JUNGCK-TYPE ITERATIVE SCHEMES FOR GENERALIZED CONTRACTIVE-LIKE OPERATORS

ABSTRACT. We introduce the Jungck-multistep iteration and show that it converges strongly to the unique common fixed point of a pair of weakly compatible generalized contractive-like operators defined on a Banach space. As corollaries, the results show that the Jungck-Mann, Jungck-Ishikawa and Jungck-Noor iterations can also be used to approximate the common fixed points of such maps. The results are improvements, generalizations and extensions of the work of Olatinwo and Imoru [13], Olatinwo [14-15]. Consequently, several results in literature are generalized.

KEY WORDS: iteration schemes, quasi-contractive operators, weakly compatible maps, coincidence points, common fixed points.

AMS Mathematics Subject Classification: 47H10.

1. Introduction

The convergence of Picard, Mann, Ishikawa, Noor and multistep iterations have been commonly used to approximate the fixed points of several classes of single quasi-contractive operators. For example see [1-3, 7, 10, 24].

Let X be a Banach space, K , a nonempty convex subset of X and $T : K \rightarrow K$ a self map of K .

Definition 1. Let $z_0 \in K$. The Picard iteration scheme $\{z_n\}_{n=0}^{\infty}$ is defined by

$$(1) \quad z_{n+1} = Tz_n, \quad n \geq 0.$$

Definition 2. For any given $u_0 \in K$, the Mann iteration scheme [8] $\{u_n\}_{n=0}^{\infty}$ is defined by

$$(2) \quad u_{n+1} = (1 - \alpha_n)u_n + \alpha_n T u_n.$$

*This author was supported by the Central Research Committee of the University of Logos

where $\{\alpha_n\}_{n=0}^{\infty}$ are real sequences in $[0, 1)$ such that $\sum_{n=0}^{\infty} \alpha_n = \infty$.

Definition 3. Let $x_0 \in K$. The Ishikawa iteration scheme [5] $\{x_n\}_{n=0}^{\infty}$ is defined by

$$(3) \quad \begin{aligned} x_{n+1} &= (1 - \alpha_n)x_n + \alpha_n T y_n \\ y_n &= (1 - \beta_n)x_n + \beta_n T x_n \end{aligned}$$

where $\{\alpha_n\}_{n=0}^{\infty}, \{\beta_n\}_{n=0}^{\infty}$ are real sequences in $[0, 1)$ such that $\sum_{n=0}^{\infty} \alpha_n = \infty$.

Observe that if $\beta_n = 0$ for each n , then the Ishikawa iteration process (3) reduces to the Mann iteration scheme (2).

Definition 4. Let $x_0 \in K$. The Noor iteration (or three-step) scheme [9] $\{x_n\}_{n=0}^{\infty}$ is defined by

$$(4) \quad \begin{aligned} x_{n+1} &= 1 - \alpha_n)x_n + \alpha_n T y_n \\ y_n &= (1 - \beta_n)x_n + \beta_n T z_n \\ z_n &= (1 - \gamma_n)x_n + \gamma_n T x_n \end{aligned}$$

where $\{\alpha_n\}_{n=0}^{\infty}, \{\beta_n\}_{n=0}^{\infty}, \{\gamma_n\}_{n=0}^{\infty}$ are real sequences in $[0, 1)$ such that $\sum_{n=0}^{\infty} \alpha_n = \infty$.

Observe that if $\gamma_n = 0$ for each n , then the Noor iteration process (4) reduces to the Ishikawa iteration scheme (3).

Definition 5. Let $x_0 \in K$. The multistep iteration scheme [19] $\{x_n\}_{n=0}^{\infty}$ is defined by

$$(5) \quad \begin{aligned} x_{n+1} &= (1 - \alpha_n)x_n + \alpha_n T y_n^1 \\ y_n^i &= (1 - \beta_n^i)x_n + \beta_n^i T y_n^{i+1}, i = 1, 2, \dots, p-2, \\ y_n^{p-1} &= (1 - \beta_n^{p-1})x_n + \beta_n^{p-1} T x_n, p \geq 2 \end{aligned}$$

where $\{\alpha_n\}_{n=0}^{\infty}, \{\beta_n^i\}, i = 1, 2, \dots, p-1$ are real sequences in $[0, 1)$ such that $\sum_{n=0}^{\infty} \alpha_n = \infty$.

Observe that the multistep iteration is a generalization of the Noor, Ishikawa and the Mann iterations. In fact, if $p = 1$ in (5), we have the Mann iteration (2); if $p = 2$ in (5), we have the Ishikawa iteration (3) and if $p = 3$, we have the Noor iterations (4).

We note that while many authors have worked on the existence of fixed points for a pair of quasicontractive maps, e.g. see [1, 4, 6, 21-22], little is known about the approximations of those common fixed points using the convergence of iteration techniques. Jungck was the first to introduce an iteration scheme, which is now called Jungck iteration scheme [6] to approximate the common fixed points of what is now called Jungck contraction

maps. Singh et al. [22] of recent introduced the Jungck-Mann iteration procedure and discussed it's stability for a pair of contractive maps. Olatinwo and Imoru [13], Olatinwo [14-15] built on that work to introduce the Jungck-Ishikawa and Jungck-Noor iteration schemes and used their convergences to approximate the coincidence points (not common fixed points) of some pairs of generalized contractive-like operators with the assumption that one of each of the pairs of maps is injective. However, a coincidence point for a pair of quasicontractive maps need not be a common fixed point. We introduce the Jungck-multistep iteration and show that its convergence can be used to approximate the common fixed points of those pairs of quasicontractive maps without assuming the injectivity of any of the operators. Hence the iterative sequence used is a generalization of the those used in [13-15]. The fact that the injectivity of any of the maps is not assumed in our results and the common fixed points of those maps are approximated and not just the coincidence points make the corollary of our results an improvement of the results of Olatinwo [14-15], Olatinwo and Imoru [13]. Consequently, a lot of results dealing with convergence of Picard, Mann, Ishikawa and multistep iterations for single quasicontractive operators on Banach spaces are generalized.

2. Preliminaries

Let X be a Banach space, Y be an arbitrary set and $S, T : Y \rightarrow X$ such that $T(Y) \subseteq S(Y)$. Then we have the following definitions.

Definition 6 (21). *For any $x_o \in Y$, there exists a sequence $\{x_n\}_{n=0}^\infty \in Y$ such that $Sx_{n+1} = Tx_n$. The Jungck iteration is defined as the sequence $\{Sx_n\}_{n=1}^\infty$ such that*

$$(6) \quad Sx_{n+1} = Tx_n, \quad n \geq 0.$$

This procedure becomes Picard iteration when $Y = X$ and $S = I_d$ where I_d is the identity map on X .

Similarly, the Jungck contraction maps are the maps S, T satisfying

$$(7) \quad d(Tx, Ty) \leq k d(Sx, Sy), \quad 0 \leq k < 1 \text{ for all } x, y \in Y.$$

If $Y = X$ and $S = I_d$, then maps satisfying (7) become the well known contraction maps.

Definition 7 (22). *For any given $u_o \in Y$, the Jungck-Mann iteration scheme $\{Su_n\}_{n=1}^\infty$ is defined by*

$$(8) \quad Su_{n+1} = (1 - \alpha_n)Su_n + \alpha_n Tu_n$$

where $\{\alpha_n\}_{n=0}^\infty$ are real sequences in $[0, 1)$ such that $\sum_{n=0}^\infty \alpha_n = \infty$.

Definition 8 (15). Let $x_o \in Y$. The Jungck-Ishikawa iteration scheme $\{Sx_n\}_{n=1}^\infty$ is defined by

$$(9) \quad \begin{aligned} Sx_{n+1} &= (1 - \alpha_n)Sx_n + \alpha_nTy_n \\ Sy_n &= (1 - \beta_n)Sx_n + \beta_nTx_n \end{aligned}$$

where $\{\alpha_n\}_{n=0}^\infty, \{\beta_n\}_{n=0}^\infty$ are real sequences in $[0, 1)$ such that $\sum_{n=0}^\infty \alpha_n = \infty$.

Definition 9 (15). Let $x_o \in Y$. The Jungck-Noor iteration (or three-step) scheme $\{Sx_n\}_{n=1}^\infty$ is defined by

$$(10) \quad \begin{aligned} Sx_{n+1} &= (1 - \alpha_n)Sx_n + \alpha_nTy_n \\ Sy_n &= (1 - \beta_n)Sx_n + \beta_nTz_n \\ Sz_n &= (1 - \gamma_n)Sx_n + \gamma_nTx_n \end{aligned}$$

where $\{\alpha_n\}_{n=0}^\infty, \{\beta_n\}_{n=0}^\infty$ and $\{\gamma_n\}_{n=0}^\infty$ are real sequences in $[0, 1)$ such that $\sum_{n=0}^\infty \alpha_n = \infty$.

Definition 10. Let $x_o \in Y$. The Jungck-multistep iteration scheme $\{Sx_n\}_{n=1}^\infty$ is defined by

$$(11) \quad \begin{aligned} Sx_{n+1} &= (1 - \alpha_n)Sx_n + \alpha_nTy_n^1 \\ Sy_n^i &= (1 - \beta_n^i)Sx_n + \beta_n^iTy_n^{i+1}, i = 1, 2, \dots, p-2, \\ Sy_n^{p-1} &= (1 - \beta_n^{p-1})Sx_n + \beta_n^{p-1}Tx_n, p \geq 2 \end{aligned}$$

where $\{\alpha_n\}_{n=0}^\infty, \{\beta_n^i\}, i = 1, 2, \dots, p-1$ are real sequences in $[0, 1)$ such that $\sum_{n=0}^\infty \alpha_n = \infty$.

Observe that that the Jungck-multistep iteration is a generalization of the Jungck-Noor, Jungck-Ishikawa and the Jungck-Mann iterations. In fact, if $p = 1$ in (11), we have the Jungck-Mann iteration (8); if $p = 2$ in (11), we have the Jungck-Ishikawa iteration (9) and if $p = 3$, we have the Jungck-Noor iterations (10).

Observe that if $X = Y$ and $S = I_d$, then the Jungck-multistep (11), Jungck-Noor (10), Jungck-Ishikawa (9) and the Jungck-Mann (8) iterations respectively become the multistep (5), Noor (4), Ishikawa (3) and the Mann (2) iterative procedures.

One of the most general contractive-like operators which has been studied by several authors is the Zamfirescu operators. Suppose X is a Banach space. The map $T : X \rightarrow X$ is called a Zamfirescu operator if

$$(12) \quad \|Tx - Ty\| \leq h \max \left\{ \|x - y\|, \frac{\|x - Tx\| + \|y - Ty\|}{2}, \frac{\|x - Ty\| + d\|y - Tx\|}{2} \right\},$$

where $0 \leq h < 1$, see [24]. It is known that the operators satisfying (12) are generalizations of Kannan maps [7] and Chatterjea maps [3]. Zamfirescu [24] proved that the Zamfirescu operator has a unique fixed point which can be approximated by Picard iteration (1). Berinde [2] showed that Ishikawa iteration can be used to approximate the fixed point of a Zamfirescu operator when X is a Banach space while it was shown by the first author [10] that if X is generalised to a complete metrizable locally convex space (which includes Banach spaces), the Mann iteration can be used to approximate the fixed point of a Zamfirescu operator. Several researchers have studied the convergence rate of these iterations with respect to the Zamfirescu operators. For example, it has been shown that the Picard iteration (1) converges faster than the Mann iteration (2) when dealing with the Zamfirescu operators. For example, see [20]. It is still a subject of research as to conditions under which the Mann iteration will converge faster than the Ishikawa or vice-versa when dealing with the Zamfirescu operators.

We now consider the following conditions. X is a Banach space and Y a nonempty set such that $T(Y) \subseteq S(Y)$ and $S, T : Y \rightarrow X$. For $x, y \in Y$ and $h \in (0, 1)$:

$$(13) \quad \|Tx - Ty\| \leq h \max \left\{ \|Sx - Sy\|, \frac{\|Sx - Tx\| + \|Sy - Ty\|}{2}, \frac{\|Sx - Ty\| + \|Sy - Tx\|}{2} \right\},$$

$$(14) \quad \|Tx - Ty\| \leq h \max \left\{ \|Sx - Sy\|, \frac{\|Sx - Tx\| + \|Sy - Ty\|}{2}, \|Sx - Ty\|, \|Sy - Tx\| \right\},$$

$$(15) \quad \|Tx - Ty\| \leq \delta \|Sx - Sy\| + L \|Sx - Tx\|, \quad L > 0, \quad 0 < \delta < 1,$$

$$(16) \quad \|Tx - Ty\| \leq \frac{\delta \|Sx - Sy\| + \varphi(\|Sx - Tx\|)}{1 + M \|Sx - Tx\|}, \quad 0 \leq \delta < 1, \quad M \geq 0.$$

$$(17) \quad \|Tx - Ty\| \leq \delta \|Sx - Sy\| + \varphi(\|Sx - Tx\|), \quad 0 \leq \delta < 1.$$

where $\varphi : \mathfrak{R}_+ \rightarrow \mathfrak{R}_+$ is a monotone increasing sequence with $\varphi(0) = 0$.

Remark 1. Observe that if $X = Y$ and $S = I_d$, (13) is the same as the Zamfirescu operator (12) already studied by several authors; (14) becomes the operator studied by Rhoades [18]; while (15) becomes the operator introduced by Osilike [16]. Operators satisfying (16) and (17) were introduced by Olatinwo [15].

A comparison of the four maps show the following.

Proposition 1. (13) \Rightarrow (14) \Rightarrow (15) \Rightarrow (16) \Rightarrow (17) *but the converses are not true.*

Proof. (13) \Rightarrow (14): This follows immediately since $\frac{\|Sx-Ty\|+\|Sy-Tx\|}{2} \leq \max\{\|Sx-Ty\|, \|Sy-Tx\|\}$. (14) \Rightarrow (15): We consider each of the possibilities.

Case 1: Suppose $\|Tx-Ty\| \leq h\|Sx-Ty\| \leq h\|Sx-Tx\| + h\|Tx-Ty\|$ and consequently, $\|Tx-Ty\| \leq \frac{h}{1-h}\|Sx-Tx\|$. Setting $L = \frac{h}{1-h}$ completes the proof.

Case 2: Suppose

$$\begin{aligned} \|Tx-Ty\| &\leq h \frac{\|Sx-Tx\| + \|Sy-Ty\|}{2} \\ &\leq h \frac{\|Sx-Tx\| + \|Sy-Sx + Sx-Tx + Tx-Ty\|}{2} \\ &\leq h\|Sx-Tx\| + \frac{h}{2}\|Sy-Sx\| + \frac{h}{2}\|Tx-Ty\|. \end{aligned}$$

After computing we have $\|Tx-Ty\| \leq \frac{h}{2-h}\|Sy-Sx\| + \frac{2h}{2-h}\|Sx-Tx\|$. Setting $\delta = \frac{h}{2-h}$ and $L = \frac{2h}{2-h}$ completes the proof.

Case 3: $\|Tx-Ty\| \leq h\|Sy-Tx\| \leq h\|Sy-Sx\| + h\|Sx-Tx\|$.

(15) \Rightarrow (16): Suppose $M = 0$ and $\varphi(t) = Lt$ in (16), we have (15).

(16) \Rightarrow (17): This follows from the fact that $\|Tx-Ty\| \leq \frac{\delta\|Sx-Sy\| + \varphi(\|Sx-Tx\|)}{1+M\|Sx-Tx\|} \leq \delta\|Sx-Sy\| + \varphi(\|Sx-Tx\|)$ ■

We need the following definition.

Definition 11 (1). *A point $x \in X$ is called a coincident point of a pair of self maps S, T if there exists a point w (called a point of coincidence) in X such that $w = Sx = Tx$. Self-maps S and T are said to be weakly compatible if they commute at their coincidence points, that is, if $Sx = Tx$ for some $x \in X$, then $STx = TStx$.*

Olatinwo and Imoru [13] proved that the Jungck-Mann and Jungck-Ishikawa converge to the *coincident point* of S, T defined by (13) when S is an injective operator. It was shown in [14] that the Jungck-Ishikawa iteration converges to the coincidence point of S, T defined by (17) when S is an injective operator while the same convergence result was proved for Jungck-Noor when S, T are defined by (16) [15]. (We note that the maps satisfying (14) and of course (15)-(17) need not have a coincidence point [22]). We rather prove the convergence of multi-step iteration to the unique *common fixed point* of S, T defined by (17), without assuming that S is injective, provided the coincident point exist for S, T .

3. Main results

The following Lemma is well known.

Lemma 1. *Let $\{a_n\}$ be a sequence of nonnegative numbers such that $a_{n+1} \leq (1 - \lambda_n)a_n$ for any n , where $\lambda_n \in [0, 1)$ and $\sum_{n=0}^{\infty} \lambda_n = \infty$. Then $\{a_n\}$ converges to zero.*

Theorem 1. *Let X be a Banach space and $S, T : Y \rightarrow X$ for an arbitrary set Y such that (17) holds and $T(Y) \subseteq S(Y)$. Assume S and T have a coincidence point z such that $Tz = Sz = p$. For any $x_o \in Y$, the Jungck-multistep iteration (11) $\{Sx_n\}_{n=1}^{\infty}$ converges to p .*

Further, if $Y = X$ and S, T commute at p (i.e. S and T are weakly compatible), then p is the unique common fixed point of S, T .

Proof. In view of (11) and (17) coupled with the fact that $Tz = Sz = p$, we have

$$\begin{aligned}
 (18) \quad \|Sx_{n+1} - p\| &\leq (1 - \alpha_n)\|Sx_n - p\| + \alpha_n\|Tz - Ty_n^1\| \\
 &\leq (1 - \alpha_n)\|Sx_n - p\| \\
 &\quad + \alpha_n[\delta\|Sz - Sy_n^1\| + \varphi(\|Sz - Tz\|)] \\
 &= (1 - \alpha_n)\|Sx_n - p\| + \delta\alpha_n\|p - Sy_n^1\|.
 \end{aligned}$$

An application of (11) and (17) give

$$\begin{aligned}
 (19) \quad \|Sy_n^1 - p\| &\leq (1 - \beta_n^1)\|Sx_n - p\| + \beta_n^1\|Tz - Ty_n^2\| \\
 &\leq (1 - \beta_n^1)\|Sx_n - p\| \\
 &\quad + \beta_n^1[\delta\|Sz - Sy_n^2\| + \varphi(\|Sz - Tz\|)].
 \end{aligned}$$

Substituting (19) in (18), we have

$$\begin{aligned}
 (20) \quad \|Sx_{n+1} - p\| &\leq (1 - \alpha_n)\|Sx_n - p\| + \delta\alpha_n(1 - \beta_n^1)\|Sx_n - p\| \\
 &\quad + \delta^2\alpha_n\beta_n^1\|Sy_n^2 - p\| \\
 &= (1 - (1 - \delta)\alpha_n - \delta\alpha_n\beta_n^1)\|Sx_n - p\| \\
 &\quad + \delta^2\alpha_n\beta_n^1\|Sy_n^2 - p\|.
 \end{aligned}$$

Similarly, an application of (11) and (17) give

$$(21) \quad \|Sy_n^2 - p\| \leq (1 - \beta_n^2)\|Sx_n - p\| + \delta\beta_n^2\|Sy_n^3 - p\|.$$

Substituting (21) in (20) we have

$$\begin{aligned}
 (22) \quad \|Sx_{n+1} - p\| &\leq (1 - (1 - \delta)\alpha_n - \delta\alpha_n\beta_n^1)\|Sx_n - p\| \\
 &\quad + \delta^2\alpha_n\beta_n^1(1 - \beta_n^2)\|Sx_n - p\| \\
 &\quad + \delta^3\alpha_n\beta_n^1\beta_n^2\|Sy_n^3 - p\| \\
 &= (1 - (1 - \delta)\alpha_n - (1 - \delta)\delta\alpha_n\beta_n^1 - \delta^2\alpha_n\beta_n^1\beta_n^2) \\
 &\quad \times \|Sx_n - p\| + \delta^3\alpha_n\beta_n^1\beta_n^2\|Sy_n^3 - p\|.
 \end{aligned}$$

Similarly, an application of (11) and (17) give

$$(23) \quad \|Sy_n^3 - p\| \leq (1 - \beta_n^3)\|Sx_n - p\| + \delta\beta_n^3\|Sy_n^4 - p\|.$$

Substituting (23) in (22) we have

$$(24) \quad \begin{aligned} \|Sx_{n+1} - p\| &\leq (1 - (1 - \delta)\alpha_n - (1 - \delta)\delta\alpha_n\beta_n^1 - \delta^2\alpha_n\beta_n^1\beta_n^2) \\ &\quad \times \|Sx_n - p\| + \delta^3\alpha_n\beta_n^1\beta_n^2(1 - \beta_n^3)\|Sx_n - p\| \\ &\quad + \delta^4\alpha_n\beta_n^1\beta_n^2\beta_n^3\|Sy_n^4 - p\| \\ &= (1 - (1 - \delta)\alpha_n - (1 - \delta)\delta\alpha_n\beta_n^1 - (1 - \delta)\delta^2\alpha_n\beta_n^1\beta_n^2 \\ &\quad - \delta^3\alpha_n\beta_n^1\beta_n^2\beta_n^3)\|Sx_n - p\| \\ &\quad + \delta^4\alpha_n\beta_n^1\beta_n^2\beta_n^3\|Sy_n^4 - p\| \\ &\leq (1 - (1 - \delta)\alpha_n - \delta^3\alpha_n\beta_n^1\beta_n^2\beta_n^3)\|Sx_n - p\| \\ &\quad + \delta^4\alpha_n\beta_n^1\beta_n^2\beta_n^3\|Sy_n^4 - p\|. \end{aligned}$$

Continuing the above process we have

$$(25) \quad \begin{aligned} \|Sx_{n+1} - p\| &\leq (1 - (1 - \delta)\alpha_n - \delta^{k-2}\alpha_n\beta_n^1\beta_n^2\beta_n^3\dots\beta_n^{k-2})\|Sx_n - p\| \\ &\quad + \delta^{k-1}\alpha_n\beta_n^1\beta_n^2\beta_n^3\dots\beta_n^{k-2}\|Sy_n^{k-1} - p\| \\ &\leq (1 - (1 - \delta)\alpha_n - \delta^{k-2}\alpha_n\beta_n^1\beta_n^2\beta_n^3\dots\beta_n^{k-2})\|Sx_n - p\| \\ &\quad + \delta^{k-1}\alpha_n\beta_n^1\beta_n^2\beta_n^3\dots\beta_n^{k-2}[(1 - \beta_n^{k-1})\|Sx_n - p\| \\ &\quad + \beta_n^{k-1}\|Tz - Tx_n\|] \\ &\leq (1 - (1 - \delta)\alpha_n - \delta^{k-2}\alpha_n\beta_n^1\beta_n^2\beta_n^3\dots\beta_n^{k-2})\|Sx_n - p\| \\ &\quad + \delta^{k-1}\alpha_n\beta_n^1\beta_n^2\beta_n^3\dots\beta_n^{k-2}[(1 - \beta_n^{k-1})\|Sx_n - p\| \\ &\quad + \delta\beta_n^{k-1}\|Sx_n - p\|] \\ &\leq (1 - (1 - \delta)\alpha_n - \delta^{k-2}\alpha_n\beta_n^1\beta_n^2\beta_n^3\dots\beta_n^{k-2}) \\ &\quad + \delta^{k-1}\alpha_n\beta_n^1\beta_n^2\beta_n^3\dots\beta_n^{k-2})\|Sx_n - p\| \\ &\leq (1 - (1 - \delta)\alpha_n)\|Sx_n - p\|. \end{aligned}$$

Hence by Lemma 1 $Sx_n \rightarrow p$. Next we show that p is unique. Suppose there exists another point of coincidence p^* . Then there is an $z^* \in X$ such that $Tz^* = Sz^* = p^*$. Hence, from (17) we have

$$\|p - p^*\| = \|Tz - Tz^*\| \leq \delta\|Sz - Sz^*\| + \varphi(\|Sz - Tz\|) = \delta\|p - p^*\|.$$

Since $\delta < 1$, then $p = p^*$ and so p is unique.

Since S, T are weakly compatible, then $TSz = STz$ and so $Tp = Sp$. Hence p is a coincidence point of S, T and since the coincidence point is unique, then $p = p^*$ and hence $Sp = Tp = p$ and therefore p is the unique common fixed point of S, T and the proof is complete. \blacksquare

Remark 2. Weaker versions of Theorem 1 are the results in [14-15] where S is assumed injective and the convergence is not to the common fixed point but to the coincidence point of S, T . Furthermore, the Jungck-multistep iteration used in Theorem 1 is more general than the Jungck-Ishikawa and the Jungck-Noor iteration used in [14-15].

It is already shown in [1] and [12] that if $S(Y)$ or $T(Y)$ is a complete subspace of X , then maps satisfying the generalized Zamfirescu operators (13) has a unique coincidence point. Hence we have the following results.

Theorem 2. *Let X be a Banach space and $S, T : X \rightarrow X$ such that $\|Tx - Ty\| \leq h \max\{\|Sx - Sy\|, \frac{\|Sx-Tx\|+\|Sy-Ty\|}{2}, \frac{\|Sx-Ty\|+\|Sy-Tx\|}{2}\}$ and $T(X) \subseteq S(X)$. Assume S and T are weakly compatible. For any $x_o \in X$, the Jungck-multistep iteration (11) $\{Sx_n\}_{n=1}^\infty$ converges to the unique common fixed point of S, T .*

Since the Jungck-Noor, Jungck-Ishikawa and Jungck-Mann iterations are special cases of Jungck-multistep iteration, then we have the following consequences.

Corollary 1. *Let X be a Banach space and $S, T : X \rightarrow X$ such that $\|Tx - Ty\| \leq h \max\{\|Sx - Sy\|, \frac{\|Sx-Tx\|+\|Sy-Ty\|}{2}, \frac{\|Sx-Ty\|+\|Sy-Tx\|}{2}\}$ and $T(X) \subseteq S(X)$. Assume S and T are weakly compatible. For any $x_o \in X$, the Jungck-Noor iteration (10) $\{Sx_n\}_{n=1}^\infty$ converges to the unique common fixed point of S, T .*

Corollary 2. *Let X be a Banach space and $S, T : X \rightarrow X$ such that $\|Tx - Ty\| \leq h \max\{\|Sx - Sy\|, \frac{\|Sx-Tx\|+\|Sy-Ty\|}{2}, \frac{\|Sx-Ty\|+\|Sy-Tx\|}{2}\}$ and $T(X) \subseteq S(X)$. Assume S and T are weakly compatible. For any $x_o \in X$, the Jungck-Ishikawa iteration (9) $\{Sx_n\}_{n=1}^\infty$ converges to the unique common fixed point of S, T .*

Remark 3. (i) A weaker version of Corollary 1 is the main result of [13] where the convergence is to the coincidence point of S, T and S is assumed injective.

(ii) If $S = I_d$ in Corollary 1, then we have the main result of [2].

Corollary 3. *Let X be a Banach space and $S, T : X \rightarrow X$ such that $\|Tx - Ty\| \leq h \max\{\|Sx - Sy\|, \frac{\|Sx-Tx\|+\|Sy-Ty\|}{2}, \frac{\|Sx-Ty\|+\|Sy-Tx\|}{2}\}$ and $T(X) \subseteq S(X)$. Assume S and T are weakly compatible. For any $x_o \in X$, the Jungck-Mann iteration (8) $\{Sx_n\}_{n=1}^\infty$ converges to the unique common fixed point of S, T .*

Remark 4. If $S = I_d$, Theorem 3.3 gives the result of [10].

It is already shown in [1] and [2] that if $S(Y)$ or $T(Y)$ is a complete subspace of X , then maps satisfying the operators (14) has a unique coincidence point. Hence we have the following results.

Theorem 3. *Let X be a Banach space and $S, T : X \rightarrow X$ such that $\|Tx - Ty\| \leq h \max\{\|Sx - Sy\|, \frac{\|Sx - Tx\| + \|Sy - Ty\|}{2}, \|Sx - Ty\|, \|Sy - Tx\|\}$ and $T(X) \subseteq S(X)$. Assume S and T are weakly compatible. For any $x_o \in X$, the Jungck-multistep iteration (11) $\{Sx_n\}_{n=1}^\infty$ converges to the unique common fixed point of S, T .*

Since the Jungck-Noor, Jungck-Ishikawa and Jungck-Mann iterations are special cases of Jungck-multistep iteration, then we have the following consequences.

Corollary 4. *Let X be a Banach space and $S, T : X \rightarrow X$ such that $\|Tx - Ty\| \leq h \max\{\|Sx - Sy\|, \frac{\|Sx - Tx\| + \|Sy - Ty\|}{2}, \|Sx - Ty\|, \|Sy - Tx\|\}$ and $T(X) \subseteq S(X)$. Assume S and T are weakly compatible. For any $x_o \in X$, the Jungck-Noor iteration (10) $\{Sx_n\}_{n=1}^\infty$ converges to the unique common fixed point of S, T .*

Corollary 5. *Let X be a Banach space and $S, T : X \rightarrow X$ such that $\|Tx - Ty\| \leq h \max\{\|Sx - Sy\|, \frac{\|Sx - Tx\| + \|Sy - Ty\|}{2}, \|Sx - Ty\|, \|Sy - Tx\|\}$ and $T(X) \subseteq S(X)$. Assume S and T are weakly compatible. For any $x_o \in X$, the Jungck-Ishikawa iteration (2.4) $\{Sx_n\}_{n=1}^\infty$ converges to the unique common fixed point of S, T .*

Corollary 6. *Let X be a Banach space and $S, T : X \rightarrow X$ such that $\|Tx - Ty\| \leq h \max\{\|Sx - Sy\|, \frac{\|Sx - Tx\| + \|Sy - Ty\|}{2}, \|Sx - Ty\|, \|Sy - Tx\|\}$ and $T(X) \subseteq S(X)$. Assume S and T are weakly compatible. For any $x_o \in X$, the Jungck-Mann iteration (8) $\{Sx_n\}_{n=1}^\infty$ converges to the unique common fixed point of S, T .*

Acknowledgement. The research is supported by the African Mathematics Millennium Science Initiative (AMMSI). The first author is grateful to the Hampton University, Virginia, USA, for hospitality.

References

- [1] ABBAS M., JUNGCK G., Common fixed point results for noncommuting mappings without continuity in cone metric spaces, *J. Math. Anal. Appl.*, 341(2008), 416-420.
- [2] BERINDE V., On the convergence of Ishikawa iteration in the class of quasicontractive operators, *Acta Math. Univ. Comenianae*, LXXIII(1)(2004), 119-126.
- [3] CHATTERJEA S.K., Fixed point theorems, *Comptes rendus de l'Academic bulgare des Sciences*, 25(6)(1972), 727-730.

- [4] DAS K.M., NAIK K.V., Common fixed point theorems for commuting maps on metric spaces, *Proc. Amer. Math. Soc.*, 77(1979), 369-373.
- [5] ISHIKAWA S., Fixed points by a new iteration method, *Proc. Amer. Math. Soc.*, 149(1974), 147-150.
- [6] JUNGCK G., Commuting mappings and fixed points, *Amer. Math. Monthly*, 83(1976), 261-263.
- [7] KANNAN R., Some results on fixed points II, *Amer. Math. Monthly*, 76(1969), 405-408.
- [8] MANN W.R., Mean value methods in iteration, *Proc. Amer. Math. Soc.*, 4(1953), 506-510.
- [9] NOOR M.A., New approximation schemes for general variational inequalities, *Journal of Mathematical Analysis and Applications*, 251(1)(2004), 217-229.
- [10] OLALERU J.O., On the convergence of the Mann iteration in locally convex spaces, *Carpathian Journal of Mathematics*, 22(1-2)(2006), 115-120.
- [11] OLALERU J.O., On the equivalence of Picard, Mann and Ishikawa iterations for a class of quasi-contractive operators, *J. Nig. Assoc. Math. Phys.*, 11(2007), 51-56.
- [12] OLALERU J.O., A new approximation method for the common fixed points of two weakly compatible mappings, (*submitted*).
- [13] OLATINWO M.O., IMORU C.O., Some convergence results for the Jungck-Mann and Jungck-Ishikawa iteration process in the class of generalized Zamfirescu operators, *Acta Math. Univ. Comenianae*, 77(2)(2008), 299-304.
- [14] OLATINWO M.O., Some stability and strong convergence results for the Jungck-Ishikawa iteration process, *Creative Math. and Info.*, 17(2008), 33-42.
- [15] OLATINWO M.O., A generalization of some convergence results using the Jungck-Noor three step iteration process in arbitrary Banach space, *Fasc. Math.*, 40(2008), 37-43.
- [16] OSILIKE M.O., Stability results for Ishikawa fixed point iteration procedure, *Indian J. Pure Appl. Math.*, 26(10)(1995), 937-941.
- [17] RAFIQ A., On the equivalence of Mann and Ishikawa iteration methods with errors, *Math. Comm.*, 11(2006), 143-152.
- [18] RHOADES B.E., Comments on two fixed points iteration methods, *J. Math. Anal. Appl.*, 56(2)(1976), 741-750.
- [19] RHOADES B.E., SOLTUZ S.M., The equivalence between Mann-Ishikawa iterations and multi-step iteration, *Nonlinear Anal.*, 58(2004), 219-228.
- [20] POPESCU O., Picard iteration converges faster than Mann iteration for a class of quasi-contractive operators, *Math. Comm.*, 12(2007), 195-202.
- [21] SINGH S.L., On Common fixed points of commuting maps, *Math. Sem. Notes Kobe Univ.*, 5(1977), 131-134.
- [22] SINGH S.L., BHATNAGAR C., MISHRA S.N., Stability of Jungck-type iterative procedures, *International J. Math. and Math. Sc.*, 19(2005), 3035-3043.
- [23] ZHIQUN X., Remarks of equivalence among Picard, Mann and Ishikawa iterations in normed space, *Fixed Point Theory and Applications*, Vol. 2007, Article ID 61434, 5 pp.
- [24] ZAMFIRESCU T., Fixed point theorems in metric spaces, *Arch. Math. (Basel)*, 23(1972), 292-298.

JOHNSON O. OLALERU
MATHEMATICS DEPARTMENT
UNIVERSITY OF LAGOS
LAGOS, NIGERIA
e-mail: olaleru1@yahoo.co.uk or jolaleru@unilag.edu.ng

HUDSON AKEWE
MATHEMATICS DEPARTMENT
UNIVERSITY OF LAGOS
LAGOS, NIGERIA
e-mail: hudsonmolas@yahoo.com.

Received on 14.08.2009 and, in revised form, on 21.11.2009.