



<b>Title</b>	<b>Gender disparity of hepatocellular carcinoma: role of hepatitis B virus X protein and androgen receptor</b>
<b>Author(s)</b>	<b>Chiu, YT; Ng, IOL</b>
<b>Citation</b>	<b>Hong Kong Medical Journal, 2016, v. 22 n. 3, suppl 4, p. 43-47</b>
<b>Issued Date</b>	<b>2016</b>
<b>URL</b>	<b><a href="http://hdl.handle.net/10722/232160">http://hdl.handle.net/10722/232160</a></b>
<b>Rights</b>	<b>Hong Kong Medical Journal. Copyright © Hong Kong Academy of Medicine Press.; This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License.</b>

# Gender disparity of hepatocellular carcinoma: role of hepatitis B virus X protein and androgen receptor

YT Chiu, IOL Ng \*

## KEY MESSAGES

1. The natural COOH-terminal truncated HBx was more frequently detected in hepatocellular carcinoma (HCC) in male than female patients.
2. There was no significant clinicopathological correlation with the HBx status or remarkable changes in androgen receptor (AR) expression in the presence of different truncated HBx mutants compared with full-length HBx.
3. Lower expression of AR was associated with

more aggressive tumour phenotype and a higher metastatic potential of HCC.

Hong Kong Med J 2016;22(Suppl 4):S43-7  
RFCID project number: 12111092

YT Chiu, IOL Ng

Department of Pathology, The University of Hong Kong

\* Principal applicant and corresponding author: iolng@hku.hk

## Introduction

Hepatocellular carcinoma (HCC) is especially prevalent in Hong Kong due to prevalent hepatitis B virus (HBV) infection. HBV-associated HCC has a male predominance, with a male-to-female ratio of about 4-5:1. This is in contrast to HCV-associated HCC that has a male-to-female ratio of 1.5-2:1. HBV is a partially double-stranded DNA virus containing four overlapping open reading frames. Among them, the X gene is most frequently integrated into the host genome naturally. Naturally integrated X gene is frequently found to be deleted at the 3'-end, leading to COOH-terminal truncated HBx protein.

In our previous study, we showed that in HBV-positive patients with HCC, full-length HBx (HBxFL) was detected in all non-tumourous livers (n=51). However, HBxFL (HBx 1-154 amino acids) was found in only 53% of the tumours, whereas natural COOH-truncated HBx (HBx 1-130 amino acids) was found in the remaining 47%. The presence of natural COOH-truncated HBx significantly correlated with the presence of venous invasion ( $P < 0.01$ ), a hallmark of metastasis, suggesting COOH-truncation of HBx plays a significant role in enhancing cell invasiveness and metastasis in HCC.<sup>1</sup> In addition, HBx mutants with different COOH-truncated length cause HCC by altering intracellular distribution and consecutive modulation of the important proliferative STAT/SOCS signalling, indicating a role of various shorter HBx mutants in tumourigenesis.<sup>2</sup> The mislocalisation of HBx truncated mutants can be explained in part by the finding that HBx protein contains a functional nuclear export sequence from amino acids 89 to 100.<sup>3</sup> Therefore, the COOH-terminal truncated form of HBx has been demonstrated to play a tumourigenic role in HCC and its oncogenicity may vary according to the truncated length.

Among male HBV carriers, the risk of HCC development is significantly increased in those with higher androgen receptor (AR) activity. HBV infection may, therefore, particularly cooperate with male-specific AR signalling to accelerate hepatocarcinogenesis. Moreover, HBx has been shown to enhance AR transcriptional activity through GSK-3 $\beta$  kinase pathways.<sup>4</sup> However, no report has shown a correlation between AR signalling and COOH-terminal truncated HBx protein.

In this study, we hypothesised that HBx cooperated with male-specific AR signalling to enhance hepatocarcinogenesis, and natural HBx truncated mutants had different effects on this AR signalling. The prevalence of natural COOH-terminal truncated HBx mutants (HBx  $\Delta$ C1 and HBx  $<$  $\Delta$ C1) was slightly higher in male than female with HCC, but no significant clinicopathological association was observed with the expression of COOH-terminal truncated HBx mutants in HCCs. The COOH-terminal truncated HBx mutants were not associated with the mRNA expression of AR in the same HCC specimens, but the mRNA expression of AR showed a significant correlation with the metastatic potential, cellular differentiation and staging of HCC.

## Methods

This study was conducted from April 2013 to April 2014.

### Patient samples

A total of 99 pairs of human HCCs from patients who underwent liver resection for HCC between 1992 and 2001 at Queen Mary Hospital, Hong Kong, were selected. All these 99 patients (77 men and 22 women) aged 19 to 74 years were positive for serum hepatitis B surface antigen (HBsAg). All specimens

were snap-frozen in liquid nitrogen and kept at -80°C. Frozen sections were cut from HCC blocks separately and stained for histological examination to ensure a homogenous cell population of tissues.

**RNA extraction and reverse-transcriptase polymerase chain reaction**

Total RNA was isolated using TRIZOL reagent (Invitrogen, Carlsbad [CA], USA) following the manufacturer’s instructions. cDNA was synthesised by using Gold RNA PCR Core Kit (Applied Biosystems, Foster City [CA], USA) following the manufacturer’s instructions.

**Semi-quantitative PCR**

For semi-quantitative PCR, the reaction was set up using AmpliTaqGOLD enzyme (Applied Biosystems, Carlsbad [CA], USA) and carried out with Takara PCR Thermal Cycler. PCR products were detected by 1.5% ethidium bromide-agarose gel electrophoresis at 100V for 15 minutes and visualised on a gel documentation system. The primers for PCR were as follows: HBx ATG: 5’-CCC AAG CTT ATG GCT GCT AGG CTG TGC T-3’; HBx full length (HBxFL): 5’-CGA ATT CTT AGG CAG AGG TGA AAA AGT TG-’; HBx ΔC1: 5’-CGA ATT CTT ACT TTA ATC TAA TCT CCT CCC C-3’; HBx <ΔC1: 5’-CTT ATG TAA GAC CTT GGG CAA CAT-3’

**Quantitative real-time PCR**

The reaction was set up using SYBR Green Real-Time PCR Master Mixes (Applied Biosystems) and was carried out with Applied Biosystems 7900 HT Fast Real-Time PCR System (Applied Biosystems) according to the manufacturer’s protocol. All reactions were performed in triplicate and the expression of targeted gene relative to hypoxanthine guanine phosphoribosyl transferase was determined using 2<sup>-ΔCT</sup> method. The primer sequences for real-time PCR were as follows: AR F: 5’-CTC ACC AAG CTC CTG GAC TC-3’; AR R: 5’-GAA AGG ATC TTG GGC ACT TG-3’; HPRT F:5’-CTTTGCTGACCTGCTGGATT-3’; HPRT R:5’-CTGCATTGTTTTGCCAGTGT -3’.

**Results**

**The prevalence of natural COOH-terminal truncated HBx mutants (HBx ΔC1 and HBx <ΔC1) was slightly higher in HCCs from male than female patients**

The HCCs were screened for the presence of COOH-terminal truncated HBx mutants, ΔC1 (HBx 1-130 amino acids) and <ΔC1 (HBx 1-95 amino acids). The expression pattern of the truncated HBx mutants between male and female patients was compared. Among the 99 HCC samples, the natural COOH-terminal truncated HBx mutants (HBx ΔC1

and HBx <ΔC1) were more frequently detected in HCCs in males than females (44% vs 37%, Fig 2). The frequency of HBx ΔC1 in HCCs in males was similar to that in females (30% vs 32%), but a more remarkable difference was observed for HBx <ΔC1 mutants (14% vs 5%, Fig 1). These indicated that the COOH-terminal truncated HBx mutants that are shorter than ΔC1 may be found more frequently in male HCCs and may be more associated than HBx ΔC1 with the male predominance in HCC.

**The expression of COOH-terminal truncated HBx mutants in human HCCs showed no significant clinicopathological or survival correlation**

To examine whether the presence of natural COOH-terminal truncated HBx mutants was associated with a particular tumour biological behaviour and consequent prognosis, the expression of full-length HBx and truncated HBx mutants was correlated with different clinicopathological features and survival rates in the 99 HCC patients. The presence of COOH-terminal truncated HBx mutants (ie the group with either HBx ΔC1 or HBx<ΔC1) did not show any significant correlation with the clinicopathological features (Fig 2a). In addition, there was no significant





FIG 1. The frequency of presence of HBx FL and COOH-terminal truncated HBxΔC1 and Δ<C1 in male and female HBV-associated HCCs (n=99). Presence of HBx FL and COOH-terminal truncated HBx mutants was detected by conventional PCR. The frequency of COOH-truncated mutants was higher in male than female HCC patients.



difference between the presence of HBx mutants and the disease free or overall survival (Fig 2b). This demonstrated that the presence of COOH-terminal truncated HBx mutants might not be associated with tumour biological behaviour.


**The COOH-terminal truncated HBx mutants had no significant correlation with the mRNA expression of AR in the same HCC specimens**

To investigate the effect of different COOH-terminal truncated HBx mutants on the mRNA expression level of AR, the expression level of AR was examined and correlated with the presence of COOH-terminal truncated HBx mutants in the 99 HCCs. There was no significant difference between the mRNA levels of AR and the presence of COOH-terminal truncated HBx mutants or the full-length HBx (Fig 3a). Although there was a trend of association between the presence of HBx $\Delta C1$  and downregulation of AR (P=0.4358), the mRNA levels of AR were not

significantly different upon further stratification of the HBx mutants into HBx $\Delta C1$  and HBx $<\Delta C1$  (Fig 3b). These indicated that the COOH-terminal truncated HBx mutants, either HBx  $\Delta C1$  or HBx $<\Delta C1$ , may not regulate mRNA expression of AR.

**The mRNA expression of AR showed a significant correlation with the metastatic potential, cellular differentiation and staging of HCC**

To evaluate the association of the mRNA expression of AR with a particular tumour biological behaviour and prognostic significance in HCC, the mRNA levels of AR correlated with different clinicopathological features. Lower expression of AR in the HCC was associated with the presence of tumour venous invasion (P=0.0213), tumour microsatellite formation (P=0.0251), poorer tumour cellular differentiation (P=0.0002) and higher tumour stage (P=0.0073) [Fig 3c]. This correlation



study demonstrated that lower expression of AR was associated with a higher metastatic potential and more aggressive tumour phenotype in HCC.

## Discussion

The frequency of natural COOH-terminal truncated HBx (HBx  $\Delta$ C1 and HBx< $\Delta$ C1) was higher in HCCs of male than female patients (44% vs 37%). More importantly, when the mutants were stratified into the group with HBx< $\Delta$ C1 only and the other one with HBx FL or HBx  $\Delta$ C1, the frequency of HBx mutants in male and female HCCs (14% vs 5%) was even more obvious. With a higher frequency of HBx< $\Delta$ C1 in HCCs of males and the differential regulatory roles of shorter truncated HBx mutants in HCC due to nuclear mislocalisation,<sup>2</sup> these imply that HBx<C1 may play a role in hepatocarcinogenesis in male HCC patients and may contribute in part to the male predominance of HCC. Nonetheless, the underlying mechanisms require further investigation. In addition, the expression status of HBx in the HCCs showed no significant correlation with any clinicopathological features or patient survival. This might have been a result of technical limitations in examining HBx gene truncation, where the co-existence of HBx FL and COOH-terminal truncated mutants in the same specimen cannot be detected by conventional PCR. The HCC tumours with HBx FL detected might also harbour HBx $\Delta$ C1 and/or HBx< $\Delta$ C1, and this may influence the comparison among groups of various parameters.

Full-length HBx protein has been shown to enhance AR transcriptional activity through modulation of c-Src and GSK-3 $\beta$  kinase pathways.<sup>5</sup> In addition, HBx protein enhances AR-responsive gene expression depending on androgen level.<sup>6</sup> Therefore, the regulatory roles of full-length HBx on AR activity are well demonstrated. In the present study, we examined the consequence of COOH-terminal truncated HBx on AR transactivation. The results showed no significant correlation between the presence of X truncation and mRNA expression of AR in HCCs when compared with the full-length HBx, suggesting that COOH-truncated HBx mutants may not exert a more potent activation on AR than the full-length HBx. Nevertheless, a lower expression of AR was associated with more aggressive tumour behaviour. This finding was consistent with that in other studies that report hepatic AR suppression of HCC metastasis through modulation of cell migration and anoikis,<sup>7</sup> enhanced cell adhesion and decreased cell migration via modulating  $\beta$ 1-integrin-AKT signalling in HCC cells.<sup>8</sup> In contrast to our results, several reports found that the activation and overexpression of AR promoted cell invasion and migration in HCC and tumour staging.<sup>9</sup> Therefore, the potential roles of AR in hepatocarcinogenesis remain elusive and warrant further investigation.

## Conclusion

The HBx truncated mutants are more frequent in HCCs of males than females, and this may implicate a role of HBx truncated mutants in the male predominance of HCC. Nonetheless, the correlation study of the presence of HBxC1 or HBx< $\Delta$ C1 and the different clinicopathological features of HCC patients and the mRNA levels of AR implied that HBx truncated mutants alone may not exert a significant effect on the tumour biological behaviour or induce a more potent transactivation of AR than HBxFL. AR may play a role in hepatocarcinogenesis, in which lower expression of AR is associated with more aggressive tumour phenotype and a higher metastatic potential of HCC.

## Acknowledgement

This study was supported by the Research Fund for the Control of Infectious Diseases, Food and Health Bureau, Hong Kong SAR Government (#12111092).

## References

1. Sze KM, Chu GK, Lee JM, Ng IO. C-terminal truncated hepatitis B virus x protein is associated with metastasis and enhances invasiveness by C-Jun/matrix metalloproteinase protein 10 activation in hepatocellular carcinoma. *Hepatology* 2013;57:131-9.
2. Bock CT, Toan NL, Koeberlein B, et al. Subcellular mislocalization of mutant hepatitis B X proteins contributes to modulation of STAT/SOCS signaling in hepatocellular carcinoma. *Intervirology* 2008;51:432-43.
3. Forgues M, Marrogi AJ, Spillare EA, et al. Interaction of the hepatitis B virus X protein with the Crm1-dependent nuclear export pathway. *J Biol Chem* 2001;276:22797-803.
4. Yang WJ, Chang CJ, Yeh SH, et al. Hepatitis B virus X protein enhances the transcriptional activity of the androgen receptor through c-Src and glycogen synthase kinase-3 $\beta$  kinase pathways. *Hepatology* 2009;49:1515-24.
5. Wu MH, Ma WL, Hsu CL, et al. Androgen receptor promotes hepatitis B virus-induced hepatocarcinogenesis through modulation of hepatitis B virus RNA transcription. *Sci Transl Med* 2010;2:32ra35.
6. Chiu CM, Yeh SH, Chen PJ, et al. Hepatitis B virus X protein enhances androgen receptor-responsive gene expression depending on androgen level. *Proc Natl Acad Sci U S A* 2007;104:2571-8.
7. Ma WL, Hsu CL, Yeh CC, et al. Hepatic androgen receptor suppresses hepatocellular carcinoma metastasis through modulation of cell migration and anoikis. *Hepatology* 2012;56:176-85.
8. Ma WL, Jeng LB, Lai HC, Liao PY, Chang C. Androgen receptor enhances cell adhesion and decreases cell migration via modulating  $\beta$ 1-integrin-AKT signaling in hepatocellular carcinoma cells. *Cancer Lett* 2014;351:64-71.
9. Zhang Y, Shen Y, Cao B, Yan A, Ji H. Elevated expression levels of androgen receptors and matrix metalloproteinase-2 and -9 in 30 cases of hepatocellular carcinoma compared with adjacent tissues as predictors of cancer invasion and staging. *Exp Ther Med* 2015;9:905-8.