

Kessar, R. & Linckelmann, M. (2006). On blocks of strongly p-solvable groups. Archiv der Mathematik, 87(6), 481 - 487.



**CITY UNIVERSITY  
LONDON**

[City Research Online](#)

**Original citation:** Kessar, R. & Linckelmann, M. (2006). On blocks of strongly p-solvable groups. Archiv der Mathematik, 87(6), 481 - 487.

**Permanent City Research Online URL:** <http://openaccess.city.ac.uk/1902/>

### Copyright & reuse

City University London has developed City Research Online so that its users may access the research outputs of City University London's staff. Copyright © and Moral Rights for this paper are retained by the individual author(s) and/ or other copyright holders. All material in City Research Online is checked for eligibility for copyright before being made available in the live archive. URLs from City Research Online may be freely distributed and linked to from other web pages.

### Versions of research

The version in City Research Online may differ from the final published version. Users are advised to check the Permanent City Research Online URL above for the status of the paper.

### Enquiries

If you have any enquiries about any aspect of City Research Online, or if you wish to make contact with the author(s) of this paper, please email the team at [publications@city.ac.uk](mailto:publications@city.ac.uk).

# ON BLOCKS OF STRONGLY $P$ -SOLVABLE GROUPS

RADHA KESSAR, MARKUS LINCKELMANN

ABSTRACT. We prove that a block of a finite strongly  $p$ -solvable group  $G$  with defect group  $P$  is Morita equivalent to its corresponding block of  $N_G(Z(J(P)))$  via a bimodule with endo-permutation source.

Let  $p$  be a prime. Following [8, Ch. 6, §5] a finite group  $G$  is called *strongly  $p$ -solvable* if  $G$  is  $p$ -solvable and either  $p \geq 5$  or  $p = 3$  and  $Sl_2(3)$  is not involved in  $G$ . If  $G$  is a strongly  $p$ -solvable finite group and  $P$  a Sylow- $p$ -subgroup of  $G$ , then  $G$  is  $p$ -constrained and  $p$ -stable (cf. [8, Ch. 8, §1]), and hence  $G = N_G(Z(J(P)))O_{p'}(G)$  by a theorem of Glauberman (cf. [8, Ch. 8, Theorem 2.11]). The purpose of this paper is to show that this theorem, which is used in the proof of the  $p$ -nilpotency theorem of Glauberman and Thompson, has a generalisation to blocks of strongly  $p$ -solvable groups. We denote by  $\mathcal{O}$  a complete discrete valuation ring having an algebraically closed residue field  $k = \mathcal{O}/J(\mathcal{O})$  of characteristic  $p$ . We state our Theorem and explain the notation below.

**Theorem.** *Let  $G$  be a strongly  $p$ -solvable finite group, let  $b$  be a block of  $\mathcal{O}G$  and let  $P$  a defect group of  $b$ . Set  $N = N_G(Z(J(P)))$  and denote by  $c$  the Brauer correspondent of  $b$  in  $\mathcal{O}N$ . Then the Brauer categories  $\mathcal{F}_{G,b}$  and  $\mathcal{F}_{N,c}$  are equal and there is an indecomposable  $\mathcal{O}Gb$ - $\mathcal{O}Nc$ -bimodule  $M$  with the following properties.*

(i) *As  $\mathcal{O}(G \times N)$ -module,  $M$  has vertex  $\Delta P = \{(u, u)\}_{u \in P}$  and an endo-permutation  $\mathcal{O}\Delta P$ -module  $W$  as source.*

(ii) *The bimodule  $M$  and its  $\mathcal{O}$ -dual  $M^*$  induce a Morita equivalence between the block algebras  $\mathcal{O}Gb$  and  $\mathcal{O}Nc$ .*

The structure of the source algebras of blocks of  $p$ -solvable finite groups has been completely determined by Puig. Once the stated equality of Brauer categories is established (using Glauberman's aforementioned theorem), the rest the proof of the Theorem consists of showing that the reduction techniques used in Puig's work "commute" with taking normalisers of  $Z(J(P))$ .

---

1991 *Mathematics Subject Classification.* 20C20.

With the notation of the Theorem, the block  $c$  of  $\mathcal{O}N$  is the unique block satisfying  $\text{Br}_P(c) = \text{Br}_P(b)$ ; this makes sense as  $C_G(P) \subseteq N$ . Fix a block  $e_P$  of  $kC_G(P)$  such that  $\text{Br}_P(b)e_P = e_P$ ; that is,  $(P, e_P)$  is a maximal  $b$ -Brauer pair and a maximal  $c$ -Brauer pair.

By [1], for any subgroup  $Q$  of  $P$  there is a unique block  $e_Q$  of  $kC_G(Q)$  such that  $(Q, e_Q) \subseteq (P, e_P)$ . Denote by  $\mathcal{F}_{G,b}$  the category whose objects are the subgroups of  $P$  and whose set of morphisms from a subgroup  $Q$  of  $P$  to another subgroup  $R$  of  $P$  is the set of group homomorphisms  $\varphi : Q \rightarrow R$  for which there exists an element  $x \in G$  satisfying  $\varphi(u) = xux^{-1}$  for all  $u \in Q$  and  ${}^x(Q, e_Q) \subseteq (R, e_R)$ . Thus the automorphism group of a subgroup  $Q$  of  $P$  as object of the category  $\mathcal{F}_{G,b}$  is canonically isomorphic to  $N_G(Q, e_Q)/C_G(Q)$ . By Alperin's fusion theorem, the category  $\mathcal{F}_{G,b}$  is completely determined by the structure of  $P$  and the groups  $N_G(Q, e_Q)/C_G(Q)$  where either  $Q = P$  or  $(Q, e_Q)$  is an essential  $b$ -Brauer pair (cf. [18, §48]). Note that  $O_p(G) \subseteq Q$  whenever the pair  $(Q, e_Q)$  is essential. The category  $\mathcal{F}_{N,c}$  is defined similarly. Since  $O_p(N)$  contains  $Z(J(P))$  and  $C_N(Q) = C_G(Q)$  for any subgroup  $Q$  of  $P$  containing  $Z(J(P))$  we have an inclusion of categories  $\mathcal{F}_{N,c} \subseteq \mathcal{F}_{G,b}$ .

If  $b$  is the principal block of  $\mathcal{O}G$ , then  $e_Q$  is the principal block of  $kC_G(Q)$ , for any subgroup  $Q$  of  $P$ . Thus the above condition  ${}^x(Q, e_Q) \subseteq (R, e_R)$  is equivalent to  ${}^xQ \subseteq R$ . Therefore, if  $b$  is the principal block of  $\mathcal{O}G$ , we write  $\mathcal{F}_G$  instead of  $\mathcal{F}_{G,b}$ .

In general, the definition of  $\mathcal{F}_{G,b}$  depends on the choice of a maximal  $b$ -Brauer pair, but since all maximal  $b$ -Brauer pairs are  $G$ -conjugate, it is easy to see that  $\mathcal{F}_{G,b}$  is unique up to isomorphism of categories. Note that we always have  $\mathcal{F}_P \subseteq \mathcal{F}_{G,b}$ .

Let  $P$  be a finite  $p$ -group. An  $\mathcal{O}$ -free  $\mathcal{O}P$ -module  $U$  is called an *endo-permutation module* if  $\text{End}_{\mathcal{O}}(U)$  has a  $P$ -stable  $\mathcal{O}$ -basis with respect to the action of  $P$  by conjugation. This concept is due to Dade [5, 6]. It is shown in [5], that an endo-permutation  $\mathcal{O}P$ -module  $U$  has at most one isomorphism class of indecomposable direct summands of vertex  $P$ . Since the restriction to any subgroup  $Q$  of  $P$  of an endo-permutation  $\mathcal{O}P$ -module  $U$  is an endo-permutation  $\mathcal{O}Q$ -module, it follows that if  $U$  has an indecomposable direct summand with vertex  $P$ , then  $\text{Res}_Q^P(U)$  has exactly one isomorphism class of indecomposable direct summands with vertex  $Q$  for any subgroup  $Q$  of  $P$ . We will say that  $U$  is  *$G$ -stable* if the indecomposable summands with vertex  $Q$  of  $\text{Res}_Q^P(U)$  are isomorphic to those of  $\text{Res}_{\varphi}(U)$  for any subgroup  $Q$  of  $P$  and any group homomorphism  $\varphi : Q \rightarrow P$  for which there is an element  $x \in G$  satisfying  $\varphi(u) = xux^{-1}$  for all  $u \in Q$ . Here  $\text{Res}_{\varphi}(U)$  is the endo-permutation  $\mathcal{O}Q$ -module obtained from restricting  $U$  via  $\varphi$ ; that is, with  $u \in Q$  acting as  $\varphi(u)$  on  $U$ . In the terminology of [17] this is the same as saying that the interior  $P$ -algebra  $\text{End}_{\mathcal{O}}(U)$  contains  $G$ -fusion. We state without proof two obvious Lemmas needed in the proof of the Theorem.

**Lemma 1.** *Let  $G, H$  be finite groups and let  $P$  be a  $p$ -subgroup of both  $G$  and  $H$ . Let  $U, V$  be endo-permutation  $\mathcal{O}P$ -modules having direct summands of vertex  $P$ .*

(i) *If  $U$  is  $G$ -stable, then  $\text{Res}_Q^P(U)$  is  $G$ -stable for any subgroup  $Q$  of  $P$ .*

(ii) If  $U$  is  $G$ -stable and  $V$  is  $H$ -stable, then the  $\mathcal{O}(P \times P)$ -module  $U \otimes_{\mathcal{O}} V$  is  $G \times H$ -stable.

**Lemma 2.** *Let  $G$  be a finite group and let  $P$  be a  $p$ -subgroup of  $G$ . Let  $U$  be an endo-permutation  $\mathcal{O}P$ -module having an indecomposable direct summand  $V$  of vertex  $P$ . If  $U$  is  $G$ -stable then  $V$  is  $G$ -stable.*

We quote the following well-known result on the local structure of certain blocks of  $p$ -constrained groups; recall that a finite group  $G$  is  $p$ -constrained if  $C_G(Q) \subseteq O_{p',p}(G)$ , where  $Q$  is a Sylow- $p$ -subgroup of  $O_{p',p}(G)$ .

**Proposition 3.** ([9, 3.1, 3.4]) *Let  $G$  be a finite  $p$ -constrained group and let  $b$  be a  $G$ -stable block of  $\mathcal{O}O_{p'}(G)$ . Then  $b$  is still a block of  $\mathcal{O}G$ , any Sylow- $p$ -subgroup  $P$  of  $G$  is a defect group of  $b$  as block of  $\mathcal{O}G$ , and for any  $p$ -subgroup  $Q$  of  $G$ , the central idempotent  $\text{Br}_Q(b)$  is a block of  $kC_G(Q)$ .*

The last statement in Proposition 3 means that  $b$  is of principal type, which in turn implies that  $\mathcal{F}_{G,b} = \mathcal{F}_G$ . We will need Puig's algebra theoretic formulation of Fong's reduction [7], as outlined in [14].

**Proposition 4.** (cf. [9, 3.1(i)]) *Let  $G$  be a finite  $p$ -solvable group and let  $b$  be a block of  $\mathcal{O}G$ . There is a subgroup  $H$  of  $G$  containing  $O_{p'}(G)$  and an  $H$ -stable block  $e$  of  $\mathcal{O}O_{p'}(H)$  such that*

$$\mathcal{O}Gb \cong \text{Ind}_H^G(\mathcal{O}He)$$

*as interior  $G$ -algebras. Moreover, every Sylow- $p$ -subgroup of  $H$  is then a defect group of  $e$  and  $b$  as blocks of  $\mathcal{O}H$  and  $\mathcal{O}G$ , respectively.*

**Proposition 5.** (cf. [10, 4.4]) *Let  $G$  be a finite  $p$ -solvable group, let  $b$  be a  $G$ -stable block of  $\mathcal{O}O_{p'}(G)$  and set  $S = \mathcal{O}O_{p'}(G)b$ . Denote by  $\hat{G}$  the  $\mathcal{O}^\times$ -group opposite to that defined by the action of  $G$  on  $S$ , and set  $\hat{L} = \hat{G}/N$ , where any element  $y$  of  $N$  is identified to its canonical image  $(y, yb)$  in  $\hat{G}$ . There is a unique algebra isomorphism*

$$\mathcal{O}Gb \cong S \otimes_{\mathcal{O}} \mathcal{O}_* \hat{L}$$

*mapping  $xb$  to  $s \otimes \overline{(x, s)}$ , where  $x \in G$  and  $s \in S^\times$  such that  $(x, s) \in \hat{G}$  and where  $\overline{(x, s)}$  is the canonical image of  $(x, s)$  in  $\hat{L}$ .*

We show next, that this reduction is compatible with taking normalisers of  $Z(J(P))$ .

**Proposition 6.** *Let  $G$  be a finite  $p$ -solvable group, let  $b$  be a block of  $\mathcal{O}G$ , let  $H$  be a subgroup of  $G$  containing  $O_{p'}(G)$  and let  $e$  be an  $H$ -stable block of  $\mathcal{O}O_{p'}(H)$  such that  $\mathcal{O}Gb \cong \text{Ind}_H^G(\mathcal{O}He)$ . Let  $P$  be a Sylow- $p$ -subgroup of  $H$  and let  $Z$  be a subgroup of  $P$  such that  $N_G(P)$  is contained in  $N_G(Z)$ . Denote by  $c$  and  $f$  the blocks of  $\mathcal{O}N_G(Z)$  and  $\mathcal{O}N_H(Z)$  which are the Brauer correspondents of  $b$  and  $e$ , respectively. Then  $f$  is an  $N_H(Z)$ -stable block of  $\mathcal{O}O_{p'}(N_H(Z))$  and we have an isomorphism of interior  $N_G(Z)$ -algebras*

$$\mathcal{O}N_G(Z)c \cong \text{Ind}_{N_H(Z)}^{N_G(Z)}(\mathcal{O}N_H(Z)f) .$$

*Proof.* By [10, 4.2], we have  $O_{p'}(N_H(Z)) = O_{p'}(C_H(Z)) = O_{p'}(H) \cap C_H(Z)$ . By Proposition 3,  $\text{Br}_Z(e)$  is a block of  $kC_H(Z)$ . Since  $e$  is  $H$ -stable and every block of  $kN_H(Z)$  is contained in  $kC_H(Z)$ , it follows that  $\text{Br}_Z(e)$  is a block of  $kN_H(Z)$ . As  $e \in \mathcal{O}O_{p'}(H)$ , it follows that  $\text{Br}_Z(e) \in kO_{p'}(N_H(Z))$ . Thus  $\text{Br}_Z(e)$  lifts to a unique  $N_H(Z)$ -stable block  $f$  of  $\mathcal{O}O_{p'}(N_H(Z))$ , which is the Brauer correspondent of  $e$ . If  $x \in N_G(Z) - N_H(Z)$ , then  ${}^x\text{Br}_Z(e)\text{Br}_Z(e) = \text{Br}_Z({}^xe \cdot e) = 0$ , because  ${}^xe \cdot e = 0$  as  $\mathcal{O}Gb \cong \text{Ind}_H^G(\mathcal{O}He)$ . Thus  ${}^xf \cdot f = 0$ . Therefore,  $\text{Ind}_{N_H(Z)}^{N_G(Z)}(\mathcal{O}N_H(Z)f) \cong \mathcal{O}N_G(Z)\text{Tr}_{N_H(Z)}^{N_G(Z)}(f)$ ; in particular, the algebras  $\mathcal{O}N_H(Z)f$  and  $\mathcal{O}N_G(Z)\text{Tr}_{N_H(Z)}^{N_G(Z)}(f)$  have isomorphic centers. As  $f$  is primitive in  $Z(\mathcal{O}N_H(Z))$ , it follows that  $\text{Tr}_{N_H(Z)}^{N_G(Z)}(f)$  is primitive in  $Z(\mathcal{O}N_G(Z))$ . But then  $c = \text{Tr}_{N_H(Z)}^{N_G(Z)}(f)$ , which implies the result.  $\square$

*Proof of the Theorem.* By Proposition 4, there is a subgroup  $H$  of  $G$  containing  $O_{p'}(G)$  and an  $H$ -stable block  $e$  of  $\mathcal{O}O_{p'}(H)$  such that  $\mathcal{O}Gb \cong \text{Ind}_H^G(\mathcal{O}He)$ . By general properties of algebra induction (cf. [15]), the blocks  $\mathcal{O}Gb$  and  $\mathcal{O}He$  have isomorphic source algebras. In view of Proposition 6, we may therefore assume that  $G = H$  and  $b = e$ , and then  $c$  is an  $N$ -stable block of  $\mathcal{O}O_{p'}(N)$ . Set  $S = \mathcal{O}O_{p'}(G)b$ . Then  $S$  is a matrix algebra of rank  $n^2$  prime to  $p$ , on which  $G$  acts. We consider the subgroup

$$\hat{G} = \{(x, s) | x \in G, s \in S^\times \text{ such that } {}^xt = sts^{-1} \text{ for all } t \in S\}$$

of  $G \times S^\times$ , endowed with the group homomorphism  $\mathcal{O}^\times \rightarrow \hat{G}$  mapping  $\lambda \in \mathcal{O}^\times$  to  $(1_G, \lambda^{-1}1_S)$ ; that is,  $\hat{G}$  is the  $\mathcal{O}^\times$ -group opposite to that defined by the action of  $G$  on  $S$  (cf. [10] for more details on this terminology). By the Skolem-Noether theorem, every algebra automorphism of  $S$  is inner, and thus, for any  $x \in G$ , there is  $s_x \in S^\times$  such that  $(x, s_x) \in \hat{G}$ . Equivalently, the canonical projection  $\hat{G} \rightarrow G$  onto the first component is surjective. We may choose the  $s_x$  in such a way that  $\det(s_x) = 1$  for all  $x \in G$ , and then  $s_x s_y = \lambda_{x,y} s_{xy}$  for any  $x, y \in G$  and some  $n^{\text{th}}$  roots of unity  $\lambda_{x,y}$  in  $\mathcal{O}^\times$  (all this is well-known; see e. g. [10]). Note that this means that the 2-cocycle  $\lambda$  representing the central extension  $\hat{G}$  of  $G$  by  $\mathcal{O}^\times$  has actually values in the canonical image of  $k^\times$  in  $\mathcal{O}^\times$  (this will be relevant when we apply [16, (e)] below, which is formulated for  $k^\times$ -groups). There is a unique group homomorphism  $\sigma : P \rightarrow S^\times$  such that  $\det(\sigma(y)) = 1$  for all  $y \in P$  and such that  $(y, \sigma(y)) \in \hat{G}$ ; in other words, the action of  $P$  on  $S$  “lifts

canonically". Since  $S$  is a matrix algebra, we may write  $S = \text{End}_{\mathcal{O}}(U)$  for some free  $\mathcal{O}$ -module  $U$  of rank  $n$ . Then  $U$  becomes an  $\mathcal{O}P$ -module via the group homomorphism  $\sigma$ . In fact,  $U$  becomes an endo-permutation  $\mathcal{O}P$ -module, because  $S$  is a direct summand of the group algebra  $\mathcal{O}O_{p'}(G)$ , which has  $O_{p'}(G)$  as  $P$ -stable  $\mathcal{O}$ -basis, and thus  $S$  has a  $P$ -stable  $\mathcal{O}$ -basis.

We observe next that  $U$  is  $G$ -stable. If  $Q$  is a subgroup of  $P$  and  $x \in G$  such that  ${}^xQ \subseteq P$ , denote by  $\varphi : Q \cong {}^xQ$  the isomorphism sending  $y \in Q$  to  ${}^xy$ . By the uniqueness of  $\sigma$  we have  $\sigma({}^xy) = {}^x\sigma(y)$  for all  $y \in Q$ ; indeed, both sides have determinant 1 and act by conjugation as  ${}^xy$  on  $S$ . Choose  $s \in S^\times$  such that  $(x, s) \in \hat{G}$ , or equivalently, such that  ${}^xt = sts^{-1}$  for all  $t \in S$ . Then the map sending  $u \in U$  to  $s(u)$  is an isomorphism  $\text{Res}_\varphi(U) \cong \text{Res}_Q^P(U)$ ; in particular,  $U$  is  $G$ -stable.

We identify  $O_{p'}(G)$  to its canonical image  $\{(y, yb) \mid y \in O_{p'}(G)\}$  in  $\hat{G}$ , and set  $\hat{L} = \hat{G}/O_{p'}(G)$ . Thus  $\hat{L}$  is a central  $\mathcal{O}^\times$ -extension of the group  $L = G/O_{p'}(G)$ .

Analogously, set  $T = \mathcal{O}O_{p'}(N)c$ . This is a matrix algebra over  $\mathcal{O}$  of rank  $m^2$  prime to  $p$  on which  $N$  acts. Set

$$\tilde{N} = \{(y, t) \in N \times T^\times \mid {}^y(t') = tt't^{-1} \text{ for all } t' \in T\}$$

endowed with the group homomorphism  $\mathcal{O}^\times \rightarrow \tilde{N}$  sending  $\lambda \in \mathcal{O}^\times$  to  $(1_N, \lambda^{-1}1_T)$ . For any  $y \in N$ , denote by  $t_y$  an element in  $T^\times$  such that  $(y, t_y) \in \tilde{N}$  and such that  $\det(t_y) = 1$ . Denote by  $\tau : P \rightarrow T^\times$  the unique group homomorphism such that  $(u, \tau(u)) \in \tilde{N}$  and such that  $\det(\tau(u)) = 1$  for all  $u \in P$ . Write  $T = \text{End}_{\mathcal{O}}(V)$  for some free  $\mathcal{O}$ -module  $V$  of rank  $m$ . As before,  $V$  becomes an endopermutation  $\mathcal{O}P$ -module via  $\tau$ , and  $V$  is  $H$ -stable by the uniqueness of  $\tau$ .

Similarly, identify  $O_{p'}(N)$  to its canonical image in  $\tilde{N}$  and set  $\tilde{L} = \tilde{N}/O_{p'}(N)$ .

Let  $W$  be an indecomposable direct summand of  $U^* \otimes_{\mathcal{O}} V$ , viewed as  $\mathcal{O}\Delta P$ -module, such that  $W$  has vertex  $\Delta P$ . It follows from Lemma 1 and Lemma 2 that  $W$  is  $G \times H$ -stable.

By Glauberman's theorem [8, Ch. 8, Theorem 2.11], we have  $G = NO_{p'}(G)$ . Since, by Proposition 4, both  $b$  and  $c$  are of principal type, we have the equality of the Brauer categories  $\mathcal{F}_{G,b} = \mathcal{F}_{N,c}$ . By [10, 4.2], we have  $O_{p'}(G) \cap N = O_{p'}(N)$ . Thus the inclusion  $N \subseteq G$  induces a group isomorphism  $N/O_{p'}(N) \cong G/O_{p'}(G) = L$ . Identify  $N/O_{p'}(N)$  to  $L$  through this isomorphism. The crucial step is to show that there is an isomorphism of  $\mathcal{O}^\times$ -groups

$$\hat{L} \cong \tilde{L}$$

which induces the identity on the canonical quotients  $L$  of  $\hat{L}$ ,  $\tilde{L}$ , and which preserves the canonical images of  $P$  in  $\hat{L}$  and  $\tilde{L}$  elementwise. Before we prove the existence of such a group isomorphism, let us show how this concludes the proof of the Theorem.

Identifying  $\mathcal{O}_* \tilde{L}$  and  $\mathcal{O}_* \hat{L}$  through the algebra isomorphism induced by this group isomorphism, together with Proposition 5, shows that we have isomorphisms of interior  $P$ -algebras

$$\mathcal{O}Gb \cong S \otimes_{\mathcal{O}} \mathcal{O}_* \hat{L} \quad \text{and} \quad \mathcal{O}Nc \cong T \otimes_{\mathcal{O}} \mathcal{O}_* \hat{L}.$$

If we consider  $U \otimes_k V^*$  as  $S$ - $T$ -bimodule, then through the above algebra isomorphisms, the bimodule  $M = U \otimes_{\mathcal{O}} \mathcal{O}_* \hat{L} \otimes_{\mathcal{O}} V^*$  has vertex  $\Delta P$  and  $W$  as source. Clearly  $M$  and its dual induce a Morita equivalence between  $\mathcal{O}Gb$  and  $\mathcal{O}Nc$ .

It remains to prove the existence of a group isomorphism  $\hat{L} \cong \tilde{L}$  as claimed. By the above remarks the central extensions  $\hat{L}, \tilde{L}$  of  $L$  by  $\mathcal{O}^\times$  can be represented by 2-cocycles with values in the canonical image of  $k^\times$  in  $\mathcal{O}^\times$ , and hence we may assume that  $\mathcal{O} = k$ . We will show, that there is an isomorphism of  $k^\times$ -groups  $\tilde{N} \cong \hat{N}$  preserving the canonical images of  $P$  and of  $O_{p'}(N)$  and inducing the identity on the canonical quotients  $N$  of  $\hat{N}, \tilde{N}$ . Since  $O_{p'}(N)$  and  $Z = Z(J(P))$  are normal subgroups of coprime order in  $N$ , they commute, and thus the Brauer construction with respect to  $Z$  applied to  $S$  yields  $S(Z) = T$ ; in particular,  $\text{Br}_Z(b) = c$ . By Puig's version [16, (e)] of Dade's splitting theorem on fusion for endo-permutation modules (applied to  $Z$  instead of  $P$ ), there is a group homomorphism  $f : N_{S^\times}(\sigma(Z)) \rightarrow T^\times$  which extends the group homomorphism  $(S^Z)^\times \rightarrow T^\times$  induced by  $\text{Br}_Z$  and which satisfies  $f(s)\text{Br}_Z(s')f(s^{-1}) = \text{Br}_Z(ss's^{-1})$  for all  $s \in N_{S^\times}(\sigma(Z))$  and all  $s' \in S^Z$ . The latter condition implies that if  $(x, s) \in \hat{N}$  then  $(x, f(s)) \in \tilde{N}$ , and the map sending  $(x, s)$  to  $(x, f(s))$  is in fact a  $k^\times$ -group isomorphism inducing the identity on  $N$ . Now  $\tau$  and  $f \circ \sigma$  are two group homomorphisms from  $P$  to  $T^\times$  lifting the action of  $P$  on  $T$ , hence they are equal, and therefore the above isomorphism  $\tilde{N} \cong \hat{N}$  preserves the canonical images of  $P$  elementwise. Finally, since  $f$  extends  $\text{Br}_Z$  we get for any  $x \in O_{p'}(N)$  that  $f(xb) = \text{Br}_Z(xb) = xc$ , and so the isomorphism  $\tilde{N} \cong \hat{N}$  preserves the canonical images of  $O_{p'}(N)$ . Taking the quotients by these images yields a  $k^\times$ -isomorphism  $\hat{L} \cong \tilde{L}$  with the required properties.  $\square$

**Remarks.** (1) If  $G$  is a strongly  $p$ -solvable finite group then  $SL_2(p)$  is not isomorphic to a subquotient of  $G$ , and hence every block of  $\mathcal{O}G$  is  $SL_2(p)$ -free in the sense of [12, 1.2]. By [12, Theorem 1.4] the equality  $\mathcal{F}_{G,b} = \mathcal{F}_{N,c}$  holds more generally for any block  $b$  of a finite group whenever the block  $b$  is  $SL_2(p)$ -free.

(2) If  $b$  is the principal block of a strongly  $p$ -solvable finite group  $G$ , then  $c$  is the principal block of  $N$  and the inclusion  $N \subseteq G$  induces an algebra isomorphism  $\mathcal{O}Nc \cong \mathcal{O}Gb$ . This in turn is equivalent to Glauberman's Theorem [8, Ch. 8, Theorem 2.11] because the kernel of the principal block  $b$  of  $\mathcal{O}G$  is well-known to be  $O_{p'}(G)$ .

(3) Glauberman's Theorem [8, Ch. 8, Theorem 2.11] is used in the proof of the  $p$ -nilpotency Theorem of Glauberman and Thompson, stating that for an odd prime  $p$  a finite group  $G$  is  $p$ -nilpotent if and only if  $N = N_G(Z(J(P)))$  is  $p$ -nilpotent, where here  $P$  is a Sylow- $p$ -subgroup of  $G$ . The latter has a block theoretic version as well; cf. [11].

(4) A. Watanabe [19] pointed out that the statements and proofs remain valid with  $Z(J(P))$  replaced by any normal subgroup  $Z$  of  $P$  with the property that  $N_G(Z)$  controls fusion in  $b$ .

## REFERENCES

1. J. L. Alperin, M. Broué, *Local methods in block theory*, Ann. Math. **110** (1979), 143–157.
2. R. Brauer, *On the structure of blocks of characters of finite groups*, Lecture Notes in Mathematics **372** (1974), 103–130.
3. M. Broué, L. Puig, *A Frobenius theorem for blocks*, Invent. Math. **56** (1980), 117–128.
4. M. Cabanes, *Extensions of  $p$ -groups and construction of characters*, Comm. Alg. **15** (1987), 1297–1311.
5. E. C. Dade, *Endo-permutation modules over  $p$ -groups I*, Ann. Math. **107** (1978), 459–494.
6. E. C. Dade, *Endo-permutation modules over  $p$ -groups II*, Ann. Math. **108** (1978), 317–346.
7. P. Fong, *On the characters of  $p$ -solvable groups*, Trans. Amer. Math. Soc. **98** (1961), 263–284.
8. D. Gorenstein, *Finite Groups*, Chelsea Publishing Company, New York, 1980.
9. M. E. Harris, M. Linckelmann, *Splendid derived equivalences for blocks of finite  $p$ -solvable groups*, J. London Math. Soc. **62** (2000), 85–96.
10. M. E. Harris, M. Linckelmann, *On the Glauberman and Watanabe correspondences for blocks of finite  $p$ -solvable groups*, Trans. Amer. Math. Soc. **354** (2002), 3435–3453.
11. R. Kessar, M. Linckelmann, *A block theoretic analogue of a theorem of Glauberman and Thompson*, Proc. Amer. Math. Soc. **131** (2003), 35–40.
12. R. Kessar, M. Linckelmann, G. R. Robinson, *Local control in fusion systems of  $p$ -blocks of finite groups*, J. Algebra **257** (2002), 393–413.
13. B. Külshammer, L. Puig, *Extensions of nilpotent blocks*, Invent. Math. **102** (1990), 17–71.
14. L. Puig, *Local block theory in  $p$ -solvable groups*, Proceedings of Symp. Pure Math. **37** (1980), 385–388.
15. L. Puig, *Pointed groups and construction of characters*, Math. Z. **176** (1981), 265–292.
16. L. Puig, *Local extensions in endo-permutation modules split: a proof of Dade’s theorem*, Séminaire sur les groupes finis, Publ. Math. Univ. Paris VII (1986), 199–205.
17. L. Puig, *Nilpotent blocks and their source algebras*, Invent. Math. **93** (1988), 77–116.
18. J. Thévenaz,  *$G$ -Algebras and Modular Representation Theory*, Oxford Science Publications, Clarendon Press, Oxford, 1995.
19. A. Watanabe, private communication (2005).

RADHA KESSAR, MARKUS LINCKELMANN  
DEPARTMENT OF MATHEMATICAL SCIENCES  
MESTON BUILDING  
ABERDEEN, AB24 3UE  
U.K.