



Original citation:

Harrison, E. et al. (2012). Selective measurement of anti-tTG antibodies in coeliac disease and IgA deficiency: an alternative pathway. *Postgraduate Medical Journal*

Permanent WRAP url:

<http://wrap.warwick.ac.uk/50135>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes the work of researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.


Publisher's statement:

Copyright © 2012 BMJ Publishing Group Ltd

A note on versions:

The version presented here may differ from the published version or, version of record, if you wish to cite this item you are advised to consult the publisher's version. Please see the 'permanent WRAP url' above for details on accessing the published version and note that access may require a subscription.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk



<http://go.warwick.ac.uk/lib-publications>

POSTGRADUATE MEDICAL JOURNAL

Selective measurement of anti-tTG antibodies in coeliac disease and IgA deficiency – an alternative pathway

Journal:	<i>Postgraduate Medical Journal</i>
Manuscript ID:	postgradmedj-2011-130188.R4
Article Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Harrison, Elizabeth; University Hospitals Coventry and Warwickshire, Department of Gastroenterology Li, Ka-Kit; University Hospital Coventry and Warwickshire, Department of Gastroenterology Petchey, Michael; University Hospital Coventry and Warwickshire, Department of Pathology Nwokolo, Chuka; University Hospital Coventry and Warwickshire, Department of Gastroenterology Loft, Duncan; University Hospital Coventry and Warwickshire, Department of Gastroenterology Arasaradnam, Ramesh; University Hospital Coventry and Warwickshire, Department of Gastroenterology
Keywords:	Coeliac disease < GASTROENTEROLOGY

SCHOLARONE™
Manuscripts

1
2
3 **Selective measurement of anti-tTG antibodies in coeliac disease and IgA**
4
5 **deficiency – an alternative pathway**
6
7

8
9 Elizabeth Harrison¹, Ka-Kit Li¹, Michael Petchey², Chuka Nwokolo¹, Duncan Loft¹,
10 Ramesh Arasaradnam¹
11
12

13
14
15
16 Department of Gastroenterology, University Hospitals Coventry and Warwickshire.¹
17

18 Department of Pathology, University Hospitals Coventry and Warwickshire.²
19

20
21
22 University Hospitals Coventry and Warwickshire.
23

24 Clifford Bridge Road, Coventry, CV2 2DX, United Kingdom.
25
26
27

28
29 Corresponding author:
30

31 Ramesh Arasaradnam, Department of Gastroenterology, University Hospitals
32 Coventry and Warwickshire, Clifford Bridge Road, Coventry, CV2 2DX, United
33 Kingdom.
34
35
36

37 E-mail: R.Arasaradnam@warwick.ac.uk
38

39 Telephone: 02476966089
40

41 Fax: 02476966090
42
43
44

45
46
47 Keywords: Coeliac disease, serology, anti-tissue transglutaminase, IgA, IgG
48
49

50
51
52 Word count = 1758
53
54
55
56
57
58
59
60

CONTRIBUTORSHIP STATEMENT

All authors contributed equally.

Confidential: For Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

ABSTRACT**Objective**

To determine the ability of selective antibody testing to screen for coeliac disease in the presence of IgA deficiency and to define the sensitivity of a pathway using this method (Figure1).

Method

All IgA and IgG anti-tTG tests performed at our centre between January 2008 and December 2009, using the Immucap 250 analyser, were retrospectively reviewed. Positive results were correlated with histology. Results were used to validate our diagnostic pathway.

Results

12,289 consecutive serological tests were reviewed. IgA deficient patients gave either an "error" reading or very low response on the Immucap 250 analyser. Subsequent testing of this sub-group demonstrated raised IgG anti-tTG antibodies in those with histologically proven coeliac disease.

Conclusions

Using our antibody screening pathway, which involves the selective use of IgG anti-tTG, sensitivity increased from 87% to 92% in those with IgA deficiency. Adoption of this pathway for coeliac screening would negate the routine screening of immunoglobulin levels, with resultant cost saving.

INTRODUCTION

Coeliac disease is a common disease that affects around 1% of the general population in the UK, and makes up a large proportion of the workload seen in general gastroenterology clinics [1]. The pathogenesis of coeliac disease is related to hypersensitivity to gluten in genetically susceptible individuals. As a result of ingestion of gluten containing products (wheat, rye and barley), damage is caused to the mucosa of the small bowel leading to villous atrophy. This can result in a wide spectrum of signs and symptoms which include lethargy, diarrhoea, bloating, weight loss and anaemia but, in some cases, patients can remain asymptomatic. However, as such symptoms and signs are non-specific, and are common to many other gastrointestinal diseases, the diagnosis of coeliac disease may be challenging. Therefore, a high index of clinical suspicion is needed to exclude coeliac disease and to prevent delays in diagnosis.

Duodenal biopsies remain the gold standard in diagnosis. However, identification of those patients to biopsy usually involves serology testing against specific antibodies. The type of antibodies being used has evolved over the years with more sensitive and specific assays replacing older ones. Currently, IgA anti-tissue transglutaminase (IgA anti-tTG) antibodies are recommended as a first choice test for the diagnosis of coeliac disease [2]. However, variation in the sensitivity and specificity of serology tests has been noted from previous studies [3]. Patients with IgA deficiency are prone to false negative antibody results [4]. To add to the problem, IgA deficiency is more prevalent in patients with coeliac disease than in those without coeliac disease [5,6]. The measurement of IgA immunoglobulin level is traditionally carried out at the same time as the IgA anti-tTG antibodies, in an attempt to identify patients with IgA deficiency [6]. However, such an approach involves carrying out two assays (anti-tTG

1
2
3 antibodies & IgA immunoglobulin) for each patient being tested for coeliac disease.

4
5 In addition, some laboratories perform Endomysial Antibody (EMA) testing which
6
7 increases the yield slightly but at an additional cost. Moreover EMA testing is subject
8
9 to the same limitation as it is IgA dependent, and is not as reproducible as an ELISA
10
11 method, as it relies on immuno-fluorescence for which there is inter-assay variation
12
13 [7].
14

15
16 The use of IgG anti-tTG antibodies in screening for coeliac disease in patients with
17
18 selective IgA deficiency has previously been demonstrated [8,9]. However, these
19
20 previous studies did not clearly define a role for the use of IgG anti-tTG antibodies in
21
22 the diagnosis of coeliac disease. It has been suggested that it may be used as a further
23
24 screening test in those patients with IgA deficiency, as shown by very low IgA anti-
25
26 tTG antibodies [10].
27

28
29 We believe that the most cost effective and practical approach, to aid in the diagnosis
30
31 of coeliac disease, is by initially testing with IgA anti-tTG antibody and then selecting
32
33 a subset to undergo IgG anti-tTG antibody testing, by an enzyme-linked
34
35 immunosorbent assay (ELISA). Based on this we devised an ELISA-based serological
36
37 pathway to simplify screening for coeliac disease and co-existent IgA deficiency
38
39 (figure 1). The primary objective of this study is to validate our coeliac disease
40
41 screening pathway, by determining its sensitivity and specificity.
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

METHOD

Subjects

Our centre is a university teaching hospital serving a population of over 450,000. Since 2007, serological screening for coeliac disease at our centre has consisted of performing ELISA testing for both IgA anti-tTG antibodies and IgG anti-tTG antibodies in all patients. All patients screened for coeliac disease by IgA and IgG anti-tTG antibodies during the 2 year study period (January 2008 and December 2009) were included. Patients diagnosed with coeliac disease prior to 2008 were excluded. Ethical approval was not required and all results in this retrospective study were anonymised.

Serology

Serology tests were performed using the commercial kits tTG-IgA Celikey and tTG-IgG Celikey (Phadia, Freiburg, Germany) and in accordance with the manufacturer's instructions the cut-off for negative results was set at <5 U/ml (the manufacturer's quote <10 U/ml). The Immunocap 250 analyser ® Phadia was used. In those patients with IgA deficiency, the Immunocap 250 will give either an 'error' reading or a very low response [11]. A very low response was defined by Response Unit (RU) of less than 50, as determined by the calibration curve on the Immunocap 250. Previous practice had been to refer members of this subgroup for duodenal biopsy, regardless of their IgG anti-tTG antibody result, to determine if they had coeliac disease.

Histology

The diagnosis of coeliac disease was defined as positive serology (IgA or IgG anti-tTG) in the presence of confirmatory histological findings (Marsh grade of 2 - 4). Latent coeliac disease was defined as positive serology in the absence of confirmatory

1
2
3 histology, while histology in the absence of serology was defined as possible
4 serology-negative coeliac disease. Our electronic histology database was reviewed
5
6 over the same 2 year period. Details were recorded of all patients diagnosed with
7
8 coeliac disease through positive histology, on duodenal biopsy, regardless of
9
10 serological result.
11
12

13 **Pathway**

14
15 A pathway for the selective use of IgG anti-tTG in serological screening for coeliac
16
17 disease was designed (Figure 1).
18
19

20 **Data-analysis**

21
22 Results of all patients screened for coeliac disease were entered into an Excel
23
24 spreadsheet. Serological and histological results were analysed to determine which
25
26 patients met the criteria for a diagnosis of coeliac disease. Results were analysed in
27
28 the context of the proposed pathway for serological screening for coeliac disease.
29
30 Sensitivity and specificity for coeliac detection, if the pathway had been used, were
31
32 calculated and used to determine its screening ability.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

RESULTS

During the study period, 12,289 patients with no prior history of coeliac disease underwent serological screening for coeliac disease by IgA anti-tTG antibodies and IgG anti-tTG antibodies. Of these 12,289 patients, 87 were classified, through either confirmatory histology or elevated anti-tTG antibodies, as possibly having coeliac disease (Table 1).

IgA anti tTG and positive histology

Further analysis showed that 66 had both an elevated IgA anti-tTG antibody and confirmatory histology (Marsh grading 2 – 4). Therefore, in this group, the diagnosis of coeliac disease was suggested by using only the IgA anti-tTG antibodies serological test and then confirmed with histology. Thus, the sensitivity for diagnosing coeliac disease with IgA anti-tTG antibodies alone was 87% (specificity 99 %) (Table 2).

Negative IgA anti tTG and positive histology

Those patients identified as having either undetectable (“error” reading) or low IgA anti-tTG levels also had their IgG anti-tTG antibodies and duodenal histology checked. Of these patients, 4 had a positive IgG anti-tTG antibodies serological test and subsequently had positive histology on biopsy (table 3). Thus, these 4 patients had IgA deficiency (with an “error” reading on the Immunocap 250), but raised IgG anti-tTG antibodies and positive histology. They were diagnosed with coeliac disease. Therefore, the use of selective IgG anti-tTG improved the sensitivity for coeliac disease detection to 92% (specificity 99%).

Latent coeliac disease

In 11 patients with positive serology, histology was reported normal. In these patients the mean IgA anti-tTG level was 43 ± 35 and 8.1 ± 20 for IgG anti-tTG. Therefore, it

1
2
3 was assumed that these patients have latent disease and that they will manifest
4
5 clinically at a later stage in their lives.
6
7

8 9 **DISCUSSION**

10 Our objective on commencement of this study was to determine the role of IgG anti-
11
12 tTG in screening for coeliac and to define the sensitivity of a pathway using such a
13
14 method. Through this study we reviewed the results of non-selective IgA and IgG anti-
15
16 tTG screening at our centre and applied these results to the proposed pathway. This
17
18 enabled us to determine the sensitivity (92%) and specificity (99%) for coeliac
19
20 detection that such a pathway would generate (Figure 1).
21
22

23 We show that this proposed Immunocap 250 pathway is able to identify coeliac
24
25 disease in spite of IgA deficiency. These individuals produced a very low or “error”
26
27 result on IgA anti-tTG testing, but were subsequently shown to have raised IgG anti-
28
29 tTG antibodies. Without testing IgG anti-tTG in these patients, a false negative result
30
31 in those with IgA deficiency would have been generated (4 patients out of 76 had IgA
32
33 deficiency, but their IgG anti-tTG was elevated).
34
35

36 Furthermore, in those patients without coeliac disease, but with IgA deficiency, an
37
38 “error” reading was also found. However, in this patient subset, the subsequent
39
40 measurement of IgG anti-tTG antibodies showed a negative result. This would have
41
42 allowed a decision to be made that a small bowel biopsy was not necessary. This
43
44 would not only have benefited the patient by avoiding an unnecessary and invasive
45
46 investigation, but would also have generated significant cost saving.
47
48

49 Additionally, the proposed approach is likely to be even more cost effective and less
50
51 labour intensive. Both IgA and IgG anti-tTG antibodies can be carried out on the
52
53 same ELISA plate. Furthermore, we believe that adhering to this pathway would
54
55
56
57
58
59
60

1
2
3 negate the routine screening of immunoglobulin levels, with resultant additional cost
4
5 saving.

6
7 Additional saving, without clinical detriment, is of great importance, given the
8
9 increasing awareness of coeliac disease leading to an increasing demand for testing.
10
11 As with all serological testing, it does not purport 100% specificity or sensitivity. In
12
13 our series we report a 92% sensitivity which must be borne in mind when requesting
14
15 such tests.
16
17

18
19 A limitation of this study was that the researchers did not have input into which
20
21 patients were selected for screening. This may have influenced sensitivities and
22
23 specificities. Decision to screen was made purely by the referring clinicians.
24
25 Duplicate screens may have been performed on patients, but as coeliac disease is a
26
27 dynamic disease it is reasonable to repeat serology over time. Our low prevalence rate
28
29 may also be explained by the mixed ethnic population in Coventry.
30
31

32 33 34 **CONCLUSION**

35
36 Previous studies have demonstrated that the addition of IgG anti-tTG antibodies is a
37
38 useful tool in diagnosing coeliac disease in patients with IgA deficiency. We expand
39
40 on this by demonstrating how it can be adopted into a specific pathway [8,9] The
41
42 benefit of this is that it offers a selective and economical approach, which can easily
43
44 be adopted in most laboratories.
45
46
47

48 49 **MAIN MESSAGES**

- 50
51 1. IgA anti-tTG identifies most people who may have coeliac disease
52
53 2. Selective use of IgG anti-tTG detects coeliac disease in those with IgA deficiency
54
55
56
57
58
59
60

1
2
3 3. Use of comprehensive clinical/biochemical pathway allows for an economical
4
5 approach to detecting coeliac disease
6
7

8
9
10 **CURRENT RESEARCH QUESTIONS**

- 11 1. To validate the proposed pathway using EMA instead of IgG in the presence of IgA
12
13 deficiency
14
15 2. To measure corresponding mucosal tTG, especially in those with IgA deficiency
16
17 3. To determine the validity of our proposed pathway in primary care setting
18
19 4. To determine the validity of our pathway using different manufactures' automated
20
21 platforms
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

TABLES

Table 1: Summary of patients with coeliac disease

Screening modality which identified patient	Number of patients
Traditionally detected coeliac disease group (raised IgA anti-tTG, histology positive)	66
IgA deficient coeliac disease group (IgA anti-tTG antibodies showed “error” but raised IgG anti-tTG and positive histology)	4
Presumed latent coeliac disease group (raised IgA and IgG anti-tTG, but histology negative)	11
Histology suggestive of coeliac disease alone (both IgA and IgG anti-tTG negative)	6
Total number of patients with manifest or latent coeliac disease	87

Table 2: IgA anti-tTG antibody results correlated with histological confirmation

	Negative biopsy	Positive biopsy	Total
Positive anti-tTG IgA	11	66	77
Negative anti-tTG IgA	12202	10	12212
Total	12213	76	12289

Sensitivity of IgA anti-tTG for detecting coeliac screening = 0.8

Specificity of IgA anti-tTG for detecting coeliac screening = 0.9

Table 3: Detection with proposed subsequent IgG anti-tTG antibody results correlated with histological confirmation

	Negative biopsy	Positive biopsy	Total
Positive IgA anti-tTG or IgG anti-tTG	11	70	81
Negative IgA with negative IgG anti-tTG	12202	6	12208
Total	12213	76	12289

Sensitivity of subsequent IgG anti-tTG for detecting coeliac screening = 0.9

Specificity of subsequent IgG anti-tTG for detecting coeliac screening = 0.9

ACKNOWLEDGMENTS / COMPETING INTERESTS /FUNDING

None.

Confidential: For Review Only

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60


REFERENCES

- [1] Leeds JS, Hopper AD, Sanders DS. Coeliac Disease. *Br Med Bull* 2008;88(1):157-70.
- [2] NICE clinical guideline 86. Coeliac disease: recognition and assessment of coeliac disease.
- [3] Hill ID. What are the sensitivity and specificity of serologic tests for celiac disease? Do sensitivity and specificity vary in different populations? *Gastroenterology*. 2005;128(4 Suppl 1):S25-32.
- [4] McGowan KE, Lyon ME, Butzner JD. Celiac disease and IgA deficiency: complications of serological testing approaches encountered in the clinic. *Clin Chem*. 2008;54(7):1203-9
- [5] Heneghan MA, Stevens FM, Cryan EM, et al. Celiac sprue and immunodeficiency states: a 25-year review. *J Clin Gastroenterol* 1997;25(2):421-5
- [6] Thomas PD, Forbes A, Green J, et al. Guidelines for the investigation of chronic diarrhoea. 2nd edn. *Gut* 2003;52(Suppl 5):v1-v15
- [7] Leffler DA, Schuppan D. Update on Serologic Testing in Celiac Disease. *Am J Gastroenterol*. 2010;105:2520–2524
- [8] Korponay-Szabó IR, Dahlbom I, Laurila K, et al. Elevation of IgG antibodies against tissue transglutaminase as a diagnostic tool for coeliac disease in selective IgA deficiency. *Gut*. 2003;52(11):1567-71
- [9] Lenhardt A, Plebani A, Marchetti Fet al. Role of human-tissue transglutaminase IgG and anti-gliadin IgG antibodies in the diagnosis of coeliac disease in patients with selective immunoglobulin A deficiency. *Dig Liver Dis*. 2004;36(11):730-4

1
2
3 [10] Fernandez E, Blanco C, Garcia S, et al. Use of low concentrations of human IgA
4 anti-tissue transglutaminase to rule out selective IgA deficiency in patients with
5 suspected coeliac disease. Clin Chem. 2005; 51(6):1014-1016
6
7

8
9
10 [11] Löwbeer C, Wallinder H. Undetectable anti-tissue transglutaminase IgA antibody
11 measured with ELISAA Celikey indicates selective IgA deficiency. Clin Chim Acta.
12 2010;411(7-8):612
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 1: Pathway of selective testing with anti-tTG for coeliac disease



RU = Response unit determined by calibrated curve on the Immunocap 250