

THE UNIVERSITY OF WARWICK

Original citation:

Baltrus, David A., Nishimura, Marc T., Dougherty, Kevin M., Biswas, Surojit, Mukhtar, M. Shahid, Vicente, Joana G., Holub, E. B. and Dangl, Jeffery L. (2012) The Molecular Basis of Host Specialization in Bean Pathovars of *Pseudomonas syringae*. *Molecular Plant-Microbe Interactions*, Vol. 25 (No. 7). pp. 877-888.

Permanent WRAP url:

<http://wrap.warwick.ac.uk/50231>

Copyright and reuse:

The Warwick Research Archive Portal (WRAP) makes the work of researchers of the University of Warwick available open access under the following conditions. Copyright © and all moral rights to the version of the paper presented here belong to the individual author(s) and/or other copyright owners. To the extent reasonable and practicable the material made available in WRAP has been checked for eligibility before being made available.

Copies of full items can be used for personal research or study, educational, or not-for-profit purposes without prior permission or charge. Provided that the authors, title and full bibliographic details are credited, a hyperlink and/or URL is given for the original metadata page and the content is not changed in any way.


Publisher's statement:

Copyright The American Phytopathological Society

A note on versions:


The version presented in WRAP is the published version or, version of record, and may be cited as it appears here.

For more information, please contact the WRAP Team at: wrap@warwick.ac.uk



<http://go.warwick.ac.uk/lib-publications>

Molecular Plant-Microbe Interactions



**The molecular basis of host specialization in bean pathogens
of *Pseudomonas syringae***

Journal:	<i>Molecular Plant-Microbe Interactions</i>
Manuscript ID:	MPMI-08-11-0218.R2
Manuscript Type:	Research
Date Submitted by the Author:	n/a
Complete List of Authors:	Baltrus, David; University of Arizona, Plant Sciences Nishimura, Marc; UNC Chapel Hill, Biology Dougherty, Kevin; University of Arizona, Plant Sciences Biswas, Surojit; UNC Chapel Hill, Biology Muhktar, Shahid; U Alabama Birmingham, Biology Vicente, Joana; University of Warwick, Warwick-HRI Holub, Eric; University of Warwick, Warwick-HRI; Dangl, Jeff; Univ. of North Carolina, Biology
Area That Best Describes Your Manuscript:	Avirulence factors, Bacterial pathogenesis, effectors < Bacterial pathogenesis, type-3 secretion < Bacterial pathogenesis, Genomics, PAMPs, MAMPs, and elicitors

The Molecular Basis of Host Specialization in Bean Pathovars of *Pseudomonas syringae*

David A. Baltrus,¹ Marc T. Nishimura,² Kevin M. Dougherty,¹ Surojit Biswas,² M. Shahid Mukhtar,² Joana Vicente,³ Eric B. Holub,³ and Jeffery L. Dangl^{2,4,5,6,7}

¹School of Plant Sciences, The University of Arizona, PO Box 210036, Forbes Building 303, Tucson, AZ 85721-0036, U.S.A.; ²Department of Biology, University of North Carolina (UNC), Chapel Hill, NC 27599, U.S.A.; ³University of Warwick, School of Life Sciences, Wellesbourne Campus, CV35 9EF, U.K.; ⁴Howard Hughes Medical Institute, ⁵Curriculum in Genetics and Molecular Biology, ⁶Carolina Center for Genome Sciences, and ⁷Department of Microbiology and Immunology, University of North Carolina, Chapel Hill, NC 27599, U.S.A.

Submitted 29 August 2011. Accepted 4 March 2012.

Biotrophic phytopathogens are typically limited to their adapted host range. In recent decades, investigations have teased apart the general molecular basis of intraspecific variation for innate immunity of plants, typically involving receptor proteins that enable perception of pathogen-associated molecular patterns or avirulence elicitors from the pathogen as triggers for defense induction. However, general consensus concerning evolutionary and molecular factors that alter host range across closely related phytopathogen isolates has been more elusive. Here, through genome comparisons and genetic manipulations, we investigate the underlying mechanisms that structure host range across closely related strains of *Pseudomonas syringae* isolated from different legume hosts. Although type III secretion-independent virulence factors are conserved across these three strains, we find that the presence of two genes encoding type III effectors (*hopCI* and *hopMI*) and the absence of another (*avrB2*) potentially contribute to host range differences between pathovars *glycinea* and *phaseolicola*. These findings reinforce the idea that a complex genetic basis underlies host range evolution in plant pathogens. This complexity is present even in host–microbe interactions featuring relatively little divergence among both hosts and their adapted pathogens.

Pseudomonas syringae is a facultative phytopathogen with the potential to cause disease in many crop species across the globe. Despite a collectively broad host range for this bacterial species, pathogen variants (pathovars) have largely been defined by their ability to cause disease in different species. Non-adapted isolates either elicit immune responses or simply fail to thrive on the incompatible host (Lindeberg et al. 2009). Individual isolates will exhibit quantitative variation in growth and symptoms both within and between host species (Fan et al. 2007; Lin and Martin 2007); thus, virulence within *P. syringae* is expected to be a complex trait. Although progress has been made in understanding the factors that limit and promote the

expansion of host range for several bacteria, there is little consensus concerning the generality and relative importance of various molecular mechanisms known to underlie host range evolution.

The host's ability to perceive and induce defense against a potential pathogen is a major barrier to plant–bacteria compatibility. For instance, the perception of pathogen- or microbe-associated molecular patterns (PAMPS or MAMPS) by matching host pattern recognition receptor (PRR) proteins is generally described as a basic layer of incompatibility because PAMPS are necessary for microbial function (e.g., flagellum or translation elongation factor) and, thus, are indispensable in the pathogen. PRR proteins induce defense responses described as PAMP- or MAMP-triggered immunity (PTI) (Jones and Dangl 2006). An additional layer of defense is provided by highly variable receptors such as nucleotide-binding leucine-rich repeat (NB-LRR) proteins, which induce defense responses described as effector-triggered immunity (ETI) (Jones and Dangl 2006). Typically, type III effectors (T3E) are the main determinants of both virulence and ETI in plant-pathogenic bacteria such as *P. syringae* (Lindeberg et al. 2009). Current opinions are divided over whether nonhost resistance (NHR), in which all tested isolates of a pathogen trigger resistance reactions in all tested cultivars of a plant species, is mechanistically different from induced immune response involving pathogen recognition (PTI and ETI) within demonstrated hosts (Abramovitch et al. 2006; Mellersh and Heath 2003; Schulze-Lefert and Panstruga 2011; Wen et al. 2010).

Mutually exclusive mechanisms for host versus NHR are difficult to prove. For instance, all bacterial cells contain multiple conserved PAMPS and, thus, are likely to trigger PTI. Virulence is expected to have evolved by an acquired ability to suppress PTI, most likely by the delivery and action of effector molecules, and specific perception of some effectors has been shown as the molecular basis for elicitation of ETI (Jones and Dangl 2006). However, plants can also contain receptors that mediate ETI to nonadapted pathogens, such as RPS4 and RPS5 proteins that could potentially protect *Arabidopsis thaliana* against legume pathovars of *P. syringae* in natural populations (Gassmann et al. 1999; Warren et al. 1999). Additionally, ETI-mediating receptors can provide an effective barrier of broad-spectrum resistance to a pathogen species (Borhan et al. 2008; Song et al. 2003; Tai et al. 1999; Vera Cruz et al. 2000), particularly if disruption or loss of the recognized effector significantly affects pathogen fitness on susceptible hosts.

Corresponding author: D. A. Baltrus; E-mail: baltrus@email.arizona.edu

Current address: of M. S. Mukhtar: MSM Department of Biology, CH106, University of Alabama at Birmingham, 1300 University Blvd., Birmingham, AL 35294, U.S.A.

*The e-Xtra logo stands for “electronic extra” and indicates that two supplementary figures and one supplementary table are published online.

Both PTI and ETI can limit the growth of *P. syringae* across a wide variety of hosts (Ferrante et al. 2009; Ham et al. 2007; Li et al. 2005; Lin and Martin 2007; Perchepped et al. 2006; Takeuchi et al. 2003; Wen et al. 2010; Wroblewski et al. 2009). For example, transfer of the EFR receptor from *Arabidopsis* to *Nicotiana benthamiana* can restrict the growth of previously pathogenic strains (Lacombe et al. 2010), while diversity in flagellar proteins across strains contributes to immune evasion (Li et al. 2005; Takeuchi et al. 2003). Similarly, for ETI, the transfer of several single genes from *P. syringae* pv. *tomato* PT23 into various strains of *P. syringae* pv. *glycinea* can restrict the growth of the normally virulent recipient strain on soybean (Kobayashi et al. 1989). However, disruption of these genes within *P. syringae* pv. *tomato* PT23 did not increase its virulence on soybean (Lorang et al. 1994), implying either the existence of additional recognized avirulence factors or the lack of necessary virulence genes. A wide variety of strains were shown to be more virulent on a *pto* background of tomato (Lin and Martin 2007), which cannot recognize the T3E families *avrPto* or *hopAB*. Conversely, recognition of HopQ1 was shown to be the primary limitation for growth of *P. syringae* pv. *tomato* DC3000 on tobacco (Wei et al. 2007), and both *AvrRpt2* and *HopAS1* can limit the growth of multiple strains on *Arabidopsis* (Almeida et al. 2009; Sohn et al. 2011). Transfer of T3E across strains has also been shown to increase the recipient's virulence on certain nonhost species (Ferrante et al. 2009; Lindeberg et al. 2009). In most if not all of these examples, the pathogen isolates used were collected from divergent host species and are distantly related. This divergence makes it difficult to place changes within an evolutionary context. Identifying molecular factors that limit growth of recently diverged strains can illuminate the molecular basis of barriers that delimit host range as well as the evolutionary path toward adaptation to new hosts.

P. syringae is a diverse bacterial species that can be phylogenetically subdivided based on DNA hybridization, multi-locus sequence typing (MLST), and core genome phylogenies (Baltrus et al. 2011; Gardan et al. 1999; Hwang et al. 2005). MLST group III is composed of pathogens of numerous plant species, with a subset of isolates displaying enough diversity from other *P. syringae* clades to suggest designation as a separate species, *P. savastanoi* (Gardan et al. 1999; Hwang et al. 2005; Qi et al. 2011). Indeed, some members of this subgroup are metabolically less diverse than other strains, which could contribute to apparent differences in dispersion capabilities among MLST group III strains compared with other *P. syringae* isolates (Morris et al. 2010; Rico and Preston 2008). Such ecological differences could also provide greater opportunity for host specialization than is found in other clades of this species.

Two important pathovars within *P. syringae* MLST group III are each agronomically relevant pathogens of different legume species: *P. syringae* pv. *glycinea* causes bacterial blight of soybean (*Glycine max*), and *P. syringae* pv. *phaseolicola* causes halo blight in French and common dry bean (*Phaseolus vulgaris*) (Arnold et al. 2011; Qi et al. 2011). The interaction of isolate *P. syringae* pv. *phaseolicola* 1448a with bean has been a longstanding model for understanding the basis of pathogenicity (Arnold et al. 2011). Whole genome sequences are publicly available for isolates from both *P. syringae* pv. *glycinea* and *P. syringae* pv. *phaseolicola*, including strains *P. syringae* pv. *glycinea* R4 and *P. syringae* pv. *phaseolicola* 1448a (Baltrus et al. 2011; Qi et al. 2011; Studholme 2011). Therefore, exploration of pathogenicity within this subgroup represents an important opportunity to understand host range evolution at a fine scale because of recent divergence of both the pathogens and their hosts.

Here, we report host specialization within three isolates of *P. syringae*, each collected from a different legume host and representing pathovars *glycinea* and *phaseolicola*. We added next-generation sequencing of isolate *P. syringae* pv. *phaseolicola* 1644R, (collected from mung bean) to enable comparisons of putative virulence gene repertoires with the genomes of *P. syringae* pv. *phaseolicola* 1448a and *P. syringae* pv. *glycinea* R4. We attempted to identify molecular determinants that contribute to quantitative differences in virulence in a cultivar of *Phaseolus vulgaris* ('Canadian Wonder') that is susceptible to all known *P. syringae* pv. *phaseolicola* isolates. By placing the evolution of host range within a more narrowly defined phylogenetic context, we sought to identify molecular factors that contribute to changes in host range and to better understand the process of host range evolution.

RESULTS

Sequencing of *P. syringae* pv. *phaseolicola* isolate 1644R.

A draft genome sequence of *P. syringae* pv. *phaseolicola* 1644R was created by assembling one lane of Illumina PE reads with SoapDenovo (Li et al. 2010). The final assembly contains 1,169 contigs with a mean contig size of 5,073, an N50 value of 13,068, and a maximum contig size of 66,841. The total size of the draft genome assembly is 5,930,644 nucleotides. Using Genemark (Besemer and Borodovsky 2005), we identified a total of 6,429 nonredundant putative proteins in the draft genome, which is likely an overestimate due to its fractured nature. In all, 5,306 putative protein sequences are shared among all three strains. There are 452 putative proteins that are uniquely present within *P. syringae* pv. *phaseolicola* 1644R compared with these other two isolates, while 333 (approximately 16%, *P. syringae* pv. *glycinea* R4) and 338 (approximately 16%, *P. syringae* pv. *phaseolicola* 1448a) are missing from only one of the other strains.

Fine-scale differences in host range.

P. syringae pv. *phaseolicola* 1448a and 1644R and *P. syringae* pv. *glycinea* R4 are closely related (Fig. 1A, phylogeny) but were collected from naturally diseased leaves of three different host species: *Phaseolus vulgaris* (French bean), *Vigna radiata* (mung bean), and *G. max* (soybean), respectively. Additionally, *P. syringae* pv. *phaseolicola* 1644R has been assigned to race 1 within pv. *phaseolicola* based on differential defense reactions for multiple cultivars of French bean (data not shown). We measured growth of each of these isolates on each of the host species in order to identify fine-scale host specialization indicative of the early stages of host range evolution (Fig. 1B). Each bacterial isolate achieved its highest population sizes on the particular host species from which it was collected and grew to lower levels on the alternative species. Within a given host species, the isolate collected from that host grew to the highest levels. Furthermore, *P. syringae* pv. *glycinea* R4 grew relatively poorly on both French and mung bean while *P. syringae* pv. *phaseolicola* 1448a and 1644R grew to intermediate levels on mung bean and French bean, respectively. All three of these isolates grew to higher levels in French bean than most other non-bean pathogens in a phylogenetically diverse panel of bacteria (Supplementary Fig. S1).

Non-type III secretion system dependent growth differences in bean pathogens.

Suppression of PTI by *hrpL*-regulated virulence proteins (a subset of which are T3E) is a necessary component for successful *P. syringae* infection. To better understand the contribution of *hrpL*- or T3E-dependent virulence effects within

these bean pathogens and across strains, we compared growth of *hrpL* and *hrcC* mutants in planta across various isolates. We found that *hrpL* and *hrcC* mutants derived from isolate *P. syringae* pv. *phaseolicola* 1448a retained the ability to grow weakly in French bean leaves, and consistently achieved significantly higher population sizes in planta than comparable mutants derived from the bean pathogen *P. syringae* B728a or the tomato pathogen *P. syringae* pv. *tomato* DC3000 (Fig. 2A and B). Additionally, although the *P. syringae* B728a *hrcC* mutant grew to higher levels than an *hrcC* mutant of *P. syringae* pv. *tomato* DC3000 (Fig. 2B), growth differences between *hrpL* mutants of these strains were not statistically distinguishable (Fig. 2A). There was no difference in growth between *hrpL* and *hrcC* mutants of *P. syringae* pv. *phaseolicola* 1448a (data not shown). We further found that *P. syringae* pv. *glycinea* R4 and *P. syringae* pv. *phaseolicola* 1644R and 1448a all shared this potential for limited growth in French bean in the absence of a functioning *hrpL* regulon (Fig. 2B).




Fig. 1. Fine-scale host specialization in three bean pathogens. **A**, Phylogenetic relationships among all three bean pathogens based on conserved sequences within the whole genomes. *Pseudomonas syringae* pv. *mori* MAFF301020 is used as an outgroup. This Bayesian *P. syringae* phylogeny was constructed by analyzing sequences of 12 proteins whose evolutionary relationships match the consensus tree from genomic comparisons. The numerical label displays the fraction of individual loci phylogenies from comparisons of proteins conserved across all four strains that support these relationships. A Bayesian phylogeny of the three bean species based on *matK* gene sequences is shown to the right. All nodes for the bean phylogeny had posterior probabilities >0.95 and are consistent with other studies (Delgado-Salinas et al. 2006; McClean et al. 2008). **B**, The growth of each bean pathogen isolate within three different host plants: French bean ‘Canadian Wonder’, mung bean, and soybean ‘Merit’. Primary leaves from 2-week-old plants were syringe inoculated with bacterial suspensions and assayed 4 days (French bean, soybean) or 5 days (mung bean) after inoculation. Letters represent significant differences with $P < 0.05$ according to Tukey’s highly significant difference and error bars display two standard errors. Separate analyses of variance were run for each host plant data set, which is reflected by differences in the labeling scheme for statistical differences.

Comparing T3E repertoires among the bean pathogens.

We identified putative T3E within *P. syringae* pv. *phaseolicola* 1644R and compared this repertoire to those previously characterized from *P. syringae* pv. *glycinea* R4 and *P. syringae* pv. *phaseolicola* 1448a (Fig. 3A) (Baltrus et al. 2011; Chang et al. 2005; Zumaquero et al. 2010). There were 13 T3E genes shared by all three isolates, as well as one conserved truncation within *hopAA1*. *P. syringae* pv. *phaseolicola* 1644R and *P. syringae* pv. *glycinea* R4 share three T3E genes and a disrup-




Fig. 2. Measureable type III secretion system independent virulence effects within group III bean pathogens. Bacterial population sizes for virulence mutants from several isolates were measured at day 0 and day 4 after syringe inoculation of primary leaves within 2-week-old French bean plants. **A**, Comparison of *hrpL* mutants within *Pseudomonas syringae* B728a, *P. syringae* pv. *phaseolicola* 1448a, and *P. syringae* pv. *tomato* DC3000. **B**, Comparison of *hrcC* mutants within *P. syringae* B728a, *P. syringae* pv. *phaseolicola* 1448a, and *P. syringae* pv. *tomato* DC3000. **C**, Comparison of *hrpL* mutants of *P. syringae* pv. *phaseolicola* 1448a and 1644R and *P. syringae* pv. *glycinea*. Letters represent significant differences with $P < 0.05$ according to Tukey’s highly significant difference and error bars display two standard errors. Separate analyses of variance were run for each host plant data set, which is reflected by differences in the labeling scheme for statistical differences.

tion within *hopV1* that was not found in *P. syringae* pv. *phaseolicola* 1448a (Fig. 3B). Conversely, *P. syringae* pv. *phaseolicola* 1644R and 1448a share four T3E genes, a truncation in *hopAB3*, and a C-terminus frame shift in *hopM1* that was not found in *P. syringae* pv. *glycinea* R4 (Fig. 3B). Only one T3E gene (*hopX1*) was truncated only in *P. syringae* pv. *phaseolicola* 1644R, but this T3E appears to be dispensable for growth within pv. *phaseolicola* on French bean Canadian Wonder and we did not further characterize this locus (Stevens et al. 1998). We additionally queried the predicted protein repertoire of *P. syringae* pv. *phaseolicola* 1644R for novel T3E using the SIEVE server (McDermott et al. 2011) but found no convincing evidence for new translocated families.

Testing T3E within *Phaseolus vulgaris* for avirulence elicitation.

To test for phenotypic differences underlying differences in growth, French bean pods from Canadian Wonder were inoculated with all three isolates and followed for disease symptoms (Fig. 4A). *P. syringae* pv. *phaseolicola* 1448a and 1644R both caused water-soaked lesions after 3 days, whereas *P. syringae* pv. *glycinea* R4 elicited a browning of tissue surrounding the inoculation point, indicative of hypersensitive cell death (Tsiamis et al. 2000).

Because pod inoculations can yield variable responses, we cloned all of the T3E genes unique to *P. syringae* pv. *glycinea* R4 (Fig. 3), expressed them in *P. syringae* pv. *phaseolicola* 1448a under a constitutive promoter, and performed growth curves in French bean Canadian Wonder to identify potential avirulence factors underlying a hypersensitive response (HR) in the bean pod (Fig. 4B). Expression of the *P. syringae* pv. *glycinea* R4 allele of *hopC1* in *P. syringae* pv. *phaseolicola* 1448a reduced growth in planta (Fig. 4B and C). This is consistent with previous reports that *hopC1* from pv. *pisi* is an

avirulence factor on all tested cultivars of French bean (Arnold et al. 2001). Similarly, expression of *P. syringae* pv. *glycinea* R4 *hopM1*, which was truncated in both *P. syringae* pv. *phaseolicola* 1448a and 1644R compared with *P. syringae* pv. *glycinea* R4 (Fig. 3B), also reduced growth in French bean (Fig. 4B and C). Although expression of *hopC1* triggered an HR in the pod assay, *hopM1* did not appear to have any effect compared with wild-type *P. syringae* pv. *phaseolicola* 1448a (data not shown) (Arnold et al. 2001). We found, surprisingly, that expression of *avrPto* from *P. syringae* pv. *glycinea* R4 increased the growth of *P. syringae* pv. *phaseolicola* 1448a on




Fig. 3. Three bean isolates share similar type III effector (T3E) suites. **A**, The T3E suite of *Pseudomonas syringae* pv. *phaseolicola* 1644R was determined by tBLASTn, as described by Baltrus and associates (2011), and compared with *P. syringae* pv. *phaseolicola* 1448a and *P. syringae* pv. *glycinea* R4. Black letters indicate shared T3E alleles, while red letters indicate truncations in T3E compared with the other genomes. **B**, Diagrams of the length in amino acids of each shared truncated T3E. *hopM1* from both *P. syringae* pv. *phaseolicola* 1448a and 1644R possesses a frame-shift mutation (represented by *) relative to the *P. syringae* pv. *glycinea* R4 nucleotide sequence, so that the C terminus of the proteins in both strains is truncated and diverged from that in *P. syringae* pv. *glycinea* R4.




Fig. 4. Type III effectors (T3E) within *Pseudomonas syringae* pv. *glycinea* R4 but not *P. syringae* pv. *phaseolicola* 1644R trigger avirulence responses in French bean ‘Canadian wonder’. **A**, The three group III bean pathogen isolates as well as an *hrcC*- version of *P. syringae* pv. *phaseolicola* 1448a were inoculated into bean pods and observed for virulence responses. **B**, T3E present in *P. syringae* pv. *glycinea* R4 but absent within *P. syringae* pv. *phaseolicola* 1448a were cloned as open reading frames (ORF) including stop codons and measured for their ability to affect growth of *P. syringae* pv. *phaseolicola* 1448a in French bean. Expression of these effectors was driven from an *nptII* promoter. **C**, T3E that modify growth of *P. syringae* pv. *phaseolicola* 1448a in French bean as ORF were cloned with their native promoters and without stop codons and measured for their effects on growth of *P. syringae* pv. *phaseolicola* 1448a in French bean. Only two independent growth curves were performed. An asterisk (*) indicates that the effector clone includes an upstream chaperone. Primary leaves from 2-week-old plants were syringe inoculated with bacterial suspensions and assayed 4 days after inoculation. Letters represent significant differences with $P < 0.05$ according to Tukey’s highly significant difference and error bars display two standard errors. Growth values are listed below statistical groups, with maximum and minimum values listed for pooled groups.

French Bean (Fig. 4B and C). We also tested two genes uniquely present in *P. syringae* pv. *glycinea* R4 known to be involved in virulence: *efe* and *iaaL* (Mazzola and White 1994; Weingart et al. 2001). The *efe* gene codes for ethylene-forming enzyme, which catalyzes the oxygenation of 2-oxoglutarate to form ethylene (Weingart et al. 2001), while the protein product of *iaaL* metabolizes indole acetic acid (auxin) to 3-indole-acetyl-L-lysine (Glickmann et al. 1998). Neither altered the growth of *P. syringae* pv. *phaseolicola* 1448a on French bean. Finally, we note that, although both *P. syringae* pv. *glycinea* R4 and *P. syringae* pv. *phaseolicola* 1644R appeared to lack the pathway for production of phaseolotoxin, absence of this virulence factor has not been shown to alter bacterial growth in planta (Patil et al. 1974).

Identifying virulence factors that enhance growth on French bean.

We generated a mutant of *P. syringae* pv. *glycinea* R4 (DAB890) in which both copies of the *hopH1/hopC1* locus and *hopM1* (the genes identified above conferring avirulence on French bean Canadian Wonder) were all deleted from the genome. This mutant did not exhibit altered growth on French bean Canadian Wonder (Fig. 5A) and still appeared to trigger an HR-like response in pod assays (data not shown). We then replaced T3E genes that were present in *P. syringae* pv. *phaseolicola* 1448a but absent from *P. syringae* pv. *glycinea* R4 (Fig. 3) into DAB890, and tested the resulting strains for

their ability to increase the growth of the mutant on French bean Canadian Wonder. Only the expression of *avrB2* increased the growth of DAB890 (Fig. 5A). A similar effect but with smaller magnitude was observed when *avrB2* from *P. syringae* pv. *phaseolicola* 1448a was expressed from its native promoter in the wild-type *P. syringae* pv. *glycinea* R4 background (Fig. 5C). We also moved T3E genes that were present in *P. syringae* pv. *phaseolicola* 1448a but not *P. syringae* pv. *phaseolicola* 1644R (Fig. 3) into *P. syringae* pv. *phaseolicola* 1644R; none consistently increased the growth of *P. syringae* pv. *phaseolicola* 1644R on French bean Canadian Wonder (Fig. 5D).

DISCUSSION

Many studies have focused attention on characterizing the molecular basis of NHR to phytopathogens (Mellersh and Heath 2003; Mysore and Ryu 2004; Schulze-Lefert and Panstruga 2011; Wen et al. 2010). There is often a continuum of quantitative differences in growth, persistence, or virulence for pathogens on plant genotypes that have been used in experiments to represent nonhost species (Lin and Martin 2007). Moreover, investigations of NHR often involve comparisons of virulence between pathogen isolates that have significantly diverged. The phylogenetic relationship among the isolates and the original hosts from which the isolates were collected is an important consideration for these studies.




Fig. 5. *avrB2* acts as a virulence factor on French bean for *Pseudomonas syringae* pv. *glycinea* R4. **A**, Deletion of *hopC1*, *hopH1*, and *hopM1* from *P. syringae* pv. *glycinea* R4 does not affect growth on French bean. **B**, Type III effectors (T3E) present in *P. syringae* pv. *phaseolicola* 1448a but absent within *P. syringae* pv. *glycinea* R4 were cloned as open reading frames (ORF) with stop codons and measured for their ability to modify growth of DAB890 in French bean. **C**, *avrB2* was cloned with its native promoter and without stop codons and measured for its ability to modify growth of *P. syringae* pv. *phaseolicola* 1448a in French bean. **D**, T3E present in *P. syringae* pv. *phaseolicola* 1448a but absent within *P. syringae* pv. *phaseolicola* 1644R were cloned as ORF with stop codons and measured for their ability to affect growth of *P. syringae* pv. *phaseolicola* 1644R in French bean. An asterisk (*) indicates that the effector clone includes an upstream chaperone. Primary leaves from 2-week-old plants were syringe inoculated with bacterial suspensions and assayed 4 days after inoculation. Letters represent significant differences with $P < 0.05$ according to Tukey's highly significant difference and error bars display two standard errors. Growth values are listed below statistical groups, with maximum and minimum values listed for pooled groups.

Given the rapid advances in DNA sequencing, it is now feasible and straightforward to compare suites of virulence factors between pathogen isolates. Such comparisons will be most informative between closely related bacterial isolates that differ quantitatively in their respective ability to grow on recently diverged host species. This approach allows one to phylogenetically trace the early steps of host range specialization and enables identification of molecular differences that may ultimately contribute to NHR. Conversely, a better understanding of molecular barriers that delimit host range will provide insight into the evolutionary potential for host shifts of these pathogens to new plant species.

With this approach in mind, we chose three recently diverged *P. syringae* isolates collected from closely related legume species to explore the evolution of pathogenicity. Three strains (*P. syringae* pv. *phaseolicola* 1448a and 1644R and *P. syringae* pv. *glycinea* R4) were originally isolated from naturally diseased leaves of *Phaseolus vulgaris* (French bean), *V. radiata* (mung bean), and *G. max* (soybean), respectively (Arnold et al. 2011; Baltrus et al. 2011; Marques et al. 2000). The legume hosts are estimated to have diverged from each other within the last 20 million years, and their genomes display a high level of similarity and colinearity (Delgado-Salinas et al. 2006; McClean et al. 2008). The demographic and evolutionary history of the three bacterial isolates is unknown; however, their phylogenetic branching pattern mirrors that of their respective host species, suggesting co-evolution of pathovars with host species (Fig. 1A). We observed quantitative differences in host specialization, such that each isolate grew best on its original host and slightly less on the other two hosts. All three bacterial strains exhibited higher levels of growth on French bean than did a majority of phylogenetically diverse *P. syringae* isolates.

Given the common occurrence of genotype–host interactions in this pathosystem (Taylor et al. 1996), the molecular mechanisms that define host range differences between pathovars are difficult to determine without extensive sampling of multiple combinations of strains and cultivars. Such studies are complicated by subjective decisions in defining host range, such as interpreting subtle phenotypic variation in symptoms that could indicate partial overlap of host ranges. However, to enable progress with the *Pseudomonas*-legume pathosystem, we chose cultivars of each host plant (except mung bean) that are fully susceptible to all defined races of each particular pathovar in order to limit the influence of ETI. Conversely, *P. syringae* pv. *phaseolicola* 1448a and *P. syringae* pv. *glycinea* R4 are also virulent on all tested cultivars of French bean and soybean, respectively. Although the magnitude of differences between these isolates may change in a cultivar-dependent way, we focused on the virulence potential of each isolate on a particular host and highlight changes with the potential to modify host range between these isolates.

As noted in the introduction, NHR is considered to be a complex phenotype involving defense induced by recognition of either PAMPs or translocated effector proteins. The fact that *hrpL* and *hrcC* mutants of *P. syringae* pv. *phaseolicola* 1448a grew to higher levels than mutant counterparts from either *P. syringae* pv. *tomato* DC3000 (a tomato pathogen) or *P. syringae* B728a (a distantly related bean pathogen) on French bean demonstrates that *P. syringae* pv. *phaseolicola* 1448a has adapted to its bean host in an *hrpL*- and type III secretion-independent manner (Fig. 2A), and also suggests a fundamental difference in virulence mechanisms on the same host between two well-studied isolates of pvs. *phaseolicola* and *syringae*. It is possible that *P. syringae* pv. *phaseolicola* 1448a PAMPs have evolved to better evade recognition by bean PRR, although the full complement of *P. syringae* pv. *phaseolicola* 1448a T3E may still be weakly recognized and generate basal

defenses on French bean. *P. syringae* pv. *phaseolicola* 1448a may also be better adapted to use nutrients or survive within the bean apoplast, even in the absence of a type III secretion system. Nevertheless, these growth differences are *hrpL* independent, likely *gacAS* independent (Ortiz-Martín et al. 2010), and at least partially conserved in *P. syringae* pv. *phaseolicola* 1644R and *P. syringae* pv. *glycinea* R4 (Fig. 2B). Comparisons of signatures of evolution within the genomes of these isolates in connection with measurement of growth *hrpL* mutants for other group III isolates could allow for identification of additional virulence factors.

Although both *P. syringae* pv. *glycinea* R4 and *P. syringae* pv. *phaseolicola* 1644R grew more poorly than *P. syringae* pv. *phaseolicola* 1448a on French bean, only *P. syringae* pv. *glycinea* R4 triggered a hypersensitive cell death response when inoculated into French bean pods (Fig. 4A). ETI has previously been shown to contribute to NHR across closely related strains (Almeida et al. 2009). Thus, we sought to identify avirulence factors from this isolate by cloning all T3E from *P. syringae* pv. *glycinea* R4 that are not found within *P. syringae* pv. *phaseolicola* 1644R or 1448a, and determining whether these T3E affected growth when expressed in *P. syringae* pv. *phaseolicola* 1448a. Arnold and associates (2001) have previously shown that the *hopC1* allele from *P. syringae* pv. *pisi* acts as an avirulence factor on all French bean cultivars tested. Therefore, it is not surprising that the *P. syringae* pv. *glycinea* R4 allele of *hopC1* was recognized by French bean Canadian Wonder. *P. syringae* pv. *glycinea* R4 contains two copies of the *hopC1/hopH1* locus (Baltrus et al. 2011), and this locus is absent from all other known group III pathovars of *P. syringae*. This is evidence for recent acquisition by *P. syringae* pv. *glycinea* R4 and suggests that *hopC1* or *hopH1* may play an important virulence role in some context during natural infection of soybean. However, we were unable to find a role for these genes after hand inoculation of soybean leaves under laboratory conditions; we found that *P. syringae* pv. *glycinea* R4 and DAB890 (in which both copies of the *hopC1/hopH1* operon were deleted) grew to similar levels on ‘Merit’ soybean (data not shown).

We also found that HopC1 can act as an avirulence factor in mung bean (Supplementary Fig. S2), suggesting conservation of a resistance gene across a wide variety of *Phaseolus vulgaris* cultivars as well as across legume species (Arnold et al. 2001). This may explain why *P. syringae* pv. *glycinea* R4 grows equally poorly on both French bean and mung bean, whereas *P. syringae* pv. *phaseolicola* 1644R grows to intermediate levels on French bean and soybean. Therefore, acquisition of *hopC1* could be an event that differentiates pv. *glycinea* from pv. *phaseolicola*. We also found that *P. syringae* pv. *glycinea* R4 contains an additional factor with potential to restrict growth in French bean, because translocation of full-length HopM1 leads to lower growth of *P. syringae* pv. *phaseolicola* 1448a. This is surprising because the *hopM1* locus is thought to be a fundamental virulence factor present within all pathogenic strains of *P. syringae*, and is not known to be recognized by any other host plants despite being independently truncated within multiple isolates, (Baltrus et al. 2011; Cai et al. 2011).

Despite being the least virulent of our three isolates on the tested cultivars of French bean and mung bean, *P. syringae* pv. *glycinea* R4 contains a T3E (*avrPto*) that increases growth of *P. syringae* pv. *phaseolicola* 1448a on French bean Canadian Wonder (Fig. 4). Because AvrPto suppresses upstream steps in plant defense by binding to and inhibiting the action of the FLS2-BAK PRR complex (Xiang et al. 2008), this could indicate that *P. syringae* pv. *phaseolicola* 1448a growth is ultimately still limited by the French bean immune system in a way that is not overcome by its repertoire of T3E. It is also possible that the *P.*

syringae pv. *glycinea* R4 allele of *avrPto* acts more rapidly than the *P. syringae* pv. *phaseolicola* 1448a T3E suite to block plant defenses, or masks “weak” ETI dependent on recognition of a T3E. The possibility of weak ETI constraining multiplication of virulent pathogens has been reported previously. For example, *P. syringae* pv. *tomato* DC3000 grows to higher levels on *rar1-21 Arabidopsis* plants and RAR1 functions only as a chaperone to stabilize steady-state levels of the NB-LRR receptors required for ETI (Holt et al. 2005).

This result is intriguing because *P. syringae* pv. *phaseolicola* 1448a contains an active version of *hopABI*, alleles of which have been shown to target the FLS2-BAK PRR complex (Shan et al. 2008). *HopABI* is considered part of the same redundant effector group (REG) as *AvrPto* and, thus, thought to suppress the same level of the plant immune response (Cunnac et al. 2011). Although components of an REG may act somewhat independently to disrupt conserved defense pathways, evolutionary divergence may also result in different roles for these T3E in different host contexts. That *avrPto-hopAB2* affect similar defense pathways in *Arabidopsis* and tomato does not necessarily mean that *avrPto-hopABI* have the same functions in French bean or soybean. Similar to *hopCI*, this allele of *avrPto* appears to be a recent import into *P. syringae* pv. *glycinea* R4 because it is not present within any other group III isolate and may signal differences in virulence strategies between pvs. *glycinea* and *phaseolicola*.

Disruption of all avirulence genes identified above from *P. syringae* pv. *glycinea* R4 for French bean Canadian Wonder did not increase growth of this isolate on this host (Fig. 5A). Therefore, growth differences between *P. syringae* pv. *glycinea* R4 and *P. syringae* pv. *phaseolicola* 1448a must be explained by the presence of additional avirulence factors or the absence of virulence factors (Lorang et al. 1994). To explore these possibilities, we singly added all T3E present in *P. syringae* pv. *phaseolicola* 1448a but absent in *P. syringae* pv. *glycinea* R4 to strain DAB890 (*P. syringae* pv. *glycinea* R4 in which the *hopCI-hopH1* operons and *hopMI* have been deleted) and measured growth on French bean Canadian Wonder.

AvrB2 from *P. syringae* pv. *phaseolicola* 1448a did increase growth of this isolate on French bean (Fig. 5B and C). This difference is consistent, although more subtle, when expression of *avrB2* was driven by its native promoter within *P. syringae* pv. *glycinea* R4 containing the resident two copies of *hopCI* as well as *hopMI*. *AvrB2* has previously been shown to mask the avirulence effect of *HopF1* in French bean within *P. syringae* pv. *phaseolicola* 1449b (Tsiamis et al. 2000) but *hopF1* is absent from *P. syringae* pv. *glycinea* R4. Furthermore, *avrB2* does not appear to increase the growth of *P. syringae* pv. *phaseolicola* 1644R on French bean (Fig. 5D), so that the virulence function of this T3E is either compensated for by other virulence genes within *P. syringae* pv. *phaseolicola* 1644R or is epistatic to an alternative *P. syringae* pv. *glycinea* R4 specific factor.

It has recently been proposed that the dominant mode of recognition progresses from ETI to PTI as the phylogenetic divergence between host species increases (Schulze-Lefert and Panstruga 2011). In this comparison between three closely related isolates, recognition of T3E plays a limited role over relatively short divergence times because there is no evidence that *P. syringae* pv. *phaseolicola* 1644R contains T3E recognized by French bean (Fig. 4A). Moreover, absence of single T3E does not explain the virulence deficit for *P. syringae* pv. *phaseolicola* 1644R (Fig. 5D). Further investigation will explore whether allelic differences and expression differences in the T3E suites or differences in metabolic capabilities between these two isolates contributes toward adaptation to their respective hosts of isolation. Over longer time scales (*P. syringae* pv. *glycinea* R4 to *P. syringae* pv. *phaseolicola* 1448a), there are multiple events

leading to recognition of *P. syringae* pv. *glycinea* R4 on French bean. Although acquisition of *HopC1* by *P. syringae* pv. *glycinea* R4 limits growth on both French bean and mung bean, the default state for many *P. syringae* strains on French bean may be NHR due to recognition of the conserved T3E *hopMI*. Taken as a whole, our study suggests that the comparisons of whole genome sequences within a phylogenetic context can provide insight into fine-scale host range differences, and that some of the functional complexity of host range evolution in plant–microbe interactions can be observed even among closely related host and pathogen species.

MATERIALS AND METHODS

Plasmids, bacterial isolates, and growth conditions.

All plasmids used or created are listed in Table 1, while the bacterial isolates are listed in Table 2. Typically, *P. syringae* isolates were grown at 27°C on King’s B (KB) media using rifampicin at 50 µg/ml. When necessary, cultures of both *P. syringae* and *Escherichia coli* were supplemented with antibiotics or sugars in the following concentrations: tetracycline at 10 µg/ml, kanamycin at 50 µg/ml, gentamycin at 25 µg/ml, and 5% sucrose.

All clones were created by first amplifying target sequences using *Pfx* polymerase (Invitrogen, Carlsbad, CA, U.S.A.), followed by recombination of these fragments into the entry vector pDONR207 using BP clonase (Invitrogen). All open reading frame (ORF) (without an *hrp*-box promoter) and gene (including the promoter) sequences were confirmed by Sanger sequencing of these pDONR207 clones. ORF clones always included stop codons at the end of the sequence but gene clones did not, and the start codons for all ORF sequences were based on previous annotations, comparison across alleles, and proximity to the *hrp* box. Known chaperones for some T3E (*shcM* and *shcV*) were included in both ORF and gene clones. Clones in entry vectors were recombined into destination vectors of interest using LR clonase (Invitrogen).

Generation of mutants.

Bacterial mutants were generated using a modified suicide plasmid protocol (Marco et al. 2005). Regions (>500 bp) upstream and downstream of the target genes were polymerase chain reaction (PCR) amplified separately and then combined into one fragment by overlap extension PCR. The bridge PCR amplicon was then cloned into pDONR207, and further moved into MTN1907 using LR clonase. MTN1907 is a modified version of the pLVC-D suicide plasmid (Marco et al. 2005), providing tetracycline resistance as well as sucrose counter-selection from *sacB* cloned downstream of a *trp* promoter (giving constitutive *sacB* expression). MTN1907 also includes an *nptII* gene providing resistance to kanamycin but we note that expression from this locus is not consistent in *P. syringae*. Once mated into *P. syringae*, single recombination of a homologous region upstream or downstream of the target region and subsequent selection on sucrose allows for screening of clean deletions. Mutants were confirmed by phenotyping for sucrose resistance, tetracycline sensitivity, PCR amplification of the deletion, and failure of PCR to amplify regions within the deletion.

Sequencing and annotation of *P. syringae* pv. *phaseolicola* 1644R.

P. syringae pv. *glycinea* R4 is a race 4 isolate of pv. *glycinea* obtained from the Staskawicz lab (Baltrus et al. 2011), while the history of *P. syringae* pv. *phaseolicola* 1448a is well documented (Arnold et al. 2011) and was originally collected by Habtu Assefa in Ethiopia in 1985. *P. syringae* pv. *phaseolicola*

1644 (NCPBB 2435) was originally isolated from *V. radiata* in the United States in 1971. A rifampicin-resistant mutant of *P. syringae* pv. *phaseolicola* 1644 was generated through selection, and one colony of this isolate was picked and grown overnight in 250 ml of KB liquid media to generate genomic DNA for sequencing. Genomic DNA was collected using the cetyltrimethylammonium bromide extraction method. One lane of sequencing on an Illumina GAI using a standard protocol for paired end library generation (yielding 18,862,298 total reads of 36 bp) was used to generate a draft genome sequence of *P. syringae* pv. *phaseolicola* 1644R. Reads were assembled de novo using SoapDenovo (Li et al. 2010) and custom Perl scripts that optimized the assembly for number of contigs and mean contig size. The best assembly was generated with a kmer size of 32. Reads were not culled for quality before the assembly. The nucleotide sequences of contigs from this assembly can be found under GenBank accession AGAS00000000.

It is difficult to precisely annotate lower-quality draft genomes, which often contain gaps and ORF fragments. There-

fore, we used a progression of scripts and filters to quantify a putative number of proteins present within *P. syringae* pv. *phaseolicola* 1644R that were absent from *P. syringae* pv. *glycinea* R4 or *P. syringae* pv. *phaseolicola* 1488a. We first used Genemark (Besemer and Borodovsky 2005) to call ORF from the draft genome, screened for ORF > 50 amino acids, and used BLASTp versus the *P. syringae* pv. *phaseolicola* 1448a proteome (Joardar et al. 2005) (screened for exact matches in amino acid identity [AAID]) to identify proteins that were duplicated or truncated due to poor assembly quality. Once these redundant features were culled, we used BLASTp (*e* value = 1e-5, AAID > 85%) to identify loci absent from the *P. syringae* pv. *glycinea* R4 and *P. syringae* pv. *phaseolicola* 1448a proteomes (Baltrus et al. 2011). We then used tBLASTn (*e* value = 1e-5, AAID > 85%) against both the *P. syringae* pv. *glycinea* R4 and *P. syringae* pv. *phaseolicola* 1448a genomes to further narrow the list of novel ORF. Finally, we used BLASTp (*e* value = 1e-5, no cutoff) against the NR database to screen out redundant *P. syringae* pv. *phaseolicola* 1644R ORF and create a final list of novel proteins. Although these ORF are poten-

Table 1. Strains

Strain	Description	Reference
<i>Pseudomonas syringae</i> pv. <i>phaseolicola</i> 1448a	<i>P. syringae</i> pv. <i>phaseolicola</i> race 6 isolate, rifampicin resistant	Joardar et al. 2005; Taylor et al. 1996
<i>P. syringae</i> pv. <i>glycinea</i> R4	<i>P. syringae</i> pv. <i>glycinea</i> race 4 isolate, rifampicin resistant	Baltrus et al. 2011; Staskawicz et al. 1984
<i>P. syringae</i> pv. <i>phaseolicola</i> 1644R	<i>P. syringae</i> pv. <i>phaseolicola</i> race 1 isolate, rifampicin resistant	This article, derived from Taylor et al. 1996
DAB007	<i>P. syringae</i> pv. <i>tomato</i> DC3000 <i>hrcC</i> mutant	Roine et al. 1997
DAB087	<i>P. syringae</i> pv. <i>syringae</i> B728a <i>hrcC</i> mutant	Hirano et al. 1999
DAB093	<i>P. syringae</i> pv. <i>phaseolicola</i> race 4 strain 1448a <i>hrcC</i> mutant, rifampicin resistant	This article, derived from Ortiz-Martin et al. 2006
DAB786	<i>P. syringae</i> pv. <i>phaseolicola</i> 1448a <i>hrpL</i> clean deletion, rifampicin resistant	This article
DAB963	<i>P. syringae</i> pv. <i>tomato</i> DC3000 <i>hrpL</i> clean deletion, rifampicin resistant	This article
<i>P. syringae</i> B728a <i>hrpL</i>	Obtained from Dr. Gwyn Beattie	Unpublished
DAB886	<i>P. syringae</i> pv. <i>glycinea</i> R4 isolate with <i>hrpL</i> clean deletion, rifampicin resistant	This article
DAB890	<i>P. syringae</i> pv. <i>glycinea</i> R4 isolate with 2x <i>hopC1/hopH1</i> and <i>shcM1/hopM1</i> clean deletion	This article
DAB907	<i>P. syringae</i> pv. <i>phaseolicola</i> 1644R with <i>hrpL</i> clean deletion, rifampicin resistant	This article
DAB748	<i>P. syringae</i> pv. <i>phaseolicola</i> 1448a with pDAB256	This article
DAB809	<i>P. syringae</i> pv. <i>phaseolicola</i> 1448a with pDAB245	This article
DAB784	<i>P. syringae</i> pv. <i>phaseolicola</i> 1448a with pDAB363	This article
DAB775	<i>P. syringae</i> pv. <i>phaseolicola</i> 1448a with pDAB267	This article
DAB771	<i>P. syringae</i> pv. <i>phaseolicola</i> 1448a with pDAB272	This article
DAB804	<i>P. syringae</i> pv. <i>phaseolicola</i> 1448a with pDAB289	This article
DAB774	<i>P. syringae</i> pv. <i>phaseolicola</i> 1448a with pDAB268	This article
DAB828	<i>P. syringae</i> pv. <i>phaseolicola</i> 1448a with pDAB296	This article
DAB831	<i>P. syringae</i> pv. <i>phaseolicola</i> 1644R with pDAB296	This article
DAB767	<i>P. syringae</i> pv. <i>phaseolicola</i> 1644R with pDAB269	This article
DAB768	<i>P. syringae</i> pv. <i>phaseolicola</i> 1644R with pDAB270	This article
DAB799	<i>P. syringae</i> pv. <i>phaseolicola</i> 1644R with pDAB271	This article
DAB806	<i>P. syringae</i> pv. <i>phaseolicola</i> 1644R with pDAB286	This article
DAB750	<i>P. syringae</i> pv. <i>phaseolicola</i> 1644R with pDAB250	This article
DAB830	<i>P. syringae</i> pv. <i>phaseolicola</i> 1644R with pDAB254	This article
DAB183	<i>P. syringae</i> pv. <i>phaseolicola</i> 1448a with pDAB040	This article
DAB524	<i>P. syringae</i> pv. <i>phaseolicola</i> 1448a with pDAB151	This article
DAB182	<i>P. syringae</i> pv. <i>phaseolicola</i> 1448a with pDAB039	This article
DAB892	<i>P. syringae</i> pv. <i>phaseolicola</i> 1448a with pDAB334	This article
DAB954	<i>P. syringae</i> pv. <i>glycinea</i> R4 with pDAB334	This article
DAB960	<i>P. syringae</i> pv. <i>glycinea</i> R4 with MTN500	This article
DAB939	DAB890 with pDAB269	This article
DAB943	DAB890 with pDAB296	This article
DAB940	DAB890 with pDAB270	This article
DAB941	DAB890 with pDAB250	This article
DAB937	DAB890 with pDAB287	This article
DAB936	DAB890 with pDAB339	This article
DAB938	DAB890 with pDAB341	This article
DAB945	DAB890 with pDAB286	This article

tially full length, we emphasize that the number of true ORF within *P. syringae* pv. *phaseolicola* 1644R may differ due to inclusions or exclusions of truncated loci as well as inclusion of noncoding regions.

Phylogenetics.

To resolve evolutionary relationships among these three isolates (and including *P. syringae* pv. *mori* MAFF301020 as an outgroup) (Baltrus et al. 2011), we created amino acid sequence-based trees using a database of 2,400 orthologous genes shared by all four isolates. We required reciprocal best hits, with 80% amino acid identity for greater than 80% of the length and an *e* value < 1 × 10⁻²⁰⁰. For each apparent ortholog, we used ProbCons (Do 2005), a probabilistic consistency algorithm that combines sum-of-pairs scoring and HMM-derived posterior probabilities, to produce a consensus alignment for all orthologs of a gene. For each gene, we performed a model test to determine the best amino acid substitution model. We then constructed individual trees for each protein sequence using RAXML. RAXML is a maximum likelihood-based tool for large phylogenetic trees,

and was optimized for running on our computers (Stamatakis 2004, 2006). A majority rules consensus tree was created from all of these individual trees for individual loci using the program Consense within the Phylip package. Consense was then used to create a majority rules consensus tree from phylogenies for all 2,400 loci. In order to display branch lengths on the phylogeny in Figure 1A, we chose a subset of loci whose phylogenetic patterns matched that of the consensus tree of all conserved genes (locus numbers in the *P. syringae* pv. *phaseolicola* 1448a genome: PSPPH_0035, PSPPH_0196, PSPPH_0405, PSPPH_0450, PSPPH_0737, PSPPH_1378, PSPPH_1507, PSPPH_2368, PSPPH_2957, PSPPH_4280, PSPPH_4340, and PSPPH_4623), concatenated these sequences together, aligned with ClustalX, and created a Bayesian phylogeny using Mr. Bayes 3.1.2. To create a phylogeny for the bean species, we obtained sequences of the *matK* gene from *Phaseolus vulgaris* (GenBank number EU196765.1), *G. max* (AF142700.1), and *V. radiata* (JN008226.1), with *Desmodium psilocarpum* (AY386896.1) as an outgroup. Once sequences were aligned with ClustalX, a Bayesian phylogeny was created using Mr. Bayes 3.1.2.

Table 2. Plasmids

Plasmid	Description	Antibiotics ^a	Reference, source
MTN1907	Gateway destination vector for making clean deletions in <i>P. syringae</i>	Tet ^r , Kan ^r , Cam ^r , Suc ^s	This article
pDONR207	Gateway entry vector from Invitrogen	Gent ^r , Cam ^r	Invitrogen ^b
MTN1970-2	3' Untranslated region of <i>Arabidopsis</i> HSP18.2 in pDONR207	Gent ^r	This article
pDAB255	<i>avrpto</i> open reading frame (ORF) with stop codon in pDONR207	Gent ^r	This article
pDAB238	<i>efe</i> ORF with stop codon in pDONR207	Gent ^r	This article
pDAB281	<i>laaL</i> ORF with stop codon in pDONR207	Gent ^r	This article
MTN1382	<i>hopH1</i> ORF with stop codon in pDONR207	Gent ^r	This article
pDAB059	<i>hopZ1</i> ORF with stop codon in pDONR207	Gent ^r	This article
pDAB283	<i>shcM/hopM1</i> ORF with stop codon in pDONR207	Gent ^r	This article
pDAB061	<i>hopC1</i> ORF with stop codon in pDONR207	Gent ^r	This article
pDAB261	<i>avrB2</i> ORF with stop codon in pDONR207	Gent ^r	This article
pDAB262	<i>avrB4-1</i> ORF with stop codon in pDONR207	Gent ^r	This article
pDAB249	<i>hopAW1</i> ORF with stop codon in pDONR207	Gent ^r	This article
pDAB285	<i>hopQ1</i> ORF with stop codon in pDONR207	Gent ^r	This article
pDAB336	<i>hopAF1</i> ORF with stop codon in pDONR207	Gent ^r	This article
pDAB297	<i>hopAF1</i> ORF with stop codon in pDONR207	Gent ^r	This article
pDAB282	<i>shcV/hopV1</i> ORF with stop codon in pDONR207	Gent ^r	This article
pDAB263	<i>hopAB1</i> ORF with stop codon in pDONR207	Gent ^r	This article
pDAB253	<i>avrRps4</i> ORF in pDONR207	Gent ^r	This article
pBV226	Gateway destination vector with expression driven by <i>nptII</i> promoter	Tet ^r , Cam ^r	Vinatzer et al. 2006
pDAB256	<i>avrpto</i> ORF with stop codon in pBV226	Tet ^r	This article
pDAB245	<i>efe</i> ORF with stop codon in pBV226	Tet ^r	This article
pDAB363	<i>laaL</i> ORF with stop codon in pBV226	Tet ^r	This article
pDAB267	<i>hopH1</i> ORF with stop codon in pBV226	Tet ^r	This article
pDAB272	<i>hopZ1</i> ORF with stop codon in pBV226	Tet ^r	This article
pDAB289	<i>shcM/hopM1</i> ORF with stop codon in pBV226	Tet ^r	This article
pDAB268	<i>hopC1</i> ORF with stop codon in pBV226	Tet ^r	This article
pDAB296	MTN1970-2 EV construct in pBV226	Tet ^r	This article
pDAB269	<i>avrB2</i> ORF with stop codon in pBV226	Tet ^r	This article
pDAB270	<i>avrB4-1</i> ORF with stop codon in pBV226	Tet ^r	This article
pDAB250	<i>hopAW1</i> ORF with stop codon in pBV226	Tet ^r	This article
pDAB287	<i>hopQ1</i> ORF with stop codon in pBV226	Tet ^r	This article
pDAB339	<i>hopAF1</i> ORF with stop codon in pBV226	Tet ^r	This article
pDAB341	<i>hopAF1</i> ORF with stop codon in pBV226	Tet ^r	This article
pDAB286	<i>shcV/hopV1</i> ORF with stop codon in pBV226	Tet ^r	This article
pDAB271	<i>hopAB1</i> ORF with stop codon in pBV226	Tet ^r	This article
pDAB254	<i>avrRps4</i> ORF with stop codon in pBV226	Tet ^r	This article
SMPgy5172	<i>hrp</i> gene with <i>hrp</i> -box promoter and no stop codon in pDONR207	Gent ^r	This article
pDAB081	<i>shcM/hopM1</i> gene with <i>hrp</i> -box promoter and no stop codon in pDONR207	Gent ^r	This article
SMPgy5168	<i>hopC1</i> gene with <i>hrp</i> -box promoter and no stop codon in pDONR207	Gent ^r	This article
MTN366	<i>avrB2</i> gene with <i>hrp</i> -box promoter and no stop codon in pDONR207	Gent ^r	This article
pJC531	Gateway destination vector based off of pBBR1-MCS2 no promoter	Kan ^r , Cam ^r	Chang et al. 2005
pDAB040	<i>avrpto</i> gene with <i>hrp</i> -box promoter and no stop codon in pJC531	Kan ^r	This article
pDAB151	<i>shcM/hopM1</i> gene with <i>hrp</i> -box promoter and no stop codon in pJC531	Kan ^r	This article
pDAB039	<i>hopC1</i> gene with <i>hrp</i> -box promoter and no stop codon in pJC531	Kan ^r	This article
pDAB334	MTN1970-2 EV construct in pJC531	Kan ^r	This article
pDAB49	<i>avrB2</i> gene with <i>hrp</i> -box promoter and no stop codon in pJC531	Kan ^r	This article

^a Tet^r, Kan^r, Cam^r, and Gent^r = resistance to tetracycline, kanamycin, chloramphenicol, and gentamycin; Suc^s = susceptible to sucrose.

^b Invitrogen, Carlsbad, CA, U.S.A.

Comparison of virulence genes across *P. syringae* pv. *phaseolicola* 1448a, *P. syringae* pv. *glycinea* R4, and *P. syringae* pv. *phaseolicola* 1644R.

We searched the draft genome sequence of *P. syringae* pv. *phaseolicola* 1644R for virulence genes by tBLASTn using protein sequences of known T3E and toxin pathways as queries (same list as given by Baltrus and associates [2011]). There were no cutoffs used for this initial search, and all potential hits were further confirmed manually. When only fragments of T3E genes could be identified within the draft genome assembly, completeness of these loci was gauged by using SOAP (Li et al. 2008) to align Illumina reads to the most closely related allele of each virulence gene, with sequences verified by hand. Truncations within candidate virulence genes were confirmed by targeted Sanger sequencing. Additionally, *P. syringae* pv. *phaseolicola* 1644R-specific proteins were bioinformatically screened for translocation potential using the SIEVE server (McDermott et al. 2011). If a duplicated member of a T3E family was absent (i.e., *avrB4-2* in *P. syringae* pv. *phaseolicola* 1644R) or there was only an allelic variant of a T3E family, these specific alleles were considered to be absent.

Growth curves and pod inoculation.

Pod inoculation was done as described by Tsiamis and associates (2000). Briefly, fresh bean pods (before seed clearly developed) were harvested from plants grown under greenhouse conditions and pricked with toothpicks that had been dipped in a 10 mM MgCl₂ solution containing bacteria at an optical density at 600 nm (OD₆₀₀) of 0.5. Pods were kept under continuous light and moderate humidity for 3 days, after which symptoms were photographed.

To measure bacterial growth in planta, French bean and soybean were grown in a growth chamber under long-day conditions (18 h of light and 6 h of darkness) while mung bean were grown in the greenhouse under similar light conditions. Growth conditions differ across bean species because we found that mung bean plants grown in the greenhouse were able to tolerate hand inoculation whereas those grown in the growth chamber under conditions used for the other legume species were easily crushed. Experiments for Figure 5A and 5 also took place under greenhouse conditions. All bacterial growth measurements on all plant species were performed on primary leaves from 2-week-old plants. In each case, bacteria were grown overnight in liquid cultures, centrifuged, rinsed with 10 mM MgCl₂, and resuspended in 10 mM MgCl₂ at OD₆₀₀ = 0.0001 (approximately 10⁴ cells/ml). Bacterial suspensions were inoculated into leaves using a blunt-end syringe in the afternoon (between 2:00 and 6:00 p.m., lights out at approximately 10:00 p.m. in both growth chambers and greenhouses). Four independent replicates of inoculum population size were measured for each strain within each experiment. For ease of comparison across strains within figures, and because day 0 bacterial population sizes were largely indistinguishable within most experiments, population sizes for inoculum are listed in Supplementary Table S1. In order to conserve plants, day 0 bacterial population sizes were measured from the inoculum for all experiments but the *hrpL* and *hrcC* analyses; levels in planta are roughly 50- to 100-fold lower than in the inoculum. Bacterial populations were measured after 4 days of growth within French bean and soybean and after 5 days of growth in mung bean because these were the days that showed maximum levels of growth for each specific pathogen in pilot experiments (data not shown). Four cores (1 cm²) were taken on each half of a leaf, with two total leaves per plant sampled (16 cores total, four independent measurements per plant). At least three independent experiments were performed

per isolate, except that only two experiments were measured for Figure 5A and C. Data for each experiment were statistically explored by analysis of variance, with isolate as a fixed factor, experiment as a random factor, and plant as a random factor nested within isolate and experiment. If there was a significant effect of isolate in the model, means were compared by post hoc analysis using Tukey's highly significant difference and an α value of 0.05.

ACKNOWLEDGMENTS

Funding was provided by National Institute of Health (NIH) grant GM066025 and Department of Energy Basic Energy Sciences grant 95ER20187 to J. L. Dangl, an NIH National Research Service Ruth Kirchstein Award GM082279-01 and startup funds from the University of Arizona to D. A. Baltrus, and from the Arnold and Mabel Beckmann foundation to S. Biswas. J. L. Dangl is an Investigator of the Howard Hughes Medical Institute. We thank C. Jones for technical support with genome assembly and processing; P. Miecskowski, H. Kelkar, and J. Walsh of the UNC-CH HTS core for technical assistance; A. Jones for providing the pLVC-D plasmid; G. Beattie for the *P. syringae* B728a *hrpL* mutant; and J. Taylor (NVRS-Wellesbourne, retired) for providing historical background on isolates from his *P. syringae* pv. *phaseolicola* collection.

LITERATURE CITED

Abramovitch, R. B., Anderson, J. C., and Martin, G. B. 2006. Bacterial elicitation and evasion of plant innate immunity. *Nat. Rev. Mol. Cell Biol.* 7:601-611.

Almeida, N. F., Yan, S., Lindeberg, M., Studholme, D. J., Schneider, D. J., Condon, B., Liu, H., Viana, C. J., Warren, A., Evans, C., Kemen, E., Maclean, D., Angot, A., Martin, G. B., Jones, J. D., Collmer, A., Setubal, J. C., and Vinatzer, B. A. 2009. A draft genome sequence of *Pseudomonas syringae* pv. *tomato* T1 reveals a type III effector repertoire significantly divergent from that of *Pseudomonas syringae* pv. *tomato* DC3000. *Mol. Plant-Microbe Interact.* 22:52-62.

Arnold, D. L., Jackson, R. W., Fillingham, A. J., Goss, S. C., Taylor, J. D., Mansfield, J. W., and Vivian, A. 2001. Highly conserved sequences flank avirulence genes: Isolation of novel avirulence genes from *Pseudomonas syringae* pv. *pisi*. *Microbiology* 147:1171-1182.

Arnold, D. L., Lovell, H. C., Jackson, R. W., and Mansfield, J. W. 2011. *Pseudomonas syringae* pv. *phaseolicola*: From 'has bean' to supermodel. *Mol. Plant Pathol.* 12:617-627.

Baltrus, D. A., Nishimura, M. T., Romanchuk, A., Chang, J. H., Mukhtar, M. S., Cherkis, K., Roach, J., Grant, S. R., Jones, C. D., and Dangl, J. L. 2011. Dynamic evolution of pathogenicity revealed by sequencing and comparative genomics of 19 *Pseudomonas syringae* isolates. *PLoS Pathog.* 7:e1002132. Published online

Besemer, J., and Borodovsky, M. 2005. GeneMark: Web software for gene finding in prokaryotes, eukaryotes and viruses. *Nucleic Acids Res.* 33:W451-W454.

Borhan, M. H., Gunn, N., Cooper, A., Gulden, S., Tör, M., Rimmer, S. R., and Holub, E. B. 2008. WRR4 encodes a TIR-NB-LRR protein that confers broad-spectrum white rust resistance in *Arabidopsis thaliana* to four physiological races of *Albugo candida*. *Mol. Plant-Microbe Interact.* 21:757-768.

Cai, R., Lewis, J., Yan, S., Liu, H., Clarke, C. R., Campanile, F., Almeida, N. F., Studholme, D. J., Lindeberg, M., Schneider, D., Zaccardelli, M., Setubal, J. C., Morales-Lizcano, N. P., Bernal, A., Coaker, G., Baker, C., Bender, C. L., Leman, S., and Vinatzer, B. A. 2011. The plant pathogen *Pseudomonas syringae* pv. *tomato* is genetically monomorphic and under strong selection to evade tomato immunity. *PLoS Pathog.* 7:e1002130. Published online.

Chang, J. H., Urbach, J. M., Law, T. F., Arnold, L. W., Hu, A., Gombar, S., Grant, S. R., Ausubel, F. M., and Dangl, J. L. 2005. A high-throughput, near-saturating screen for type III effector genes from *Pseudomonas syringae*. *Proc. Natl. Acad. Sci. U.S.A.* 102:2549-2554.

Cunnac, S., Chakravarthy, S., Kvitko, B. H., Russell, A. B., Martin, G. B., and Collmer, A. 2011. Genetic disassembly and combinatorial reassembly identify a minimal functional repertoire of type III effectors in *Pseudomonas syringae*. *Proc. Natl. Acad. Sci. U.S.A.* 108:2975-2980.

Delgado-Salinas, A., Bibler, R., and Lavin, M. 2006. Phylogeny of the genus *Phaseolus* (Leguminosae): A recent diversification in an ancient landscape. *Syst. Bot.* 31:779-791.

Do, C. B. 2005. ProbCons: Probabilistic consistency-based multiple sequence alignment. *Genome Res.* 15:330-340.

Fan, J., Crooks, C., and Lamb, C. 2007. High-throughput quantitative lu-

- minescence assay of the growth in planta of *Pseudomonas syringae* chromosomally tagged with *Photorhabdus luminescens luxCDABE*. *Plant J.* 53:393-399.
- Ferrante, P., Clarke, C. R., Cavanaugh, K. A., Michelmore, R. W., Buonaurio, R., and Vinatzer, B. A. 2009. Contributions of the effector gene *hopQ1-1* to differences in host range between *Pseudomonas syringae* pv. *phaseolicola* and *P. syringae* pv. *tabaci*. *Mol. Plant Pathol.* 10:837-842.
- Gardan, L., Shafik, H., Belouin, S., Broch, R., Grimont, F., and Grimont, P. A. 1999. DNA relatedness among the pathovars of *Pseudomonas syringae* and description of *Pseudomonas tremae* sp. nov. and *Pseudomonas cannabina* sp. nov. (ex Sutic and Dowson 1959). *Int. J. Syst. Bacteriol.* 492:469-478.
- Gassmann, W., Hirsch, M. E., and Staskawicz, B. J. 1999. The *Arabidopsis* RPS4 bacterial-resistance gene is a member of the TIR-NBS-LRR family of disease-resistance genes. *Plant J.* 20:265-277.
- Glickmann, E., Gardan, L., Jacquet, S., Hussain, S., Elasmri, M., Petit, A., and Dessaux, Y. 1998. Auxin production is a common feature of most pathovars of *Pseudomonas syringae*. *Mol. Plant-Microbe Interact.* 11:156-162.
- Ham, J. H., Kim, M. G., Lee, S. Y., and Mackey, D. 2007. Layered basal defenses underlie non-host resistance of *Arabidopsis* to *Pseudomonas syringae* pv. *phaseolicola*. *Plant J.* 51:604-616.
- Hirano, S. S., Charkowski, A. O., Collmer, A., Willis, D. K., and Upper, C. D. 1999. Role of the Hrp type III protein secretion system in growth of *Pseudomonas syringae* pv. *syringae* B728a on host plants in the field. *Proc. Natl. Acad. Sci. U.S.A.* 96:9851-9856.
- Holt, B. F., Belkhadir, Y., and Dangl, J. L. 2005. Antagonistic control of disease resistance protein stability in the plant immune system. *Science* 309:929-932.
- Hwang, M. S. H., Morgan, R. L., Sarkar, S. F., Wang, P. W., and Guttman, D. S. 2005. Phylogenetic characterization of virulence and resistance phenotypes of *Pseudomonas syringae*. *Appl. Environ. Microbiol.* 71:5182-5191.
- Joardar, V., Lindeberg, M., Jackson, R. W., Selengut, J., Dodson, R., Brinkac, L. M., Daugherty, S. C., Deboy, R., Durkin, A. S., Giglio, M. G., Madupu, R., Nelson, W. C., Rosovitz, M. J., Sullivan, S., Crabtree, J., Creasy, T., Davidsen, T., Haft, D. H., Zafar, N., Zhou, L., Halpin, R., Holley, T., Khouri, H., Feldblyum, T., White, O., Fraser, C. M., Chatterjee, A. K., Cartinhour, S., Schneider, D. J., Mansfield, J., Collmer, A., and Buell, C. R. 2005. Whole-genome sequence analysis of *Pseudomonas syringae* pv. *phaseolicola* 1448A reveals divergence among pathovars in genes involved in virulence and transposition. *J. Bacteriol.* 187:6488-6498.
- Jones, J. D. G., and Dangl, J. L. 2006. The plant immune system. *Nature* 444:323-329.
- Kobayashi, D. Y., Tamaki, S. J., and Keen, N. T. 1989. Cloned avirulence genes from the tomato pathogen *Pseudomonas syringae* pv. *tomato* confer cultivar specificity on soybean. *Proc. Natl. Acad. Sci. U.S.A.* 86:157-161.
- Lacombe, S. E. V., Rougon-Cardoso, A., Sherwood, E., Peeters, N., Dahlbeck, D., van Esse, H. P., Smoker, M., Rallapalli, G., Thomma, B. P. H. J., Staskawicz, B., Jones, J. D. G., and Zipfel, C. 2010. Interfamily transfer of a plant pattern-recognition receptor confers broad-spectrum bacterial resistance. *Nat. Biotechnol.* 28:365-369.
- Li, R., Li, Y., Kristiansen, K., and Wang, J. 2008. SOAP: Short oligonucleotide alignment program. *Bioinformatics* 24:713-714.
- Li, R., Zhu, H., Ruan, J., Qian, W., Fang, X., Shi, Z., Li, Y., Li, S., Shan, G., Kristiansen, K., Li, S., Yang, H., Wang, J., and Wang, J. 2010. De novo assembly of human genomes with massively parallel short read sequencing. *Genome Res.* 20:265-272.
- Li, X., Lin, H., Zhang, W., Zou, Y., Zhang, J., Tang, X., and Zhou, J.-M. 2005. Flagellin induces innate immunity in nonhost interactions that is suppressed by *Pseudomonas syringae* effectors. *Proc. Natl. Acad. Sci. U.S.A.* 102:12990-12995.
- Lin, N.-C., and Martin, G. B. 2007. Pto- and Prf-mediated recognition of AvrPto and AvrPtoB restricts the ability of diverse *Pseudomonas syringae* pathovars to infect tomato. *Mol. Plant-Microbe Interact.* 20:806-815.
- Lindeberg, M., Cunnac, S., and Collmer, A. 2009. The evolution of *Pseudomonas syringae* host specificity and type III effector repertoires. *Mol. Plant Pathol.* 10:767-775.
- Lorang, J., Shen, H., and Kobayashi, D. 1994. *avrA* and *avrE* in *Pseudomonas syringae* pv. *tomato* PT23 play a role in virulence on tomato plants. *Mol. Plant-Microbe Interact.* 7:508-515.
- Marco, M. L., Legac, J., and Lindow, S. E. 2005. *Pseudomonas syringae* genes induced during colonization of leaf surfaces. *Environ. Microbiol.* 7:1379-1391.
- Marques, A., Corbiere, R., Gardan, L., Tourte, C., Manceau, C., Taylor, J., and Samson, R. 2000. Multiphasic approach for the identification of the different classification levels of *Pseudomonas savastanoi* pv. *phaseolicola*. *Eur. J. Plant Pathol.* 106:715-734.
- Mazzola, M., and White, F. F. 1994. A mutation in the indole-3-acetic acid biosynthesis pathway of *Pseudomonas syringae* pv. *syringae* affects growth in *Phaseolus vulgaris* and syringomycin production. *J. Bacteriol.* 176:1374-1382.
- McClellan, P., Lavin, M., Gepts, P., and Jackson, S. 2008. *Phaseolus vulgaris*: A diploid model for soybean. Pages 55-78 in: *Soybean Genomics*. Springer, Berlin.
- McDermott, J., Corrigan, A., and Peterson, E. 2011. Computational prediction of type III and IV secreted effectors in gram-negative bacteria. *Infect. Immun.* 79:23-32.
- Mellersh, D. G., and Heath, M. C. 2003. An investigation into the involvement of defense signaling pathways in components of the nonhost resistance of *Arabidopsis thaliana* to rust fungi also reveals a model system for studying rust fungal compatibility. *Mol. Plant-Microbe Interact.* 16:398-404.
- Morris, C. E., Sands, D. C., Vanneste, J. L., Montarry, J., Oakley, B., Guilbaud, C., and Glaux, C. 2010. Inferring the evolutionary history of the plant pathogen *Pseudomonas syringae* from its biogeography in headwaters of rivers in North America, Europe, and New Zealand. *mBio* 1:e00107-00110-e00107-00120. Published online.
- Mysore, K. S., and Ryu, C.-M. 2004. Nonhost resistance: How much do we know? *Trends Plant Sci.* 9:97-104.
- Ortiz-Martín, I., Macho, A. P., Lambersten, L., Ramos, C., and Beuzón, C. R. 2006. Suicide vectors for antibiotic marker exchange and rapid generation of multiple knockout mutants by allelic exchange in gram-negative bacteria. *J. Microbiol. Methods* 67:395-407.
- Ortiz-Martín, I., Thwaites, R., Macho, A. P., Mansfield, J. W., and Beuzón, C. R. 2010. Positive regulation of the Hrp type III secretion system in *Pseudomonas syringae* pv. *phaseolicola*. *Mol. Plant-Microbe Interact.* 23:665-681.
- Patil, S. S., Hayward, A., and Emmons, R. 1974. An ultraviolet-induced nontoxic mutant of *Pseudomonas phaseolicola* of altered pathogenicity. *Phytopathology* 64:590-595.
- Perchepped, L., Kroj, T., Tronchet, M., Loudet, O., and Roby, D. 2006. Natural variation in partial resistance to *Pseudomonas syringae* is controlled by two major QTLs in *Arabidopsis thaliana*. *PLoS ONE* 1:e123. Published online.
- Qi, M., Wang, D., Bradley, C. A., and Zhao, Y. 2011. Genome sequence analyses of *Pseudomonas savastanoi* pv. *glycinea* and subtractive hybridization-based comparative genomics with nine pseudomonads. *PLoS ONE* 6:e16451. Published online.
- Rico, A., and Preston, G. M. 2008. *Pseudomonas syringae* pv. *tomato* DC3000 uses constitutive and apoplast-induced nutrient assimilation pathways to catabolize nutrients that are abundant in the tomato apoplast. *Mol. Plant-Microbe Interact.* 21:269-282.
- Roine, E., Wei, W., Yuan, J., Nurmiaho-Lassila, E. L., Kalkkinen, N., Romantschuk, M., and He, S. Y. 1997. Hrp pilus: An hrp-dependent bacterial surface appendage produced by *Pseudomonas syringae* pv. *tomato* DC3000. *Proc. Natl. Acad. Sci. U.S.A.* 94:3459-3464.
- Schulze-Lefert, P., and Panstruga, R. 2011. A molecular evolutionary concept connecting nonhost resistance, pathogen host range, and pathogen speciation. *Trends Plant Sci.* 16:117-125.
- Shan, L., He, P., Li, J., Heese, A., Peck, S. C., Nürnberger, T., Martin, G. B., and Sheen, J. 2008. Bacterial effectors target the common signaling partner BAK1 to disrupt multiple MAMP receptor-signaling complexes and impede plant immunity. *Cell Host Microbe* 4:17-27.
- Sohn, K. H., Saucet, S. B., Clarke, C. R., Vinatzer, B. A., O'Brien, H. E., Guttman, D. S., and Jones, J. D. G. 2011. HopAS1 recognition significantly contributes to *Arabidopsis* nonhost resistance to *Pseudomonas syringae* pathogens. *New Phytol.* 193:58-66.
- Song, J., Bradeen, J. M., Naess, S. K., Raasch, J. A., Wielgus, S. M., Haberland, G. T., Liu, J., Kuang, H., Austin-Phillips, S., Buell, C. R., Helgeson, J. P., and Jiang, J. 2003. Gene RB cloned from *Solanum bulbocastanum* confers broad spectrum resistance to potato late blight. *Proc. Natl. Acad. Sci. U.S.A.* 100:9128-9133.
- Stamatakis, A. 2004. RAXML-III: A fast program for maximum likelihood-based inference of large phylogenetic trees. *Bioinformatics* 21:456-463.
- Stamatakis, A. 2006. RAXML-VI-HPC: Maximum likelihood-based phylogenetic analyses with thousands of taxa and mixed models. *Bioinformatics* 22:2688-2690.
- Staskawicz, B. J., Dahlbeck, D., and Keen, N. T. 1984. Cloned avirulence gene of *Pseudomonas syringae* pv. *glycinea* determines race-specific incompatibility on *Glycine max* (L.) Merr. *Proc. Natl. Acad. Sci. U.S.A.* 81:6024-6028.
- Stevens, C., Bennett, M. A., Athanassopoulos, E., Tsiamis, G., Taylor, J. D., and Mansfield, J. W. 1998. Sequence variations in alleles of the avirulence gene *avrPphE.R2* from *Pseudomonas syringae* pv. *phaseo-*

- licola* lead to loss of recognition of the AvrPphE protein within bean cells and a gain in cultivar-specific virulence. *Mol. Microbiol.* 29:165-177.
- Studholme, D. J. 2011. Application of high-throughput genome sequencing to intrapathovar variation in *Pseudomonas syringae*. *Mol. Plant Pathol.* 12:829-838.
- Tai, T. H., Dahlbeck, D., Clark, E. T., Gajiwala, P., Pasion, R., Whalen, M. C., Stall, R. E., and Staskawicz, B. J. 1999. Expression of the Bs2 pepper gene confers resistance to bacterial spot disease in tomato. *Proc. Natl. Acad. Sci. U.S.A.* 96:14153-14158.
- Takeuchi, K., Taguchi, F., Inagaki, Y., Toyoda, K., Shiraishi, T., and Ichinose, Y. 2003. Flagellin glycosylation island in *Pseudomonas syringae* pv. *glycinea* and its role in host specificity. *J. Bacteriol.* 185:6658-6665.
- Taylor, J., Teverson, D., and Allen, D. 1996. Identification and origin of races of *Pseudomonas syringae* pv. *phaseolicola* from Africa and other bean growing areas. *Plant Pathol.* 45:469-478.
- Tsiamis, G., Mansfield, J. W., Hockenhull, R., Jackson, R. W., Sesma, A., Athanassopoulos, E., Bennett, M. A., Stevens, C., Vivian, A., Taylor, J. D., and Murillo, J. 2000. Cultivar-specific avirulence and virulence functions assigned to *avrPphF* in *Pseudomonas syringae* pv. *phaseolicola*, the cause of bean halo-blight disease. *EMBO (Eur. Mol. Biol. Organ.) J.* 19:3204-3214.
- Vera Cruz, C. M., Bai, J., Ona, I., Leung, H., Nelson, R. J., Mew, T. W., and Leach, J. E. 2000. Predicting durability of a disease resistance gene based on an assessment of the fitness loss and epidemiological consequences of avirulence gene mutation. *Proc. Natl. Acad. Sci. U.S.A.* 97:13500-13505.
- Vinatzer, B. A., Teitzel, G. M., Lee, M.-W., Jelenska, J., Hotton, S., Fairfax, K., Jenrette, J., and Greenberg, J. T. 2006. The type III effector repertoire of *Pseudomonas syringae* pv. *syringae* B728a and its role in survival and disease on host and non-host plants. *Mol. Microbiol.* 62:26-44.
- Warren, R. F., Merritt, P. M., Holub, E., and Innes, R. W. 1999. Identification of three putative signal transduction genes involved in R gene-specified disease resistance in *Arabidopsis*. *Genetics* 152:401-412.
- Wei, C.-F., Kvitko, B. H., Shimizu, R., Crabill, E., Alfano, J. R., Lin, N.-C., Martin, G. B., Huang, H.-C., and Collmer, A. 2007. A *Pseudomonas syringae* pv. *tomato* DC3000 mutant lacking the type III effector HopQ1-1 is able to cause disease in the model plant *Nicotiana benthamiana*. *Plant J.* 51:32-46.
- Weingart, H., Ullrich, H., Geider, K., and Völksch, B. 2001. The role of ethylene production in virulence of *Pseudomonas syringae* pvs. *glycinea* and *phaseolicola*. *Phytopathology* 91:511-518.
- Wen, C., Tang, X., Su, Z., and Zhou, J. 2010. Effector-triggered and pathogen-associated molecular pattern-triggered immunity differentially contribute to basal resistance to *Pseudomonas syringae*. *Mol. Plant-Microbe Interact.* 23:940-948.
- Wroblewski, T., Caldwell, K. S., Piskurewicz, U., Cavanaugh, K. A., Xu, H., Kozik, A., Ochoa, O., McHale, L. K., Lahre, K., Jelenska, J., Castillo, J. A., Blumenthal, D., Vinatzer, B. A., Greenberg, J. T., and Michelmore, R. W. 2009. Comparative large-scale analysis of interactions between several crop species and the effector repertoires from multiple pathovars of *Pseudomonas* and *Ralstonia*. *Plant Physiol.* 150:1733-1749.
- Xiang, T., Zong, N., Zou, Y., Wu, Y., Zhang, J., Xing, W., Li, Y., Tang, X., Zhu, L., Chai, J., and Zhou, J.-M. 2008. *Pseudomonas syringae* effector AvrPto blocks innate immunity by targeting receptor kinases. *Curr. Biol.* 18:74-80.
- Zumaquero, A., Macho, A. P., Rufián, J. S., and Beuzón, C. R. 2010. Analysis of the role of the type III effector inventory of *Pseudomonas syringae* pv. *phaseolicola* 1448a in interaction with the plant. *J. Bacteriol.* 192:4474-4488.

AUTHOR-RECOMMENDED INTERNET RESOURCES

Phylip package: evolution.genetics.washington.edu/phylip.html
Mr. Bayes database: mrbayes.csit.fsu.edu