




<b>Title</b>	Romidepsin induces caspase-dependent cell death in human neuroblastoma cells
<b>Author(s)</b>	Hegarty, Shane V.; Togher, Katie L.; O'Leary, Eimear; Solger, Franziska; Sullivan, Aideen M.; O'Keeffe, Gerard W.
<b>Publication date</b>	2017-05-12
<b>Original citation</b>	Hegarty, S. V., Togher, K. L., O'Leary, E., Solger, F., Sullivan, A. M. and O'Keeffe, G. W. (2017) 'Romidepsin induces caspase-dependent cell death in human neuroblastoma cells', <i>Neuroscience Letters</i> , 653, pp. 12-18. doi:10.1016/j.neulet.2017.05.025
<b>Type of publication</b>	Article (peer-reviewed)
<b>Link to publisher's version</b>	<a href="http://dx.doi.org/10.1016/j.neulet.2017.05.025">http://dx.doi.org/10.1016/j.neulet.2017.05.025</a> Access to the full text of the published version may require a subscription.
<b>Rights</b>	© 2017, Elsevier Ltd. All rights reserved. This manuscript version is made available under the CC-BY-NC-ND 4.0 license. <a href="http://creativecommons.org/licenses/by-nc-nd/4.0/">http://creativecommons.org/licenses/by-nc-nd/4.0/</a>
<b>Embargo information</b>	Access to this article is restricted until 12 months after publication by request of the publisher.
<b>Embargo lift date</b>	2018-05-12
<b>Item downloaded from</b>	<a href="http://hdl.handle.net/10468/4602">http://hdl.handle.net/10468/4602</a>

Downloaded on 2018-08-23T19:46:27Z

## Accepted Manuscript

Title: Romidepsin induces caspase-dependent cell death in human neuroblastoma cells

Authors: Shane V. Hegarty, Katie L. Togher, Eimear O'Leary, Franziska Solger, Aideen M. Sullivan, Gerard W. O'Keefe



PII: S0304-3940(17)30413-5  
DOI: <http://dx.doi.org/doi:10.1016/j.neulet.2017.05.025>  
Reference: NSL 32832

To appear in: *Neuroscience Letters*

Received date: 27-1-2017  
Revised date: 2-5-2017  
Accepted date: 12-5-2017

Please cite this article as: Shane V.Hegarty, Katie L.Togher, Eimear O'Leary, Franziska Solger, Aideen M.Sullivan, Gerard W.O'Keefe, Romidepsin induces caspase-dependent cell death in human neuroblastoma cells, *Neuroscience Letters*<http://dx.doi.org/10.1016/j.neulet.2017.05.025>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

## **Romidepsin induces caspase-dependent cell death in human neuroblastoma cells.**

**Shane V. Hegarty<sup>1#</sup>, Katie L. Togher<sup>1,2,3#</sup>, Eimear O’Leary<sup>1</sup>, Franziska Solger<sup>1</sup>, Aideen M. Sullivan<sup>1,2</sup>, Gerard W. O’Keeffe<sup>1,2,3</sup>.**

<sup>1</sup> Department of Anatomy and Neuroscience, Western Gateway Building, University College Cork (UCC), Cork, Ireland.

<sup>2</sup> APC Microbiome Institute, UCC, Cork, Ireland.

<sup>3</sup> INFANT Centre, Cork University Maternity Hospital and UCC, Cork, Ireland.

Address correspondence to

Dr. Gerard O’Keeffe

Phone (+353) 21 420 5570

Fax (+353) 21 420 5471

Email g.okeeffe@ucc.ie

or

Prof. Aideen Sullivan

Phone (+353) 21 420 5427


Fax (+353) 21 420 5471

Email a.sullivan@ucc.ie

#These authors contributed equally to this work.

<sup>1</sup> **Abbreviations:** 6-OHDA - 6-hydroxydopamine; pACh3 - p-acetylated-histone H3; DIV – day(s) *in vitro*; HDAC - histone deacetylase; LDH - lactate dehydrogenase; MTT - Thiazolyl Blue Tetrazolium Bromide; N - number of repetitions

## Graphical abstract



## Highlights

- Romidepsin is an FDA-approved small molecule drug and a selective HDAC1/2 inhibitor
- Romidepsin potently induces caspase-dependent cell death in human neuroblastoma cells
- MYCN-amplified neuroblastoma cells are more sensitive to Romidepsin-induced death
- The cytotoxic dose of Romidepsin can be lowered by combination with other cytotoxins
- Romidepsin may be a promising mono-/combination- chemotherapeutic for neuroblastoma

## Abstract

Neuroblastoma is the most common extracranial pediatric solid tumor, arising from the embryonic sympathoadrenal lineage of the neural crest, and is responsible for 15% of childhood cancer deaths. Although survival rates are good for some patients, those children diagnosed with high-risk neuroblastoma have survival rates as low as 35%. Thus, neuroblastoma remains a significant clinical challenge and the development of novel

therapeutic strategies is essential. Given that there is widespread epigenetic dysregulation in neuroblastoma, epigenetic pharmacotherapy holds promise as a therapeutic approach. In recent years, histone deacetylase (HDAC) inhibitors, which cause selective activation of gene expression, have been shown to be potent chemotherapeutics for the treatment of a wide range of cancers. Here we examined the ability of the FDA-approved drug Romidepsin, a selective HDAC1/2 inhibitor, to act as a cytotoxic agent in neuroblastoma cells. Treatment with Romidepsin at concentrations in the low nanomolar range induced neuroblastoma cell death through caspase-dependent apoptosis. Romidepsin significantly increased histone acetylation, and significantly enhanced the cytotoxic effects of the cytotoxic agent 6-hydroxydopamine, which has been shown to induce cell death in neuroblastoma cells through increasing reactive oxygen species. Romidepsin was also more potent in MYCN-amplified neuroblastoma cells, which is an important prognostic marker of poor survival. This study has thus demonstrated that the FDA-approved chemotherapeutic drug Romidepsin has a potent caspase-dependent cytotoxic effect on neuroblastoma cells, whose effects enhance cell death induced by other cytotoxins, and suggests that Romidepsin may be a promising chemotherapeutic candidate for the treatment of neuroblastoma.

**Key words:**

Romidepsin; Neuroblastoma; Cell Death; Epigenetic Regulation; Chemotherapeutic Drug.

## **Introduction**

Neuroblastoma is the most common extracranial pediatric solid tumor, and arises from the sympathoadrenal lineage of the neural crest. The incidence rate has been reported to be 10.9 cases per million children [1,2]. Neuroblastoma accounts for 8 to 10% of pediatric cancers and is responsible for 15% of childhood deaths from cancer [3]. Although survival rates for patients with loco-regional tumours are greater than 90%, the prognosis for those less than 18

months of age at diagnosis with metastatic disease remains dismal, and can be as low as 35% [2]. Therefore, neuroblastoma remains a significant clinical challenge, especially for those children with poorer prognoses, and the development of novel therapeutic strategies is essential.

Cancer is now understood to be a disease of widespread epigenetic dysregulation, which contributes to almost every step of tumour progression and interacts extensively with underlying genetic mutations [4-6]. Therefore, epigenetic modulators may offer an effective means to therapeutically alter the regulation of proto-oncogenes and tumor suppressor genes in cancer cells. One key epigenetic process is histone acetylation which regulates the accessibility of genes for transcription factor binding, thereby controlling the levels of gene expression [7,8]. Histone acetylation involves a dynamic interplay between histone acetyltransferase and histone deacetylase (HDAC) enzymes, which are responsible for histone acetylation and deacetylation respectively [9]. In recent years, small molecule HDAC inhibitors have been shown to be potent anti-cancer agents, several of which are now FDA-approved cancer therapies. Indeed, HDAC inhibitors are being developed as drugs for the treatment of a wide range of cancers including: glioblastoma; leukemia; lymphoma; myeloma; and breast, colorectal, gastrointestinal, lung, ovarian, pancreatic, and prostate cancer [4,10-20]. In neuroblastoma, HDAC inhibitors have been shown to induce apoptosis in a number of neuroblastoma cell lines, and thus have promise for treating high-risk neuroblastoma [21-25].

Romidepsin (FK228; FR901228 or Istodax®) is a depsipeptide small molecule (MW=540.7) that belongs to bicyclic peptide selective inhibitors of HDAC1 and HDAC2. It was approved by FDA for the treatment of refractory cutaneous T-cell lymphoma based on evidence from phase II clinical trials [20,26-28]. Romidepsin is being used/developed as a monotherapy or combination therapy for a number of cancers, in particular T-cell lymphoma [10,12,13,18,20,26,27,29,30-32]. Romidepsin is well tolerated in pediatric patients with refractory solid tumors, as well as patients with advanced cancers [33,34], meaning that it may be promising for clinical use in neuroblastoma patients. In the present study, we investigated the ability of Romidepsin to induce cytotoxic effects in the human neuroblastoma cells.

## **Materials and Methods**

### **Cell Culture**

SH-SY5Y cells were cultured in Dulbecco's Modified Eagle Medium Nutrient Mixture F-12, supplemented with 10% foetal calf serum (FCS), 100 nM L-Glutamine, 100 U/ml Penicillin/Streptomycin. IMR-32 cells were cultured in Eagle's Minimum Essential Medium with 2 mM L-Glutamine, 1% Non-Essential Amino Acids (NEAA) with 10% FCS. SK-N-BE cells were cultured in Eagle's Minimum Essential Medium with 2mM L-Glutamine, 1% NEAA: Hams F12 (1:1) with 15% FCS (all from Sigma). All cells were cultured in a humidified atmosphere containing 5% CO<sub>2</sub> at 37°C. For cell viability assays, cells were plated at a density of  $1 \times 10^5$  cells/well in a 24-well plate. For western blotting,  $2 \times 10^6$  cells were plated per well of a 6-well plate. For RNA extraction  $1.5 \times 10^6$  cells were plated per well in a 12-well plate. Where indicated, cells were treated with 5–5,000 nM of Romidepsin (MedChem Express) and/or 1 µg/ml caspase-3 inhibitor (Calbiochem) or with 15 µM 6-hydroxydopamine (6-OHDA; Sigma) for 24 h or daily for 4 days *in vitro* (DIV).

### **Cell Viability and Cell Death Assessment**

Thiazolyl Blue Tetrazolium Bromide (MTT) assays were performed to assess cell viability [35]. MTT was added to cells at a concentration of 0.5 mg/ml. Lactate dehydrogenase (LDH) activity was measured in 100 µl of the cell culture medium of each treatment group using an LDH Activity Assay Kit (Sigma), according to the manufacturer's instructions. Fresh cell culture medium, which had no contact with cultured cells, was used as the negative control.

### **Immunocytochemistry**

Cultures were immunocytochemically stained for p-acetylated-histone H3 (Ser 11/Lys 15) (pAcH3; 1:200; rabbit polyclonal IgG; Santa Cruz) and β-actin (1:200; mouse monoclonal IgG; Sigma) [36]. Cells were imaged with an Olympus IX70 inverted microscope fitted with an Olympus DP70 camera and AnalysisD™ software. The fluorescence intensity of individual cells stained for pAcH3 was measured using Image J and the relative fluorescence intensity was calculated as the intensity of each individual cell after subtraction of the background noise.

### **Measurement of Cellular Morphology**

Neurite growth analysis as a proxy measure differentiation of SH-SY5Y cells was performed as previously described [36], using the using the formula: neurite length =  $\alpha \times T \times (\pi/2)$ , where  $\alpha$  is the number of times the neurite intersects the grid lines, and T is the distance between the gridlines on the magnified image.

### Western Blotting

Western blotting was carried out as described [37]. Cells were lysed in RIPA buffer for 1h on ice, and insoluble debris was removed by centrifugation. 15 µg of protein was run by SDS-PAGE and transferred to nitrocellulose membranes using a Mini Trans-Blot Electrophoretic Transfer Cell (Bio-Rad, CA, USA). The membranes were incubated with primary antibodies against pAcH3 (1:1000; rabbit polyclonal IgG; Santa Cruz) or β-actin (1:1000; mouse monoclonal IgG; Sigma) overnight at 4°C, washed, incubated with the appropriate horseradish peroxidase-labelled secondary antibodies (1:2000; Promega), washed and developed with enhanced chemiluminescence (GE Healthcare). Protein expression was normalised to β-actin by densitometry using Image J.

### RNA extraction, cDNA synthesis and real-time PCR

RNA was extracted from the cells using Trizol reagent as per the manufactures instructions (Invitrogen). RNA concentration was quantified using a Spectrophotometer (NanoDrop 1000). Reverse transcription was performed using a high capacity cDNA Reverse Transcription Kit (Applied Biosystems) under the following parameters: 25 °C for 10 min; 37 °C for 120 min; 85 °C for 5 min; 4 °C for 10 min. The reaction mix for real-time PCR consisted of 5 µl TaqMan® Gene Expression Master Mix (Applied Biosystems), 4 µl of RNase-free H<sub>2</sub>O (Applied Biosystems), 0.5 µl 20× TaqMan® Gene Expression Assays (*ACTB*, *Bax*, *Bcl2*, *caspase-8*, *caspase-3*, *PPAR-γ*; Applied Biosystems) and 0.5 µl of cDNA (10 ng/ml). Each sample was run in duplicate under the following cycling parameters: 50 °C for 2 min; 95 °C for 10 min; 40 repetitions of 95 °C for 15 s and 60 °C for 1 min. Expression levels were calculated using the 2- $\delta$ CT method (Schmittgen and Livak, 2008).

### Statistical Analysis

Unpaired Student's t-test or one-way ANOVA with a *post hoc* Tukey's or Bonferroni's test was performed, as appropriate, to determine significant differences between groups. For non-parametric data, Kruskal-Wallis ANOVA with Dunn's *post hoc* test was used. Outliers were removed prior to analysis if Grubb's Test  $p < 0.05$ . Results were expressed as means with SEM and deemed significant when  $p < 0.05$ .

## Results



**Romidepsin induces cell death of human SH-SY5Y neuroblastoma cells.**

We first examined the effects of Romidepsin using human SH-SY5Y neuroblastoma cells, which were treated with 5 – 500 nM of Romidepsin daily for 4 DIV. These concentrations were selected based on IC<sub>50</sub> values of Romidepsin being in the nanomolar range (36 and 47 nM for HDAC1 and HDAC2, respectively), and that Romidepsin has been shown to reduce viability at 10 nM in the HL-60 cell line and at 500 nM in the HP100 cell line [38]. An MTT assay revealed that Romidepsin had no effect on cell viability at concentrations  $\leq$  40 nM when compared to control (Fig. 1A), but significantly reduced cell viability at concentrations  $\geq$  100 nM (Fig. 1A). Phase contrast microscopy confirmed that Romidepsin treatment led to a decrease in cell number and an increase in dead, non-adherent cells (Fig. 1B). We further examined the effects of the non-cytotoxic doses of Romidepsin on cellular morphology by immunostaining for  $\beta$ -actin. Treatment with 40 nM of Romidepsin significantly reduced the length of SH-SY5Y cell neurites (Fig. 1C), indicating a neuritotoxic effect. These data demonstrate that Romidepsin exerts potent cytotoxic effects in SH-SY5Y neuroblastoma cells in the low nano-molar range.

**Romidepsin increases histone acetylation in human SH-SY5Y neuroblastoma cells.**

As Romidepsin is a selective HDAC1 and HDAC2 inhibitor, we next assessed the ability of Romidepsin to increase histone acetylation. Given that 40 nM Romidepsin had a neuritoxic effect, and to eliminate the confounding effects of low cell numbers at the higher doses, cells were treated with 20 nM and 40 nM of Romidepsin and the levels of acetylated histones (pACh3) were first measured by western blotting. 20 nM and 40 nM Romidepsin significantly increased the levels of pACh3 relative to control, which was measured by densitometric quantification of pACh3 protein levels relative to  $\beta$ -actin loading control (Fig. 2A). Immunocytochemistry was used to confirm these findings and Romidepsin treatment resulted in a significant increase in the levels of pACh3 relative to control, as quantified by densitometry (Fig. 2B). Taken together, these results suggest that Romidepsin leads to increases in histones in acetylation in SH-SY5Y neuroblastoma cells.

**The effects of Romidepsin on SH-SY5Y cells are caspase-dependent.**

To determine whether Romidepsin induces apoptosis in SH-SY5Y cells, cultures were co-treated with a broad-spectrum caspase-3 inhibitor, which inhibits to the pro-apoptotic caspase-3 that functions upstream of other pro-apoptotic caspases; caspase-6 and caspase-7 [39]. An MTT assay revealed that caspase-3 inhibition significantly inhibited Romidepsin-

induced cell death in the SH-SY5Y neuroblastoma cells at 4 DIV (Fig. 3A). We next aimed to investigate whether the cytotoxic dose of Romidepsin could be reduced by combination therapy, in which neuroblastoma cells are exposed to another cytotoxic agent. To provide proof of principle for this, SH-SY5Y cells were co-treated with 6-OHDA, which is a potent cytotoxin that has been shown to induce SH-SY5Y cell death through increasing reactive oxygen species [40,41]. SH-SY5Y cells were treated with 15  $\mu$ M of 6-OHDA [36] with or without 20 nM of Romidepsin (a concentration which does not adversely affect cell viability or morphology, but induces hyperacetylation) for 24 h at which time LDH assays were performed to assess cell death. Interestingly, 20 nM Romidepsin significantly enhanced the cytotoxic effects of 6-OHDA (Fig. 3B). These data suggest that it may be possible for the therapeutic dose of Romidepsin to be lowered when used as an add-on pharmacotherapy in combination therapy.

#### **Romidepsin exerts cytotoxic effects in MYCN amplified cells.**

As SH-SY5Y cells express wild-type p53 and are not MYCN amplified, we carried out additional experiments in IMR-32 (MYCN amplified, p53 wild-type) and SK-N-BE(2) (MYCN amplified, p53 mutant) human neuroblastoma cell lines. MYCN amplification is the primary and most important prognostic marker of poor survival in neuroblastoma [54]. Based on the dose-response experiments carried out in SH-SY5Y cells, these cells were treated with 40 nM or 100 nM of Romidepsin for 4 DIV. An MTT assay revealed that both MYCN amplified neuroblastoma cell lines were more sensitive to the cytotoxic effects of Romidepsin than SH-SY5Y cells (Fig. 1A and 4A). Indeed, Romidepsin doses of 40 nM and 100 nM significantly reduced IMR-32 and SK-N-BE cell viability (Fig. 4A). Real-time PCR revealed that 40 nM Romidepsin treatment significantly increased the expression levels of *caspase-3* mRNA (Fig. 4B), and increased the ratio of *Bax* mRNA to *Bcl2* mRNA in all the neuroblastoma cell lines tested (Fig. 4C). These data show that Romidepsin is a potent cytotoxin that leads to pro-apoptotic transcriptional changes in MYCN amplified and non-amplified human neuroblastoma cells.

## **Discussion**

The selective inhibitor of HDAC1 and HDAC2, Romidepsin, is an FDA-approved chemotherapeutic drug that is being used/developed as a monotherapy or combination therapy for a number of cancers [10,12,13,18,20,26-29,30-32]. To investigate the potential of

Romidepsin for the treatment of neuroblastoma, we examined the cytotoxic effects of Romidepsin in human neuroblastoma cell lines. In the present study, we first demonstrated that Romidepsin induces cell death of SH-SY5Y cells at concentrations in the low nanomolar range through caspase-dependent apoptosis. We next found that Romidepsin significantly increases histone acetylation in these neuroblastoma cells, most likely through inhibition of HDAC1 and HDAC2 activity, and that the cytotoxic dose of Romidepsin can be lowered when combined with other cytotoxic agents. Finally, we demonstrated that Romidepsin is a potent cytotoxin that leads to pro-apoptotic transcriptional changes in MYCN amplified and non-amplified human neuroblastoma cell lines. As a result, we believe that Romidepsin may be useful in monotherapy or combination therapy approaches for neuroblastoma.

In the present study, following a dose-response experiment, we initially found that concentrations >100 nM of Romidepsin induce a significant reduction in cell viability of SH-SY5Y cells. We observed extensive cell death by phase contrast microscopy, and then showed that a non-cytotoxic dose of Romidepsin (40 nM) significantly reduced the length of SH-SY5Y cell neurites, while lower doses had no effect on neurite length. These data indicate that Romidepsin has a neuritotoxic effect on SH-SY5Y cells. Moreover, Romidepsin did not increase the length of SH-SY5Y cell neurites, and thus did not induce their neuronal differentiation. Taken together, we concluded that Romidepsin has potent cytotoxic effects on SH-SY5Y neuroblastoma cells. In support of our findings, HDAC inhibitors have been shown to induce cell death in a number of neuroblastoma cell lines [21-25].

We subsequently confirmed the ability of Romidepsin to induce hyperacetylation of histone H3 in the SH-SY5Y cells, which was most likely due to selective inhibition of HDAC1 and HDAC2. Romidepsin induced significant increases in the levels of acetylated histone H3 at 20 nM and 40 nM. In support of this, the IC<sub>50</sub> values of Romidepsin are 36 nM and 47 nM for HDAC1 and HDAC2, respectively. Furthermore, Romidepsin has been consistently demonstrated to inhibit HDAC1 and HDAC2 activity *in vitro*, *in vivo* and in cancer patients, in which histone acetylation levels were assessed [28,38,42]. This suggests that the Romidepsin-induced cytotoxic effects observed in this study may be mediated by hyperacetylation, following inhibition of HDAC1 and HDAC2 activity. Indeed, epigenetic dysregulation has been reported to contribute to a cancer phenotype [4-6].

To determine the cellular mechanisms through which Romidepsin induced SH-SY5Y cell death, we next showed that Romidepsin-induced cell death in SH-SY5Y neuroblastoma cells was caspase-dependent. In support of these findings, Romidepsin has previously been reported to induce apoptosis in a number of cancer cell lines [24,38,43]. SH-SY5Y cell death

induced by treatment with 100 nM of Romidepsin was prevented by co-treatment with a caspase-3 inhibitor. Similarly, Romidepsin has been shown to induce apoptosis at 10 nM in the HL-60 cell line and at 500 nM in the HP100 cell line [38]. Taken together, these results suggest that Romidepsin induces caspase-dependent apoptosis in neuroblastoma cells.

We next sought to determine if the cytotoxic dose of Romidepsin in neuroblastoma cells could be reduced by combination therapy with another cytotoxic agent. Using a concentration of Romidepsin which does not adversely affect cell viability or morphology but induces hyperacetylation (20 nM), we showed that Romidepsin significantly increases the SH-SY5Y cell death induced by the cytotoxin 6-OHDA, which has been shown to induce SH-SY5Y cell death through increasing reactive oxygen species [40,41]. These data suggest that it may be possible for the therapeutic dose of Romidepsin to be lowered when used as an add-on pharmacotherapy in combination therapies for neuroblastoma.

Finally, we examined the effects of Romidepsin in MYCN amplified human neuroblastoma cell lines, IMR-32 and SK-N-BE, as MYCN amplification is the primary and most important prognostic marker of poor survival in neuroblastoma [54]. MYCN amplified neuroblastoma cell lines were significantly more sensitive to the cytotoxic effects of Romidepsin than the non-MYCN amplified SH-SY5Y cells. This suggests that Romidepsin may be particularly effective in neuroblastoma cases with MYCN amplification, which accounts for ~25% of cases and correlates with high-risk disease and poor prognosis [54]. Lastly, Romidepsin was shown to cause significant pro-apoptotic transcriptional changes in both MYCN amplified and non-amplified human neuroblastoma cells.

Collectively these *in vitro* data demonstrate the potential of Romidepsin to be used as chemotherapy for neuroblastoma patients. Indeed, the FDA-approved Romidepsin is being used/developed as a monotherapy or combination therapy for various other cancer subtypes [10,12,13,18,20,26,27,29,30-32]. Furthermore, Romidepsin has been demonstrated to be well tolerated in pediatric patients with refractory solid tumors, as well as patients with advanced cancers [33,34], which gives it promise for clinical use in neuroblastoma patients. Before Romidepsin is considered for clinical use, it should be examined in animal models of neuroblastoma (for examples see [47]) as the next stage in rationalizing its use as a potential therapy for this cancer. Despite the use, effectiveness and promise of Romidepsin as a chemotherapy, Romidepsin has been ineffective in its treatment of some cancers in clinical trials, including glioblastomas and colorectal cancer, with some patients experiencing adverse effects [42,48-52]. Thus, preclinical efficacy with Romidepsin does not necessarily predict effectiveness in clinical trials. In the event of Romidepsin-related clinical complications,

Romidepsin has been shown to be metabolized by cytochrome P450 enzymes [53], which provides a potential avenue to counteract any adverse effects of Romidepsin. However, clinical trials aimed at harnessing these enzymes to metabolise Romidepsin were unsuccessful [52]. Given the dismal survival rates in children with high-risk neuroblastoma [2], it is important that novel therapeutic approaches are developed for these patients. Romidepsin has demonstrated promise in this regard, and warrants further investigation in this context.

### **Acknowledgements**

Studies in the authors' laboratories are supported by grants from the Irish Research Council (R15897; SVH/AS/G'OK) the National University of Ireland (R16189; SVH/AS/G'OK). This publication has also emanated from research supported in part by a research grant from Science Foundation Ireland (SFI) under the Grant Number 15/CDA/3498 (G'OK).

### **References**

- [1] Kaatsch P (2010) Epidemiology of childhood cancer. *Cancer Treat Rev* 36: 277-285.
- [2] Modak S, Cheung NK (2010) Neuroblastoma: Therapeutic strategies for a clinical enigma. *Cancer Treat Rev* 36: 307-317.
- [3] Maris JM (2010) Recent advances in neuroblastoma. *N Engl J Med* 362: 2202-2211.
- [4] Abdelfatah E, Kerner Z, Nanda N, Ahuja N (2016) Epigenetic therapy in gastrointestinal cancer: the right combination. *Therap Adv Gastroenterol* 9: 560-579.
- [5] Kumar R, Li DQ, Muller S, Knapp S (2016) Epigenomic regulation of oncogenesis by chromatin remodeling. *Oncogene*.
- [6] Shinjo K, Kondo Y (2015) Targeting cancer epigenetics: Linking basic biology to clinical medicine. *Adv Drug Deliv Rev* 95: 56-64.
- [7] Allfrey VG, Faulkner R, Mirsky AE (1964) ACETYLATION AND METHYLATION OF HISTONES AND THEIR POSSIBLE ROLE IN THE REGULATION OF RNA SYNTHESIS. *Proc Natl Acad Sci U S A* 51: 786-794.
- [8] Serrano L, Vazquez BN, Tischfield J (2013) Chromatin structure, pluripotency and differentiation. *Exp Biol Med (Maywood)* 238: 259-270.
- [9] Yang XJ, Seto E (2007) HATs and HDACs: from structure, function and regulation to novel strategies for therapy and prevention. *Oncogene* 26: 5310-5318.

- [10] Kaushik D, Vashistha V, Isharwal S, Sediqe SA, Lin MF (2015) Histone deacetylase inhibitors in castration-resistant prostate cancer: molecular mechanism of action and recent clinical trials. *Ther Adv Urol* 7: 388-395.
- [11] Juo YY, Gong XJ, Mishra A, Cui X, Baylin SB, et al. (2015) Epigenetic therapy for solid tumors: from bench science to clinical trials. *Epigenomics* 7: 215-235.
- [12] Barbarotta L, Hurley K (2015) Romidepsin for the Treatment of Peripheral T-Cell Lymphoma. *J Adv Pract Oncol* 6: 22-36.
- [13] Damaskos C, Karatzas T, Nikolidakis L, Kostakis ID, Karamaroudis S, et al. (2015) Histone Deacetylase (HDAC) Inhibitors: Current Evidence for Therapeutic Activities in Pancreatic Cancer. *Anticancer Res* 35: 3129-3135.
- [14] Zhu L, Wu K, Ma S, Zhang S (2015) HDAC inhibitors: a new radiosensitizer for non-small-cell lung cancer. *Tumori* 101: 257-262.
- [15] Bian J, Zhang L, Han Y, Wang C (2015) Histone deacetylase inhibitors: potent anti-leukemic agents. *Curr Med Chem* 22: 2065-2074.
- [16] Nwabo Kamdje AH, Seke Etet PF, Vecchio L, Tagne RS, Amvene JM, et al. (2014) New targeted therapies for breast cancer: A focus on tumor microenvironmental signals and chemoresistant breast cancers. *World J Clin Cases* 2: 769-786.
- [17] Tampakis A, Tampaki EC, Nebiker CA, Kouraklis G (2014) Histone deacetylase inhibitors and colorectal cancer: what is new? *Anticancer Agents Med Chem* 14: 1220-1227.
- [18] Marsh DJ, Shah JS, Cole AJ (2014) Histones and their modifications in ovarian cancer - drivers of disease and therapeutic targets. *Front Oncol* 4: 144.
- [19] Bezecny P (2014) Histone deacetylase inhibitors in glioblastoma: pre-clinical and clinical experience. *Med Oncol* 31: 985.
- [20] McGraw AL (2013) Romidepsin for the treatment of T-cell lymphomas. *Am J Health Syst Pharm* 70: 1115-1122.
- [21] Wang G, Edwards H, Caldwell JT, Buck SA, Qing WY, et al. (2013) Panobinostat synergistically enhances the cytotoxic effects of cisplatin, doxorubicin or etoposide on high-risk neuroblastoma cells. *PLoS One* 8: e76662.
- [22] de Ruijter AJ, Kemp S, Kramer G, Meinsma RJ, Kaufmann JO, et al. (2004) The novel histone deacetylase inhibitor BL1521 inhibits proliferation and induces apoptosis in neuroblastoma cells. *Biochem Pharmacol* 68: 1279-1288.
- [23] Muhlethaler-Mottet A, Flahaut M, Bourlout KB, Auderset K, Meier R, et al. (2006) Histone deacetylase inhibitors strongly sensitise neuroblastoma cells to TRAIL-induced

apoptosis by a caspases-dependent increase of the pro- to anti-apoptotic proteins ratio. *BMC Cancer* 6: 214.

[24] Panicker J, Li Z, McMahon C, Sizer C, Steadman K, et al. (2010) Romidepsin (FK228/depsipeptide) controls growth and induces apoptosis in neuroblastoma tumor cells. *Cell Cycle* 9: 1830-1838.

[25] Witt O, Deubzer HE, Lodrini M, Milde T, Oehme I (2009) Targeting histone deacetylases in neuroblastoma. *Curr Pharm Des* 15: 436-447.

[26] Piekarz RL, Frye R, Turner M, Wright JJ, Allen SL, et al. (2009) Phase II multi-institutional trial of the histone deacetylase inhibitor romidepsin as monotherapy for patients with cutaneous T-cell lymphoma. *J Clin Oncol* 27: 5410-5417.

[27] Whittaker SJ, Demierre MF, Kim EJ, Rook AH, Lerner A, et al. (2010) Final results from a multicenter, international, pivotal study of romidepsin in refractory cutaneous T-cell lymphoma. *J Clin Oncol* 28: 4485-4491.

[28] Furumai R, Matsuyama A, Kobashi N, Lee KH, Nishiyama M, et al. (2002) FK228 (depsipeptide) as a natural prodrug that inhibits class I histone deacetylases. *Cancer Res* 62: 4916-4921.

[29] Haigentz M, Jr., Kim M, Sarta C, Lin J, Keresztes RS, et al. (2012) Phase II trial of the histone deacetylase inhibitor romidepsin in patients with recurrent/metastatic head and neck cancer. *Oral Oncol* 48: 1281-1288.

[30] Jones SF, Infante JR, Spigel DR, Peacock NW, Thompson DS, et al. (2012) Phase 1 results from a study of romidepsin in combination with gemcitabine in patients with advanced solid tumors. *Cancer Invest* 30: 481-486.

[31] Tinari N, De Tursi M, Grassadonia A, Zilli M, Stuppia L, et al. (2012) An epigenetic approach to pancreatic cancer treatment: the prospective role of histone deacetylase inhibitors. *Curr Cancer Drug Targets* 12: 439-452.

[32] Harrison SJ, Quach H, Link E, Seymour JF, Ritchie DS, et al. (2011) A high rate of durable responses with romidepsin, bortezomib, and dexamethasone in relapsed or refractory multiple myeloma. *Blood* 118: 6274-6283.

[33] Fouladi M, Furman WL, Chin T, Freeman BB, 3rd, Dudkin L, et al. (2006) Phase I study of depsipeptide in pediatric patients with refractory solid tumors: a Children's Oncology Group report. *J Clin Oncol* 24: 3678-3685.

[34] Amiri-Kordestani L, Luchenko V, Peer CJ, Ghafourian K, Reynolds J, et al. (2013) Phase I trial of a new schedule of romidepsin in patients with advanced cancers. *Clin Cancer Res* 19: 4499-4507.

- [35] Hegarty SV, Sullivan AM, O'Keeffe GW (2013) BMP2 and GDF5 induce neuronal differentiation through a Smad dependant pathway in a model of human midbrain dopaminergic neurons. *Mol Cell Neurosci* 56C: 263-271.
- [36] Hegarty SV, O'Leary E, Solger F, Stanicka J, Sullivan AM, et al. (2016) A Small Molecule Activator of p300/CBP Histone Acetyltransferase Promotes Survival and Neurite Growth in a Cellular Model of Parkinson's Disease. *Neurotox Res*.
- [37] Crampton SJ, Collins LM, Toulouse A, Nolan YM, O'Keeffe GW (2012) Exposure of foetal neural progenitor cells to IL-1beta impairs their proliferation and alters their differentiation - a role for maternal inflammation? *J Neurochem* 120: 964-973.
- [38] Mizutani H, Hiraku Y, Tada-Oikawa S, Murata M, Ikemura K, et al. (2010) Romidepsin (FK228), a potent histone deacetylase inhibitor, induces apoptosis through the generation of hydrogen peroxide. *Cancer Sci* 101: 2214-2219.
- [39] Siegel RM (2006) Caspases at the crossroads of immune-cell life and death. *Nat Rev Immunol* 6: 308-317.
- [40] Yamamuro A, Yoshioka Y, Ogita K, Maeda S (2006) Involvement of endoplasmic reticulum stress on the cell death induced by 6-hydroxydopamine in human neuroblastoma SH-SY5Y cells. *Neurochem Res* 31: 657-664.
- [41] Kwon SH, Ma SX, Lee SY, Jang CG (2014) Sulfuretin inhibits 6-hydroxydopamine-induced neuronal cell death via reactive oxygen species-dependent mechanisms in human neuroblastoma SH-SY5Y cells. *Neurochem Int* 74: 53-64.
- [42] Schrump DS, Fischette MR, Nguyen DM, Zhao M, Li X, et al. (2008) Clinical and molecular responses in lung cancer patients receiving Romidepsin. *Clin Cancer Res* 14: 188-198.
- [43] Karthik S, Sankar R, Varunkumar K, Ravikumar V (2014) Romidepsin induces cell cycle arrest, apoptosis, histone hyperacetylation and reduces matrix metalloproteinases 2 and 9 expression in bortezomib sensitized non-small cell lung cancer cells. *Biomed Pharmacother* 68: 327-334.
- [44] Toulouse A, Collins GC, Sullivan AM (2012) Neurotrophic effects of growth/differentiation factor 5 in a neuronal cell line. *Neurotox Res* 21: 256-265.
- [45] Sullivan AM, Opacka-Juffry J, Hotten G, Pohl J, Blunt SB (1997) Growth/differentiation factor 5 protects nigrostriatal dopaminergic neurones in a rat model of Parkinson's disease. *Neurosci Lett* 233: 73-76.




- [46] Hurley FM, Costello DJ, Sullivan AM (2004) Neuroprotective effects of delayed administration of growth/differentiation factor-5 in the partial lesion model of Parkinson's disease. *Exp Neurol* 185: 281-289.
- [47] Teitz T, Stanke JJ, Federico S, Bradley CL, Brennan R, et al. (2011) Preclinical models for neuroblastoma: establishing a baseline for treatment. *PLoS One* 6: e19133.
- [48] Whitehead RP, Rankin C, Hoff PM, Gold PJ, Billingsley KG, et al. (2009) Phase II trial of romidepsin (NSC-630176) in previously treated colorectal cancer patients with advanced disease: a Southwest Oncology Group study (S0336). *Invest New Drugs* 27: 469-475.
- [49] Stadler WM, Margolin K, Ferber S, McCulloch W, Thompson JA (2006) A phase II study of depsipeptide in refractory metastatic renal cell cancer. *Clin Genitourin Cancer* 5: 57-60.
- [50] Molife LR, Attard G, Fong PC, Karavasilis V, Reid AH, et al. (2010) Phase II, two-stage, single-arm trial of the histone deacetylase inhibitor (HDACi) romidepsin in metastatic castration-resistant prostate cancer (CRPC). *Ann Oncol* 21: 109-113.
- [51] Otterson GA, Hodgson L, Pang H, Vokes EE (2010) Phase II study of the histone deacetylase inhibitor Romidepsin in relapsed small cell lung cancer (Cancer and Leukemia Group B 30304). *J Thorac Oncol* 5: 1644-1648.
- [52] Iwamoto FM, Lamborn KR, Kuhn JG, Wen PY, Yung WK, et al. (2011) A phase I/II trial of the histone deacetylase inhibitor romidepsin for adults with recurrent malignant glioma: North American Brain Tumor Consortium Study 03-03. *Neuro Oncol* 13: 509-516.
- [53] Shiraga T, Tozuka Z, Ishimura R, Kawamura A, Kagayama A (2005) Identification of cytochrome P450 enzymes involved in the metabolism of FK228, a potent histone deacetylase inhibitor, in human liver microsomes. *Biol Pharm Bull* 28: 124-129.
- [54] Huang A, Weiss WA (2013) Neuroblastoma and MYCN. *Cold Spring Harb Perspect Med*. 2013 Oct; 3(10): a014415.

## Figures and Figure Legends

### Figure 1: Romidepsin induces cell death of human SH-SY5Y neuroblastoma cells.


(A) MTT assay with (B) representative photomicrographs of Romidepsin-treated SH-SY5Y cells treated daily for 4 DIV with 0 - 500 nM of Romidepsin (\*\*\*)  $p < 0.001$ ; ANOVA with post-hoc Tukey's test;  $N = 4$ ). (C) Graph with (D) representative photomicrographs (Green =

$\beta$ -actin) showing the average total neurite length of cells in the control or 40 nM Romidepsin group (\*\*  $p < 0.01$ ; 20 images analysed per group per experiment.  $N = 3$ ). Scale bar = 50  $\mu\text{m}$ .




**Figure 2: Romidepsin increases histone acetylation in SH-SY5Y neuroblastoma cells.**

Western blots of (A) pACh3 and  $\beta$ -actin levels in SH-SY5Y cells cultured for 24 h with 0 to 40 nM of Romidepsin, and (B) normalised to  $\beta$ -actin levels ( $N = 3$ ). (C) Quantification of the relative fluorescence intensity of (D) pACh3 staining in SH-SY5Y cultured for 24 h with 0 to 40 nM of Romidepsin (\*  $p < 0.05$ , \*\*\*  $p < 0.001$  v control; One-way ANOVA with post-hoc Tukey's test; 50 cells analysed per group per experiment.  $N = 3$ ). Scale bar = 100  $\mu\text{m}$ .




**Figure 3: Caspase-dependent cytotoxic effects of Romidepsin in neuroblastoma cells.**

(A) Representative photomicrographs and (B, C) MTT assay of SH-SY5Y cells treated daily with without or with 1  $\mu\text{g/ml}$  of caspase-3 inhibitor and/or 40 or 100 nM of Romidepsin for 4 DIV (\*\*  $p < 0.01$  v control; One-way ANOVA with post-hoc Tukey's test;  $N = 3$ ). Scale bar = 50  $\mu\text{m}$ . (D) Standardised LDH assay of SH-SY5Y cells treated with 15  $\mu\text{M}$  6-OHDA in the presence or absence of 20 nM of Romidepsin for 24 h (\*\*\*  $p < 0.001$  v control; ++  $p < 0.01$  v 6-OHDA alone; One-way ANOVA with post-hoc Tukey's test;  $N = 6$ ).



**Figure 4: Effects of Romidepsin in MYCN-amplified and non-amplified neuroblastoma cells.**

(A) MTT assay of Romidepsin-treated IMR-32 and SK-N-BE(2) cells treated daily for 4 DIV with 0, 40 or 100 nM of Romidepsin (\*\*\*)  $p < 0.001$  vs control; ANOVA with post-hoc Tukey's test. Number of repetitions (N) = 4). (B) Real-time PCR of *caspase-3* mRNA in all three neuroblastoma cells lines treated for 24 h with 40 nM Romidepsin (\*  $p < 0.05$  vs control; Student's t-test). (C) Graphical representation of real-time PCR showing the ratio of *Bax* to *Bcl2* mRNA in all three cells lines treated for 24 h with 40 nM Romidepsin.



C