

## TROPHIC LEVELS AND METHODS FOR STOMACH CONTENT ANALYSIS OF FISHES

**P. U. Zacharia**

*Demersal Fisheries Division, CMFRI, Kochi*

ICAR- Central Marine Fisheries Research Institute

# 28

Investigation of food and feeding of fishes has traditionally been an important field of activity in fisheries biology, but it is one in which there are great difficulties in correlating the results with the research made in the other fields (FAO, 1974). Investigations of the food of the fish cannot be considered in isolation but have to be discussed in relation to the whole marine environment, of which the fish constitute single elements.

### Food Chains and Trophic Levels

The production of organic substances (food) by photosynthesis is a process involving transformation of light energy into potential chemical energy. The transfer of this food energy from the producers through a series of consumers is called a food chain, each organism through which it is passed being a link in the chain.

Three different food chains may be recognized.

1	2	3
<ul style="list-style-type: none"><li>•The carnivore chain, where the energy is passed from smaller to larger organisms</li></ul>	<ul style="list-style-type: none"><li>•The parasite chain, where the energy is passed from larger to smaller organisms</li></ul>	<ul style="list-style-type: none"><li>•The saprophyte chain, where the energy is passed from dead organic matter to micro-organism in most cases</li></ul>

In reality food may be passed through parts of all three chains before it is finally decomposed into inorganic nutrients by the bacteria and fungi found at the end of every food chain. In other words, the species population within a community or ecosystem form many food chains which interconnect, anastomose or cross each other in a complex pattern, which is usually referred to as the **food web**.

Organisms which belong to the same link of the food chain as counted from the producer level are said to belong to the same trophic level. Thus the plants constitute the first trophic level, the herbivores the second, and the carnivores feeding on herbivores the third trophic level. Secondary

---

Reprinted from the CMFRI, FRAD. 2014. Training Manual on Fish Stock Assessment and Management, p.150.



carnivores feeding on third level carnivores belong to the fourth trophic level and so forth. However, there is a very definite limit to the number of possible links in a food chain, and consequently also to the number of trophic levels in any ecosystem. The reason for this is that only about 10 percent of the available energy is assimilated in passing from one trophic level to the next. At the top of the food chain there are usually only one or two major predators. The number of species in each trophic layer increases with approach to the first layer, giving rise to what is called a pyramid of numbers. For the major predators introduction of small amounts of pollutants into the first trophic layer can have fatal consequences because it is eventually concentrated in them.

### **Gross Production and Net Production**

Only a very small portion of the light energy absorbed by green plants that is transformed into food energy (gross production) because most of it is dispersed as heat. Furthermore, some of the synthesized gross production is used by the plants in their own respiratory processes, leaving a still smaller amount of potential energy (the net production) available for transfer to the next trophic level.

### **The Loss of Energy**

True production of organic matter takes place only in the chlorophyll-possessing plants and certain synthetic bacteria, and this has been referred to as the primary production. Copepods and euphausiids, convert plant material into protein that can be assimilated by the animals which eat them but which themselves could not exist on plant material. In reality, of course, they only assimilate and store energy derived from the primary producers. They are called secondary producers, a term which of course fits animals at higher trophic levels just as well because they too - although indirectly - utilize the primary production of the plants. The loss of energy is generally referred to as the respiratory loss because the organisms utilize the food energy by oxidizing it. Because of the respiratory losses the food chains cannot be very long and the number of trophic levels in natural communities is therefore seldom more than four or five and often only three. It also means that the total amount of food available decreases with increasing trophic level. For this reason, the largest animals are found feeding on either plants or other animals which are in a low trophic level as, for example, whales on krill and elephants on plants.

### **Studying Food and Feeding of Fishes**

The study of the feeding habits of fish and other animals based upon analysis of stomach content has become a standard practice (Hyslop 1980). Stomach content analysis provides important insight into fish feeding patterns and quantitative assessment of food habits is an important aspect of fisheries management. Lagler (1949) pointed out that the gut contents only indicate what the fish would feed on. Accurate description of fish diets and feeding habits also provides the basis for understanding trophic interactions in aquatic food webs. Diets of fishes represent an integration of many important ecological components that included behavior, condition, habitat use, energy intake and inter/intra specific interactions. A food habit study might be conducted to determine the most



frequently consumed prey or to determine the relative importance of different food types to fish nutrition and to quantify the consumption rate of individual prey types. Each of these questions requires information on fish diets and necessitates different approaches in how one collects and analyzes data. Here, we outline qualitative and quantitative techniques used to describe food habits and feeding patterns of fishes. For a better understanding of diet data and for accurate interpretation of fish feeding patterns, time of day, sampling location, prey availability and even the type of collecting gear used need to be considered before initiating a diet study or analyzing existing diet data.

Stomach contents can be collected either from the live or fresh died fish. Regardless of the method, investigators should ensure that the removal technique effectively samples all items in the gut. Other wise data will be skewed toward items that are more easily displaced from the stomach. Alternatively, live fish can be sacrificed and stomach contents removed for analysis. If fish are to be sacrificed, they should be preserved immediately either by freezing or by fixing in formalin. Stomach contents will continue to digest, rendering rapid preservation of the fish or removed contents necessary to prevent loss of resolution. As in most fish groups feeding behavior of juveniles and adults vary distinctly attention should be taken to encounter more samples which will include all size groups of the particular fish. The specimens either from live or preserved should be measured to its total length to the nearest 1mm and weight to the nearest 0.1 g. Cut open the fish and record the sex and maturity stage of the fish. Remove the stomach and preserve them in 5% neutralized formalin for further analysis. For the analysis, a longitudinal cut must be made across the stomach and the contents are transferred into a petri dish. The contents then keep for five minutes to remove excess formalin and then examine under binocular microscope. Identify the gut content up to the genus and if possible up to species level depending up on the state of digestion. Various taxa digest at different rates. As such, all recently consumed taxa may be present in the foregut but only resistant items remain in the hindgut. To avoid bias when both easily digested prey and resistant prey are present, only the immediate foregut (e.g., stomach) should be sampled.

Prey items in fish stomachs are often not intact. Hard parts such as otoliths, scales, cleithra or backbones have diagnostic, species specific characteristics useful for identifying prey. Alternatively, partially digested prey may be identified using unique biochemical methods such as allozyme electrophoresis, or immunoassays. An important fact assessed by the examination of the stomach is the state or the intensity of feeding. This is judged by the degree of distension of the stomach or by the quantity of food that is contained in it. The distension of the stomach is judged and classified as 'gorged or distended', 'full', '3/4full', '1/2full' etc by eye estimation.

Fish diets can be measured in a variety of ways. Methods of gut contents analysis are broadly divisible into two, viz., qualitative and quantitative. The qualitative analysis consists of a complete identification of the organisms in the gut contents. Only with extensive experience and with the aid of good references it is possible to identify them from digested, broken and finely comminuted



materials. Quantitative methods of analysis are three types, viz., numerical, gravimetric and volumetric. All these types of analysis are widely employed by different workers. The following outline of methods is based mainly on the reviews by Hynes (1950), Pillay (1952), Windell (1968), Hyslop (1980) and Chipps et al (2002 ).

### 1. Numerical Methods

The numerical methods are based on the counts of constituent items in the gut contents. The numerical methods have been adapted in different ways to assess the relative importance of food items and these can be classified under four distinct heads, viz., a) Occurrence, b) Dominance, c) Number and d) Point (Numerical) methods.

**a) Frequency of Occurrence:** Stomach contents are examined and the individual food organisms sorted and identified. The number of stomachs in which each item occurs is recorded and expressed as a percentage of the total number of stomachs examined.

Frequency of Occurrence, =  $O_i = \frac{J_i}{P}$ , where,  $J_i$  is number of fish containing prey  $i$  and  $P$  is the number of fish with food in their stomach.

This method demonstrates what organisms are being fed upon, but it gives no information on quantities or numbers and does not take in to consideration the accumulation of food organisms resistant to digestion. For instance, three organisms in a stomach, say, prawn, rotifers and diatoms, present in the ratio of 1:200:2000 would all be treated by this method as 1:1:1 with reference to the stomach in question. This method holds good even when there is differential distribution of various food organisms in the water for the same reason that it is not biased by size or numbers of organism comprising the food. Many have used this method as an indicator of inter-specific competition while some utilized this method to illustrate the seasonal changes in diet composition.

**b) Number Method:** The number of individual of each food type in each stomach is counted and expressed as a percentage of the total number of food items in the sample studied, or as a percentage of the gut contents of each specimen examined, from which the total percentage composition is estimated.

Percent by number,  $N_i = \frac{N_i}{\sum_{i1}^Q N_i}$ , where,  $N_i$  is the number of food category  $i$

This method has been employed successfully by several workers in studies on the food of plankton feeding fishes where the items can be counted with ease. In the basic number method, no allowance is made for the differences in size of food items. So in the studies on the food of fishes other than plankton feeders, the number method has very limited use. The counting of comminuted plant matter in the stomach of fish is impracticable and will not yield correct



evaluations. So also in the analysis of the gut contents of a carnivore which may consist of only one large sized fish and a couple of small larvae, the counting are of little value computations. These are summed to give totals for each kind of food item in the whole sample, and then a grand total of all items. The quotient of these gives the percentage representation, by number, of each type of food item.

- c) Dominance Method:** Essentially the dominance method is a partial improvement of the occurrence method, viz., the lack of consideration of the quantities of the food items present in the stomach, sought to be remedied. The stomach contents comprising the main bulk of the food materials present, is determined and the number of fish in which each such dominant food material is present is expressed as a percentage of the total number of fishes examined. The percentage composition of the dominant food materials can also be expressed by this method as in the occurrence method.

Though in an analysis of dominance the bulk of the food material is taken in to account, it can yield only a very rough picture of the dietary of a fish. Moreover, items which are less dominant due to environmental reasons may escape notice. Though this defect can also be remedied to a certain extent by the examination of large samples spread over a long period of time, a system of assay that takes in to account the relative importance of food constituents will obviously be more suitable in gut content analysis.

- d) Points (Numerical) Method:** The points method is an improvement on the numerical method where consideration is given to the bulk of the food items. The simple form of points method is the one in which the counts are computed falling a certain organisms as the unit. In a more modified form, the food items are classified as 'very common', 'common', 'frequent', 'rare', etc., based on rough counts and judgments by the eye. In this arbitrary classification the size of the individual organisms is also given due consideration. The contents of all stomachs are then tabulated and as a further approximation, different categories are allotted a certain number of points and the summations of the points for each food item are reduced to percentages to show the percentage composition of the diet. This method is essentially a numerical one; the volume being only a secondary consideration and it is only in the counts that a certain amount of accuracy can be claimed.

## **2. Volumetric Methods**

Many workers consider the volume as a more satisfactory method for quantitative analysis of gut contents. As Hynes (1950) pointed out, volume forms a very suitable means of assessment, this is especially so in the case of herbivorous and mud feeding fishes where the numerical methods "become meaningless as well as inaccurate". Even in cases where the numerical methods are suitable, volume has been considered as an essential factor to be reckoned with, and in all improved numerical methods the volume of the food items is taken in to consideration in some way or other.



The chief methods that are employed in assessing the volume of food items in the gut contents of fishes are:

- a) Eye estimation Method:** This is probably the simplest and easiest means of determining the volume of food constituents. In this method the contents of each sample is considered as unity, the various items being expressed in terms of percentage by volume as estimated by inspection. This method of analysis is subjective in nature and the investigator's personal bias is likely to influence the results very greatly. This defect can be minimized to a great extent by the examination of large samples conducted over a long period.
- b) Points (Volumetric) Method:** This method is a variation of the eye estimation method. Here instead of directly assessing the volume by sight as in the previous method, each food item in the stomach is allotted a certain number of points based on its volume. Certain workers have taken into account both the size of the fish and the fullness of the stomach in the allotment of points. The diet component with highest volume was given 16 points. Every other component was awarded 16, 8, 4, 2, 1 and 0 points depending on the volume relative to the component with the highest volume. Percentage volumes within each subsample were calculated as:

$$\alpha = \frac{\text{Number of points allocated to component a}}{\text{Total points allocated to sub sample}} \times 100$$

where,  $\alpha$  is the percentage volume of the prey component a

This method is quite useful for analyzing omnivorous and herbivores where measuring volumes of microscopic organisms such as diatoms and filamentous algae are very difficult.

- c) Displacement Method:** The displacement method is probably the most accurate one for assessing the volume. The volume of each food item is measured by displacement in a graduated container such as a cylinder with the smallest possible diameter for accuracy. This method is eminently suited in the estimation of the food of carnivorous fishes. But the differential rate of digestion of the food items may sometimes affect the accuracy of the observations. However, if the collections are made when the fish are on feed, this defect can be easily overcome. A knowledge of the volumes of the different size groups of the food items may be of great help in estimating the volume of the whole item from the semi digested fragments

### 3. Gravimetric Method

The gravimetric method consists of the estimation of the weight of each of the food items, which is usually expressed as percentages of the weight of the total gut contents as in other quantitative methods.

$$\text{Percent by weight, } w_i = \frac{W_i}{\sum_{i=1}^Q W_i},$$

Where,  $W_i$  is the weight of the prey i



Generally the wet weigh of the food after removing superfluous water buy pressing it dry between filter papers is taken for this purpose. Dry weight estimation is more time consuming and is usually employed where accurate determinations of calorific intake is required. The limitation of weight as a criterion of analysis has already been referred in the consideration of the method of assessing the condition of feed. Besides these, the accurate weighing of small quantities of food matter is extremely difficult and impracticable in studies of large collections. This method is, therefore generally employed only in conjunction with other methods to demonstrate seasonal variations in the intensity of feeding.

**Table:** Example of results obtained using different methods of estimation of stomach contents for two numbers of *Lactariuslactarius* (L1)

*L. lactarius 1* (L1). 1. *Stolephorusbataviensis*, 9 cm long, weight 5 g, volume 7 ml, 6 *Acetes* each 3.0cm long, weight 300mg vol. 2ml, 1 *Bregmaceros* ,4cm, 1 g, vol. 1 ml.

*L. lactarius2* (L2). 1. *Stolephoursbataviensis*, 7 cm long, weight 3 g, volume 4 ml, 4 *Acetes* 2.5 cm long, weight 250 mg, vol.1 ml.

Food	Method	Fish			%	Total of which % expressed
		LI1	LI2			
<i>S. bataviensis</i>	Occurrence	1	1	2	40	All food occurrences
<i>Acetes</i>		1	1	2	40	
<i>Bregmaceros</i>		1	0	1	20	
<i>S. bataviensis</i>	Numerical	1	1	2	15.4	All food organisms
<i>Acetes</i>		6	4	10	76.9	
<i>Bregmaceros</i>		1	0	1	7.7	
<i>S. bataviensis</i>	Dominance	1	1	2	100	All fish
<i>Acetes</i>		1	1	2	100	
<i>Bregmaceros</i>		1	0	1	50	
Food	Method	Fish			%	Total of which % expressed
		LL1	LL2			
<i>S. bataviensis</i>	Total Volume	7	4	11	73.3	Total food volume
<i>Acetes</i>		2	1	3	20	
<i>Bregmaceros</i>		1	0	1	6.7	
<i>S. bataviensis</i>	% volume	70	80	75	75	Food volume
<i>Acetes</i>		20	20	20	20	
<i>Bregmaceros</i>		10	0	5	5	
<i>S. bataviensis</i>	Gravimetric	5	3	8	67.8	Total weight of food
<i>Acetes</i>		1.8	1	2.8	23.7	
<i>Bregmaceros</i>		1	0	1	8.5	



## Food Analysis Indices

### A. Simple Indices

- 1) Index of Fullness:** This is measured as the ratio of food weight to body weight as an index of fullness, which is very widely employed. (The ratio of corresponding volume can also be used.) This index can be applied to the food in the stomach, or to that in the whole digestive tract. It is usually expressed as parts per 10,000 (%00, or parts per decimile); that is:

$$\text{Fullness index} = \frac{\text{weight of the stomach contents} \times 10,000}{\text{weight of fish}}$$

- 2) Index of Selection or Forage Ratio:** Most fishes have a scale of preference for the organisms in their environment, so that some are consumed in large numbers, others moderately, some not at all. A quantitative index of such differences called as the forage ratio. A study of the quantities of different organisms available to the fish is made, and also of the various items in their stomachs; then;

$$\text{Selection index} = \text{forage ratio} = \frac{s}{b}$$

where,  $s$  = percentage representation by weight, of a food organism in the stomach and  $b$  = percentage representation of the same organism in the environment. The lower limit for this index is 0; its upper limit is indefinitely large.

- 3) Index of Electivity:** Ivlev (1961) proposed a somewhat different quantitative measure of selection which has been widely used as mean of comparing the feeding habits of fishes and other aquatic organisms with the availability of potential food resources in natural habitats. The relationship is defined as

$$\text{Electivity index} = E = \frac{s - b}{s + b}$$

The index has a possible range of -1 to +1, with negative values indicating avoidance or inaccessibility of the prey item, zero indicating random selection from the environment, and positive values indicating active selection.

### B. Compound Indices

In an attempt to consolidate the desirable properties of individual diet measures (e.g.,  $N_i$ ,  $W_i$ ,  $F_i$ ), compound indices were developed that combine two or more measures into a single index. The belief is that compound indices capture more information than do single component measures (Chippset *al* 2002).





### 1) Index of Preponderance: (Natarajan and Jhingran, 1961)

This index gives a summary picture of frequency of occurrence as well as bulk of various food items. It provides a definite and measurable basis of grading the various food elements. The bulk of food items can be evaluated by 1) Numerical 2) volumetric and 3) Gravimetric methods. As the numerical method is not suited to the index with the frequency of occurrence it magnifies the importance of smaller organisms which may appear in enormous numbers. Therefore either volumetric or gravimetric are best to assess the food items quantitatively. If we  $V_i$  and  $O_i$  are the volume and occurrence index of food item i. then,

$$\text{Index of preponderance } I_i = \frac{V_i O_i}{\sum V_i O_i} \times 100$$

Example: The 'Index of Preponderance' of food items of *Catlacatla* (Ham.) is given in the table 2 with rankings in brackets.

**Table 2 :** Index of Preponderance (Natarajan and Jhingran, 1961) of adult *Catlacatla*

Food items	Percentage of Occurrence ( $O_i$ )	Percentage of volume ( $V_i$ )	$V_i O_i$	$\frac{V_i O_i}{\sum V_i O_i} \times 100$
Crustaceans	24.5	57.1	1398.95	64.50 (1)
Algae	27.3	24.0	655.20	30.06 (2)
Plants	6.4	8.2	52.48	2.41 (3)
Rotifers	10.8	2.4	25.92	1.19 (4)
Insects	3.6	6.0	21.60	0.99 (5)
Protozoa	0.6	0.3	0.18	0.01 (8)
Molluscs	....	....	.....	.....
Polyzoa	....	....	.....	. ....
Detritus	10.0	1.3	13.00	0.60 (6)
Sand and mud	16.8	0.7	11.76	0.54 (7)
$\Sigma$	<b>100</b>	<b>100</b>	<b>2179.09</b>	<b>100</b>

According to the index crustaceans and algae constitute 1 and 2 ranks in *Catla catla*. While third, fourth and fifth places are held by plants, rotifers and insects. In grading the food elements accidental and incidental inclusions like sand, mud, etc., may be left out of consideration.



## 2) Index of Relative Importance (IRI):- Leo Pinkas et al (1971)

This index is an integration of measurement of number, volume and frequency of occurrence to assist in evaluating the relationship of the various food items found in the stomach. It is calculated by summing the numerical and volumetric percentages values and multiplying with frequency of occurrence percentage value.;

$$\text{Index of relative importance, } IRI_i = (\% N_i + \% V_i) \% O_i,$$

where,  $N_i$ ,  $V_i$  and  $O_i$  represent percentages of number, volume and frequency of occurrence prey  $i$  respectively.

*Example:* Index of Relative Importance of pelagic preflexion summer flounder, *Paralichthysdentatus* larvae (Grover, 1998) with ranking in brackets

Prey	% $N_i$	% $V_i$	% $O_i$	(% $N_i$ + % $V_i$ ) % $O_i$	%IRI
Tintinnids	28.7	3.3	37.6	1203.2	19.3 (3)
Copepod	20.0	10.2	41.2	1244.24	20.0 (2)
nauplii	16.0	61.4	30.0	2322	37.3 (1)
Copepodites	0.6	4.9	2.0	11	0.2 (8)
Calanoids	0.6	2.0	2.4	6.24	0.1 (9)
Cyclopoids	16.0	1.2	34.8	598.56	9.6 (5)
Copepod eggs	12.1	14.8	28.0	753.2	12.1 (4)
Bivalve larvae	3.7	0.9	11.6	53.36	0.9 (6)
Invertebrate	2.3	1.3	9.2	33.12	0.5 (7)
eggs					
Other					

In pelagic preflexion summer (*Paralichthy dentatus*) larvae, copepodites composed the bulk of the diet (61.4% Vol, 37.3 % IRI) and formed the most important prey. Copepod nauplii, the second most important prey, composed 20.0% (N and IRI). Tintinnids, despite being the most abundantly ingested prey (28.7% N); ranked third in importance at 19.3% (IRI). Bivalve larvae and copepod eggs were the only other prey that accounted for >1% of the diet, and together they composed 21.7% (IRI).



## Suggested Reading

- Chipps S. R. and E. J. Garvey 2002. *Assessment of Food Habits and Feeding Patterns*, USGS South Dakota Cooperative Fish and Wildlife Research Unit, Department of Wildlife and Fisheries Sciences, South Dakota State University, Brookings, SD 57007.
- FAO, 1974. *Manual of Fisheries Science Part 2 - Methods of Resource Investigation and their Application*. Fish.Tech paper 115.p.255.
- Grover, J. J. 1998. Feeding habits of pelagic summer flounder, *Paralichthys dentatus*, larvae in oceanic and estuarine habitats. Fish.Bull, 90 (2): 248-257.
- Hynes, H. B. N. 1950. The food of the freshwater sticklebacks (*Gastrosteus aculeatus*) and *Pygosteus pungitius*) with a review of methods used in studies of the food of fishes. J. Anim. Ecol., 19: 36-58.
- Hyslop, E. J. 1980. Stomach contents analysis: a review of methods and their application. *J.Fish. Biol*, 17:411-429.
- Ivlev, V. S. 1961. Experimental ecology of the feeding of fishes. Yale University Press, New Haven, Conn.
- Lagler, K. F. 1949. *Studies in freshwater biology*, Ann Arbor, Michigan.
- Natarajan, A. V. and A. C. Jhingran. 1961. 'Index of preponderance'-a method of grading the food elements in the stomach analysis of fishes. *Indian J. Fish*, 8: 54-59.
- Pillay, T. V. R. 1952. A critique of the methods of study of food of fishes. *J. zool. Soc. India.*, 4: 1885-200.
- Pinkas, L., M. S. Olipahnt, and I. L. K. Iverson 1971. Food habits of albacore, bluefin tuna, and bonito in Californian waters. Calif. Dep.Fish Game, *Fish. Bull*, 152: 1-105.
- Seaburg, K. G. 1957. A stomach sampler for live fish. *Progre. Fish. Cult.* 19: 137- 144
- Shchoener, T. W. 1970. Non synchronous spatial over lap of lizards in patchy habitats. *Ecol*, 51: 408-418.
- Strauss, R. E. 1979. Reliability estimates for Ivlev's electivity index, the forage ratio, and a proposed linear index of food selection. *Transactions of the American Fisheries Society* 108:344-352.
- Windell, J. T. 1968. Food analysis and rate of digestion. In. W. E. Ricker (editor), methods of assessment of fish production in fresh waters, 2nd ed., P. 215-226. IBP (Int. Biol. Programme). Handb.3.

