



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
Main Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2017

Delayed cerebral ischemia after subarachnoid hemorrhage: comparing and integrating classification systems

Kubben, Pieter L; Germans, Menno R; Jabbarli, Ramazan

DOI: <https://doi.org/10.4103/2152-7806.208805>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-139588>

Published Version



Originally published at:

Kubben, Pieter L; Germans, Menno R; Jabbarli, Ramazan (2017). Delayed cerebral ischemia after subarachnoid hemorrhage: comparing and integrating classification systems. *Surgical Neurology International*, 8(1):121.

DOI: <https://doi.org/10.4103/2152-7806.208805>

Letter to the Editor

Delayed cerebral ischemia after subarachnoid hemorrhage: Comparing and integrating classification systems

Pieter L. Kubben, Menno R. Germans¹, Ramazan Jabbarli²Department of Neurosurgery, Maastricht University Medical Center, Maastricht, ¹Department of Neurosurgery, University Hospital Zurich, Switzerland, ²Department of Neurosurgery, University Hospital of Essen, Essen, GermanyE-mail: *Pieter L. Kubben - p.kubben@mumc.nl; Menno R. Germans - Menno.Germans@usz.ch;
Ramazan Jabbarli - ramazan.jabbarli@uniklinik-essen.de
*Corresponding author

Received: 06 April 16 Accepted: 12 December 16 Published: 21 June 17

Sir,

Subarachnoid hemorrhage (SAH) is a significant healthcare problem. It is estimated that 30000 Americans suffer from aneurysmal SAH each year. Delayed cerebral ischemia (DCI) is one of the major causes of morbidity and mortality after SAH, and occurs in 20–40% of patients.^[4,5,8] To better predict and diagnose DCI after SAH, multiple grading scales have been published, of which the Fisher score is the most well-known.^[3] More recently, this score has been modified to predict symptomatic vasospasm after SAH more accurately.^[4] Many other prediction models have been developed and published, and some of them use different parameters to calculate various outcomes that are associated with DCI and vasospasm.^[1,2,7] Here, we present the SAH DCI app, which was developed to facilitate comparison between the various grading scales and to aid physicians in the use of prediction models. It incorporates automated calculation of a maximum of seven SAH grading scales, namely, the WFNS SAH grade, the Hunt and Hess grade, the original and modified Fisher grade, the early prediction scale by De Rooij *et al.*, the BEHAVIOR score, and the VASOGRADE score.^[1-4,6,7,9] The WFNS SAH grade and the Hunt and Hess grade are general outcome predictors, and the other classification systems are specific predictors for DCI.

We developed and evaluated a mobile medical app for iOS (Apple Inc, Mountain View, CA) using class 1 CE-marked technology for risk minimization. The app, called SAH DCI, has been evaluated by all authors independently, where the results of the mobile clinical decision support system were compared with manually calculated scores, which were based on the original classification systems. The app provided identical results compared to manual calculation of the relevant classification systems.




Figure 1: SAH DCI basic module (3 classification systems)

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Access this article online	
Quick Response Code:	Website: www.surgicalneurologyint.com
	DOI: 10.4103/2152-7806.208805

How to cite this article: Kubben PL, Germans MR, Jabbarli R. Delayed cerebral ischemia after subarachnoid hemorrhage: Comparing and integrating classification systems. *Surg Neurol Int* 2017;8:121.
<http://surgicalneurologyint.com/Delayed-cerebral-ischemia-after-subarachnoid-hemorrhage--Comparing-and-integrating-classification-systems/>

DCI risk after SAH Info < Back **Advanced Results**

Risks for delayed cerebral ischemia (DCI) after SAH according to different grading systems.

Age < 55 years

Age ≥ 55 years ✓

BEHAVIOR Score

- Score: 3
- Risk percentage cerebral infarction: 49.6 (95% CI: 40.8–58.4)

VASOGRADE

- Grade: Red
- Odds ratio DCI: 3.19 (95% CI 2.07 - 4.50)

Risk Chart (from De Rooij, et al.)

- Risk percentage DCI: 26%.

Modified Fisher

- Score: 4
- Odds ratio vasospasm: 2.2 (95% CI 1.6 - 3.1)

Hunt & Hess grade

- Grade: 4

EARLY VASOSPASM ON DSA

No ✓

Yes

INTRACRANIAL PRESSURE

ICP ≤ 20mm Hg ✓

ICP > 20mm Hg

TREATMENT OF MULTIPLE ANEURYSMS

No ✓

Yes

HUNT & HESS GRADE

Basic Advanced Basic Advanced

Figure 2: SAH DCI advanced module (7 classification systems)

The app consists of two screens, a basic module and an advanced module. The basic module asks 4 questions to calculate 3 grading scales (WFNS SAH Grade,^[9] Modified Fisher,^[4] and VASOGRADE^[11]) An example is illustrated in Figure 1. The advanced module asks 10 questions to calculate 7 grading scales (the WFNS SAH grade,^[9] the Hunt and Hess grade,^[6] original and modified Fisher grade,^[3,4] early prediction scale by De Rooij *et al.*,^[2] the BEHAVIOR score,^[7] and the VASOGRADE score^[11]). An example is illustrated in Figure 2. Both the modules offer the ability to export all user input and provide recommendation by email for later reference [Figure 3]. Email is sent unencrypted, which is clearly displayed. No patient identification details are necessary to use the app.

In conclusion, the SAH DCI app offers clinicians a user-friendly and accurate tool for mobile clinical decision support on DCI after SAH. It combines multiple classification systems for risk calculation that may help to guide treatment, as well as ease and quicken the clinical implication of different SAH scores. Further research and comparative effectiveness studies focusing on different classification systems are needed to extract the main parameters that predict DCI after SAH.

Financial support and sponsorship

Nil.

Cancel SAH DCI advanced results re... Send

Input parameters

- Glasgow Coma Scale: 7 - 12
- Motor deficit: Yes
- Subarachnoid hemorrhage: Thick
- Intraventricular hemorrhage: Thin
- Age: Age ≥ 55 years
- Early vasospasm on DSA: No
- Intracranial pressure: ICP ≤ 20mm Hg
- Treatment of multiple aneurysms: No
- Hunt & Hess grade: Stuporous; moderate to severe hemiparesis; possibly early decerebrate rigidity and vegetative disturbances
- Need for external ventricular shunt: No

Advanced results

Risks for delayed cerebral ischemia (DCI) after SAH according to different grading systems.

BEHAVIOR Score

Figure 3: Export results by email dialog

Conflicts of interest

The first author is the developer of the app. Other authors report no conflict of interest.

REFERENCES

- de Oliveira Manoel AL, Jaja BN, Germans MR, Yan H, Qian W, Kouzmina E, *et al.* The VASOGRADE: A Simple Grading Scale for Prediction of Delayed Cerebral Ischemia After Subarachnoid Hemorrhage. *Stroke* 2015;46:1826-31.
- de Rooij NK, Greving JP, Rinkel GJE, Frijns CJM. Early Prediction of Delayed Cerebral Ischemia After Subarachnoid Hemorrhage: Development and Validation of a Practical Risk Chart. *Stroke* 2013;44:1288-94.
- Fisher CM, Kistler JP, Davis JM. Relation of Cerebral Vasospasm to Subarachnoid Hemorrhage Visualized by Computerized Tomographic Scanning. *Neurosurgery* 1980;6:1-9.
- Frontera JA, Claassen J, Schmidt JM, Wartenberg KE, Temes R, Connolly ES, *et al.* Prediction of Symptomatic Vasospasm after Subarachnoid Hemorrhage: The Modified Fisher Scale. *Neurosurgery* 2006;59:21-7.
- Harrod CG, Bendok BR, Batjer HH. Prediction of Cerebral Vasospasm in Patients Presenting with Aneurysmal Subarachnoid Hemorrhage: A Review. *Neurosurgery* 2005;56:633-54.
- Hunt WE, Hess RM. Surgical risk as related to time of intervention in the repair of intracranial aneurysms. *J Neurosurg* 1968;28:14-20.
- Jabbarli R, Reinhard M, Roelz R, Shah M, Niesen W-D, Kaier K, *et al.* Early identification of individuals at high risk for cerebral infarction after aneurysmal subarachnoid hemorrhage: The BEHAVIOR score. *J Cereb Blood Flow Metab* 2015;35:1587-92.
- MacDonald RL. Delayed neurological deterioration after subarachnoid haemorrhage. *Nature* 2014;10:44-58.
- Teasdale GM, Drake CG, Hunt W, Kassell N, Sano K, Pertuiset B, *et al.* A universal subarachnoid hemorrhage scale: Report of a committee of the World Federation of Neurosurgical Societies. *J Neurol Neurosurg Psychiatr* 1988;51:1457-7.