

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26

The primosomal protein DnaD inhibits cooperative DNA binding by the  
replication initiator DnaA in *Bacillus subtilis*

Carla Y. Bonilla and Alan D. Grossman\*

Department of Biology

Massachusetts Institute of Technology

Cambridge, MA 02139

Running title: Primosomal protein DnaD affects DNA binding by DnaA

\*Corresponding author:  
Department of Biology  
Building 68-530  
Massachusetts Institute of Technology  
Cambridge, MA 02139  
phone: 617-253-1515  
fax: 617-253-2643  
E-mail: [adg@mit.edu](mailto:adg@mit.edu)

27 **Abstract**

28 DnaA is a AAA+ ATPase and the conserved replication initiator in bacteria. Bacteria control  
29 the timing of replication initiation by regulating the activity of DnaA. DnaA binds to multiple  
30 sites in the origin of replication (*oriC*) and is required for recruitment of proteins needed to load  
31 the replicative helicase. DnaA also binds to other chromosomal regions and functions as a  
32 transcription factor at some of these sites. *Bacillus subtilis* DnaD is needed during replication  
33 initiation for assembly of the replicative helicase at *oriC* and during replication restart at stalled  
34 replication forks. DnaD associates with DnaA at *oriC* and at other chromosomal regions bound  
35 by DnaA. Using purified proteins, we found that DnaD inhibited the ability of DnaA to bind  
36 cooperatively to DNA and caused a decrease in the apparent dissociation constant. These effects  
37 of DnaD were independent of the ability of DnaA to bind or hydrolyze ATP. Other proteins  
38 known to regulate *B. subtilis* DnaA also affect DNA binding, whereas much of the regulation of  
39 *E. coli* DnaA affects nucleotide hydrolysis or exchange. We found that the rate of nucleotide  
40 exchange for *B. subtilis* DnaA was rapid and not affected by DnaD. The rapid exchange is  
41 similar to that of *Staphylococcus aureus* DnaA and in contrast to the slow exchange rate of  
42 *Escherichia coli* DnaA. We suggest that organisms in which DnaA has a rapid rate of nucleotide  
43 exchange predominantly regulate the DNA binding activity of DnaA and those with slow rates of  
44 exchange regulate hydrolysis and exchange.

45

46

47

## 48        **Introduction**

49        Accurate and complete replication of DNA is essential for the propagation of genomic  
50 information. DNA replication in bacteria initiates from a single origin of replication (*oriC*) and  
51 depends on the conserved AAA+ ATPase DnaA (reviewed in 13, 25, 27, 29, 35, 45). DnaA  
52 binds both ATP and ADP, and DnaA-ATP is required for replication initiation (2, 17, 33, 47,  
53 57). The ATP bound form of DnaA forms oligomers that are important for promoting replication  
54 initiation (14, 18, 35, 41, 58). DnaA-ATP binds to sites in *oriC* and promotes the unwinding of  
55 the DNA Unwinding Element (DUE), which serves as a platform for assembly of the replicative  
56 helicase and the rest of the replication machinery. Wherever examined, the nucleotide bound  
57 state of DnaA controls its activity (e.g., 39, 42, 57, 63), and this can be affected by factors that  
58 alter nucleotide hydrolysis and exchange (16, 26, 28, 64). The rate of nucleotide exchange for  
59 purified *E. coli* DnaA is relatively slow, with a half-life of 45 min (57). This slow inherent rate  
60 of exchange enables the modulation of DnaA activity by factors that stimulate the rate of  
61 nucleotide hydrolysis and/or exchange (11, 16, 26, 64).

62        In contrast to the widespread conservation of DnaA, other proteins required for replication  
63 initiation are less conserved. For example, steps involved in loading the replicative helicase at  
64 *oriC* are different between *E. coli* and *Bacillus subtilis*. In *B. subtilis* and other low G+C Gram  
65 positive bacteria, helicase loading requires the essential primosomal proteins DnaD, DnaB, and  
66 DnaI, in addition to DnaA (5-8, 12, 36, 37, 54, 59, 62, 65). DnaD is associated with *oriC* and this  
67 association depends on DnaA (54, 59). Association of DnaB with *oriC* depends on DnaD, and  
68 finally, DnaI-mediated assembly of the helicase at *oriC* depends on DnaB. DnaD and DnaB bind  
69 both double and single stranded DNA, which may help stabilize opening up of the origin of  
70 replication (7, 38, 68).

71 In addition to binding to sites in *oriC*, *B. subtilis* DnaA binds to sites in chromosomal regions  
72 outside of *oriC* (4, 20, 24). DnaA functions as a transcription factor at some of these secondary  
73 binding regions (4, 9, 20, 24). Where tested, DnaD and DnaB are also found associated with  
74 these secondary DnaA binding regions, and this association depends on DnaA (60). However, in  
75 contrast to *oriC*, these secondary DnaA-binding regions do not function as origins of replication  
76 and there is no indication that they are capable of loading the replicative helicase (60).

77 Because of the association of DnaD with DnaA at multiple regions throughout the  
78 chromosome, we hypothesized that DnaD modulates the activity of DnaA. Some factors that  
79 affect the activity of DnaA, predominantly in *E. coli*, are known to alter its nucleotide-bound  
80 state (16, 26, 64). In contrast, regulators of *B. subtilis* DnaA, (e.g., YabA, Soj, and SirA) are  
81 known to alter its DNA binding properties (40, 53, 55, 66). Using purified proteins, we tested for  
82 effects of DnaD on both the ability of DnaA to bind DNA and on nucleotide exchange.

83 We found that the rate of nucleotide exchange for *B. subtilis* DnaA was relatively rapid,  
84 similar to that of DnaA from *Staphylococcus aureus* (32), and in contrast to the slow rate of  
85 exchange for DnaA from *E. coli* (57). DnaD had no effect on the rate of exchange of *B. subtilis*  
86 DnaA. In contrast, DnaD had a marked effect on the ability of DnaA to bind DNA. Binding of  
87 DnaA-ATP to DNA fragments that contain multiple binding sites is normally highly cooperative  
88 (40, 41). We found that in the presence of DnaD, binding of DnaA to DNA was no longer  
89 cooperative and the apparent dissociation constant for DnaA and DNA was reduced. We found  
90 that the ATPase activity of DnaA was not needed for these effects by DnaD, indicating that  
91 DnaD is not regulating the ATPase activity of DnaA. These effects of DnaD on the ability of  
92 DnaA to bind DNA are similar to the effects of YabA (40) and Soj (55), two other regulators of  
93 *B. subtilis* DnaA and replication initiation and further substantiate the notion that modulation of

94 cooperative binding and oligomerization of DnaA to DNA might be a common mechanism of  
95 regulation (40, 55).

96

97

## 98 **Materials and Methods**

### 99 **Purification of DnaA and DnaD**

100 *B. subtilis* DnaA (no tag) was produced in and purified from an *E. coli dnaA* null mutant,  
101 using a clone and strain provided by A. Albuzzi and W. F. Burkholder, essentially as described  
102 (17, 60). Protein was stored frozen (-80 °C) in buffer containing 45 mM Hepes pH 7.6, 0.5 mM  
103 EDTA, 10 mM magnesium acetate, 1 mM DTT, 700 mM potassium glutamate, and 20%  
104 sucrose. DnaD-his6 was produced in and purified from *E. coli*, essentially as described (60).  
105 Protein was stored at -80°C in buffer containing 50 mM Tris pH 8, 0.1 mM EDTA, 1 mM DTT,  
106 500 mM NaCl and 10% Glycerol. Proteins were quantified using absorbance at 280 nm.

### 107 **Nucleotide exchange reactions**

108 Nucleotide exchange was measured using alpha-<sup>32</sup>P-ATP or <sup>14</sup>C-ADP. Exchange reactions  
109 contained 40 mM Hepes pH 7.5, 10 mM magnesium acetate, 0.5 mM EDTA, 1 mM DTT, 150  
110 mM potassium glutamate, 100 µg/ml BSA, 10% glycerol and 1 µM alpha-<sup>32</sup>P-ATP or 1 µM <sup>14</sup>C-  
111 ADP in the absence or presence of DnaD (600 nM). DnaA (300 nM) was incubated with either  
112 nucleotide for 2 hours on ice in exchange buffer. Fifty microliters were removed at time zero to  
113 measure binding and unlabeled ATP was added in excess (2 mM) and incubated at 37°C. Filter  
114 binding was used to measure the amount of radio-labeled nucleotide still bound to DnaA at each  
115 time point. Fifty microliter aliquots were removed and placed on equilibrated nitrocellulose  
116 membranes (Millipore), washed with buffer (40 mM Hepes pH 7.5, 150 mM KCl, 10 mM

117 magnesium acetate, 0.5 EDTA, 10ug/ml BSA) and the amount of  $^{32}\text{P}$ -ATP or  $^{14}\text{C}$ -ADP  
118 remaining on the filter was measured in triplicate and averaged. The half-life was calculated  
119 using an exponential decay formula and plotted using GraphPad Prism 5 software.

## 120 **Gel Shift Assays**

121 The DNA template from the *oriC* region (*dnaA* promoter region) used for the gel shift assays  
122 was an end-labeled 400 bp fragment, 382 bp of which correspond to chromosomal DNA from  
123 the part of the *oriC* region that is upstream from *dnaA*. The fragment was generated by PCR  
124 using primers OCB23 (5'-CCGGAATTCTTTTTTTAGTATCCACAGAGG-3') and OCB24 (5'-  
125 CGCGGATCCCTTTTCTTAGAAAATGGC-3') and *B. subtilis* chromosomal DNA as template.  
126 Allowing for one mismatch from the DnaA binding site consensus sequence (5'-TTATNCACA-  
127 3'), this fragment contains eight consensus DnaA binding sites. The DNA template from  
128 upstream of *yydA* used for gel shift assays was an end-labeled 228 bp fragment generated by  
129 PCR using primers WKS167 (5'-CCCACAGCCTGTGAATTATG-3') and WKS168 (5'-  
130 CGTAGGCCGAAAGTCGTTTG--3'). Allowing for 1 mismatch, this fragment contains four  
131 consensus DnaA binding sites. It is important to note that the sequence requirements for binding  
132 DnaA are not well defined and this estimate of the number of potential binding sites is likely an  
133 underestimate as DnaA is also likely to bind sequences with more than one mismatch from  
134 consensus (17).

135 The PCR products were purified on columns (Qiagen) and end-labeled with gamma- $^{32}\text{P}$ -ATP  
136 using T4 polynucleotide kinase. The labeled DNA fragment was separated from free ATP with a  
137 G50 Column (GE). DnaA was incubated with 2.5 mM ATP for two hours on ice before being  
138 used in gel shift reactions containing 40 mM Hepes pH 7.6, 10 mM KCl, 140 mM potassium  
139 glutamate, 10 mM magnesium acetate, 2.5 mM ATP, 0.5 mM EDTA, 1 mM DTT, 50  $\mu\text{g}/\text{ml}$

140 BSA, 20% glycerol, and 50 pM DNA probe in the presence or absence of DnaD-his6 (300 nM)  
141 for 20 minutes at room temperature.

142 To determine an appropriate concentration of DnaD to use, we measured the effects of  
143 different concentrations of DnaD-his6 on the electrophoretic mobility of the DNA fragment from  
144 the *oriC* region in the presence of 10 nM DnaA-ATP, or with no DnaA (Fig. 1). DnaD-his6 was  
145 used at 25, 50, 100, 200, 300 nM. At these concentrations of DnaD-his6, there was little or no  
146 change in electrophoretic mobility of the DNA fragment in the absence of DnaA. However, in  
147 the presence of DnaA, there was a change in the gel shift beginning at 50 nM DnaD-his6. We  
148 chose to use 300 nM DnaD because there was a large change in the gel shift in the presence of  
149 DnaA, but little or no change in its absence.

150 The binding reactions were run on a 5% polyacrylamide gel with 2.5% glycerol run in 0.5X  
151 TBE at approximately 12 volts/cm for 3 hours. Gels were imaged on a Typhoon scanner (GE  
152 Healthcare) and GraphPad Prism 5 software used to plot binding curves. Data were plotted and  
153 fitted to the Hill equation  $\{y=(m1*x^n)/(Kd^n+x^n)\}$  where y is the % DNA bound at any given  
154 DnaA concentration, x is the DnaA concentration, m1 is maximal binding (100%), Kd is the  
155 apparent dissociation constant (concentration at which 50% of DNA is bound determined from  
156 data) and n is the Hill coefficient. All experiments were done in triplicate. Data presented are  
157 averages of triplicates  $\pm$  standard error.

### 158 **ATPase Assays**

159 ATPase assays were carried out using gamma-<sup>32</sup>P-ATP as substrate and products were  
160 separated by thin layer chromatography (TLC). Reactions (50  $\mu$ l) contained 100 nM DnaA, 50  
161 mM Tris pH 7, 5 mM magnesium acetate, 1 mM DTT, 100 ng/ml BSA, 10% glycerol, and 1  $\mu$ M  
162 ATP (1/1000 gamma-<sup>32</sup>P-ATP) and, where added, 1  $\mu$ M PCR product from *oriC* fragment

163 containing eight DnaA binding sites. Time points were stopped with 2 volumes of stop buffer  
164 (0.5% SDS, 250 mM NaCl, 25 mM EDTA) and spotted on cellulose TLC plates. Products were  
165 separated with 0.5 M LiCl, 1M formic acid. Plates were dried and exposed to a phosphostorage  
166 screen. Free radiolabeled orthophosphate and ATP were measured and percent hydrolysis  
167 calculated. All experiments were done in triplicate and data are presented as the averages  $\pm$   
168 standard error.

### 169 **Nucleotide Binding Assays**

170 Nucleotide binding was measured using alpha-<sup>32</sup>P-ATP. DnaA (100 nM) and ATP (1  $\mu$ M)  
171 were incubated for 30 min at room temperature in 50  $\mu$ l reactions containing 40 mM Hepes pH  
172 7.5, 10 mM Mg Acetate, 0.5 mM EDTA, 1 mM DTT, 150 mM potassium glutamate, 100  $\mu$ g/ml  
173 BSA, and 10% glycerol. The reactions were placed on equilibrated nitrocellulose membranes  
174 (Millipore), washed with buffer (40 mM Hepes pH 7.5, 150 mM KCl, 10 mM magnesium  
175 acetate, 0.5 mM EDTA, 10  $\mu$ g/ml BSA) and radioactivity was measured by filter binding as  
176 described for the nucleotide exchange assay. All experiments were done in triplicate and data are  
177 presented as the averages  $\pm$  standard error. The apparent Kd for ATP binding to DnaA was 29  
178 nM (data not shown) similar to previous reports (17).

179

180

181



## 182 **Results**

### 183 **DnaD does not affect nucleotide exchange for DnaA**

184 We found that the rate of nucleotide exchange for DnaA was relatively rapid and that DnaD  
185 had no detectable effect on this rate. To measure nucleotide exchange, we incubated purified  
186 DnaA with radio-labeled ADP or ATP. The amount of radioactive nucleotide that remained  
187 associated with DnaA was measured at various times after addition of excess unlabeled ATP at  
188 37°C, and the amount of nucleotide that was released was calculated. For both DnaA-ADP (Fig.  
189 2A) and DnaA-ATP (Fig. 2B), the radioactive nucleotide was released with a half-life of 5  
190 minutes. The addition of DnaD-his6 (Materials and methods) had no detectable effect on this  
191 half-life (Fig. 2). Based on these results, we conclude that the rate of nucleotide exchange for *B.*  
192 *subtilis* DnaA is relatively rapid compared to the 45 min half-life of exchange for *E. coli* DnaA  
193 (57) and that DnaA-ATP is regenerated from DnaA-ADP in the absence of any other cellular  
194 factors. The relatively rapid rate of nucleotide exchange is similar to that of DnaA from *S. aureus*  
195 (32).

### 196 **DnaD increases affinity and reduces cooperativity of DnaA-ATP binding to DNA**

197 Given the in vivo association of DnaD to DnaA binding sites around the chromosome (60),  
198 we tested for effects of DnaD on the ability of DnaA to bind DNA using gel electrophoretic  
199 mobility shift assays. DnaA-ATP bound to a DNA fragment from the *oriC* region with an  
200 apparent dissociation constant ( $K_d$ ) of 27 nM (Fig. 3A, C). Binding was highly cooperative and  
201 had a Hill coefficient of 8. These results are consistent with previous findings (40).

202 We measured the effects of DnaD-his6 on the ability of DnaA to bind to DNA. DnaD-his6  
203 alone (300 nM) did not have detectable binding activity under these assay conditions (Fig. 1,  
204 3B), as previously reported (60). However, addition of DnaD-his6 to DNA and DnaA-ATP

205 substantially altered the binding properties of DnaA-ATP to DNA. The apparent dissociation  
206 constant in the presence of DnaD was approximately 7 nM, compared to 27 nM in the absence of  
207 DnaD. There was a concomitant loss of cooperative binding as the Hill coefficient decreased  
208 from 8 in the absence of DnaD to 1 in its presence (Fig. 3, Table 1). This decrease in the Hill  
209 coefficient and decrease in the apparent binding constant is consistent with DnaA-ATP binding  
210 independently to multiple sites in the DNA fragment.

211 The DNA fragment used in these experiments was derived from sequences upstream of *dnaA*  
212 in the *oriC* region. We found that DnaD had a similar effect on the ability of DnaA-ATP to bind  
213 to a DNA fragment from a different chromosomal region. In addition to the *oriC* region, DnaA is  
214 found associated with several chromosomal regions in vivo (20, 24), including the region  
215 between *yydA* and *yydS* (4, 20, 24), two genes of unknown function. This region has also been  
216 implicated in regulating DNA replication by recruiting DnaA away from *oriC* (51). Therefore,  
217 we isolated a DNA fragment from the region upstream of *yydA* that contains four DnaA binding  
218 sites with  $\leq 1$  mismatch to the consensus and tested the ability of DnaD to affect DnaA binding to  
219 this region. DnaA-ATP bound to this fragment with an apparent dissociation constant of 25 (Fig.  
220 3D). Binding was cooperative with a Hill coefficient of 6, indicative of binding to 6 possible  
221 DnaA sites. Addition of DnaD-his6 (300 nM) to these reactions decreased the apparent  
222 dissociation constant to 5 nM and reduced the Hill coefficient to 1, indicating that there was  
223 essentially no cooperative binding in the presence of DnaD. These effects are similar to those on  
224 the binding of DnaA-ATP to DNA fragment from the *oriC* region upstream from *dnaA* (Fig. 3).  
225 Together, these results indicate that DnaD affects binding of DnaA to DNA fragments from the  
226 *oriC* region and at least one origin-distal region. Since DnaD is found at multiple chromosomal

227 regions bound by DnaA in vivo (60), we suspect that DnaD similarly affects DnaA binding at  
228 these regions.

229 We have not been able to detect any changes in the footprint of DnaA on DNA in the  
230 presence compared to the absence of DnaD (unpublished results). This is likely because at the  
231 high concentrations of DnaA needed to observe a footprint (17), the addition of DnaD has no  
232 detectable effect on binding (Fig. 3). At lower concentrations of DnaA, where DnaD does  
233 influence binding, we suspect that there is a population of DNA molecules with different sites  
234 occupied by DnaA, thereby not producing any obvious protection in a footprint experiment, but  
235 still capable of generating a change in electrophoretic mobility.

#### 236 **Characterization of DnaA mutants defective in ATPase activity**

237 Since the rate of ADP exchange for ATP of *B. subtilis* DnaA is relatively rapid and  
238 unaffected by DnaD, and DnaD affects the ability of DnaA-ATP to bind DNA, we hypothesized  
239 that DnaD would affect DnaA mutants that are defective in nucleotide hydrolysis and/or binding.  
240 To test this, we made two different mutations in *dnaA*. One mutation is in the conserved Walker  
241 A motif and changes the lysine at amino acid 157 to alanine, DnaA(K157A), and is predicted to  
242 reduce nucleotide binding (21). We also made a mutation in the conserved Walker B motif that  
243 changes the glutamate at amino acid 215 to alanine, DnaA(D215A), and is predicted to alter  
244 nucleotide hydrolysis (21). We purified the mutant proteins and tested them in vitro.

245 Both DnaA(K157A) and DnaA(D215A) were defective in ATP hydrolysis. We measured the  
246 rate of ATP hydrolysis using gamma-<sup>32</sup>P-ATP and measuring the release of ortho-phosphate  
247 (Materials and Methods). The rate of ATP hydrolysis by wild type DnaA was 1.8 moles of ATP  
248 hydrolyzed per mole of DnaA per hour. Upon addition of DNA, the rate of hydrolysis increased  
249 approximately 6-fold to 12 moles of ATP hydrolyzed per mole of DnaA per hour. These rates of

250 ATP hydrolysis and the effects of DNA are consistent with previously published data for DnaA  
251 from *E. coli* and *S. aureus* (32, 57). In contrast to the wild type protein, DnaA(K157A) and  
252 DnaA(D215A) had rates of ATP hydrolysis of approximately 0.07 and 0.1 moles of ATP per  
253 mole of DnaA per hour, respectively (Table 2).

254 As expected, the DnaA(K157A) mutant was defective and the DnaA(D215A) mutant had  
255 normal ATP binding. At saturating ATP concentrations, we found that wild type protein bound  
256 0.4 molecules of ATP per molecule of DnaA (Table 2), consistent with previous reports for  
257 DnaA from *E. coli* (0.48 and 0.55) (10, 57) but greater than a previous report for *B. subtilis*  
258 DnaA (0.17) (17). The DnaA(D215A) mutant had ATP binding (Table 2) that was  
259 indistinguishable from that of the wild type protein. In contrast, the DnaA(K157A) mutant  
260 appeared to bind 0.02 molecules of ATP per molecule of DnaA (Table 2), consistent with little or  
261 no ATP binding.

#### 262 **DnaD affects DnaA mutants defective in ATPase activity and nucleotide binding**

263 We determined the effects of DnaD on the ability of the mutant DnaA proteins to bind DNA  
264 and compared the binding properties to those of wild type DnaA. The mutant DnaA that binds  
265 ATP but is defective in hydrolysis {DnaA(D215A)} bound DNA with an apparent  $K_d$  of 12 nM,  
266 compared to 27 nM for the wild type protein (Fig. 4A). Binding to DNA was cooperative and  
267 had a Hill coefficient of approximately 5 (Fig. 4A). Addition of DnaD reduced the apparent  $K_d$   
268 to approximately 6 nM and the Hill coefficient to approximately 2 (Fig. 4A, Table 1). The  
269 mutant DnaA that is defective in binding ATP {DnaA(K157A)} has an apparent  $K_d$  of 36 nM  
270 and a Hill coefficient of approximately 5 (Fig. 4B). Addition of DnaD reduced the apparent  $K_d$   
271 to 17 nM and the Hill coefficient to approximately 3 (Fig. 4B, Table 1). Together, these results

272 indicate that the effects of DnaD on the ability of DnaA to bind DNA do not require ATP  
273 binding or hydrolysis.

274 Previously, we found that YabA, a negative regulator of replication initiation, inhibited  
275 cooperativity of DnaA while reducing the apparent K<sub>d</sub> (40). We tested the effects of YabA on  
276 the DnaA(K157A) mutant and found that the addition of YabA (700 nM) reduced the apparent  
277 K<sub>d</sub> from 36 nM to 14.7 nM and the Hill coefficient from 5.6 to 2 (Fig. 4C). These results are  
278 consistent with the previous reports of the effects of YabA on the ability of DnaA to bind DNA  
279 (40) and are comparable to the effects of DnaD, and suggest that they may regulate DnaA by  
280 similar mechanisms.

281

282

## 283 **Discussion**

284 In many bacteria, the conserved replication initiator DnaA is a target for the control of  
285 replication initiation. DnaA is also a transcription factor, and many of the factors that modulate  
286 its activity in replication initiation are likely to affect its activity as a transcription factor.  
287 Mechanisms regulating DnaA have been most studied with *E. coli* and its close relatives.  
288 However, the proteins and mechanisms used by *E. coli* are largely limited to the proteobacteria  
289 and are not found in Gram positive organisms like *B. subtilis*. Likewise, some of the proteins and  
290 mechanisms used by *B. subtilis* are not found in *E. coli* and other proteobacteria. Results  
291 presented here indicate that *B. subtilis* DnaD is a regulator of DnaA. Below, we discuss the  
292 possible role of DnaD in regulation of DnaA and the different properties of DnaA that might lead  
293 to differences in its regulation in different organisms.

294 **Role of DnaD in replication initiation and regulation of DnaA**

295 DnaD is found in *B. subtilis* and other low G+C-content Gram positive bacteria, but not in *E.*  
296 *coli* and other Gram negative bacteria. DnaD is required for replication initiation (5, 6), interacts  
297 with DnaA (23, 60), and is needed to recruit the helicase loading protein DnaB to *oriC* (7, 59).  
298 DnaD is found associated with the *oriC* region of the chromosome (54, 59) and many other  
299 chromosomal regions that also bind DnaA (60). The association of DnaD with these regions is  
300 dependent on DnaA (59, 60).

301 We found that DnaD decreases the apparent K<sub>d</sub> of DnaA-ATP for DNA, and also decreases  
302 the cooperativity of DnaA binding to DNA. These effects could indicate that DnaD functions  
303 either as an activator or repressor of DnaA, or both. For many regulators, the phenotype caused  
304 by a null mutation typically indicates regulation is positive or negative. Unfortunately, *dnaD* is  
305 essential and null mutations are not viable. Temperature sensitive and other conditional loss of  
306 function *dnaD* mutants result in increased association of DnaA with chromosomal regions (4,  
307 20), consistent with DnaD normally functioning to reduce binding of DnaA to DNA. However,  
308 these mutations also lead to a decrease in replication initiation. Analyzing effects of  
309 overexpressing DnaD is also problematic because overexpression of DnaD causes a severe  
310 growth defect that is independent of replication initiation from *oriC* (38). Mutations in other  
311 genes that cause a decrease in replication initiation also cause an increase in DnaA activity (4,  
312 20) making it difficult to discern if the effects of DnaD are direct, due to changes in replication  
313 initiation, or both.

314 It is well established that DnaD is essential and has a positive role in replication initiation.  
315 During replication initiation, DnaA-ATP binds cooperatively to many sites in *oriC* (e.g., 18, 39,  
316 41). By analogy to *E. coli*, it is likely that the temporal order of binding of DnaA-ATP to sites in

317 *oriC* is important for open complex formation (39, 41). In *B. subtilis*, association of DnaD with  
318 the *oriC* region requires DnaA. DnaD is then required for association of DnaB and subsequent  
319 loading of the replicative helicase (6, 7, 54, 59). It is possible that the effects of DnaD on the  
320 binding of DnaA to *oriC* (increase in apparent affinity) could also stimulate replication initiation  
321 by maintaining DnaA bound to *oriC*. However, we think that this is unlikely if ordered and  
322 cooperative binding of DnaA is important for replication initiation.

323 In addition to its known positive role in replication initiation, we postulate that DnaD also  
324 serves to negatively regulate replication initiation through its effects on the ability of DnaA to  
325 bind DNA. The ability of DnaD to inhibit cooperative binding of DnaA-ATP to DNA is similar  
326 to the effect of two other negative regulators of replication initiation, YabA (40) and Soj (55).

327 We postulate that immediately before or after replication initiation, DnaD helps keep DnaA  
328 inactive at *oriC* by inhibiting cooperative binding. This activity of DnaD as a negative regulator  
329 of DnaA could be modulated by changes in the amount of available DnaD during the replication  
330 cycle. For example, the amount of DnaD available to interact with DnaA at *oriC* could change  
331 during a replication cycle, perhaps due to association of DnaD with other proteins or  
332 chromosomal regions (60), or possible changes in its synthesis or stability. We have not yet  
333 tested these possibilities.

334 It is also possible that inhibition of DnaA by DnaD is relieved by the replication initiation  
335 protein DnaB. That is, DnaD might be keeping DnaA inactive at *oriC* until proper assembly of  
336 additional parts of the replication initiation complex. For example, DnaD is needed to recruit  
337 DnaB (part of the helicase loader) to *oriC* (59) and other chromosomal regions bound by DnaA  
338 (60). Association of DnaB might alter interactions between DnaA and DnaD, relieving the  
339 putative inhibitory effect mediated by DnaD, enabling replication initiation. Preliminary

340 attempts to test this in vitro have not been successful, perhaps because of a possible role of the  
341 membrane in interactions between DnaB and DnaD (54).

342

### 343 **Emerging theme in the regulation of DnaA**

344 One of the emerging themes of regulation of *B. subtilis* DnaA and replication initiation is the  
345 role of regulators that directly alter the ability of DnaA to bind DNA. Including DnaD, there are  
346 now at least four regulators of this type. The production and activities of the regulators are  
347 differentially controlled and each regulator is likely to be important at different times during the  
348 growth and replication cycles. The regulators are also likely to be partly redundant.

349 SirA. SirA is a negative regulator of DnaA that is produced during entry into stationary phase  
350 and the initiation of sporulation (52, 66). SirA likely interacts with domain I of DnaA, and it  
351 inhibits the ability of DnaA to bind sequences in *oriC* in vivo (53).

352 YabA. YabA was identified in a yeast two-hybrid screen for interactors with replication  
353 proteins (48). YabA is produced during growth and interacts with both DnaA and DnaN, the  
354 processivity clamp of DNA polymerase (48, 49). Like DnaD, YabA reduces the apparent K<sub>d</sub> and  
355 cooperativity of DnaA binding to DNA in vitro, and these effects are independent of the ATPase  
356 activity of DnaA (40). Also like DnaD, YabA is found associated with chromosomal regions that  
357 are bound by DnaA in vivo and this association is DnaA-dependent (40). In addition, YabA is  
358 found associated with replication forks during ongoing replication (19, 49, 61), and this  
359 association is likely due to interaction between YabA and DnaN. The interaction between DnaN  
360 and YabA likely functions to reduce the ability of YabA to negatively regulate DnaA and  
361 replication initiation (40, 49, 61) and could couple relief of YabA-mediated inhibition to the  
362 release of DnaN from the replisome during replication termination (40).



363 Soj. *B. subtilis* Soj is expressed during growth and is a member of the ParA family of  
364 chromosome partitioning proteins involved in chromosome and plasmid partitioning. Soj is a  
365 negative regulator of replication initiation (34) and DnaA (46, 55, 56). Soj inhibits the ability of  
366 DnaA to form a helix on DNA, independently of the ATPase activity of DnaA (55). The  
367 inhibitory effects of Soj on replication initiation appear to be relieved by Spo0J (56), perhaps  
368 coupling an aspect of replication control to chromosome organization or partitioning (46, 56).

### 369 **Diverse mechanisms controlling DnaA in different organisms**

370 The mechanisms used to control DnaA are diverse. Using *E. coli* and *B. subtilis* as examples,  
371 there are some common mechanisms and some striking differences. In both of these organisms,  
372 and many others, DnaA represses its own transcription (e.g., 1, 3, 20, 50, 67), establishing a  
373 homeostatic regulatory loop. In addition, there are DnaA binding sites outside of *oriC* that  
374 function to titrate DnaA away from *oriC* (51, 60). In *E. coli*, the *datA* locus binds DnaA and  
375 appears to help limit the amount of DnaA available for replication initiation (31, 43, 44).  
376 Similarly, in *B. subtilis*, there are six chromosomal regions outside of *oriC* that have clusters of  
377 DnaA binding sites (20, 24, 51). At least one of these clusters seems to function to help limit the  
378 amount of DnaA available for replication initiation (51).

379 One of the most notable differences between regulation of *E. coli* and *B. subtilis* DnaA is the  
380 stimulation of nucleotide binding and hydrolysis in *E. coli*. One of the primary mechanisms used  
381 by *E. coli* to inhibit the activity of DnaA is called RIDA (Regulatory Inactivation of DnaA) and  
382 uses a protein called Hda (26, 28). Hda interacts with *E. coli* DnaN ( $\beta$ -clamp) and stimulates  
383 nucleotide hydrolysis by DnaA, thereby stimulating conversion of the replication-competent  
384 DnaA-ATP to the inactive DnaA-ADP (64). *E. coli* also has specific DnaA-reactivating  
385 sequences that directly promote nucleotide exchange to generate DnaA-ATP from DnaA-ADP

386 (16). The stimulated rate of nucleotide exchange for *E. coli* DnaA (15) is about the same as the  
387 basal rate for *B. subtilis* DnaA. *E. coli* also has a protein called DiaA that stimulates replication  
388 initiation by stimulating binding by DnaA-ATP (22, 30).

389 In contrast to the mechanisms used by *E. coli* to regulate DnaA and replication initiation, *B.*  
390 *subtilis* is not known to regulate nucleotide hydrolysis or exchange. Rather, the primary  
391 mechanisms for controlling *B. subtilis* DnaA affect its binding to DNA (40, 53, 55), probably by  
392 inhibiting formation of multimeric DnaA structures (helix formation) on the DNA and  
393 preventing cooperative binding to sites in *oriC* (40, 55). No known regulator of DnaA in *B.*  
394 *subtilis* affects nucleotide hydrolysis or exchange.

395 Clearly, different organisms use different mechanisms to control the activity of DnaA and  
396 replication initiation. We suggest that the multiple mechanisms may have evolved in different  
397 organisms, in part, due to the different rates of nucleotide exchange. For organisms like *E. coli*  
398 where DnaA has a relatively slow rate of nucleotide exchange, stimulation of nucleotide  
399 hydrolysis and exchange is likely to be a predominant mode of regulation. In contrast, for  
400 organisms like *B. subtilis* and *S. aureus* where DnaA has a relatively rapid rate of nucleotide  
401 exchange, the predominant modes of regulation of DnaA affect DNA binding and cooperativity.

402

403

#### 404 **Acknowledgments**

405 We thank T. Baker, A. Olivares, and members of the Grossman lab for useful discussions,  
406 and C. Lee, J.L. Smith, C. Seid, M. Laub, and L. Simmons for comments on the manuscript.

407 This work was supported, in part, by Public Health Service grant GM41934 to A.D.G.

408

409 **References**

- 410 1. **Atlung, T., E. S. Clausen, and F. G. Hansen.** 1985. Autoregulation of the *dnaA* gene of  
411 *Escherichia coli* K12. *Mol Gen Genet* **200**:442-450.
- 412 2. **Bramhill, D., and A. Kornberg.** 1988. Duplex opening by *dnaA* protein at novel  
413 sequences in initiation of replication at the origin of the *E. coli* chromosome. *Cell*  
414 **52**:743-755.
- 415 3. **Braun, R. E., K. O'day, and A. Wright.** 1985. Autoregulation of the DNA replication  
416 gene *dnaA* in *E. coli* K-12. *Cell* **40**:159-169.
- 417 4. **Breier, A. M., and A. D. Grossman.** 2009. Dynamic association of the replication  
418 initiator and transcription factor DnaA with the *Bacillus subtilis* chromosome during  
419 replication stress. *J Bacteriol* **191**:486-493.
- 420 5. **Bruand, C., S. D. Ehrlich, and L. Janniere.** 1995. Primosome assembly site in *Bacillus*  
421 *subtilis*. *EMBO J* **14**:2642-2650.
- 422 6. **Bruand, C., M. Farache, S. Mcgovern, S. D. Ehrlich, and P. Polard.** 2001. DnaB,  
423 DnaD and DnaI proteins are components of the *Bacillus subtilis* replication restart  
424 primosome. *Mol Microbiol* **42**:245-255.
- 425 7. **Bruand, C., M. Velten, S. Mcgovern, S. Marsin, C. Serena, S. D. Ehrlich, and P.**  
426 **Polard.** 2005. Functional interplay between the *Bacillus subtilis* DnaD and DnaB  
427 proteins essential for initiation and re-initiation of DNA replication. *Mol Microbiol*  
428 **55**:1138-1150.
- 429 8. **Bruck, I., and M. O'donnell.** 2000. The DNA replication machine of a gram-positive  
430 organism. *J Biol Chem* **275**:28971-28983.
- 431 9. **Burkholder, W. F., I. Kurtser, and A. D. Grossman.** 2001. Replication initiation  
432 proteins regulate a developmental checkpoint in *Bacillus subtilis*. *Cell* **104**:269-279.
- 433 10. **Carr, K. M., and J. M. Kaguni.** 1996. The A184V missense mutation of the *dnaA5* and  
434 *dnaA46* alleles confers a defect in ATP binding and thermolability in initiation of  
435 *Escherichia coli* DNA replication. *Mol Microbiol* **20**:1307-1318.
- 436 11. **Crooke, E., C. E. Castuma, and A. Kornberg.** 1992. The chromosome origin of  
437 *Escherichia coli* stabilizes DnaA protein during rejuvenation by phospholipids. *J Biol*  
438 *Chem* **267**:16779-16782.
- 439 12. **Davey, M. J., and M. O'donnell.** 2003. Replicative helicase loaders: ring breakers and  
440 ring makers. *Curr Biol* **13**:R594-596.
- 441 13. **Duderstadt, K. E., and J. M. Berger.** 2008. AAA+ ATPases in the initiation of DNA  
442 replication. *Crit Rev Biochem Mol Biol* **43**:163-187.
- 443 14. **Felczak, M. M., and J. M. Kaguni.** 2004. The box VII motif of *Escherichia coli* DnaA  
444 protein is required for DnaA oligomerization at the *E. coli* replication origin. *J Biol Chem*  
445 **279**:51156-51162.
- 446 15. **Fujimitsu, K., and T. Katayama.** 2004. Reactivation of DnaA by DNA sequence-  
447 specific nucleotide exchange in vitro. *Biochem Biophys Res Commun* **322**:411-419.
- 448 16. **Fujimitsu, K., T. Senriuchi, and T. Katayama.** 2009. Specific genomic sequences of *E.*  
449 *coli* promote replicational initiation by directly reactivating ADP-DnaA. *Genes Dev*  
450 **23**:1221-1233.
- 451 17. **Fukuoka, T., S. Moriya, H. Yoshikawa, and N. Ogasawara.** 1990. Purification and  
452 characterization of an initiation protein for chromosomal replication, DnaA, in *Bacillus*  
453 *subtilis*. *J Biochem (Tokyo)* **107**:732-739.

- 454 18. **Fuller, R. S., B. E. Funnell, and A. Kornberg.** 1984. The dnaA protein complex with  
455 the *E. coli* chromosomal replication origin (*oriC*) and other DNA sites. *Cell* **38**:889-900.
- 456 19. **Goranov, A. I., A. M. Breier, H. Merrikh, and A. D. Grossman.** 2009. YabA of  
457 *Bacillus subtilis* controls DnaA-mediated replication initiation but not the transcriptional  
458 response to replication stress. *Mol Microbiol* **74**:454-466.
- 459 20. **Goranov, A. I., L. Katz, A. M. Breier, C. B. Burge, and A. D. Grossman.** 2005. A  
460 transcriptional response to replication status mediated by the conserved bacterial  
461 replication protein DnaA. *Proc Natl Acad Sci U S A* **102**:12932-12937.
- 462 21. **Hanson, P. I., and S. W. Whiteheart.** 2005. AAA+ proteins: have engine, will work.  
463 *Nat Rev Mol Cell Biol* **6**:519-529.
- 464 22. **Ishida, T., N. Akimitsu, T. Kashioka, M. Hatano, T. Kubota, Y. Ogata, K. Sekimizu,  
465 and T. Katayama.** 2004. DiaA, a novel DnaA-binding protein, ensures the timely  
466 initiation of *Escherichia coli* chromosome replication. *J Biol Chem* **279**:45546-45555.
- 467 23. **Ishigo-Oka, D., N. Ogasawara, and S. Moriya.** 2001. DnaD protein of *Bacillus subtilis*  
468 interacts with DnaA, the initiator protein of replication. *J Bacteriol* **183**:2148-2150.
- 469 24. **Ishikawa, S., Y. Ogura, M. Yoshimura, H. Okumura, E. Cho, Y. Kawai, K.  
470 Kurokawa, T. Oshima, and N. Ogasawara.** 2007. Distribution of stable DnaA-binding  
471 sites on the *Bacillus Subtilis* genome detected using a modified ChIP-chip method. *DNA*  
472 *Res* **14**:155-168.
- 473 25. **Kaguni, J. M.** 2006. DnaA: controlling the initiation of bacterial DNA replication and  
474 more. *Annu Rev Microbiol* **60**:351-375.
- 475 26. **Katayama, T., T. Kubota, K. Kurokawa, E. Crooke, and K. Sekimizu.** 1998. The  
476 initiator function of DnaA protein is negatively regulated by the sliding clamp of the *E.*  
477 *coli* chromosomal replicase. *Cell* **94**:61-71.
- 478 27. **Katayama, T., S. Ozaki, K. Keyamura, and K. Fujimitsu.** 2010. Regulation of the  
479 replication cycle: conserved and diverse regulatory systems for DnaA and *oriC*. *Nat Rev*  
480 *Microbiol* **8**:163-170.
- 481 28. **Kato, J., and T. Katayama.** 2001. Hda, a novel DnaA-related protein, regulates the  
482 replication cycle in *Escherichia coli*. *EMBO J* **20**:4253-4262.
- 483 29. **Kawakami, H., and T. Katayama.** 2010. DnaA, ORC, and Cdc6: similarity beyond the  
484 domains of life and diversity. *Biochem Cell Biol* **88**:49-62.
- 485 30. **Keyamura, K., N. Fujikawa, T. Ishida, S. Ozaki, M. Su'etsugu, K. Fujimitsu, W.  
486 Kagawa, S. Yokoyama, H. Kurumizaka, and T. Katayama.** 2007. The interaction of  
487 DiaA and DnaA regulates the replication cycle in *E. coli* by directly promoting ATP  
488 DnaA-specific initiation complexes. *Genes Dev* **21**:2083-2099.
- 489 31. **Kitagawa, R., T. Ozaki, S. Moriya, and T. Ogawa.** 1998. Negative control of  
490 replication initiation by a novel chromosomal locus exhibiting exceptional affinity for  
491 *Escherichia coli* DnaA protein. *Genes Dev* **12**:3032-3043.
- 492 32. **Kurokawa, K., H. Mizumura, T. Takaki, Y. Ishii, N. Ichihashi, B. L. Lee, and K.  
493 Sekimizu.** 2009. Rapid exchange of bound ADP on the *Staphylococcus aureus*  
494 replication initiation protein DnaA. *J Biol Chem* **284**:34201-34210.
- 495 33. **Kurokawa, K., S. Nishida, A. Emoto, K. Sekimizu, and T. Katayama.** 1999.  
496 Replication cycle-coordinated change of the adenine nucleotide-bound forms of DnaA  
497 protein in *Escherichia coli*. *EMBO J* **18**:6642-6652.
- 498 34. **Lee, P. S., and A. D. Grossman.** 2006. The chromosome partitioning proteins Soj  
499 (ParA) and Spo0J (ParB) contribute to accurate chromosome partitioning, separation of

- 500 replicated sister origins, and regulation of replication initiation in *Bacillus subtilis*. Mol  
 501 Microbiol **60**:853-869.
- 502 35. **Leonard, A. C., and J. E. Grimwade.** 2010. Regulating DnaA complex assembly: it is  
 503 time to fill the gaps. Curr Opin Microbiol **13**:766-772.
- 504 36. **Li, Y., K. Kurokawa, M. Matsuo, N. Fukuhara, K. Murakami, and K. Sekimizu.**  
 505 2004. Identification of temperature-sensitive *dnaD* mutants of *Staphylococcus aureus* that  
 506 are defective in chromosomal DNA replication. Mol Genet Genomics **271**:447-457.
- 507 37. **Marsin, S., S. Mcgovern, S. D. Ehrlich, C. Bruand, and P. Polard.** 2001. Early steps  
 508 of *Bacillus subtilis* primosome assembly. J Biol Chem **276**:45818-45825.
- 509 38. **Marston, F. Y., W. H. Grainger, W. K. Smits, N. H. Hopcroft, M. Green, A. M.**  
 510 **Hounslow, A. D. Grossman, C. J. Craven, and P. Sultanas.** 2010. When simple  
 511 sequence comparison fails: the cryptic case of the shared domains of the bacterial  
 512 replication initiation proteins DnaB and DnaD. Nucleic Acids Res.
- 513 39. **Megarry, K. C., V. T. Ryan, J. E. Grimwade, and A. C. Leonard.** 2004. Two  
 514 discriminatory binding sites in the *Escherichia coli* replication origin are required for  
 515 DNA strand opening by initiator DnaA-ATP. Proc Natl Acad Sci U S A **101**:2811-2816.
- 516 40. **Merrikh, H., and A. D. Grossman.** 2011. Control of the replication initiator DnaA by  
 517 an anti-cooperativity factor. Mol Microbiol **82**:434-446.
- 518 41. **Miller, D. T., J. E. Grimwade, T. Betteridge, T. Rozgaja, J. J. Torgue, and A. C.**  
 519 **Leonard.** 2009. Bacterial origin recognition complexes direct assembly of higher-order  
 520 DnaA oligomeric structures. Proc Natl Acad Sci U S A **106**:18479-18484.
- 521 42. **Mizushima, T., T. Takaki, T. Kubota, T. Tsuchiya, T. Miki, T. Katayama, and K.**  
 522 **Sekimizu.** 1998. Site-directed mutational analysis for the ATP binding of DnaA protein.  
 523 Functions of two conserved amino acids (Lys-178 and Asp-235) located in the ATP-  
 524 binding domain of DnaA protein in vitro and in vivo. J Biol Chem **273**:20847-20851.
- 525 43. **Morigen, A. Lobner-Olesen, and K. Skarstad.** 2003. Titration of the *Escherichia coli*  
 526 DnaA protein to excess *datA* sites causes destabilization of replication forks, delayed  
 527 replication initiation and delayed cell division. Mol Microbiol **50**:349-362.
- 528 44. **Morigen, F. Molina, and K. Skarstad.** 2005. Deletion of the *datA* site does not affect  
 529 once-per-cell-cycle timing but induces rifampin-resistant replication. J Bacteriol  
 530 **187**:3913-3920.
- 531 45. **Mott, M. L., and J. M. Berger.** 2007. DNA replication initiation: mechanisms and  
 532 regulation in bacteria. Nat Rev Microbiol **5**:343-354.
- 533 46. **Murray, H., and J. Errington.** 2008. Dynamic control of the DNA replication initiation  
 534 protein DnaA by Soj/ParA. Cell **135**:74-84.
- 535 47. **Nishida, S., K. Fujimitsu, K. Sekimizu, T. Ohmura, T. Ueda, and T. Katayama.**  
 536 2002. A nucleotide switch in the *Escherichia coli* DnaA protein initiates chromosomal  
 537 replication: evidence from a mutant DnaA protein defective in regulatory ATP hydrolysis  
 538 in vitro and in vivo. J Biol Chem **277**:14986-14995.
- 539 48. **Noirot-Gros, M. F., E. Dervyn, L. J. Wu, P. Mervelet, J. Errington, S. D. Ehrlich,**  
 540 **and P. Noirot.** 2002. An expanded view of bacterial DNA replication. Proc Natl Acad  
 541 Sci U S A **99**:8342-8347.
- 542 49. **Noirot-Gros, M. F., M. Velten, M. Yoshimura, S. Mcgovern, T. Morimoto, S. D.**  
 543 **Ehrlich, N. Ogasawara, P. Polard, and P. Noirot.** 2006. Functional dissection of  
 544 YabA, a negative regulator of DNA replication initiation in *Bacillus subtilis*. Proc Natl  
 545 Acad Sci U S A **103**:2368-2373.

- 546 50. **Ogura, Y., Y. Imai, N. Ogasawara, and S. Moriya.** 2001. Autoregulation of the *dnaA-*  
547 *dnaN* operon and effects of DnaA protein levels on replication initiation in *Bacillus*  
548 *subtilis*. *J Bacteriol* **183**:3833-3841.
- 549 51. **Okumura, H., M. Yoshimura, M. Ueki, T. Oshima, N. Ogasawara, and S. Ishikawa.**  
550 2012. Regulation of chromosomal replication initiation by oriC-proximal DnaA-box  
551 clusters in *Bacillus subtilis*. *Nucleic Acids Res* **40**:220-234.
- 552 52. **Rahn-Lee, L., B. Gorbatyuk, O. Skovgaard, and R. Losick.** 2009. The conserved  
553 sporulation protein YneE inhibits DNA replication in *Bacillus subtilis*. *J Bacteriol*  
554 **191**:3736-3739.
- 555 53. **Rahn-Lee, L., H. Merrikh, A. D. Grossman, and R. Losick.** 2011. The sporulation  
556 protein SirA inhibits the binding of DnaA to the origin of replication by contacting a  
557 patch of clustered amino acids. *J Bacteriol* **193**:1302-1307.
- 558 54. **Rokop, M. E., J. M. Auchtung, and A. D. Grossman.** 2004. Control of DNA  
559 replication initiation by recruitment of an essential initiation protein to the membrane of  
560 *Bacillus subtilis*. *Mol Microbiol* **52**:1757-1767.
- 561 55. **Scholefield, G., J. Errington, and H. Murray.** 2012. Soj/ParA stalls DNA replication  
562 by inhibiting helix formation of the initiator protein DnaA. *Embo J*.
- 563 56. **Scholefield, G., R. Whiting, J. Errington, and H. Murray.** 2011. Spo0J regulates the  
564 oligomeric state of Soj to trigger its switch from an activator to an inhibitor of DNA  
565 replication initiation. *Mol Microbiol* **79**:1089-1100.
- 566 57. **Sekimizu, K., D. Bramhill, and A. Kornberg.** 1987. ATP activates dnaA protein in  
567 initiating replication of plasmids bearing the origin of the *E. coli* chromosome. *Cell*  
568 **50**:259-265.
- 569 58. **Simmons, L. A., M. Felczak, and J. M. Kaguni.** 2003. DnaA Protein of *Escherichia*  
570 *coli*: oligomerization at the *E. coli* chromosomal origin is required for initiation and  
571 involves specific N-terminal amino acids. *Mol Microbiol* **49**:849-858.
- 572 59. **Smits, W. K., A. I. Goranov, and A. D. Grossman.** 2010. Ordered association of  
573 helicase loader proteins with the *Bacillus subtilis* origin of replication in vivo. *Mol*  
574 *Microbiol* **75**:452-461.
- 575 60. **Smits, W. K., H. Merrikh, C. Y. Bonilla, and A. D. Grossman.** 2011. Primosomal  
576 proteins DnaD and DnaB are recruited to chromosomal regions bound by DnaA in  
577 *Bacillus subtilis*. *J Bacteriol* **193**:640-648.
- 578 61. **Soufo, C. D., H. J. Soufo, M. F. Noirot-Gros, A. Steindorf, P. Noirot, and P. L.**  
579 **Graumann.** 2008. Cell-cycle-dependent spatial sequestration of the DnaA replication  
580 initiator protein in *Bacillus subtilis*. *Dev Cell* **15**:935-941.
- 581 62. **Soultanas, P.** 2002. A functional interaction between the putative primosomal protein  
582 DnaI and the main replicative DNA helicase DnaB in *Bacillus*. *Nucleic Acids Res*  
583 **30**:966-974.
- 584 63. **Speck, C., C. Weigel, and W. Messer.** 1999. ATP- and ADP-dnaA protein, a molecular  
585 switch in gene regulation. *EMBO J* **18**:6169-6176.
- 586 64. **Su'etsugu, M., T. R. Shimuta, T. Ishida, H. Kawakami, and T. Katayama.** 2005.  
587 Protein associations in DnaA-ATP hydrolysis mediated by the Hda-replicase clamp  
588 complex. *J Biol Chem* **280**:6528-6536.
- 589 65. **Velten, M., S. Mcgovern, S. Marsin, S. D. Ehrlich, P. Noirot, and P. Polard.** 2003. A  
590 two-protein strategy for the functional loading of a cellular replicative DNA helicase.  
591 *Mol Cell* **11**:1009-1020.

- 592 66. **Wagner, J. K., K. A. Marquis, and D. Z. Rudner.** 2009. SirA enforces diploidy by  
593 inhibiting the replication initiator DnaA during spore formation in *Bacillus subtilis*. Mol  
594 Microbiol **73**:963-974.
- 595 67. **Wang, Q. P., and J. M. Kaguni.** 1987. Transcriptional repression of the *dnaA* gene of  
596 *Escherichia coli* by dnaA protein. Mol Gen Genet **209**:518-525.
- 597 68. **Zhang, W., S. Allen, C. J. Roberts, and P. Sultanas.** 2006. The *Bacillus subtilis*  
598 primosomal protein DnaD untwists supercoiled DNA. J Bacteriol **188**:5487-5493.  
599  
600

601

602 **Table 1. Summary of DNA binding by wild type and mutant DnaA<sup>1</sup>.**

603

DnaD <sup>2</sup>	DnaA		DnaA(K157A)		DnaA(D215A)	
	-	+	-	+	-	+
apparent Kd	27 ± 0.4	6.6 ± 1	36 ± 0.8	17 ± 1	12 ± 0.3	5.7 ± 0.3
Hill coefficient	8.6 ± 0.9	1.0 ± 0.1	5.3 ± 0.5	2.7 ± 0.4	5.2 ± 0.4	2.2 ± 0.2

604

605

606 <sup>1</sup> Results presented in Fig. 3C, 4A, and 4B are summarized. The DNA template was a fragment607 containing sequences upstream from *oriC* with eight DnaA binding sites (≤1 mismatch from

608 consensus).

609 <sup>2</sup> DnaD-his6 was either added at 300 nM (+) or absent (-).

610

611



612  
613  
614

**Table 2. Summary of ATP hydrolysis and binding by wild type and mutant DnaA<sup>1</sup>.**

protein	ATPase	ATPase + DNA	ATP binding
DnaA	$1.8 \pm 0.25$	$11.9 \pm 2.2$	$0.4 \pm 0.05$
DnaA(K157A)	$0.07 \pm 0.002$	$0.3 \pm 0.04$	$0.02 \pm 0.01$
DnaA(D215A)	$0.1 \pm 0.2$	$0.3 \pm 0.01$	$0.4 \pm 0.11$

615  
616  
617  
618  
619  
620  
621  
622

<sup>1</sup> The rate of ATP hydrolysis (ATPase) is presented as the number of moles of ATP hydrolyzed per mole of DnaA per hour. Where indicated, the 400 bp DNA fragment from the *oriC* region that was used for the gel shift assays was added (1  $\mu$ M). The amount of ATP bound (ATP binding) is presented as moles of ATP per mole of DnaA.

623 **Figure legends**

624

625 **Figure 1. Effect of different concentrations of DnaD on DnaA binding to *oriC*.**

626 Representative gel of the radiolabeled DNA probe from the *oriC* region with different amount of  
627 DnaD-his, in the absence of DnaA (six lanes on the left) or presence of 10 nM DnaA-ATP (six  
628 lanes on the right). Concentrations of DnaD-his are 0 (-), 25, 50, 100, 200, or 300 nM and are  
629 indicated below each lane.

630

631

632 **Figure 2. DnaD does not affect nucleotide exchange by DnaA.** The amount of  $^{14}\text{C}$ -ADP  
633 (A) or  $^{32}\text{P}$ -ATP (B) bound to DnaA (300 nM) at various times after addition of unlabeled ATP (2  
634 mM) at 37°C was measured by filter binding in the absence (open circles) and presence (filled  
635 squares) of DnaD-his6 (600 nM). Data are averages of triplicates  $\pm$  standard error and are  
636 normalized to the starting amount of radioactivity in the absence of unlabeled ATP.

637

638

639 **Figure 3. DnaD inhibits cooperative binding of DnaA to DNA.** Representative gels and  
640 binding curves measuring binding of DnaA-ATP to DNA (50 pM) with and without purified  
641 DnaD-his6 (300 nM) are shown. DnaA concentrations used were: 0, 1, 2, 5, 10, 20, 30, 40, 50,  
642 60, 80, 100, and 200 nM.

643 **A, B.** Representative gels with increasing concentrations of DnaA-ATP incubated with  
644 template DNA from the *oriC* region in the absence (A) or presence (B) of DnaD-his6. Probe with  
645 no added protein is shown in the first lane (A) or first lane (B). Probe with DnaD-his6 and no  
646 DnaA is shown in the second lane of panel B.

647 **C, D.** Data from three independent gel shift assays using template DNA from the *oriC* region  
648 (C) or the *yydA* region (D) are plotted as percent DNA bound vs. the concentration of DnaA-  
649 ATP, in the absence (open circles) and presence (filled squares) of DnaD-his6.

650 In experiments with the DNA fragment from the *oriC* region (C), the calculated Hill  
651 coefficient for DnaA-ATP was 8.6 in the absence of DnaD-his6 and 1 in the presence of DnaD-  
652 his6. The apparent K<sub>d</sub> for DnaA-ATP was 27 nM in the absence and 6.6 nM in the presence of  
653 DnaD-his6. In experiments with the DNA fragment from the *yydA* region (D), the calculated Hill  
654 coefficient for DnaA-ATP was 6 in the absence of DnaD-his6 and 1 in the presence of DnaD-  
655 his6. The apparent K<sub>d</sub> for DnaA-ATP was 25 nM in the absence and 5 nM in the presence of  
656 DnaD-his6.

657

658

659

660 **Figure 4. Effects of DnaD and YabA on DnaA binding to DNA are independent of**  
661 **ATPase activity.** Binding curves of DnaA mutants defective in ATP hydrolysis, DnaA(D215A)-  
662 ATP (A) and ATP binding, DnaA(K157A) (B, C) within the DNA fragment from the *oriC*  
663 region. DnaA concentrations tested were: 0, 1, 2, 5, 10, 20, 30, 40, 50, 60, 80, 100, and 200 nM.  
664 **A, B.** Binding in the absence (open circles) and presence (filled squares) of DnaD-his6 (300  
665 nM).  
666 **C.** Binding in the absence (open circles) and presence (filled diamonds) of his6-YabA (700  
667 nM). For the DnaA mutant defective in ATP binding, DnaA(K157A), the presence of YabA  
668 reduced the Hill coefficient from 5.6 to 2 and the apparent Kd from 36 nM to 14.7 nM.  
669











