

Professor Anthony J Avery

Professor of Primary Care, Division of Primary Care

School of Medicine, University of Nottingham, Nottingham.

Email: tony.avery@nottingham.ac.uk

Will pharmacists working in general practice help tackle the current workload crisis?

It is widely acknowledged that there is a workload crisis in UK general practice. In addition to the impact of population growth and an increase in the number of older people with complex health needs, GPs have also taken on a considerable amount of work previously undertaken in secondary care. The result has been a 16% increase in overall GP workload, over a seven-year period up to 2014, according to a recent large-scale English study.¹ So far, there is little sign that the promised increase in resource heralded in the five-year forward review² is making a difference on the front line.

One approach to tackling the workload crisis is to draw upon the skills of other healthcare professionals, with pharmacists being one example; particularly those that are able to prescribe. In this context, the investment from NHS England in expanding the number of pharmacists working in general practices is welcome. In addition to the pilot wave investment of £31 million for the first recruitment wave of 490,³ there is a commitment to invest an additional £100 million to part-fund 1500 pharmacists to work in English general practices over the next 3-4 years.⁴

There is a reasonable body of evidence confirming the benefits that pharmacists can bring by working in general practices. For example, pharmacists working in primary care can help with chronic disease management, and there is evidence that they can be effective for a wide variety of conditions^{5,6,7}. They can also help more broadly with medicines optimisation⁵ and can work with GPs to reduce hazardous prescribing.⁸ From my experience of working with a practice-based pharmacist there are many other ways in which they can ease the burden on GPs, including taking on difficult prescribing challenges (titrating up or withdrawing medicines), undertaking complex medication reviews, taking on medicines reconciliation for patients discharged from hospital, and negotiating with patients about medicines changes for increased effectiveness and cost-control.

Also, as highlighted in this edition of the BJGP, pharmacists are enthusiastic about their new roles in general practice.⁹ In a qualitative study in the South West of England, pharmacists were interviewed before and after a training programme aimed at equipping them for working in general practice. While recognising their ongoing training needs, and the challenges and uncertainties around their new roles, the pharmacists were willing to take some of the load off GPs.

So, what is not to like about this expansion of pharmacists in English general practice? The main problem, is the funding model. The NHS England contribution to the costs of recruitment and employment is tapered from 60% in year one to 20% in year three.⁴ This means that general practices that have taken on a pharmacist are now having to give serious consideration to the opportunity costs of continuing to employ them in the longer term. When faced with 80% of the annual salary of a pharmacist, GPs in a fully staffed general practice (or federation of practices) will be asking whether any benefits of the pharmacist on their workload are sufficient for them to be prepared to reduce their drawings. In practices with staff shortages they will be asking whether they might be better off investing in more GP sessions or a full-time nurse. The problem is that however

good pharmacists are, it is unlikely that they will reduce GP workload in the way that an additional GP or nurse would. Also, as far as I am aware, there is little or no evidence available on the impact of practice-based pharmacists on GP workload to enable GPs to make a fully informed decision on this issue.

Therefore, I worry that unless the funding formula is changed we may find that this admirable scheme to expand the pharmacy workforce in general practice may run into the sand. I suspect that many general practices will be prepared to make a financial contribution to employing pharmacists, particularly where they genuinely reduce GP workload. However, previous studies suggest that the majority of the impact of practice-based pharmacists will be on quality and safety,⁵ rather than on GP workload, and it is reasonable to ask whether GPs or the government should be paying for this? If NHS England is serious about putting more resources into general practice and easing the burden on GPs a more generous approach is needed to the ongoing funding of practice-based pharmacists.

References

1. Hobbs FDR, Bankhead C, Mukhtar T, et al, on behalf of the National Institute for Health Research School for Primary Care Research. Clinical workload in UK primary care: a retrospective analysis of 100 million consultations in England, 2007–14. *Lancet* 2016; published online April 5. [http://dx.doi.org/10.1016/S0140-6736\(16\)00620-6](http://dx.doi.org/10.1016/S0140-6736(16)00620-6).
2. NHS England. General Practice Forward View, April 2016. Available at: <https://www.england.nhs.uk/wp-content/uploads/2016/04/gp-fv.pdf> [accessed 2 July 2017]
3. More than 400 pharmacists to be recruited to GP surgeries by next year. Available at: <https://www.england.nhs.uk/2015/11/pharmacists-recruited/> [accessed 16 July 2017]
4. NHS England. Clinical Pharmacists in General Practice. Available at: <https://www.england.nhs.uk/gp/gp-fv/workforce/building-the-general-practice-workforce/cp-gp/> [accessed 2 July 2017]
5. Nkansah N, Mostovetsky O, Yu C, Chheng T, Beney J, Bond CM, Bero L. Effect of outpatient pharmacists' non-dispensing roles on patient outcomes and prescribing patterns. *Cochrane Database of Systematic Reviews* 2010, Issue 7. Art. No.: CD000336. DOI: 10.1002/14651858.CD000336.pub2.
6. Santschi V, Chiolero A, Burnand B, Colosimo AL, Paradis G. Impact of pharmacist care in the management of cardiovascular disease risk factors: a systematic review and meta-analysis of randomized trials. *Arch Intern Med*. 2011 Sep 12;171(16):1441-53. doi: 10.1001/archinternmed.2011.399.
7. Weeks G, George J, Maclure K, Stewart D. Non-medical prescribing versus medical prescribing for acute and chronic disease management in primary and secondary care. *Cochrane Database of Systematic Reviews* 2016, Issue 11. Art. No.: CD011227. DOI: 10.1002/14651858.CD011227.pub2
8. Avery AJ, Rodgers S, Cantrill JA, et al. A pharmacist-led information technology intervention for medication errors (PINCER): a multicentre, cluster randomised, controlled trial and cost-effectiveness analysis. *The Lancet* 2012; 379(9823): 1310-1319. doi:10.1016/S0140-6736(11)61817-5.
9. Butterworth J, Sansom A, Sims L, Healey M, Kingsland E, Campbell J. Pharmacist perceptions of their emerging general practice roles in UK primary care: a qualitative interview study. *Br J Gen*

Pract 2017; DOI: <https://doi.org/10.3399/bjgpxxxxxx>. [Note to editorial team: please update this reference]