



Hdud, Ismail M. and Loughna, Paul (2014) Influence of $1\alpha, 25$ -dihydroxyvitamin D₃ [$1, 25(\text{OH})_2\text{D}_3$] on the expression of Sox 9 and the transient receptor potential vanilloid 5/6 ion channels in equine articular chondrocytes. *Journal of Animal Science and Technology*, 56 . 33/1-33/8. ISSN 2055-0391

Access from the University of Nottingham repository:

<http://eprints.nottingham.ac.uk/46623/1/SOX%20s40781-014-0033-1.pdf>

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the Creative Commons Attribution licence and may be reused according to the conditions of the licence. For more details see: <http://creativecommons.org/licenses/by/2.5/>

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

RESEARCH

Open Access

Influence of 1 α , 25-dihydroxyvitamin D₃ [1, 25(OH)₂D₃] on the expression of Sox 9 and the transient receptor potential vanilloid 5/6 ion channels in equine articular chondrocytes

Ismail M Hdud^{1,3} and Paul T Loughna^{1,2*}

Abstract

Background: Sox 9 is a major marker of chondrocyte differentiation. When chondrocytes are cultured *in vitro* they progressively de-differentiate and this is associated with a decline in Sox 9 expression. The active form of vitamin D, 1, 25 (OH)₂D₃ has been shown to be protective of cartilage in both humans and animals. In this study equine articular chondrocytes were grown in culture and the effects of 1, 25 (OH)₂D₃ upon Sox 9 expression examined. The expression of the transient receptor potential vanilloid (TRPV) ion channels 5 and 6 in equine chondrocytes *in vitro*, we have previously shown, is inversely correlated with de-differentiation. The expression of these channels in response to 1, 25 (OH)₂D₃ administration was therefore also examined.

Results: The active form of vitamin D (1, 25 (OH)₂D₃) when administered to cultured equine chondrocytes at two different concentrations significantly increased the expression of Sox 9 at both. In contrast 1, 25 (OH)₂D₃ had no significant effect upon the expression of either TRPV 5 or 6 at either the protein or the mRNA level.

Conclusions: The increased expression of Sox 9, in equine articular chondrocytes *in vitro*, in response to the active form of vitamin D suggests that this compound could be utilized to inhibit the progressive de-differentiation that is normally observed in these cells. It is also supportive of previous studies indicating that 1 α , 25-dihydroxyvitamin D₃ can have a protective effect upon cartilage in animals *in vivo*. The previously observed correlation between the degree of differentiation and the expression levels of TRPV 5/6 had suggested that these ion channels may have a direct involvement in, or be modulated by, the differentiation process *in vitro*. The data in the present study do not support this.

Keywords: Cartilage, Vitamin D, Sox 9, TRPV, Chondrocyte

Background

The active metabolite of vitamin D, 1, 25 (OH)₂D₃ has been reported to be involved in the regulation of cellular functions such as differentiation, proliferation, and the immune system [1]. Similar to other steroids, 1, 25 (OH)₂D₃ exerts physiological actions at two levels by genomic and non-genomic mechanisms. The genomic mechanism

exerts its effect on the cells by binding to specific receptors known as vitamin D receptors, which in turn bind to retinoid X receptors to form heterodimeric complexes [2]. These heterodimeric complexes in turn bind to VDREs (vitamin D responsive elements) of the upstream region of the responsive gene leading to activation/suppression of the target gene. In contrast, the non-genomic effect of 1, 25 (OH)₂D₃ is mediated through plasma membrane receptors of the target cells which are involved in ion channel activity regulation and also activate signal transduction pathways [3,4]. Exogenous 1, 25 (OH)₂D₃ in mice increased chondrocyte proliferation as well as enhancing

* Correspondence: paul.loughna@nottingham.ac.uk

¹School of Veterinary Medicine and Science, Faculty of Medicine and Health Sciences, University of Nottingham, Sutton Bonington Campus, Leicestershire LE12 5RD, UK

²Medical Research Council-Arthritis Research UK Centre for Musculoskeletal Ageing Research, University of Nottingham, Leicestershire, UK

Full list of author information is available at the end of the article

cartilage matrix mineralization [5]. The profound effects of vitamin D on calcium metabolism are well established and it has been reported that it can protect against cartilage loss in osteoarthritis in humans and rats [6].

Sox 9 (Sry-type high mobility group domain) transcription factor is expressed in articular chondrocytes, central nervous and urogenital systems [7,8]. It is expressed in all primordial cartilage tissues during embryogenesis [8,9] and functions to induce differentiation of stem cells to chondrocytes [10], regulate cartilage development and phenotypic maintenance during embryonic development [11]. Studies indicate that Sox 9 is involved in chondrocyte differentiation by regulating the expression of cartilage-specific genes such as collagen type II α 1 [8,12], XI α 2 [13], aggrecan [14], collagen link protein [15], cartilage oligomeric matrix protein and Cd-rap [16]. It has been shown that Sox 9 expression is maintained in articular chondrocytes and reduced in osteoarthritic chondrocytes [17].

Haploinsufficiency of Sox 9 results in campomelic dysplasia, a severe syndrome resulting in inadequate cartilage formation during development leading to severe dwarfism [18]. In chimaeric mice, Sox 9^{-/-} the prechondrogenic mesenchyme cells were prevented from differentiating into chondrocytes and lost their ability to express chondrocyte specific genes [19,20]. Sox 9 expression is also significantly reduced in osteoarthritic cartilage compared to normal healthy cartilage [17,21].

In vitro propagation of articular chondrocytes results in de-differentiation that is characterized by gradual loss of chondrocytic phenotype and acquisition of a fibroblastic phenotype [22] which is associated with a rapid decline in Sox 9 expression [23]. We have previously shown in equine chondrocytes that there is an association between the degree of de-differentiation and the expression and the transient receptor potential vanilloid (TRPV) channels 5 and 6 [24].

The transient receptor potential (TRP) superfamily is a non-selective cation ion channels with relative calcium selectivity. The TRPV sub-family is divided into two groups. TRPV1-4 channels are non-selective ion channels with modest permeation to calcium. This group can be activated by different stimuli such as heat/cold, chemical/mechanical stresses and binding to second messengers [25,26]. The other group comprises of TRPV5 and TRPV6 channels that are highly calcium selective and tightly regulated by cytosolic Ca²⁺ concentration [27,28]. The TRPV5 channel is implicated in Ca²⁺ reabsorption from the kidney, whereas TRPV6 channel is involved in Ca²⁺ absorption in the intestine [29]. Expression of both channels in human articular chondrocytes at mRNA level has been reported [30]. We also demonstrated their expression at the protein levels in equine articular chondrocytes (EAC) [24]. A correlation between the expression of TRPV5/6 channels and administered 1, 25 (OH)₂D₃ concentration has

also reported in intestinal endothelial cells [31,32], renal cells [33], osteoblasts [34,35]. Calbindin-D9K is a cytosolic Ca²⁺ binding protein, a member of cellular proteins found in the cells with high affinity for Ca²⁺ ions. Calbindin-D9k knockout mice models demonstrated that 1, 25 (OH)₂D₃ intake increased the expression of both channels [36]. The aim of the present study was to examine the effects of 1, 25 (OH)₂D₃ on the expression of Sox 9 and TRPV 5 and 6 in cultured equine chondrocytes.

Methods

Chondrocyte isolation

Articular chondrocytes were isolated from equine articular cartilage removed from load bearing synovial joints (metacarpophalangeal joints) of skeletally mature animals obtained on the day of slaughter from a local abattoir; these animals were euthanized for purposes other than research. All experiments were performed with local institutional ethical approval (University of Nottingham, School of Veterinary Medicine and Science Ethical Committee), in strict accordance with national guidelines. Articular cartilage slices were collected in serum free DMEM medium supplemented with 2% antibiotics (50 U/ml penicillin and 50 μ g/ml streptomycin) (Invitrogen, UK). Cartilage slices were washed in phosphate buffer saline (PBS) supplemented with 10% antibiotics (50 U/ml penicillin and 50 μ g/ml) for 30 min with agitation, followed by enzymatic digestion in freshly prepared 0.1% (v/w) collagenase type I from *clostridium histolyticum* enzyme dissolved in serum free medium supplemented with 2% antibiotics at relative humidity of 95%, 5% CO₂, 37C for 18 h. Undigested cartilage debris were removed from the cell/medium suspension by filtering the mixture through a nylon filter strainer of 70 μ m pore size (BD Bioscience, Europe). Supernatant was spun to isolate the chondrocytes, followed by three washes using PBS containing 10% antibiotics. Finally cells were suspended in DMEM contains 2% antibiotics and 10% FCS and cultured at 37C, 95% humidity and 5% CO₂ until confluent. Cells viability was assessed by the trypan blue exclusion test (Sigma Aldrich, UK).

Vitamin D treatment

To explore the influence of the active form of vitamin D (1, 25, α -dihydroxy vit D) on equine articular chondrocytes, cells were cultivated at 2 $\times 10^5$ cells/well until sub-confluence. Chondrocytes were treated with different concentrations of 1, 25 (OH)₂D₃ (1 $\times 10^{-9}$ and 1 $\times 10^{-12}$) [37] in serum free-medium for 24 h. The concentration of DMSO was below 0.1%. At the end of 1, 25 (OH)₂D₃ treatment, cells were washed three times sterile PBS and whole cell lysate was collected for protein expression using western blotting.

Western blotting

Total cellular protein lysate was isolated using radio-immunoprecipitation assay (RIPA) buffer (150 mM NaCl, 50 mM Tris HCl, pH 7.5, 5 mM ethylene glycol tetraacetic acid (EGTA), 1% Triton, 0.5% sodium deoxycholate and 0.1% sodium dodecyl sulphate) supplemented with phosphatase and protease inhibitors cocktail (Roche Diagnostic, Mannheim, Germany) on ice. Protein concentration was quantified by the Bradford method. 25 µg of total protein lysate was mixed with sample buffer (0.5 M Tris HCl, pH 6.8, 100% glycerol, 20% SDS, 0.5% bromophenol blue and 5% β-mercaptoethanol), separated on a 4/1-% polyacrylamide gel, then electrically transferred to PVDF membrane (Invitrogen, UK) by semi-dry apparatus (Bio-Rad, UK). Transferred proteins were blocked for non-specific binding in 5% (w/v) non-fat milk diluted in Tris base buffer saline with 0.1% tween 20 (TBS-T), followed by incubation with designated antibodies overnight. Membranes were washed five times in TBST, followed by 1 h incubation with goat anti-rabbit IgG conjugated with horseradish peroxidase (HRP) (Dako, UK) secondary antibody at room temperature. Finally, five washes were carried out for 5 min each followed by developing the membrane with the Amersham ECL western blot enhanced chemiluminescence kit (GE Healthcare, UK) and visualized by exposing to X-ray film.

Quantitative PCR

The quantitative real time PCR was carried out using a LightCycler 480 PCR System (Roche Diagnostics) using SYBR green DNA-binding fluorescent dye. 20 µl reactions were made in an optical 96-well reaction plate in triplicate and contained: a mixture of template cDNA (5 µl), sense and anti-sense gene specific primers (0.8 µl) (20 pmol), SYBR Green detection reagent and 3.4 µl RNA-free water. Reaction plate was sealed with ABI-prism optical adhesive cover, and spun at 2000 rpm for 2 min. The expression of target genes were normalised against GAPDH (Glyceraldehyde 3-phosphate dehydrogenase) and HPRT (Hypoxanthine-guanine phosphoribosyltransferase) using comparative cycle of threshold (Ct) value method.

Statistical analysis

Data values are presented as the mean ± SEM. Each experiment was performed in triplicate. The relative expression on the graphs represents the mean of a combination of three experiments. The differences between animals were analyzed utilizing Students *t*-test followed by the Bonferroni correction. *P* values less than or equal to 0.05 were considered statically significant.

Results and discussion

The active form of vitamin D (1, 25 (OH)₂D₃) has been shown to play an important role in Ca²⁺ homeostasis.

Intracellular Ca²⁺ concentration is involved in several chondrocyte functions including ECM biosynthesis. This study was designed to explore the possible influence of the active form of vitamin D (1, 25 (OH)₂D₃) treatment on the expression of Sox 9 and members of the TRP subfamily that are known as epithelial Ca²⁺ channels (TRPV5 and TRPV6 channels).

The influence of two doses (1 10⁻⁹ and 1 10⁻¹²) of 1, 25 (OH)₂D₃ treatment for 24 h on the expression of the transcription factor (Sox 9) at the protein level in equine chondrocytes was examined. Incubation of chondrocytes for 24 h with 1 10⁻¹² of 1, 25 (OH)₂D₃ significantly increased the expression of Sox 9 by nearly 3 fold, whereas treatment with 1 10⁻⁹ of 1, 25 (OH)₂D₃ induced more than 2 fold increase (*P* < 0.001) compared to the non-treated cells (Figure 1).

In chondrocytes, the mechanisms by which Sox 9 expression is regulated is of great interest due to the critical role played by this transcription factor in controlling the chondrocyte phenotype. The findings of the present study shows an elevation of Sox 9 protein in response to treatment with 1, 25 (OH)₂D₃. Isolation of chondrocytes results in their de-differentiation and loss of chondrocyte phenotype to fibroblast phenotype [22]. Recent studies have reported changes in the pattern of expression of some proteins during the course of chondrocytes de-differentiation. Cultivation of de-differentiated chondrocytes in alginate gel results in restoring the expression of transcription factor Sox 9 and chondrocyte differentiation markers such as collagen type II and aggrecan [38]. The current study indicated that treatment of cultured chondrocytes with 1, 25 (OH)₂D₃ results in restoring of the transcription factor Sox 9, which could indicate chondrocyte re-differentiation and restoration of the chondrocyte phenotype.

The physiological activities of Sox 9 protein were demonstrated by mouse genetic studies which indicated the importance of Sox 9 expression level in determining chondrocyte phenotype during development [39,40]. These studies demonstrated that chondrocyte differentiation and cartilage development were severely affected by knock-out or knock-in of a single allele of Sox 9. Moreover, the campomelic dysplasia is a genetic disorder characterized by multiple developmental abnormalities including cartilage induced by haplo-insufficiency of Sox 9 [18].

In human normal articular chondrocytes, Sox 9 expression was progressively reduced by passage in monolayer cell culture [41]. Moreover its expression is reduced in osteoarthritic chondrocytes compared to the normal articular chondrocytes [23] and has therefore been suggested to contribute in osteoarthritis disease processes by altering the ECM gene expression [42]. Thus, improving the expression of Sox 9 by 1, 25 (OH)₂D₃ treatment could




Figure 1 The effect of 1, 25 (OH)₂D₃ on protein expression of Sox 9 in equine articular chondrocytes following 24 h of incubation. **(A)** Western blot analysis of protein expression levels using an antibody raised against the Sox 9 protein. Freshly isolated equine whole chondrocyte lysate was used as a positive control C. β-actin was used as an internal load control. **(B)** the level of expression of the protein following 24 h of incubation with different doses (1 10⁻⁹ and 1 10⁻¹²) of 1, 25 (OH)₂D₃ treatment in addition to the solvent (DMSO). Values are presented as means ± S.E. ***Indicates statistically significant at P < 0.001.

provide a new insight into the prevention and/or treatment of osteoarthritis.

In fetal cartilage as well as in mature cartilage, the main role of Sox 9 is to maintain the chondrocyte phenotype in addition to inhibition of hypertrophic chondrocyte differentiation [43]. Therefore, down regulation of Sox 9 expression was suggested to be a precondition for hypertrophic alteration occurring in degenerative cartilage [42,44]. As Sox 9 protein is augmented by 1, 25 (OH)₂D₃ treatment, this study suggests that 1, 25 (OH)₂D₃ treatment could be utilized to improve the ECM proteins biosynthesis by enhancing the anabolic activities of articular chondrocytes. After further passages, chondrocytes have been shown to exhibit a more pronounced de-differentiated phenotype and lower levels of Sox 9 [41]. Further studies on the effects of 1, 25 (OH)₂D₃ upon these cells would be of great interest.

We have previously shown that the expression of both TRPV5 and 6 channels, at the protein level, are inversely related to chondrocyte de-differentiation [24]. In this study we therefore examined the influence of 1, 25 (OH)₂D₃ treatment on the expression of TRPV5/6 channels on primary equine articular chondrocytes isolated at passage two. The findings of the current study indicated that no

statistically significant changes were observed on the expression of TRPV5 and TRPV6 channels at the protein level following treatment with different doses of 1, 25 (OH)₂D₃ (Figure 2). To ensure that 1, 25 (OH)₂D₃ had no effect on the expression of these channels we also examined its effect upon their transcript levels but again no significant changes were observed (Figure 3). Previous studies conducted on mouse investigated the effect of 1, 25 (OH)₂D₃ on epithelial Ca²⁺ transport including TRPV5 and 6 channels following 1, 25 (OH)₂D₃ treatment [45,46]. These studies demonstrated that the expression of TRPV5 and TRPV6 channels in kidney and duodenum were stimulated by binding of VDR (vitamin D receptor) at transcriptional level. TRPV6 was reported to have several classes of VDREs (vitamin D response elements) in humans and the mouse [45,47,48].

In the vitamin D receptor knockout mice model, the level of gene expression pattern of the TRPV6 and 5 channels in duodenum is dramatically down-regulated, in contrast no changes are observed in gene expression of other Ca²⁺ transporters [49,50]. The role of vitamin D in the stimulation of Ca²⁺ transporters has emerged recently. Therefore, the correlation between 1, 25 (OH)₂D₃ treatment and Ca²⁺ transport proteins were investigated





Figure 2 Expression of TRPV5 and TRPV6 channels in equine articular chondrocytes at the protein level. The changes in expression of TRPV5 (A) and TRPV6 (B) channels following treatment with 1, 25 vit D were assessed by densitometry analysis of western blots and normalized to the expression of housekeeping protein (β -Tubulin). Equine kidney (K) lysate was used as a positive control. Levels of expression at the protein level of the TRPV5 (C) and TRPV6 (D) channels following 24 h of incubation with different doses of 1, 25 (OH) $_2$ D $_3$ in addition to the solvent (DMSO). Data presented as a mean \pm S.E.

in several cell types. 1, 25 (OH) $_2$ D $_3$ was demonstrated to increase the expression of TRPV6 channel, calbindin-D9k and PMCA1b genes in Caco-2 cell lines [36,51]. This finding was not consistent with the current study, where no changes were observed on the expression of either TRPV5 or 6 channels at mRNA level or protein levels. As stated above we have previously observed an inverse correlation between TRPV 5 and 6 levels and differentiation state. The current study suggests that the changes in levels may not be directly linked to the de-differentiation process. This however, needs further study as does the mechanism by which Sox-9 is regulated by 1,

25 (OH) $_2$ D $_3$. Ca $^{2+}$ ions and Ca $^{2+}$ channels have been implicated in chondrocyte metabolism [52,53]. In vitro studies show that augmented extracellular Ca $^{2+}$ promotes differentiation toward a hypertrophic phenotype, in contrast Ca $^{2+}$ reduction improves ECM protein synthesis including collagen type II and aggrecan and delays hypertrophy [54]. Therefore, investigating the effect of the 1, 25 (OH) $_2$ D $_3$ on the expression of Ca $^{2+}$ channels could be considered as a candidate for treatment of joint diseases.

The results of the current study do not, however, suggest an involvement of TRPV5/6 ion channels in the regulation of vitamin D mediated up-regulation of sox-9.



Conclusions

This study examined the effects of the active form of vitamin D (1, 25 (OH)₂D₃) upon the expression of transcription factor Sox 9 and the calcium sensitive TRPV 5 and 6 channels, in equine articular chondrocytes, *in vitro*. An increased expression of Sox 9, in these chondrocytes, in response to the active form of vitamin D suggests

that this compound acts to inhibit the progressive de-differentiation that is normally observed in these cells. It is also supportive of previous studies indicating that 1 α , 25-dihydroxyvitamin D₃ can have a protective effect upon cartilage in animals *in vivo*. There was no effect of vitamin D (1, 25 (OH)₂D₃) upon the expression of either TRPV channel at either the protein or mRNA

level. We have previously observed a correlation between the degree of differentiation and the expression levels of TRPV 5/6 and suggested that these ion channels may have a direct interaction with the differentiation process *in vitro*. The data in the present study do not support this.

Competing interests

The authors declare that they have no competing interests.

Authors contributions

ISM and PTL designed the experiment and analyzed the results. ISM carried out cell culture work. Both authors wrote the manuscript and made subsequent changes.

Author details

¹School of Veterinary Medicine and Science, Faculty of Medicine and Health Sciences, University of Nottingham, Sutton Bonington Campus, Leicestershire LE12 5RD, UK. ²Medical Research Council-Arthritis Research UK Centre for Musculoskeletal Ageing Research, University of Nottingham, Leicestershire, UK. ³School of Veterinary Medicine and Science, Tripoli University, Tripoli, Libya.

Received: 26 June 2014 Accepted: 19 November 2014

Published online: 17 December 2014

References

1. Boyan BD, Sylvia VL, Dean DD, Del Toro F, Schwartz Z: **Differential regulation of growth plate chondrocytes by 1 α ,25-(OH) $_2$ D $_3$ and 24R,25-(OH) $_2$ D $_3$ involves cell-maturation-specific membrane-receptor-activated phospholipid metabolism.** *Crit Rev Oral Biol Med* 2002, **13**(2):143–154.
2. Tetlow LC, Smith SJ, Mawer EB, Woolley DE: **Vitamin D receptors in the rheumatoid lesion: expression by chondrocytes, macrophages, and synoviocytes.** *Ann Rheum Dis* 1999, **58**(2):118–121.
3. Lehmann B, Meurer M: **Vitamin D metabolism.** *Dermatol Ther* 2010, **23**(1):2–12.
4. Fleet JC: **Genomic and proteomic approaches for probing the role of vitamin D in health.** *Am J Clin Nutr* 2004, **80**(6 Suppl):1730S–1734S.
5. Xue Y, Karaplis AC, Hendy GN, Goltzman D, Miao D: **Exogenous 1,25-dihydroxyvitamin D $_3$ exerts a skeletal anabolic effect and improves mineral ion homeostasis in mice that are homozygous for both the 1 α -hydroxylase and parathyroid hormone null alleles.** *Endocrinology* 2006, **147**(10):4801–4810.
6. Castillo EC, Hernandez-Cueto MA, Vega-Lopez MA, Lavalle C, Kouri JB, Ortiz-Navarrete V: **Effects of vitamin D supplementation during the induction and progression of osteoarthritis in a rat model.** *Evid Based Complement Altern Med* 2012, **2012**:156563.
7. Morais Da Silva S, Hacker A, Harley V, Goodfellow P, Swain A, Lovell-Badge R: **Sox9 expression during gonadal development implies a conserved role for the gene in testis differentiation in mammals and birds.** *Nat Genet* 1996, **14**(1):62–68.
8. Zhao Q, Eberspaecher H, Lefebvre V, De Crombrugge B: **Parallel expression of Sox9 and Col2a1 in cells undergoing chondrogenesis.** *Dev Dyn* 1997, **209**(4):377–386.
9. Wright E, Hargrave MR, Christiansen J, Cooper L, Kun J, Evans T, Gangadharan U, Greenfield A, Koopman P: **The Sry-related gene Sox 9 is expressed during chondrogenesis in mouse embryos.** *Nat Genet* 1995, **9**(1):15–20.
10. Shintani N, Hunziker EB: **Chondrogenic differentiation of bovine synovium: bone morphogenetic proteins 2 and 7 and transforming growth factor beta1 induce the formation of different types of cartilaginous tissue.** *Arthritis Rheum* 2007, **56**(6):1869–1879.
11. Sive JJ, Baird P, Jeziorski M, Watkins A, Hoyland JA, Freemont AJ: **Expression of chondrocyte markers by cells of normal and degenerate intervertebral discs.** *Mol Pathol* 2002, **55**(2):91–97.
12. Lefebvre V, Huang W, Harley VR, Goodfellow PN, de Crombrugge B: **SOX 9 is a potent activator of the chondrocyte-specific enhancer of the pro alpha1(II) collagen gene.** *Mol Cell Biol* 1997, **17**(4):2336–2346.
13. Bridgewater LC, Lefebvre V, de Crombrugge B: **Chondrocyte-specific enhancer elements in the Col11a2 gene resemble the Col2a1 tissue-specific enhancer.** *J Biol Chem* 1998, **273**(24):14998–15006.
14. Sekiya I, Tsuji K, Koopman P, Watanabe H, Yamada Y, Shinomiya K, Nifuji A, Noda M: **SOX9 enhances aggrecan gene promoter/enhancer activity and is up-regulated by retinoic acid in a cartilage-derived cell line, TC6.** *J Biol Chem* 2000, **275**(15):10738–10744.
15. Kou I, Ikegawa S: **SOX9-dependent and -independent transcriptional regulation of human cartilage link protein.** *J Biol Chem* 2004, **279**(49):50942–50948.
16. Xie WF, Zhang X, Sakano S, Lefebvre V, Sandell LJ: **Trans-activation of the mouse cartilage-derived retinoic acid-sensitive protein gene by Sox 9.** *J Bone Miner Res* 1999, **14**(5):757–763.
17. Brew CJ, Clegg PD, Boot-Handford RP, Andrew JG, Hardingham T: **Gene expression in human chondrocytes in late osteoarthritis is changed in both fibrillated and intact cartilage without evidence of generalised chondrocyte hypertrophy.** *Ann Rheum Dis* 2010, **69**(1):234–240.
18. Wagner T, Wirth J, Meyer J, Zabel B, Held M, Zimmer J, Pasantes J, Bricarelli FD, Keutel J, Hustert E, Wolf U, Tommerup N, Schempp W, Scherer G: **Autosomal sex reversal and campomelic dysplasia are caused by mutations in and around the SRY-related gene SOX9.** *Cell* 1994, **79**(6):1111–1120.
19. Bi W, Deng JM, Zhang Z, Behringer RR, de Crombrugge B: **Sox 9 is required for cartilage formation.** *Nat Genet* 1999, **22**(1):85–89.
20. Hargus G, Kist R, Kramer J, Gerstel D, Neitz A, Scherer G, Rohwedel J: **Loss of Sox 9 function results in defective chondrocyte differentiation of mouse embryonic stem cells in vitro.** *Int J Dev Biol* 2008, **52**(4):323–332.
21. Cucchiariini M, Thurn T, Weimer A, Kohn D, Terwilliger EF, Madry H: **Restoration of the extracellular matrix in human osteoarthritic articular cartilage by overexpression of the transcription factor SOX9.** *Arthritis Rheum* 2007, **56**(1):158–167.
22. Benya PD, Padilla SR, Nimmi ME: **Independent regulation of collagen types by chondrocytes during the loss of differentiated function in culture.** *Cell* 1978, **15**(4):1313–1321.
23. Tew SR, Clegg PD, Brew CJ, Redmond CM, Hardingham TE: **SOX9 transduction of a human chondrocytic cell line identifies novel genes regulated in primary human chondrocytes and in osteoarthritis.** *Arthritis Res Ther* 2007, **9**(5):R107.
24. Hdud IM, El-Shafei AA, Loughna P, Barrett-Jolley R, Mobasheri A: **Expression of transient receptor potential vanilloid (TRPV) channels in different passages of articular chondrocytes.** *Int J Mol Sci* 2012, **13**(4):4433–4445.
25. Hdud IM, Mobasheri A, Loughna PT: **Effect of osmotic stress on the expression of TRPV4 and BKCa channels and possible interaction with ERK1/2 and p38 in cultured equine chondrocytes.** *Am J Physiol Cell Physiol* 2014, **306**(11):C1050–C1057.
26. Hdud IM, Mobasheri A, Loughna PT: **Effects of cyclic equibiaxial mechanical stretch on alpha-BK and TRPV4 expression in equine chondrocytes.** *SpringerPlus* 2014, **3**:59.
27. Vennekens R, Hoenderop JG, Prenen J, Stuiver M, Willems PH, Droogmans G, Nilius B, Bindels RJ: **Permeation and gating properties of the novel epithelial Ca(2+) channel.** *J Biol Chem* 2000, **275**(6):3963–3969.
28. Nilius B, Vennekens R, Prenen J, Hoenderop JG, Bindels RJ, Droogmans G: **Whole-cell and single channel monovalent cation currents through the novel rabbit epithelial Ca2+ channel ECaC.** *J Physiol* 2000, **527**(Pt 2):239–248.
29. Hoenderop JG, Nilius B, Bindels RJ: **Calcium absorption across epithelia.** *Physiol Rev* 2005, **85**(1):373–422.
30. Gavenis K, Schumacher C, Schneider U, Eisfeld J, Mollenhauer J, Schmidt-Rohlfing B: **Expression of ion channels of the TRP family in articular chondrocytes from osteoarthritic patients: changes between native and in vitro propagated chondrocytes.** *Mol Cell Biochem* 2009, **321**(1–2):135–143.
31. Tapania S, Fleet JC, Peng JB, Wang XD, Wood RJ: **1,25-Dihydroxyvitamin D and 25-hydroxyvitamin D mediated regulation of TRPV6 (a putative epithelial calcium channel) mRNA expression in Caco-2 cells.** *Eur J Nutr* 2006, **45**(4):196–204.
32. Teerapornpuntakit J, Dorkkam N, Wongdee K, Krishnamra N, Charoenphandhu N: **Endurance swimming stimulates transepithelial calcium transport and alters the expression of genes related to calcium absorption in the intestine of rats.** *Am J Physiol Endocrinol Metab* 2009, **296**(4):E775–E786.

33. Embark HM, Setiawan I, Poppendieck S, van de Graaf SF, Boehmer C, Palmada M, Wieder T, Gerstberger R, Cohen P, Yun CC, Bindels RJ, Lang F: Regulation of the epithelial Ca²⁺ channel TRPV5 by the NHE regulating factor NHERF2 and the serum and glucocorticoid inducible kinase isoforms SGK1 and SGK3 expressed in *Xenopus* oocytes. *Cell Physiol Biochem* 2004, **14**(4-6):203-212.
34. Zanello LP, Norman AW: Multiple molecular mechanisms of 1 α , 25(OH)₂-vitamin D₃ rapid modulation of three ion channel activities in osteoblasts. *Bone* 2003, **33**(1):71-79.
35. Zhang M, Wang JJ, Chen YJ: Effects of mechanical pressure on intracellular calcium release channel and cytoskeletal structure in rabbit mandibular condylar chondrocytes. *Life Sci* 2006, **78**(21):2480-2487.
36. Lee GS, Jung EM, Choi KC, Oh GT, Jeung EB: Compensatory induction of the TRPV6 channel in a calbindin-D9k knockout mouse: Its regulation by 1,25-hydroxyvitamin D₃. *J Cell Biochem* 2009, **108**(5):1175-1183.
37. Fernandez-Cancio M, Audi L, Carrascosa A, Toran N, Andaluz P, Esteban C, Granada ML: Vitamin D and growth hormone regulate growth hormone/insulin-like growth factor (GH-IGF) axis gene expression in human fetal epiphyseal chondrocytes. *Growth Horm IGF Res* 2009, **19**(3):232-237.
38. Kumar D, Lassar AB: The transcriptional activity of Sox9 in chondrocytes is regulated by RhoA signaling and actin polymerization. *Mol Cell Biol* 2009, **29**(15):4262-4273.
39. Akiyama H, Lyons JP, Mori-Akiyama Y, Yang X, Zhang R, Zhang Z, Deng JM, Taketo MM, Nakamura T, Behringer RR, McCrea PD, de Crombrughe B: Interactions between Sox9 and beta-catenin control chondrocyte differentiation. *Genes Dev* 2004, **18**(9):1072-1087.
40. Kawakami Y, Rodriguez-Leon J, Izpisua Belmonte JC: The role of TGF β s and Sox9 during limb chondrogenesis. *Curr Opin Cell Biol* 2006, **18**(6):723-729.
41. Tew SR, Hardingham TE: Regulation of SOX9 mRNA in human articular chondrocytes involving p38 MAPK activation and mRNA stabilization. *J Biol Chem* 2006, **281**(51):39471-39479.
42. Aigner T, Gebhard PM, Schmid E, Bau B, Harley V, Poschl E: SOX9 expression does not correlate with type II collagen expression in adult articular chondrocytes. *Matrix Biol* 2003, **22**(4):363-372.
43. Bi W, Huang W, Whitworth DJ, Deng JM, Zhang Z, Behringer RR, de Crombrughe B: Haploinsufficiency of Sox9 results in defective cartilage primordia and premature skeletal mineralization. *Proc Natl Acad Sci U S A* 2001, **98**(12):6698-6703.
44. Gebhard PM, Gehrsitz A, Bau B, Soder S, Eger W, Aigner T: Quantification of expression levels of cellular differentiation markers does not support a general shift in the cellular phenotype of osteoarthritic chondrocytes. *J Orthop Res* 2003, **21**(1):96-101.
45. Nijenhuis T, Hoenderop JG, van der Kemp AW, Bindels RJ: Localization and regulation of the epithelial Ca²⁺ channel TRPV6 in the kidney. *J Am Soc Nephrol* 2003, **14**(11):2731-2740.
46. Song Y, Peng X, Porta A, Takanao H, Peng JB, Hediger MA, Fleet JC, Christakos S: Calcium transporter 1 and epithelial calcium channel messenger ribonucleic acid are differentially regulated by 1, 25 dihydroxyvitamin D₃ in the intestine and kidney of mice. *Endocrinology* 2003, **144**(9):3885-3894.
47. Haussler MR, Whitfield GK, Kaneko I, Haussler CA, Hsieh D, Hsieh JC, Jurutka PW: Molecular mechanisms of vitamin D action. *Calcif Tissue Int* 2013, **92**(2):77-98.
48. Hoenderop JG, Muller D, Van Der Kemp AW, Hartog A, Suzuki M, Ishibashi K, Imai M, Sweep F, Willems PH, Van Os CH, Bindels RJ: Calcitriol controls the epithelial calcium channel in kidney. *J Am Soc Nephrol* 2001, **12**(7):1342-1349.
49. Bouillon R, Van Cromphaut S, Carmeliet G: Intestinal calcium absorption: molecular vitamin D mediated mechanisms. *J Cell Biochem* 2003, **88**(2):332-339.
50. Van Cromphaut SJ, Dewerchin M, Hoenderop JG, Stockmans I, Van Herck E, Kato S, Bindels RJ, Collen D, Carmeliet P, Bouillon R, Carmeliet G: Duodenal calcium absorption in vitamin D receptor-knockout mice: functional and molecular aspects. *Proc Natl Acad Sci U S A* 2001, **98**(23):13324-13329.
51. Wood RJ, Tchack L, Tapania S: 1,25-Dihydroxyvitamin D₃ increases the expression of the CaT1 epithelial calcium channel in the Caco-2 human intestinal cell line. *BMC Physiol* 2001, **1**:11.
52. Kirsch T, Swoboda B, von der Mark K: Ascorbate independent differentiation of human chondrocytes in vitro: simultaneous expression of types I and X collagen and matrix mineralization. *Differentiation* 1992, **52**(1):89-100.
53. Bonen DK, Schmid TM: Elevated extracellular calcium concentrations induce type X collagen synthesis in chondrocyte cultures. *J Cell Biol* 1991, **115**(4):1171-1178.
54. Koyano Y, Hejna M, Flechtenmacher J, Schmid TM, Thonar EJ, Mollenhauer J: Collagen and proteoglycan production by bovine fetal and adult chondrocytes under low levels of calcium and zinc ions. *Connect Tissue Res* 1996, **34**(3):213-225.

doi:10.1186/s40781-014-0033-1

Cite this article as: Hdud and Loughna: Influence of 1 α , 25-dihydroxyvitamin D₃ [1, 25(OH)₂D₃] on the expression of Sox 9 and the transient receptor potential vanilloid 5/6 ion channels in equine articular chondrocytes. *Journal of Animal Science and Technology* 2014 **56**:33.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

