



The University of
Nottingham

UNITED KINGDOM · CHINA · MALAYSIA

Burroughs, Laurence and Eccleshare, Lee and Ritchie, John and Kulkarni, Omkar and Lygo, Barry and Woodward, Simon and Lewis, William (2015) One-pot Cannizzaro cascade synthesis of ortho-Fused Cycloocta-2,5-dien-1-ones from 2-Bromo(hetero)aryl aldehydes. *Angewandte Chemie International Edition*, 54 (36). pp. 10648-10651. ISSN 1521-3773

Access from the University of Nottingham repository:

http://eprints.nottingham.ac.uk/46633/1/Burroughs_et_al-2015-Angewandte_Chemie_International_Edition.pdf

Copyright and reuse:

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the Creative Commons Attribution licence and may be reused according to the conditions of the licence. For more details see:
<http://creativecommons.org/licenses/by/2.5/>

A note on versions:

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact eprints@nottingham.ac.uk

One-Pot Cannizzaro Cascade Synthesis of *ortho*-Fused Cycloocta-2,5-dien-1-ones from 2-Bromo(hetero)aryl Aldehydes

Laurence Burroughs, Lee Eccleshare, John Ritchie, Omkar Kulkarni, Barry Lygo, Simon Woodward,* and William Lewis


Abstract: An intramolecular Cannizzaro-type hydride transfer to an *in situ* prepared allene enables the synthesis of *ortho*-fused 4-substituted cycloocta-2,5-dien-1-ones with unprecedented technical ease for an eight-ring carboannulation. Various derivatives could be obtained from commercially available (hetero)aryl aldehydes, trimethylsilylacetylene, and simple propargyl chlorides in good yields.

Readily (often commercially) available 2-bromo(hetero)aryl aldehydes (**1**, Scheme 1) constitute powerful and popular units for the construction of *ortho*-fused carbocyclic rings.^[1] Two strategies commonly employed in the multitudinous syntheses described in the literature involve 1) the derivatization of **1** to intermediates of type **A**, with a π -system attacking an electrophile that is ultimately derived from the aldehyde by functional-group manipulations (e.g., an alcohol, acetal, or even the aldehyde itself); or 2) formyl group homologation to an anion-stabilizing unit that attacks a suitably activated π -bond (**B**). Whereas many elegant catalytic systems that use intermediates **A** or **B** have already been described,^[2,3] the preparation of the suitable precursors is often step-intensive, raising questions on the overall process

efficiency in the “age of sustainability”.^[4] We speculated that a new annulation strategy, based on a Cannizzaro-type reaction via intermediate **C**, might be possible. As the hydride transfer simultaneously exposes a powerful Michael acceptor (ynone or equivalent) and a carbanion in close proximity, efficient annulation might be expected to occur (especially if **C** is directly attained *in situ*). Herein, we describe the use of such a strategy for the formation of eight-membered rings. The paucity of direct single-pot/step procedures to such medium rings attests to the known issues associated with their synthesis;^[5] they therefore provide a stringent test for strategy **C**. Although redox-based Cannizzaro reactions, including asymmetric and triggered C–C bond-forming versions, have recently been developed,^[6] to the best of our knowledge, they have not been employed for carboannulation reactions to medium rings. Genuine single-pot/step procedures to eight-membered rings are very limited—Repe’s historical catalytic access to cyclooctatetraene^[7] and Murakami’s [4+2+2] approach^[8] being notable exceptions. Modern alternatives frequently describe catalytic cyclizations that proceed with exquisitely high yields (for the final step). Unfortunately, their overall synthetic efficiency is often compromised by the multistep syntheses required to obtain the cyclization precursors.^[9]

Initial investigation of our own proposal (**C**) centered on the combination of readily available 2-bromobenzaldehyde (**1a**), trimethylsilylacetylene (**2a**), and propargyl chloride **3a**. Encouragingly, even an initial run returned significant amounts of benzo[8]annulene derivative **4aaa** (Table 1, entry 1, ca. 7%) whose identity and regiochemistry was confirmed by X-ray crystallography (see the Supporting Information) as initial NMR data was not conclusive.^[10] Unfortunately, under the conditions of entry 1, multiple side products were also produced. Upon changing from an organocopper to an organocuprate formulation (entry 2), the reaction became much cleaner, but the yield of **4aaa** remained low. The recovered side products of entry 2 included significant amounts of the alcohols **5a** and **6a** together with unreacted **3a**. We could confirm that formation of **D** was almost quantitative in < 20 min under all conditions tried (Et₂O or THF, –78 to 0°C, based on recovered **5a** in independent reactions). We therefore suspected issues associated with both *n*BuLi–halogen exchange from **D** and the copper-promoted S_N2’ reaction leading to the progenitor for eight-ring formation. Further experiments were designed to probe these hypotheses (Table 1).

Organometallic compounds derived from **D** were poorly soluble in Et₂O so the reaction solvent was switched to THF (entry 3). This led to an improved reaction that enabled us to




Scheme 1. Proposed use of 2-bromo(hetero)aryl aldehydes in Cannizzaro-triggered annulation cascades compared to traditional approaches. EWG = electron-withdrawing group, LA = Lewis acid, R¹–R³ are generic groups, *n* and *m* are typically 0–1.

[*] Dr. L. Burroughs, L. Eccleshare, J. Ritchie, O. Kulkarni, Prof. Dr. B. Lygo, Prof. Dr. S. Woodward, Dr. W. Lewis
School of Chemistry, University of Nottingham
University Park, Nottingham NG7 2RD (UK)
E-mail: simon.woodward@nottingham.ac.uk

Supporting information for this article is available on the WWW under <http://dx.doi.org/10.1002/anie.201505347>.

© 2015 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.


Table 1: Optimization of the Cannizzaro-triggered cascade reaction to eight-ring compound **4aaa**.^[a]


Entry	Solvent	Halogen exchange of intermediate D	Transmetalation with CuBr-SMe ₂	Addition of 3a and cyclization	Yield 4aaa [%]
1	Et ₂ O	<i>t</i> BuLi (2 equiv), -50°C, 20 min	1 equiv, -50°C, 20 min	1 equiv, -50°C to RT over 5 h	< 10
2	Et ₂ O	<i>n</i> BuLi (1 equiv), 0°C, 10 min	0.5 equiv, -50°C, 5 min	0.5 equiv, -50°C to +10°C over 1.2 h	14 ^[b]
3	THF	<i>n</i> BuLi (1 equiv), -50°C, 15 min	0.5 equiv, -50°C, 10 min	0.5 equiv, -50°C to 10°C over 1.2 h	53
4	THF	<i>n</i> BuLi (1 equiv), -50°C, 15 min	0.5 equiv, -50°C, 1 h	0.5 equiv, -50°C to -10°C over 1.5 h; kept at -10°C for 1 h	70 (69) ^[c]
5	THF	<i>n</i> BuLi (1 equiv), -50°C, 15 min	0.5 equiv, -50°C, 1 h	0.5 equiv, -50°C to -10°C over 1.5 h; kept at -10°C for 2 h	35

[a] Reaction conditions: **1a** (0.71 mmol), **2a** (0.77 mmol)/*n*BuLi (0.72 mmol), CuBr-SMe₂ (0.36–0.71 mmol), **3a** (0.35–0.70 mmol) in Et₂O or THF (1.0 mL). The yields of **4aaa** were determined by GC analysis against genuine samples and an internal standard (1-methylnaphthalene, 50 μL). [b] The mass balance of the reaction included **3a**, **5a**, and **6a** in a 2.4:1.0:1.8 ratio. [c] The yield of isolated product for a reaction performed with 1.39 mmol of **3a** is shown in parentheses.

identify that the onset of the Cannizzaro-type cascade reaction occurs at -20 to -10°C for this substrate. Independent treatment of THF solutions containing **D** with *n*BuLi followed by quenching with D₂O confirmed that complete Br/Li exchange had occurred within 15 min at -40 to -60°C, and that the resulting organolithium species was rather stable at these temperatures. However, above approximately -15°C, the same species began to decompose rapidly. Thus, -40 to -50°C was selected as the temperature range for optimizing the transmetalation to CuBr-SMe₂. When the nominal diorganocuprate was prepared under the conditions of entry 3 (-50°C, 10 min, from the aryl lithium intermediate with CuBr-SMe₂) and then treated with methyl vinyl ketone, very poor conversions into the 1,4-addition product were attained. This strongly suggested that transmetalation to Cu^I is rather slow for this system. Gratifyingly, extending the time for Li/Cu exchange to one hour (entry 4) significantly increased the quantity of **4aaa** formed, and the GC yields were comparable to those attained through isolation (70% vs. 69%). Further extending the transmetalation time to two hours at -10°C had a detrimental effect on the yield of **4aaa** (entry 5).

With an optimized procedure for the preparation of parent compound **4aaa** in hand, we studied the scope of the reaction with respect to the 2-bromo(hetero)aryl aldehyde component **1**. Using the conditions of entry 4 (Table 1) and aldehydes **1b–1j**, the eight-ring compounds that are shown in Scheme 2 were attained. Substitution at the 3-position was better tolerated than at the 1- and 4-positions (products **4baa/4caa** vs. **4daa/4faa**). Monosubstitution at the 2-position was also tolerated. A comparison of the yields of **4baa–4gaa** indicates that steric factors are more important than electronic ones; for example, intramolecular OMe steric contacts increase the *ortho* steric profile of **4eaa** versus **4jaa**. The yields of compounds **4haa–4jaa** also support this trend. Whereas the yields of **4aaa–4jaa** are modest, it should be borne in mind that they represent one-pot processes to analytically pure crystalline products on up to multigram scales, using only inexpensive commercial starting materials. The new process compares very favorably to the consolidated




Scheme 2. Variation of the 2-bromo(hetero)aryl aldehyde; yields starting from the commercial aldehydes are given.^[9] [a] The precursor aldehyde was obtained from isatin (3 steps).


yields and time efficiencies of previously reported eight-ring syntheses.^[9]

By varying the acetylene component using the readily attained starting materials **2b–2f**, five further examples of eight-ring compounds **4** were synthesized (Scheme 3). The terminal alkyne substituent could encompass alkyl, vinyl, or aryl units, and the corresponding products were all obtained in acceptable yields. Finally, variation of the propargyl chloride was investigated using simple **3b–3i** (Scheme 3). The aryl moiety could accommodate both electron-donating and -withdrawing substituents and some functional groups (**4aae** and **4aaf**).

Whereas the full mechanistic features of our Cannizzaro-triggered cascade are still under investigation, a working hypothesis is shown in Scheme 4. Sequential 1,2-acetylide addition, halogen exchange, and transmetalation to copper affords intermediate **E**. This copper(I) species undergoes a γ -selective S_N2' addition to propargyl chloride **3** affording the non-isolable allene **C'**, which undergoes the Cannizzaro-type hydride shift proposed in Scheme 1. The resulting allyl anion




Scheme 3. Variation of the terminal acetylene and propargyl chloride. [a] Run at -50°C .



Scheme 4. Mechanistic proposal for the formation of eight-ring compounds **4**; L is indicative of the presence of a generic organocopper or -cuprate species.

is presumably placed in close proximity to the potent ynone Michael acceptor, which leads to efficient ring closure. The use of an aryl moiety as the R^2 substituent is apparently sufficient to stabilize the allyl anion and allow the necessary *trans* to *cis* isomerization. Protonation of the non-classical enolate **G** results in the formation of **4**. Several pieces of evidence support this chain of events. First, attempts to use 2-thienylcyanocuprates derived from a related aza substrate led to the isolation of **7** (X-ray structure) where the intermediate allene has been carbometalated rather than



Scheme 5. The computational models *anti-C'* and *syn-C'* and indirect evidence for the formation of **G**.

having accepted a hydride. For the formation of **4iaa**, we could isolate appreciable amounts of **8**, which is the expected product of a Cannizzaro-type hydride transfer without subsequent annulation. Use of the deuterated aldehyde 2-BrC₆H₄CDO ([D]-**1a**) led to the formation of the expected product [D]-**4aaa**, confirming the participation of the aldehyde in the Cannizzaro process.

The mechanistic proposal shown in Scheme 4 was also probed by DFT calculations at the M06-2X/6-31G(d) and CBS-QB3 levels of theory^[11] on the “metal-free” anions *anti-C'* and *syn-C'* (Scheme 5; see also the Supporting Information). Whereas the relationship of *anti-C'* and *syn-C'* to the reaction coordinate of the true organometallic species involved in the cascade process needs to be treated with caution, it does provide further evidence for the proposed mechanism. The calculated barriers for the Cannizzaro-type hydride transfer and the rearrangement of the allyl anion (the equivalent of the transformation of **C'** into **F** in Scheme 4) were approximately 5 and 7 kcal mol⁻¹, respectively. The calculations also indicate that non-classical enolates of type **G** are favored in the reaction over the equivalent allenolate structures. This idea was further reinforced by the observation that trapping of the reaction intermediates with iodine, iodomethane, or carbon dioxide, as opposed to a proton, leads to the formation of derivatives **9–11** (Scheme 5).

To the best of our knowledge, this Cannizzaro-triggered cascade process is unique. The nearest analogues that we could find are cyclizations reported by the Oonishi and Sato groups;^[9b,d] however, in these cases, the allene intermediate must be separately synthesized, and no Cannizzaro-type process is involved. Similarly, a single eight-ring side product (formed in 10% yield) was obtained by Alajarin and Vidal through a 1,5-hydride shift/ 8π electrocyclization from a neutral 1,2-allenylacetal at 110°C .^[12] Again, no Cannizzaro-type trigger was proposed. Whereas Cannizzaro shifts have been used for oxidation-state manipulations in *ortho*-fused systems, in these processes, no annulation steps were involved.^[13] The most similar ynone eight-ring closure process that could be identified was developed by Schreiber and Kelly,^[14] but this transformation involves an alkoxide nucleophile.

In conclusion, we believe that this very simple formation of *ortho*-fused cycloocta-2,5-dien-1-one units demonstrates the potential of the developed method for rapid and efficient annulations. Investigations into extending the scope of both the present reaction and other variants are ongoing.

Acknowledgements

This study was initiated under funding from the European Union's Seventh Programme for research, technological

development and demonstration (308768). J.R. is grateful to European Thermodynamics Ltd and the Engineering Physical Sciences Research Council (EPSRC) for provision of a studentship. L.B. thanks the EPSRC's National Service for Computational Chemistry (NSCCS) for providing computational facilities, and O.K. acknowledges a Kishore Vaigyanik Protsahan Yojana (KVPJ) Fellowship.

Keywords: aldehydes · annulation · carbocycles · medium-ring compounds · synthetic methods

How to cite: *Angew. Chem. Int. Ed.* **2015**, *54*, 10648–10651
Angew. Chem. **2015**, *127*, 10794–10797

- [1] a) *Science of Synthesis—Compounds with two carbon-heteroatom bonds, aldehydes*, Vol. 25 (Ed.: R. Brückner), Georg Thieme, Stuttgart, **2007**, p. 901, also Sections 25.4–25.6, pp. 463–710; b) P. Merino in *Science of Synthesis—Monocyclic Arenes, Quasiarenes, and Annulenes*, Vol. 45a (Eds.: J. S. Siegel, Y. Tobe), Georg Thieme, Stuttgart, **2009**, pp. 253–327, also Sections 45.6.3–45.6.4.
- [2] For selected recent examples of strategy A, see: a) A. R. Jagdale, J. H. Park, S. W. Youn, *J. Org. Chem.* **2011**, *76*, 7204–7215; b) D. Das, S. Pratihari, S. Roy, *Org. Lett.* **2012**, *14*, 4870–4873; c) W. Zi, F. D. Toste, *J. Am. Chem. Soc.* **2013**, *135*, 12600–12603; d) P. Sarnpitak, K. Trongchit, Y. Kostenko, S. Sathalalai, M. P. Gleeson, S. Ruchirawat, P. Ploypradith, *J. Org. Chem.* **2013**, *78*, 8281–8296; e) M. Chen, N. Sun, Y. Liu, *Org. Lett.* **2013**, *15*, 5574–5577; f) A. S. K. Hashmi, J. Hofmann, S. Shi, A. Schütz, M. Rudolph, C. Lotzschütz, M. Wietek, M. Bührle, M. Wölfle, F. Rominger, *Chem. Eur. J.* **2013**, *19*, 382–389 (also for reactivity of type B).
- [3] For selected recent examples of strategy B, see: a) H. Zhang, R. C. Larock, *J. Org. Chem.* **2002**, *67*, 7048–7056; b) M. Tian, P. Belmont, *J. Org. Chem.* **2008**, *73*, 4101–4109; c) S. P. Shukla, R. Tiwari, A. K. Verma, *Tetrahedron* **2012**, *68*, 9035–9044; d) X.-C. Wang, M.-J. Zhong, Y.-M. Liang, *Org. Biomol. Chem.* **2012**, *10*, 3636–3641; e) C. H. Oh, J. H. Kim, B. K. Oh, J. R. Park, J. H. Lee, *Chem. Eur. J.* **2013**, *19*, 2592–2596; f) S. Manojveer, R. Balamurugan, *Org. Lett.* **2014**, *16*, 1712–1715.
- [4] For relevant opinions on reaction efficiency, see: a) R. A. Sheldon, *Chem. Soc. Rev.* **2012**, *41*, 1437–1451; b) *Green Chemistry in the Pharmaceutical Industry* (Eds.: P. J. Dunn, A. S. Wells, M. T. Williams), Wiley-VCH, Weinheim, **2010**, pp. 370; c) P. A. Clarke, A. V. Zaytzev, A. C. Whitwood, *Tetrahedron Lett.* **2007**, *48*, 5209–5212.
- [5] For reviews on the synthesis of medium-ring carbocycles, see: a) L. Yet, *Tetrahedron* **1999**, *55*, 9349–9403 (via radicals); b) *Topics in Organometallic Chemistry—Alkene Metathesis in Organic Synthesis*, Vol. 1 (Ed.: A. Fürstner), Springer, New York, USA, **1998**, p. 231 (by metathesis); c) C. Galli, L. Mandolini, *Eur. J. Org. Chem.* **2000**, 3117–3125 (ring-strain effects); d) K. C. Majumdar, B. Chattopadhyay, *Curr. Org. Chem.* **2009**, *13*, 731–757 (by palladium catalysis).
- [6] For selected “modern era” Cannizzaro reactions, see: a) A. E. Russell, S. P. Miller, J. P. Morken, *J. Org. Chem.* **2000**, *65*, 8381–8383; b) M. S. Abaee, R. Sharifi, M. M. Mojtahedi, *Org. Lett.* **2005**, *7*, 5893–5895; c) K. Ishihara, T. Yano, M. Fushimi, *J. Fluorine Chem.* **2008**, *129*, 994–997; d) C. Wuensch, H. Lechner, S. M. Glueck, K. Zangger, M. Hall, K. Faber, *ChemCatChem* **2013**, *5*, 1744–1748; e) P. Wang, W.-J. Tao, X.-L. Sun, S. Liao, Y. Tang, *J. Am. Chem. Soc.* **2013**, *135*, 16849–16852; f) Q. Xu, J. Chen, H. Tian, X. Yuan, S. Li, C. Zhou, J. Liu, *Angew. Chem. Int. Ed.* **2014**, *53*, 225–229; *Angew. Chem.* **2014**, *126*, 229–233.
- [7] A. S. Travis, W. J. Hornix, R. Bud, P. J. T. Morris, *Br. J. Hist. Sci.* **2009**, *25*, 145–167, and references therein.
- [8] M. Murakami, S. Ashida, T. Matsuda, *J. Am. Chem. Soc.* **2006**, *128*, 2166–2167.
- [9] For a selection of representative recent examples (the mode of cyclization, the number of steps required to prepare the cyclization precursor from inexpensive starting materials, and the overall yield for the total synthesis of the eight-ring product (synthesis of the precursors and subsequent cyclization) are given in parentheses); see: a) M. Utsugi, Y. Kamada, M. Nakada, *Tetrahedron Lett.* **2008**, *49*, 4754–4757 (Pd-cat. Heck reaction, 20, 36%); b) Y. Oonishi, A. Hosotani, Y. Sato, *J. Am. Chem. Soc.* **2011**, *133*, 10386–10389 (Rh-cat. [6+2], 7, 21–23%); c) C. S. Penkett, J. A. Woolford, *Org. Lett.* **2012**, *14*, 5704–5707 (photochemistry, 1, <5%); d) Y. Oonishi, A. Hosotani, Y. Sato, *Angew. Chem. Int. Ed.* **2012**, *51*, 11548–11551; *Angew. Chem.* **2012**, *124*, 11716–11719 (Rh-cat. [6+2], 4–7, 8–25%); e) T. Iwai, H. Okochi, H. Ito, M. Sawamura, *Angew. Chem. Int. Ed.* **2013**, *52*, 4239–4242; *Angew. Chem.* **2013**, *125*, 4333–4336 (Au-cat. alkyne cyclization, 5–7 and 3 for the catalyst synthesis, 14–15%); f) C. B. Nielsen, T. Brock-Nannestad, P. Hammershøj, T. K. Reenberg, M. Schau-Magnussen, D. Trpevski, T. Hensel, R. Salcedo, G. V. Baryshnikov, B. F. Minaev, M. Pittelkow, *Chem. Eur. J.* **2013**, *19*, 3898–3904 (Friedel–Crafts, 5, 34–39%); g) T. Nishinaga, T. Ohmae, K. Aita, M. Takase, M. Iyoda, T. Araib, Y. Kunugi, *Chem. Commun.* **2013**, *49*, 5354–5356 (Cu-cat. oxidation, 3, 16%); h) K. Aita, T. Ohmae, M. Takase, K. Nomura, H. Kimura, T. Nishinaga, *Org. Lett.* **2013**, *15*, 3522–3525 (Ru-cat. metathesis, 7, <5%); i) Y. Wang, J. Song, L. Xu, Y. Kann, J. Shi, H. Wang, *J. Org. Chem.* **2014**, *79*, 2255–2262 (Cu-cat. oxidation, 3–5, 12–21%); j) Y. Wang, D. Gao, J. Shi, Y. Kan, J. Song, C. Li, H. Wang, *Tetrahedron* **2014**, *70*, 631–636 (Cu-cat. oxidation, 1, 44%); k) S. Hirai, M. Utsugi, M. Iwamoto, M. Nakada, *Chem. Eur. J.* **2015**, *21*, 355–359 (Pd-cat. alkenylation, >10, <40%); l) C. Hulot, S. Amiri, G. Blond, P. R. Schreiner, J. Suffert, *J. Am. Chem. Soc.* **2009**, *131*, 13387–13398 (Ni-cat. electrocyclization, 8, 28–57%); m) M. Charpenay, A. Boudhar, C. Hulot, G. Blond, J. Suffert, *Tetrahedron* **2013**, *69*, 7568–7591 (Ni-cat. electrocyclization, 6, 39–75%). According to this analysis (excluding **4iaa**), the performance of our own system is: Cannizzaro-triggered, 2–3, 16–62%, average: 42%, 19 examples with >40%. For the preparation of our own starting materials, costs were in the range of €0.3–8 mmol⁻¹. Compound **4iaa** is included to demonstrate the need to avoid rather long starting-material syntheses.
- [10] For these derivatives, the proton attached to C4 gives rise to a characteristic low-field resonance at approximately 5.0 ppm, presumably owing to its proximity to the aryl ring current. Compounds **4** are often highly crystalline, and in a number of cases, they could be isolated by a simple recrystallization of the crude reaction material on gram scale. The structures of **4haa** and **4jaa** were confirmed by X-ray crystallography (see the Supporting Information).
- [11] Y. Zhao, D. G. Truhlar, *Theor. Chem. Acc.* **2008**, *120*, 215–241; J. A. Montgomery, M. J. Frisch, J. W. Ochterski, G. A. Petersson, *J. Chem. Phys.* **1999**, *110*, 2822–2827; J. A. Montgomery, Jr., M. J. Frisch, J. W. Ochterski, G. A. Petersson, *J. Chem. Phys.* **2000**, *112*, 6532–6542.
- [12] M. Alajarin, B. Bonillo, M. Marin-Luna, P. Sanchez-Andrada, A. Vidal, *Chem. Eur. J.* **2013**, *19*, 16093–16103.
- [13] Y. Zhang, F. Yang, L. Zheng, Q. Dang, X. Bai, *Org. Lett.* **2014**, *16*, 6041–6043.
- [14] S. L. Schreiber, S. E. Kelly, *Tetrahedron Lett.* **1984**, *25*, 1757–1760. We thank a referee for this suggestion.

Received: June 11, 2015

Published online: July 29, 2015