



Delforce, Sarah J. and Wang, Yu and Van-Aalst, Meg E. and Corbisier de Meaultsart, Celine and Morris, Brian J. and Broughton Pipkin, Fiona and Roberts, Claire T. and Lumbers, Eugenie R. and Pringle, Kirsty G. (2016) Effect of oxygen on the expression of renin-angiotensin system components in a human trophoblast cell line. *Placenta*, 37 . pp. 1-6. ISSN 1532-3102

**Access from the University of Nottingham repository:**

<http://eprints.nottingham.ac.uk/47007/9/2015%20PL-15-100105R1%282%29%20final.pdf>

**Copyright and reuse:**

The Nottingham ePrints service makes this work by researchers of the University of Nottingham available open access under the following conditions.

This article is made available under the University of Nottingham End User licence and may be reused according to the conditions of the licence. For more details see: [http://eprints.nottingham.ac.uk/end\\_user\\_agreement.pdf](http://eprints.nottingham.ac.uk/end_user_agreement.pdf)

**A note on versions:**

The version presented here may differ from the published version or from the version of record. If you wish to cite this item you are advised to consult the publisher's version. Please see the repository url above for details on accessing the published version and note that access may require a subscription.

For more information, please contact [eprints@nottingham.ac.uk](mailto:eprints@nottingham.ac.uk)

Manuscript Number: PL-15-100105R1

Title: Effect of oxygen on the expression of renin-angiotensin system components in a human trophoblast cell line

Article Type: Full Length Article

Keywords: renin-angiotensin system; pregnancy; placenta; hypoxia; gene expression

Corresponding Author: Dr. Kirsty G Pringle, PhD

Corresponding Author's Institution: University of Newcastle

First Author: Sarah J Delforce

Order of Authors: Sarah J Delforce; Yu Wang; Meg E Van-Aalst; Celine Corbisier de Meaultsart; Brian J Morris; Fiona Broughton-Pipkin; Claire T Roberts; Eugenie R Lumbers; Kirsty G Pringle, PhD

Abstract: During the first trimester, normal placental development occurs in a low oxygen environment that is known to stimulate angiogenesis via upregulation of vascular endothelial growth factor (VEGF). Expression of the placental renin-angiotensin system (RAS) is highest in early pregnancy. While the RAS and oxygen both stimulate angiogenesis, how they interact within the placenta is unknown. We postulated that low oxygen increases expression of the proangiogenic RAS pathway and that this is associated with increased VEGF in a first trimester human trophoblast cell line (HTR-8/SVneo). HTR-8/SVneo cells were cultured in one of three oxygen tensions (1%, 5% and 20%). RAS and VEGF mRNA expression were determined by qPCR. Prorenin, angiotensin converting enzyme (ACE) and VEGF protein levels in the supernatant, as well as prorenin and ACE in cell lysates, were measured using ELISA. Low oxygen significantly increased the expression of both angiotensin II type 1 receptor (AGTR1) and VEGF (both  $P < 0.05$ ). There was a positive correlation between AGTR1 and VEGF expression at low oxygen ( $r = 0.64$ ,  $P < 0.005$ ). Corresponding increases in VEGF protein were observed with low oxygen ( $P < 0.05$ ). Despite no change in ACE1 mRNA expression, ACE levels in the supernatant increased with low oxygen (1% and 5%,  $P < 0.05$ ). Expression of other RAS components did not change. Low oxygen increased AGTR1 and VEGF expression, as well as ACE and VEGF protein levels, suggesting that the proangiogenic RAS pathway is activated. This highlights a potential role for the placental RAS in mediating the proangiogenic effects of low oxygen in placental development.

## RESPONSE TO REVIEWERS

### Manuscript # PL-15-100105

**Title:** ‘Effect of oxygen on the expression of renin-angiotensin system (RAS) components in a human trophoblast cell line’

**Authors:** Sarah J Delforce; Yu Wang; Meg E Van-Aalst; Celine Corbisier de Meaultsart; Brian J Morris; Fiona Broughton-Pipkin; Claire T Roberts; Eugenie R Lumbers; Kirsty G. Pringle

We would like to thank the Editor and reviewers for their generally favorable assessment of our manuscript entitled: ‘Effect of oxygen on the expression of renin-angiotensin system (RAS) components in a human trophoblast cell line’. We consider that we have now addressed all of the reviewers’ concerns. We have displayed our text corrections in red throughout the manuscript and described them in our response to reviewers (see below). The reviewers’ comments are stated first, in italics, followed by our response.

#### **Reviewer #1:**

*In this study the authors have shown that a low oxygen environment stimulates the expression of AGTR1 and VEGF mRNAs in HTR-8/SVneo cells in vitro. Both AT1R and VEGF are involved in angiogenesis and proliferation, which are key properties of first trimester EVT. We showed that incubation in a low oxygen environment increased production of ACE and VEGF protein by HTR-8/SVneo cells and increased expression of the Ang II/AT1R pathway. They proposed that the placental RAS is under the influence of many factors including endocrine, molecular and environmental ones.*

No response required.

*General comment: This is a well-written and designed article by experimented team. I have concerns about the reliability of the use of HTR-8/SVneo cells instead primary trophoblasts or other lines (i.e. BeWo, Jeg-3, JAr ...etc). How the authors justify the choice of this line and not another? This justification must be added to the section of discussion.*

AUTHORS’ RESPONSE: The following passage has now been added to the Materials and methods (page 2, lines 48-55) “HTR-8/SVneo trophoblast cells were chosen for the study as they are a transformed first trimester human extravillous trophoblast cell line. Because of this they are an ideal tool for investigating the effect of low oxygen on the proliferative/angiogenic RAS pathways in placental development and are superior to other cell lines such as choriocarcinoma BeWo cells. We have demonstrated previously that the pro-angiogenic/proliferative pathway of the RAS is expressed in the HTR-8/SVneo cell line [14] as occurs in the first trimester placenta *in vivo* [9], but which is not seen in BeWo cells [14].”

#### **Reviewer #2:**

*This study by Delforce and colleagues characterize the role of varying oxygen tension on the expression of the renin-angiotensin system in trophoblast cell line HTR-8/SVneo. The main findings of this paper are that the mRNA expression of AGTR1 and VEGF at low oxygen level and the increased VEGF secretion from cells. The methodological approach is sound, although first trimester primary trophoblast cells or explants would have been preferred. The overall conclusion that low oxygen regulates the RAS may be overstated given that only one gene from the RAS is altered (AGTR1) and this was not confirmed at the protein level. This is important since although REN mRNA expression was altered, there were no changes at the protein level. Changes in VEGF expression with oxygen tension are well established and therefore these findings are more confirmatory than novel insights.*

AUTHORS' RESPONSE: We agree that the conclusion that low oxygen regulates the RAS was overstated and have revised this section in the Discussion (page 6, lines 164-167): "The study also showed that even small reductions in O<sub>2</sub> have substantial effects on the expression of the AT<sub>1</sub>R and pro-angiogenic factors such as VEGF. This was based on our finding that culture in 1% oxygen had a much more powerful effect than culture in 5% oxygen"

*Minor comments:*

*Cell culture: Please provide the passage number(s) of the cell line if possible.*

AUTHORS' RESPONSE: Cells were between passages 10-20. This information has now been added to the Materials and methods section of the manuscript (page 2, lines 58-59).

*Line 53-54: RNA quality measurements are not appropriate indications of cell viability. Trypan blue exclusion, MTT, apoptosis assays may be required.*

AUTHORS' RESPONSE: We have rephrased the text to clarify that RNA quantity, not quality, was used as an indicator of cell viability, page 3 lines 66-69: "RNA quantity was used as an indicator of cell viability and was assessed using the Nanodrop spectrophotometer, no differences in RNA quantity were detected between the treatment groups (data not shown). RNA quality was determined by agarose gel electrophoresis."

*Line 73: Please check if this is supposed to be "one microliter" or "one milliliter"*

AUTHORS' RESPONSE: We thank the reviewer for noting this error. This sentence was supposed to say "one millilitre" and has now been corrected (page 3, line 89).

*Please provide details for regression analysis. Also please provide the R<sup>2</sup> value.*

AUTHORS' RESPONSE: The following passage has now been added to the Materials and methods (page 4, lines 115-116): "Pearson correlation coefficients were calculated to determine any relationships between the mRNA expression of genes." R<sup>2</sup> and r values are now present in the figure legends and have been added to the text (page 5, lines 137-138).

**Reviewer #3:** *In this manuscript, the authors describe in vitro experiments demonstrating that low oxygen influence RAS and VEGF in HTR-8/SVneo cells. The authors have shown by different techniques that culture in low oxygen tension (1%) increased AGTR1 and VEGF mRNAs as well as ACE and VEGF proteins expression in trophoblast cells. Data are interesting and add knowledge to the area, however some points should be elucidate before manuscript publication.*

AUTHORS' RESPONSE: No response required.

*Specific comments:*

*1- How did the authors define oxygen tension (1%, 5% or 20%) for culture? As the effects on gene and protein expression were observed with very low oxygen tension (1%) it is important to clarify whether this oxygen tension reflects the physiological environment of early gestation.*

AUTHORS' RESPONSE: We chose 1% and 5% oxygen tensions to closely reflect the environment experienced by the early gestation trophoblasts within the placental villi as well as the extravillous trophoblasts invading into the endometrium. We have now included a section in the introduction outlining the physiological environment in the first trimester (page 1, lines 7-12): "Rodesch *et al.* [2] established that the oxygen tension in the intervillous space at 8 weeks to be 17.9 mmHg (~2.5%) with a range of 5–30 mmHg (~0.7– 4.3%) while the oxygen tension in the endometrium is higher at 39.6 mmHg (~5.7%) with a range of 25–70 mmHg (~3.5–10%). Thus there is an oxygen gradient experienced by the first trimester trophoblasts that can range from ~1% (in the placental villi) to up to 10% in endometrium."

*2- Material and Methods- Are VEGF primers also described by Marques et al. (2011)? Did the authors investigate VEGF-A? There are multiple isoforms of VEGF-A. What was the VEGF isoform(s) investigated?*

AUTHORS' RESPONSE: The VEGF primers described target VEGF-A and will detect all transcript variants of VEGF-A. VEGF primers were described in Pringle *et al.* (2011); this reference has been added to the text (page 3, line 81).

*3- Material and Methods - Lines 53-54 - There was influence of oxygen tension in cell viability?*

AUTHORS' RESPONSE: No influence of oxygen tension on cell viability was observed. This information has now been added to the Materials and methods (page 3, 66-69). Also see response to reviewer 1, above.

*4- Material and Methods- Did the authors measure protein content (VEGF, prorenin and ACE) at 24hs of culture?*

AUTHORS' RESPONSE: We have measured VEGF protein, as well as prorenin and ACE protein at 24 hours of culture. No effect of oxygen tension on the levels of these proteins was, however, observed. The Materials and methods section has been modified to reflect this (lines 98-99) and these results have now been included in the manuscript:

- Page 5, lines 142-144: "Although measurable levels of *REN* mRNA were found (Fig. 1), prorenin protein was low in both the culture medium and cell lysate of HTR-8/SVneo cells at 24 and 48 h incubation irrespective of the prevailing O<sub>2</sub> (only 48 h data shown; Fig. 3)."
- Page 5, lines 147-148: "No effect of oxygen tension on VEGF protein was observed at 24 h incubation (data not shown)."
- Page 5 lines 153-155: "No effect of oxygen tension on ACE protein in culture medium or cell lysates was observed at 24 h incubation (data not shown)."

*5- Did the authors expect to identify ACE2, AGTR2 and MAS1 gene expression in HTR-8/SVneo cells. How to explain the observed data? Could this lack of expression alter expression of other RAS members in HTR-8/SVneo cells.*

AUTHORS' RESPONSE: We did not expect to identify *ACE2*, *AGTR2* or *MAS1* gene expression in HTR-8/SVneo cells as we have previously demonstrated that this cell line does not express these mRNAs (Wang et al. 2012). A statement outlining the similarities between these 2 studies has been included in the Results (page 4, lines 126 – 127).

*6- Results - Lines 117-118 -What is the physiological meaning of the correlations? What is the molecular correlation between these genes. Manuscript could be improved with more discussion about a hypothesized mechanisms related to the data.*

AUTHORS' RESPONSE: The following passage has now been added to the Discussion (page 7, lines 203-209) "We found that *AGT1R* mRNA abundance was highly correlated with levels of both *VEGF* and *ATP6AP2* mRNA. The association between *AGT1R* and *ATP6AP2* mRNA suggests that increased levels of the (pro)renin receptor (*ATP6AP2*) may stimulate the activation of the placental RAS cascade and in turn increase AT<sub>1</sub>R, while the association between *AGT1R* and *VEGF* mRNA suggests that increased AT<sub>1</sub>R in first trimester placenta may increase expression of VEGF mRNA and protein, and thereby stimulate angiogenesis within the developing placenta."

*7- Result - Lines 123 - How much prorenin was expected in the cell cultures? Why the values are low?*

AUTHORS' RESPONSE: The values observed in the study were expected and were similar to those obtained in previous studies in which increases in prorenin protein were only observed when HTR-8/SVneo cells were stimulated with cAMP (see Wang *et al.* 2012). This was addressed in the Discussion (page 6, lines 168-176) where we suggest that the expression of prorenin (mRNA and protein) may be influenced by factors other than low oxygen including hormonal activation and/or activation by cAMP: "A low oxygen environment was not associated with increased expression of prorenin mRNA or

its protein. This suggests that a low oxygen tension alone is not responsible for the high levels of *REN* mRNA found in first trimester placentae [10]. It is more likely that increased *REN* mRNA expression in the placenta early in pregnancy is the result of hormonal activation and/or activation by cAMP of *REN* mRNA expression. Human chorionic gonadotropin (hCG) has been recognized as a stimulus for prorenin production by villous placenta [19], as have  $\beta$ -adrenoceptor agonists [19]. We have also shown previously that cAMP increases *REN* mRNA expression and prorenin protein secretion by HTR-8/SVneo cells [14].”

*8- Discussion - Please check this statement in lines 141-145.*

AUTHORS' RESPONSE: We have adjusted the text because we agree that the effect of 1% O<sub>2</sub> was only seen for *AGT1R* mRNA and not the whole RAS cascade. Please also see our response to Reviewer 1, above.

**Conflict of Interest**

The authors have no competing interests to declare.



## **Effect of oxygen on the expression of renin-angiotensin system components in a human trophoblast cell line**

Sarah J. Delforce<sup>a\*</sup>, Yu Wang<sup>a\*</sup>, Meg E. Van-Aalst<sup>a</sup>, Celine Corbisier de Meaultsart<sup>a</sup>, Brian. J. Morris<sup>b</sup>, Fiona Broughton-Pipkin<sup>c</sup>, Claire T. Roberts<sup>d</sup>, Eugenie R. Lumbers<sup>a</sup>, Kirsty G. Pringle<sup>a†</sup>

<sup>a</sup>School of Biomedical Sciences and Pharmacy, Mothers and Babies Research Centre, Hunter Medical Research Institute, University of Newcastle, Newcastle, New South Wales, Australia

<sup>b</sup>School of Medical Sciences and Bosch Institute, University of Sydney, Sydney, New South Wales, Australia

<sup>c</sup>Department of Obstetrics & Gynaecology, School of Medicine, University of Nottingham, NG5 1PB, UK

<sup>d</sup>Discipline of Obstetrics and Gynaecology, School of Paediatrics and Reproductive Health, Robinson Research Institute, University of Adelaide, Adelaide, South Australia, Australia.

\*These authors contributed equally to this work.

†Corresponding author. Mothers and Babies Research Centre, Hunter Medical Research Institute, Level 3 East, 1 Kookaburra Circuit, New Lambton Heights, NSW 2305, Australia. Tel: +61-2-4985-5643; fax: +61-2-9421-4394

*E-mail address:* [kirsty.pringle@newcastle.edu.au](mailto:kirsty.pringle@newcastle.edu.au)

## ABSTRACT

During the first trimester, normal placental development occurs in a low oxygen environment that is known to stimulate angiogenesis via upregulation of vascular endothelial growth factor (VEGF). Expression of the placental renin-angiotensin system (RAS) is highest in early pregnancy. While the RAS and oxygen both stimulate angiogenesis, how they interact within the placenta is unknown. We postulated that low oxygen increases expression of the proangiogenic RAS pathway and **that** this is associated with increased VEGF in a first trimester human trophoblast cell line (HTR-8/SVneo). HTR-8/SVneo cells were cultured in one of three oxygen tensions (1%, 5% and 20%). RAS and *VEGF* mRNA expression were determined by qPCR. Prorenin, angiotensin converting enzyme (ACE) and VEGF protein levels in the supernatant, as well as prorenin and ACE in cell lysates, were measured using ELISA. Low oxygen significantly increased the expression of both angiotensin II type 1 receptor (*AGTR1*) and *VEGF* (both  $P < 0.05$ ). There was a positive correlation between *AGTR1* and *VEGF* expression at low oxygen ( $r = 0.64$ ,  $P < 0.005$ ). Corresponding increases in VEGF protein were observed with low oxygen ( $P < 0.05$ ). Despite no change in *ACE1* mRNA expression, ACE levels in the supernatant increased with low oxygen (1% and 5%,  $P < 0.05$ ). Expression of other RAS components did not change. Low oxygen increased *AGTR1* and *VEGF* expression, as well as ACE and VEGF protein levels, suggesting that the proangiogenic RAS pathway is activated. This highlights a potential role for the placental RAS in mediating the proangiogenic effects of low oxygen in placental development.

**Keywords:** renin-angiotensin system; pregnancy; placenta; hypoxia; gene expression

## Abbreviations

ACE, angiotensin-converting enzyme; **ACE1, angiotensin-converting enzyme 1**; ACE2, angiotensin-converting enzyme 2; AGT, angiotensinogen; **AT<sub>1</sub>R, angiotensin II type 1 receptor**; **AGTR1 mRNA**, angiotensin II type 1 receptor mRNA; Ang, angiotensin; ATP6AP2, **ATPase, H<sup>+</sup> transporting, lysosomal accessory protein 2** / (pro)renin receptor; BCA, bicinchoninic acid; EVT, extravillous trophoblast; HIFs, hypoxia inducible factors; iNOS, inducible nitric oxide synthase; RAS, renin-angiotensin system; **REN mRNA**; (pro)renin **mRNA**; VEGF, vascular endothelial growth factor.

## Highlights

- The human placenta develops in a low oxygen environment
- The placental renin-angiotensin system (**RAS**) is highly expressed in early pregnancy
- Culture in 1% O<sub>2</sub> increased **AGTRI** mRNA and ACE protein in HTR-8/SVneo cells
- **The latter** was associated with increased **VEGF** mRNA and VEGF protein
- Stimulation of the placental RAS by low oxygen may be proliferative and angiogenic

## 1 Introduction

2 A low oxygen environment during the first trimester of pregnancy is required for  
3 optimal placental development that is essential for supplying fetal demand in late  
4 gestation. This early gestation low oxygen environment occurs as a result of extravillous  
5 trophoblast (EVT) cell proliferation and subsequent invasion of the decidua and its  
6 vasculature, initially occluding maternal spiral arterioles from about two weeks after  
7 implantation [1]. Rodesch *et al.* [2] established that the oxygen tension in the  
8 intervillous space at 8 weeks is 17.9 mmHg (~2.5%) with a range of 5–30 mmHg  
9 (~0.7–4.3%) while the oxygen tension in the endometrium is higher at 39.6 mmHg  
10 (~5.7%) with a range of 25–70 mmHg (~3.5–10%). Thus there is an oxygen gradient  
11 experienced by first trimester trophoblasts that can range from ~1% (in the placental  
12 villi) to up to 10% in the decidua. The low oxygen environment stimulates angiogenesis  
13 and vascularization of the placenta. Poor placental development, characterized by  
14 insufficient decidual invasion by EVTs, incomplete occlusion of maternal arterioles  
15 with inadequate remodeling of maternal spiral arterioles, and early onset of maternal  
16 blood flow to the conceptus, ultimately results in poor nutrient and oxygen exchange  
17 during the 2<sup>nd</sup> and 3<sup>rd</sup> trimesters. These are associated with intrauterine growth  
18 restriction and preeclampsia [3, 4].

19 A low oxygen tension stabilizes hypoxia inducible factors (HIFs) which promote  
20 angiogenesis and vascularization by activating pro-angiogenic factors such as vascular  
21 endothelial growth factor (VEGF), angiopoietins, factors involved in regulation of  
22 vascular tone such as inducible nitric oxide synthase (iNOS) and proteins involved in  
23 nutrient transfer such as transferrin and glycolytic enzymes [5-8].

24 Another system that might regulate placental angiogenesis is the renin-angiotensin  
25 system (RAS). Tissue RASs have been shown to be involved in the regulation of  
26 angiogenesis, as well as cell proliferation and apoptosis [9]. We have shown that mRNA  
27 expression of prorenin (*REN mRNA*), (pro)renin receptor (*ATP6AP2 mRNA*),  
28 angiotensinogen (*AGT mRNA*), angiotensin (Ang) II type 1 (*AT<sub>1</sub>R*) (*AGTR1 mRNA*)  
29 and Ang converting enzyme 2 (*ACE2*) (*ACE2 mRNA*) are all very high in early  
30 gestation placentae compared with term [10]. In addition, we have shown that the  
31 mRNA expression of placental *VEGF* is correlated with those of *REN*, *ATP6AP2* and  
32 *AGTR1* mRNAs [10]. Thus the placental RAS is most active during the first trimester

33 and therefore could stimulate angiogenesis, as it does in other tissues. The ocular RAS  
34 is stimulated by ischemia. The increased activity of the ocular RAS is associated with a  
35 potent angiogenic response mediated via the Ang II/AT<sub>1</sub>R pathway [11]. In addition,  
36 early renal development requires activation of the RAS by a low oxygen milieu [12].

37 While a low oxygen environment regulates placental development [13], the extent to  
38 which the RAS is essential for normal placental development has not been established.  
39 To investigate interactions between a low oxygen milieu and the placental RAS, we  
40 examined the effects of low oxygen on the expression of the RAS and VEGF in a first  
41 trimester human trophoblast cell line, HTR-8/SVneo, which we have previously shown  
42 expresses mRNAs encoding those RAS pathway components that stimulate  
43 angiogenesis in the eye and kidney [14, 15].

## 45 **Materials and methods**

### 46 *Cell culture*

47 HTR-8/SVneo cells are an immortalized first trimester trophoblast cell line (a kind  
48 gift from Prof. Charles Graham, Queens University, Ontario). HTR-8/SVneo  
49 trophoblast cells were chosen for the study as they are a transformed first trimester  
50 human extravillous trophoblast cell line. Because of this they are an ideal tool for  
51 investigating the effect of low oxygen on the proliferative/angiogenic RAS pathways in  
52 placental development and are superior to other cell lines such as choriocarcinoma  
53 BeWo cells. We have demonstrated previously that the pro-angiogenic/proliferative  
54 pathway of the RAS is expressed in the HTR-8/SVneo cell line [14] as occurs in the  
55 first trimester placenta *in vivo* [9], but which is not seen in BeWo cells [14].

56 HTR-8/SVneo cells were cultured in RPMI-1640 medium (HyClone), supplemented  
57 with 10% fetal bovine serum (SAFC Biosciences), 1 mg/ml antibiotic-antimycotic  
58 (Gibco) and 1% L-glutamine in 5% CO<sub>2</sub> in room air at 37°C (cells were between  
59 passages 10–20). Cells were seeded at a density of 200,000 or 400,000 cells per well for  
60 24 h or 48 h incubation, respectively. They were seeded in 6 well plates with 2 ml of  
61 incubation medium per well and allowed to settle for 24 h, after which time the medium  
62 was changed. Cells were then transferred to sealed oxygen chambers containing either  
63 1%, 5% or 20% O<sub>2</sub> and 5% CO<sub>2</sub> in N<sub>2</sub> and cultured for 24 or 48 h, with chambers  
64 flushed every 24 h. Cells were harvested and the incubation medium was collected at 24

65 and 48 h, then snap frozen in liquid nitrogen at  $-80^{\circ}\text{C}$  for subsequent protein and  
66 mRNA analyses. Three experiments were conducted in triplicate. RNA quantity was  
67 used as an indicator of cell viability and was assessed using the Nanodrop  
68 spectrophotometer, no differences in RNA quantity were detected between the treatment  
69 groups (data not shown). RNA quality was determined by agarose gel electrophoresis.

70

#### 71 *Semi-quantitative real-time reverse transcriptase polymerase chain reaction (qPCR)*

72 Total RNA was isolated using the RNeasy mini kit according to the manufacturer's  
73 instructions (Qiagen). In addition, we examined the integrity of the total RNA in each  
74 sample using gel electrophoresis. RNA samples were DNase treated (Qiagen) and total  
75 RNA was spiked with a known amount of Alien RNA (Stratagene),  $10^7$  copies per  $\mu\text{g}$  of  
76 total RNA, before the RNA was reverse transcribed using a Superscript III RT kit with  
77 random hexamers (Invitrogen). The Alien qRT PCR inhibitor alert system serves as a  
78 reference for internal standardization [16]. qPCR was performed in an Applied  
79 Biosystems 7500 Real Time PCR System using SYBR Green for detection. Each  
80 reaction contained 5  $\mu\text{l}$  of SYBR Green PCR master mix (Applied Biosystems), RAS  
81 and VEGF primers that we have described previously [10, 17], cDNA reversed  
82 transcribed from 10 ng total RNA, and water to 10  $\mu\text{l}$ . Messenger RNA abundance was  
83 calculated as described previously, using the  $2^{-\Delta\Delta\text{CT}}$  method and expressed relative to  
84 Alien mRNA and a calibrator sample (a term placental sample collected at elective  
85 Cesarean section) [10].

86

#### 87 *Extraction and quantification of total protein from cells*

88 Protein was extracted from cells using a radioimmunoprecipitation assay (RIPA)  
89 lysis and extraction buffer. One milliliter of RIPA buffer (50 mM Tris-HCl, 150 mM  
90 NaCl, 1 mM EDTA, 1% NP-40, 0.5% sodium deoxycholate, 100 nM sodium  
91 orthovanadate and Complete Mini Protease Inhibitor Cocktail tablets (Roche  
92 Diagnostics Australia) and 10  $\mu\text{L}$  of 100 nM PMSF were added to each sample.  
93 Samples were incubated on ice for 30 min then centrifuged at 13,000 rpm at  $4^{\circ}\text{C}$  for 10  
94 min. Supernatants were collected. Protein was quantified using the Pierce BCA Protein  
95 assay kit (Life Technologies) according to the manufacturer's instructions.

96

### 97 *Measurement of prorenin, VEGF and ACE by ELISA*

98 Prorenin, VEGF and ACE concentrations in culture medium and cell lysates (at both  
99 24 and 48 h) were measured using the Human Prorenin ELISA kit (Molecular  
100 Innovations), Human VEGF DuoSet ELISA kit (R&D systems) and the Human ACE  
101 DuoSet ELISA kit (R&D Systems), respectively, according to the manufacturers'  
102 instructions, as described previously [18]. For prorenin, VEGF and ACE proteins in  
103 culture medium and prorenin in cell lysates, all samples were assayed in duplicate on  
104 one ELISA plate. Therefore there was no inter-assay variability. For ACE protein in cell  
105 lysates, samples were assayed in duplicate over two plates and inter-assay variability  
106 was 9.4%. Intra-assay coefficients of variation were 13.1% and 3.2% for prorenin  
107 culture medium and cell lysate, 6.8% for VEGF culture medium and 14.2% and 2.9%  
108 for ACE culture medium and cell lysate, respectively.

109

### 110 *Data analysis*

111 Two-way ANOVA with Tukey's multiple comparisons test was used to determine  
112 the effects of differing oxygen tensions on the relative abundance of RAS mRNAs,  
113 prorenin, ACE and VEGF protein levels. A Kruskal-Wallis non-parametric test was  
114 used to determine the effects of differing oxygen tensions on prorenin concentration in  
115 both the culture medium and cell lysate at 48 h. Pearson correlation coefficients were  
116 calculated to determine any relationships between the various mRNA levels. GraphPad  
117 Prism (Prism version 6.0) was used for all graphs and correlation analyses and the SPSS  
118 statistical package (SPSS for Windows, Release 17.0.0. Chicago) was used for all other  
119 analyses. Significance was set at  $P < 0.05$ .

120

## 121 **Results**

### 122 *Effect of O<sub>2</sub> on RAS and VEGF mRNA expression*

123 At all oxygen tensions, HTR-8/SVneo cells expressed genes of the pro-angiogenic  
124 pathway of the RAS, i.e., *AGT*, *REN*, *ATP6AP2*, *ACE1* and *AGTR1*, as well as *VEGF*,  
125 but expression of genes of the anti-angiogenic RAS pathway, i.e., *ACE2*, *AGT2R*, *MAS1*  
126 was not detected (Figure 1); this is consistent with previously published data on the  
127 expression of RAS genes in this cell line [14].

128 Expression of *AGT*, *REN*, *ATP6AP2* and *ACE1* mRNAs was not affected by the  
129 prevailing oxygen tension, but *AGTR1* and *VEGF* mRNA expression were significantly  
130 higher in HTR-8/SVneo cells cultured in low oxygen (1% O<sub>2</sub>) compared with 5% or  
131 20% O<sub>2</sub> at 48 h (all  $P < 0.01$ ). A similar trend was seen in 24 h incubates. This did not,  
132 however, reach statistical significance (Fig. 1).

133 There was a significant effect of incubation time on the expression of *AGT*, *REN*,  
134 *ATP6AP2*, *ACE1* and *AGTR1* mRNAs, in that expression of these mRNAs was  
135 significantly higher at 48 h compared with 24 h (all  $P < 0.05$ ; Fig. 1).

136 A significant correlation was found between *AGTR1* and *VEGF* mRNA and between  
137 *AGTR1* and *ATP6AP2* mRNA ( $r = 0.6378$  and  $0.6500$  respectively,  $R^2 = 0.4067$  and  
138  $0.4733$  respectively, both  $P < 0.01$ ; Fig. 2).

139

140 *Effect of O<sub>2</sub> on the concentration on prorenin and VEGF protein concentration in the*  
141 *culture medium*

142 Although measurable levels of *REN* mRNA were found (Fig. 1), prorenin protein  
143 was low in both the culture medium and cell lysate of HTR-8/SVneo cells at 24 and 48  
144 h incubation irrespective of the prevailing O<sub>2</sub> (only 48 h data shown; Fig. 3).

145 After 48 h VEGF levels in the culture medium of HTR-8/SVneo cells grown in 1%  
146 O<sub>2</sub> were much higher than levels in medium from samples cultured in 5% and 20% O<sub>2</sub>  
147 ( $P < 0.05$  and  $P < 0.01$ , respectively; Fig. 4). No effect of oxygen tension on VEGF  
148 protein was observed at 24 h incubation (data not shown).

149

150 *Effect of O<sub>2</sub> on the concentration of ACE in the culture medium*

151 After 48 h culture, HTR-8/SVneo cells cultured in 1% and 5% O<sub>2</sub> had significantly  
152 higher levels of ACE protein in the supernatant than cells cultured in 20% O<sub>2</sub> ( $P < 0.05$ ;  
153 Fig. 5). ACE levels in the cell lysate were not affected by O<sub>2</sub>. No effect of oxygen  
154 tension on ACE protein in culture medium or cell lysates was observed at 24 h  
155 incubation (data not shown).

156

## 157 Discussion

158 Incubation of primary HTR-8/SVneo trophoblast cells at a low (1%) oxygen tension  
159 increased the expression of *AGTR1* and *VEGF* mRNA and stimulated the production of



160 ACE and VEGF proteins. We propose, therefore, that a low oxygen milieu activates the  
161 pro-angiogenic pathway of the RAS by increasing AT<sub>1</sub>R and ACE protein levels and  
162 that these changes are associated with increases in VEGF, an important angiogenic  
163 factor.

164 ~~The study also showed that even small reductions in O<sub>2</sub> have substantial effects on~~  
165 ~~the expression of the AT<sub>1</sub>R and pro-angiogenic factors such as VEGF. This was based~~  
166 ~~on our finding that culture in 1% oxygen had a much more powerful effect than culture~~  
167 ~~in 5% oxygen. Thus, increased AGTR1 and VEGF mRNA expression and VEGF protein~~  
168 ~~only occurred at 1% O<sub>2</sub>.~~ A low oxygen environment was not associated with increased  
169 expression of prorenin mRNA or its protein. This suggests that a low oxygen tension  
170 alone is not responsible for the high levels of *REN* mRNA found in first trimester  
171 placentae [10]. It is more likely that increased *REN* mRNA expression in the placenta  
172 early in pregnancy is the result of hormonal activation and/or activation by cAMP of  
173 *REN* mRNA expression. Human chorionic gonadotropin (hCG) has been recognized as  
174 a stimulus for prorenin production by villous placenta [19], as have β-adrenoceptor  
175 agonists [19]. We have shown ~~previously~~ that cAMP increases *REN* mRNA expression  
176 and prorenin protein secretion by HTR-8/SVneo cells [14]. Expression of *AGT* and  
177 *ATP6AP2* mRNAs were not affected by oxygen. This suggests that their high  
178 expression in early gestation placentae [10] is not due to the low oxygen milieu.

179 Although the expression of *ACE1* was not stimulated by culturing cells in a low  
180 oxygen environment, we did observe increased ACE protein in medium from  
181 incubations carried out in low oxygen. Goyal *et al.* have observed a similar effect [20].  
182 In animal studies they showed that maternal hypoxia was associated with upregulation  
183 of placental ACE protein without a corresponding increase in *ACE1* mRNA levels [20].  
184 They concluded that this was because hypoxia-regulated miR-27, which regulates ACE  
185 translation, was down-regulated [20]. It is possible that a hypoxia-regulated miRNA in  
186 the HTR-8/SVneo cell cultures could be similarly affected by oxygen.

187 In a previous publication we reviewed evidence suggesting that the low oxygen  
188 milieu of early placental development activated the placental RAS, stimulating  
189 angiogenesis and cell proliferation [15]. The RAS pathway most likely to be involved in  
190 inducing these effects is the prorenin/AGT/ACE/Ang II/AT<sub>1</sub>R pathway because the Ang  
191 II/AT<sub>1</sub>R blocking drug, candesartan, blocks Ang II-induced endothelial proliferation

192 [21]. Hypoxia inducible factor (HIF-1 $\alpha$ ) is essential for normal trophoblast  
193 differentiation, since both HIF-1 $\alpha$  and AT<sub>1</sub>R gene ablation in mice impairs placental  
194 vascularization [22, 23]. Ang II acting via the AT<sub>1</sub>R causes cell proliferation [24].

195 We were surprised that a low oxygen milieu had no effect on prorenin expression  
196 and secretion. There may, however, be a number of factors acting on the placental  
197 renin/AGT/ACE/Ang II/AT<sub>1</sub>R pathway **that** upregulate different components, so  
198 ensuring increased placental Ang II production. As mentioned above, hCG and cAMP  
199 are potent stimuli for placental prorenin release. **cAMP also stimulates AGT mRNA**  
200 **expression [25].** MicroRNAs (miRNAs) may affect the stability or translation of *ACE1*  
201 mRNA as **has been observed** in animal experiments [20]. **AGTR1 mRNA** expression in  
202 HTR8/SVneo cells is, moreover, increased by low oxygen.

203 **We found that AGTR1 mRNA abundance was highly correlated with levels of both**  
204 **VEGF and ATP6AP2 mRNA. The association between AGTR1 and ATP6AP2 mRNA**  
205 **suggests that increased levels of the (pro)renin receptor (ATP6AP2) may stimulate the**  
206 **activation of the placental RAS cascade and in turn increase AT<sub>1</sub>R, while the**  
207 **association between AGTR1 and VEGF mRNA suggests that increased AT<sub>1</sub>R in first**  
208 **trimester placenta may increase expression of VEGF mRNA and protein, and thereby**  
209 **stimulate angiogenesis within the developing placenta.**

210 In conclusion, we have shown that a low oxygen environment stimulates the  
211 expression of *AGTR1* and *VEGF* mRNAs in HTR-8/SVneo cells *in vitro*. Both AT<sub>1</sub>R  
212 and VEGF are involved in angiogenesis and proliferation, which are key properties of  
213 first trimester EVT. We showed that incubation in a low oxygen environment increased  
214 production of ACE and VEGF protein by HTR-8/SVneo cells and increased expression  
215 of the Ang II/AT<sub>1</sub>R pathway. We propose that the placental RAS is under the influence  
216 of a number of endocrine, molecular and environmental **factors**.

217

### 218 **Author Contributions**

219 E.R. Lumbers and K.G. Pringle were responsible for designing the experiments.

220 C.T. Roberts, B.J. Morris and F. Broughton-Pipkin assisted in developing the  
221 experimental concepts that were the basis of the experiments.

222 Y. Wang, S.J. Delforce, M.E. Van-Aalst and C. Corbisier de Meaultsart carried out the  
223 experiments. Wang and Delforce analysed the data.

224 Delforce, Wang, Lumbers, Morris and Pringle interpreted the data and drafted the  
225 manuscript.

226

227 **Funding**

228 This research was funded by a National Health and Medical Research Council of  
229 Australia (NHMRC) project grant GNT1043537 awarded to ERL, CTR, BJM and FBP.  
230 CTR is supported by a NHMRC Senior Research Fellowship, GNT1020749.

231 **References**

- 232 [1] Jaffe R, Jauniaux E, Hustin J. Maternal circulation in the first-trimester human  
233 placenta – myth or reality? *Am J Obstet Gynecol.* 1997;176:695-705.
- 234 [2] Rodesch F, Simon P, Donner C, Jauniaux E. Oxygen measurements in  
235 endometrial and trophoblastic tissues during early pregnancy. *Obstet Gynecol.*  
236 1992;80:283-5.
- 237 [3] Salafia CM, Charles AK, Maas EM. Placenta and fetal growth restriction. *Clin*  
238 *Obstet Gynecol.* 2006;49:236-56.
- 239 [4] Silasi M, Cohen B, Karumanchi SA, Rana S. Abnormal placentation, angiogenic  
240 factors, and the pathogenesis of preeclampsia. *Obstet Gynecol Clin North Am.*  
241 2010;37:239-53.
- 242 [5] O'Donnell JL, Joyce MR, Shannon AM, Harmey J, Geraghty J, Bouchier-Hayes  
243 D. Oncological implications of hypoxia inducible factor-1alpha (HIF-1alpha)  
244 expression. *Cancer Treat Rev.* 2006;32:407-16.
- 245 [6] Pugh CW, Ratcliffe PJ. Regulation of angiogenesis by hypoxia: role of the HIF  
246 system. *Nat Med.* 2003;9:677-84.
- 247 [7] Schaffer L, Vogel J, Breymann C, Gassmann M, Marti HH. Preserved placental  
248 oxygenation and development during severe systemic hypoxia. *Am J Physiol Regul*  
249 *Integr Comp Physiol.* 2006;290:R844-51.
- 250 [8] Wang GL, Jiang BH, Rue EA, Semenza GL. Hypoxia-inducible factor 1 is a  
251 basic-helix-loop-helix-PAS heterodimer regulated by cellular O<sub>2</sub> tension. *Proc Natl*  
252 *Acad Sci USA.* 1995;92:5510-4.
- 253 [9] Paul M, Poyan Mehr A, Kreutz R. Physiology of local renin-angiotensin  
254 systems. *Physiol Rev.* 2006;86:747-803.
- 255 [10] Pringle KG, Tadros MA, Callister RJ, Lumbers ER. The expression and  
256 localization of the human placental prorenin/renin-angiotensin system throughout  
257 pregnancy: roles in trophoblast invasion and angiogenesis? *Placenta.* 2011;32:956-62.
- 258 [11] Wilkinson-Berka JL. Diabetes and retinal vascular disorders: role of the renin-  
259 angiotensin system. *Expert Rev Mol Med.* 2004;6:1-18.
- 260 [12] Tufro-McReddie A, Norwood VF, Aylor KW, Botkin SJ, Carey RM, Gomez  
261 RA. Oxygen regulates vascular endothelial growth factor-mediated vasculogenesis and  
262 tubulogenesis. *Dev Biol.* 1997;183:139-49.
- 263 [13] Genbacev O, Zhou Y, Ludlow JW, Fisher SJ. Regulation of human placental  
264 development by oxygen tension. *Science.* 1997;277:1669-72.
- 265 [14] Wang Y, Pringle KG, Chen YX, Zakar T, Lumbers ER. Regulation of the renin-  
266 angiotensin system (RAS) in BeWo and HTR-8/SVneo trophoblast cell lines. *Placenta.*  
267 2012;33:634-9.
- 268 [15] Lumbers ER, Pringle KG, Wang Y, Gibson KJ. The renin-angiotensin system  
269 from conception to old age: the good, the bad and the ugly. *Clin Exp Pharmacol*  
270 *Physiol.* 2013;40:743-52.
- 271 [16] Gilsbach R, Kouta M, Bonisch H, Bruss M. Comparison of in vitro and in vivo  
272 reference genes for internal standardization of real-time PCR data. *Biotechniques.*  
273 2006;40:173-7.
- 274 [17] Marques FZ, Pringle KG, Conquest A, Hirst JJ, Markus MA, Sarris M, et al.  
275 Molecular characterization of renin-angiotensin system components in human  
276 intrauterine tissues and fetal membranes from vaginal delivery and cesarean section.  
277 *Placenta.* 2011;32:214-21.

- 278 [18] Wang Y, Pringle KG, Sykes SD, Marques FZ, Morris BJ, Zakar T, et al. Fetal  
279 sex affects expression of renin-angiotensin system components in term human decidua.  
280 *Endocrinology*. 2011;153:462-8.
- 281 [19] Poisner AM. Regulation of utero-placental prorenin. *Adv Exp Med Biol*.  
282 1995;377:411-26.
- 283 [20] Goyal R, Lister R, Leitzke A, Goyal D, Gheorghe CP, Longo LD. Antenatal  
284 maternal hypoxic stress: adaptations of the placental renin-angiotensin system in the  
285 mouse. *Placenta*. 2011;32:134-9.
- 286 [21] Herr D, Rodewald M, Fraser H, Hack G, Konrad R, Kreienberg R, et al.  
287 Regulation of endothelial proliferation by the renin-angiotensin system in human  
288 umbilical vein endothelial cells. *Reproduction*. 2008;136:125-30.
- 289 [22] Cowden Dahl KD, Fryer BH, Mack FA, Compernelle V, Maltepe E, Adelman  
290 DM, et al. Hypoxia-inducible factors 1 $\alpha$  and 2 $\alpha$  regulate trophoblast differentiation. *Mol*  
291 *Cell Biol*. 2005;25:10479-91.
- 292 [23] Walther T, Jank A, Heringer-Walther S, Horn LC, Stepan H. Angiotensin II type  
293 1 receptor has impact on murine placentation. *Placenta*. 2008;29:905-9.
- 294 [24] George AJ, Thomas WG, Hannan RD. The renin-angiotensin system and cancer:  
295 old dog, new tricks. *Nat Rev Cancer*. 2010;10:745-59.
- 296 [25] Wang Y, Pringle KG, Lumbers ER. The effects of cyclic AMP, sex steroids and  
297 global hypomethylation on the expression of genes controlling the activity of the renin-  
298 angiotensin system in placental cell lines. *Placenta*. 2013;34:275-80.  
299

## 300 LEGENDS TO FIGURES

301 **Fig. 1.** Effect of O<sub>2</sub> on expression of RAS components in HTR-8/SVneo cells at 24 and  
302 48 h. (A) *AGTR1* and (B) *VEGF* mRNA were significantly upregulated by low oxygen  
303 (1% O<sub>2</sub>) at 48 h when compared with 5% and 20% O<sub>2</sub> (\*, both  $P<0.01$ ). (B) *VEGF*  
304 mRNA was significantly increased at 48 h compared with 24 h by 1% O<sub>2</sub> (#,  $P<0.05$ ).  
305 (A) *AGTR1*, (C) *ATP6AP2*, (D) *REN*, (E) *ACE1* and (F) *AGT* mRNA expression  
306 significantly increased with incubation time (•, all  $P<0.05$ ). Data are presented as mean  
307  $\pm$  SEM.

308

309 **Fig. 2.** The correlations between *VEGF* and *AGTR1* mRNA abundance and between  
310 *AGTR1* and *ATP6AP2* mRNA abundance in HTR-8/SVneo cells cultured at 1% O<sub>2</sub>. (A)  
311 *VEGF* and *AGTR1* ( $r=0.638$ ,  $R^2=0.407$  and  $P=0.0044$ ); (B) *AGTR1* and *ATP6AP2*  
312 ( $r=0.650$ ,  $R^2=0.473$  and  $P=0.0035$ ).

313

314 **Fig. 3.** Levels of prorenin protein in culture medium and cell lysates of HTR-8/SVneo  
315 cells cultured in varying oxygen tensions for 48 h. Neither culture medium (A) nor cell  
316 lysate (B) prorenin protein levels were affected by oxygen tension. Data are presented  
317 as mean  $\pm$  SEM.

318

319 **Figure 4.** The effect of O<sub>2</sub> on VEGF protein concentration in the culture medium of  
320 HTR-8/SVneo cells at 48 h. Culture in 1% O<sub>2</sub> for 48 h significantly increased the  
321 concentration of VEGF compared with culture in either 5% or 20% O<sub>2</sub> (\*,  $P<0.05$  and  
322  $P<0.01$  respectively). Data are presented as mean  $\pm$  SEM.

323

324 **Fig. 5.** The effect of O<sub>2</sub> on ACE protein concentration in the culture medium and cell  
325 lysate of HTR-8/SVneo cells at 48 h. (A) The concentration of ACE in the culture  
326 medium of HTR-8/SVneo cells was significantly increased after 48 h of culture in both  
327 1% and 5% O<sub>2</sub> when compared with the concentration measured in medium cultured in  
328 20% O<sub>2</sub> (\*, both  $P<0.05$ ). (B) ACE protein levels in cell lysates were unaffected by  
329 oxygen. Data are presented as mean  $\pm$  SEM.

Figure 1





Figure 2

A



B






Figure 3






Figure 4



**Figure 5**

