



Coulthard, JD; Palla, L; Pot, GK (2017) Breakfast consumption and nutrient intakes in 4-18-year-olds: UK National Diet and Nutrition Survey Rolling Programme (2008-2012). *The British journal of nutrition*. pp. 1-11. ISSN 0007-1145 DOI: <https://doi.org/10.1017/S0007114517001714>

Downloaded from: <http://researchonline.lshtm.ac.uk/4237547/>

DOI: [10.1017/S0007114517001714](https://doi.org/10.1017/S0007114517001714)

Usage Guidelines

Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact researchonline@lshtm.ac.uk.

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

1 **Breakfast consumption and nutrient intakes in 4-18 year olds –**
2 **UK National Diet and Nutrition Survey Rolling Programme**
3 **(2008-2012)**

4 Janine D Coulthard¹, Luigi Palla², Gerda K Pot^{1,3*}

5 ¹Faculty of Life Sciences and Medicine, Diabetes and Nutritional Sciences Division, King's College London, Franklin-
6 Wilkins Building, 150 Stamford Street, SE1 9NH, London, UK

7 ²Faculty of Epidemiology and Population Health, Department of Medical Statistics, London School of Hygiene and Tropical
8 Medicine, Keppel St, WC1E 7HT, London, UK

9 ³Faculty of Earth and Life Sciences, Section of Health and Life, Vrije University Amsterdam, de Boelelaan 1085, 1081 HV
10 Amsterdam, The Netherlands

11 *Corresponding author: G.K Pot, email Gerda.Pot@kcl.ac.uk, telephone +44 (0)20 7848 4437

12

13 **Running title:** Breakfast and nutrient intakes in children. [42 letters & spaces (max 45)]

14

15 **Key words:** Children, dietary patterns, Breakfast, Nutrient Intakes, National Diet and
16 Nutrition Survey Rolling Programme

17

18 **Journal:** British Journal of Nutrition

19 **Word Count:** 5140

20 **Number of tables/figures:** 5

21 **Number of references:** 72

22 **Conflicts of interest:** none

23 **Abstract**

24 **Word Count:** 250 (max 250)

25 Although breakfast consumption is widely considered to be an important component of a
26 healthy lifestyle, few UK studies have examined differences in nutrient intakes between
27 breakfast consumers and breakfast skippers among children and adolescents. We investigated
28 associations between breakfast skipping in 4-18 year olds and their nutrient intakes using data
29 from the UK's National Diet and Nutrition Survey Rolling Programme. Dietary data were
30 derived from 4-day estimated food diaries of 802 children aged 4-10 years and 884 children
31 aged 11-18 years (*n* 1686 in total). Daily nutrient intakes of children with different breakfast
32 habits were compared by one-way ANCOVA adjusting for relevant covariates (sex, age,
33 ethnicity, equivalised household income and BMI). Within-person analysis was carried out
34 on children with an irregular breakfast habit (*n* 879) comparing nutrient intakes on breakfast
35 days with those on non-breakfast days using repeated measures ANCOVA. We observed that
36 the overall nutritional profile of the children in terms of fibre and micronutrient intake was
37 superior in frequent breakfast consumers (micronutrients: folate, calcium, iron and iodine
38 ($P < 0.01$)) and, for the 4-10 years age group, on breakfast days (micronutrients: folate,
39 vitamin C, calcium and iodine ($P < 0.01$)). Also, significantly higher proportions of breakfast-
40 consuming children met their Reference Nutrient Intakes of folate, vitamin C, calcium, iron
41 and iodine compared to breakfast skippers (chi-square analysis, $P < 0.001$). Our study adds to
42 the body of data linking breakfast consumption with higher quality dietary intake in school-
43 age children, supporting the promotion of breakfast as an important element of a healthy
44 dietary pattern in children.

45

46 **Introduction**

47 Breakfast is widely considered to be an important component of a healthy lifestyle for both
48 adults and children. Its consumption is promoted by the UK Government's Change4Life
49 public health campaign in England⁽¹⁾. One of the suggested benefits of breakfast for children
50 and adolescents is that eating breakfast regularly protects against overweight and obesity. To
51 date, evidence for this consists largely of epidemiological studies⁽²⁻⁹⁾ and intervention studies
52 have so far failed to demonstrate a causal relationship between breakfast habit and
53 adiposity⁽¹⁰⁻¹³⁾. Another proposed benefit of breakfast consumption in children and
54 adolescents is that it improves overall nutrient intakes⁽²⁾.

55 Previous studies have been carried out into variations in nutritional intakes dependent on
56 child breakfast habits, but many of these focus on consumption of a specific type of
57 breakfast, for example breakfast cereal⁽¹⁴⁻¹⁶⁾ or a "good-quality" breakfast⁽¹⁷⁾. Some have
58 examined overall differences in nutritional profile between children who eat breakfast and
59 those that do not, including recent studies involving North American, Mexican and
60 Australian populations⁽¹⁸⁻²²⁾, and have reported that breakfast consumption compared with
61 breakfast skipping was associated with improved nutrient intakes^(18,20-30). However, the
62 authors are not aware of any recent studies of UK or other European children examining
63 differences in nutrient intakes between breakfast consumers and breakfast skippers.
64 Therefore, our main aim in this study was to investigate associations between breakfast
65 skipping in 4-18 year olds and their nutrient intakes using data from the UK's National Diet
66 and Nutrition Survey Rolling Programme (NDNS RP). We compared daily nutrient intakes
67 between children with different breakfast habits and also carried out a within-person analysis,
68 comparing intakes of children on breakfast days with their intakes on non-breakfast days.
69 This latter approach was adopted in order to minimise the impact of residual confounding
70 inherent in cross-sectional studies.

71 Many epidemiological studies comparing the characteristics of individuals with different
72 breakfast habits ask participants (or their parents) how frequently they (or their children)
73 consume breakfast, but do not include a definition of this meal^(4,27,31,32). Where breakfast is
74 defined, this definition varies widely^(2,33,34). Some studies classify any energy intake within a
75 specified time period as breakfast^(19,21), others include only solid foods and not beverages in
76 the definition, regardless of the calorie content of the latter⁽³⁴⁾. We employed an objective
77 definition of breakfast, based on a minimum energy intake within a specified time period. We
78 hypothesized that intakes of micronutrients of public health interest, which have a key role in
79 children's healthy development and growth, may be particularly affected by breakfast
80 skipping, as many breakfast items consumed widely by UK children, such as breakfast
81 cereals, fruit juice and dairy products, are important sources of micronutrients for this age

82 group⁽³⁵⁾.

83

84 **Methods**

85 *Population*

86 The data analysed in this study were collected between 2008 and 2012 as part of the NDNS
87 RP⁽³⁶⁾. Each year the NDNS RP gathers dietary and nutritional data from approximately
88 1,000 randomly sampled individuals living in private UK households, comprising equal
89 numbers of adults (aged 19 years and over) and children (aged 1.5-18 years)⁽³⁵⁾. These
90 individuals are sampled using The Royal Mail's Postcode Address File, comprising all UK
91 addresses, to randomly select addresses. Where there is more than one household at a
92 selected address, one of these households is randomly selected. At each selected household
93 either one adult and one child (if present) are selected, or one child, resulting in roughly equal
94 numbers of adults and children in the final sample. Chapter 2 of the report on the NDNS RP
95 results from years 1-4 (combined)⁽³⁵⁾ gives fuller details of its methodology. This study
96 focused on the sub-population of children in the NDNS RP 2008-2012 sample aged 4-18
97 years, consisting of 802 children aged 4-10 years and 884 children aged 11-18 years (*n* 1686
98 in total). Ethics approval for the NDNS RP was obtained from Oxfordshire A Research
99 Ethics Committee⁽³⁵⁾.

100

101 *Dietary assessment*

102 Each survey participant was visited in their home by a survey interviewer, who placed a 4-d
103 estimated (unweighed) food diary (with written instructions) to be completed on 4
104 consecutive days by the participant, or their parent for those aged 11 years and under⁽³⁷⁾⁽³⁸⁾.
105 Follow-up checks were made by the interviewer to optimise completeness of record keeping
106 in the diary⁽³⁵⁾. Within the sample analysed for years 2008 to 2012 of the NDNS RP 1686
107 children aged 4-18 years completed at least 3 diary days (98.2% of these completed the full 4
108 diary days). Home visits were carried out continuously throughout each year, from February
109 2008 to August 2012, thus allowing for seasonal variations in diet⁽³⁹⁾.

110 The diary entries were then recorded and analysed by a dietary assessment system using food
111 composition data from the Department of Health's NDNS Nutrient Databank to estimate
112 energy and nutrient intakes⁽³⁷⁾. The NDNS Nutrient Databank is based on data from McCance
113 and Widdowson's "The Composition of Foods" series^(37,40). Non-milk extrinsic sugars
114 (NMES) were defined as comprising all free sugars (added monosaccharides and
115 disaccharides, together with naturally occurring sugars in honey, syrups and fruit juices) and
116 50% of fruit sugars from stewed, dried or canned fruit⁽⁴¹⁾. Dietary fibre intakes were of non-

117 starch polysaccharides (NSP), defined by the Englyst method⁽⁴²⁾. Mean daily energy intakes
118 were expressed as a percentage of the Estimated Average Requirement (EAR) for each child,
119 as specified by the Scientific Advisory Committee on Nutrition (SACN)⁽⁴³⁾. Mean daily
120 intakes of the micronutrients folate, vitamin C, calcium, iron and iodine were compared to
121 Dietary Reference Values (DRV) (Reference Nutrient Intakes (RNI) and Lower Reference
122 Nutrient Intakes (LRNI)) for each child, as set by the Committee on Medical Aspects of Food
123 Policy (COMA)⁽⁴⁴⁾. The results of the above analysis of the diary entries form part of the
124 published core sample data for the NDNS RP 2008-2012⁽³⁶⁾ and were further analysed in this
125 study as described below.

126 For each day the food diary entries had been split into 7 different time periods: 06.00 to 08.59
127 hours, 09.00 to 11.59 hours, 12.00 to 13.59 hours, 14.00 to 16.49 hours, 17.00 to 19.59 hours,
128 20.00 to 21.59 hours and 22.00 to 05.59 hours. Microsoft Excel for Mac 2011 (version
129 14.4.6) and IBM SPSS Statistics (version 23) were used to calculate the total energy intake
130 for each diary day between 06.00 and 8.59 hours and identify those days on which at least
131 100 kcal were consumed between 06.00 and 8.59 hours (breakfast days) and those on which
132 less than 100 kcal were consumed (non-breakfast days). The threshold of 100 kcal was
133 chosen to allow for consumption of a milky drink which was not considered sufficient to be
134 classed as breakfast⁽⁴⁵⁾. Using these data, the children were split into 3 categories: those
135 consuming breakfast every diary day, those consuming breakfast on at least one but not all
136 diary days, and those not consuming breakfast on any diary day. A subset of the group of
137 children with an irregular breakfast habit i.e. who consumed breakfast on at least one but not
138 all diary days was created, by eliminating all diary days that comprised a Saturday or a
139 Sunday and then identifying all children who still had an irregular breakfast habit based on
140 weekday diary days alone. This was to correct for a possible shift to later breakfast time
141 and/or a different pattern in nutrient intakes at the weekend. For each child with an irregular
142 breakfast habit mean daily nutrient intakes for breakfast days and non-breakfast days
143 respectively were calculated.

144 The nutrients selected for analysis in this study were the macronutrients protein, fat and
145 carbohydrate plus a selection of other nutrients linked with an unhealthy (NMES and sodium)
146 or a healthy diet (dietary fibre). We also analysed intakes of a selection of micronutrients that
147 Public Health England has identified as being of particular interest to public health, namely:
148 folate, iron, vitamin C and calcium⁽³⁵⁾. To this list we added iodine: there is some evidence of
149 iodine deficiency in UK adolescent girls giving rise to public health concerns due to its vital
150 role in foetal neurodevelopment⁽⁴⁶⁾. The mean daily nutrient intake values were expressed as
151 a percentage of total energy intake for protein, fat, carbohydrate and NMES and as mean

152 intakes per 1000kcal of energy intake for the remaining nutrients. This was to allow for
153 possible differences in daily energy intakes between breakfast and non-breakfast eaters.

154

155 *Other measures*

156 During the home visit, the interviewer measured the weight and height of the participant so
157 that their BMI could be calculated (weight in kilograms divided by the square of height in
158 metres)⁽³⁵⁾. The calculated BMI and the British 1990 growth reference (UK90) charts⁽⁴⁷⁾ were
159 used to categorise the children as normal weight, overweight (85th centile cut-off) or obese
160 (95th centile cut-off). The interviewer also conducted a computer-assisted personal interview
161 (CAPI) to collect further data on the individual and their household, including age, ethnicity
162 (five main categories: white, mixed, black or black British, Asian or Asian British, and
163 other), whether the individual was currently dieting to lose weight (a “yes” or “no” response,
164 years 3 and 4 of the NDNS RP, for individuals 11 years and older) and household income
165 (choice of 13 income bands, ranging from under £5,000 to £100,000 to more)⁽⁴⁸⁾.

166

167 *Statistical analysis*

168 The energy-adjusted daily nutrient intake, equivalised household income (equivalised for
169 different household sizes and composition using the McClements equivalence scale⁽³⁵⁾) and
170 BMI variables were checked for normality by inspecting frequency distribution histograms
171 and skew and kurtosis values and transformed as necessary. The differences between the
172 children grouped by three categories of breakfast habit (consumption of breakfast on every,
173 some or no diary days) with respect to sex, age, ethnicity, weight status and equivalised
174 household income were assessed by chi-square analysis and ANOVA. Chi-square analysis
175 was carried out on the dieting variable and used to examine the proportions of children
176 meeting the RNI and LRNI for daily intakes of the selected micronutrients. Chi-square
177 analysis was also used to compare the characteristics of children with missing BMI and/or
178 income data to assess whether the relatively high incidence of missing data (n 270, 16% of
179 sample) might affect the results adjusted for these variables.

180 Daily nutrient intakes for the three groups of children with different breakfast habits were
181 compared by one-way ANOVA and then ANCOVA, with adjustments made for the
182 following covariates: sex, age, ethnicity, equivalised household income and BMI. Energy
183 intake for this analysis was expressed as a proportion of EAR, to allow for variations in
184 energy intake with age and sex. To investigate the potential effect of under- or over-reporting
185 of dietary intakes a sensitivity analysis was carried out, rerunning the chi-square analysis of
186 children meeting the DRVs and the ANCOVA analysis of energy intake after eliminating
187 those children with energy intake as a proportion of EAR more than two SDs from 100% (as

188 described by McCrory *et al.*⁽⁴⁹⁾). This was not considered necessary for the nutrient intakes,
189 as these were all expressed as a proportion of energy intake. Due to inequality of variance
190 (assessed using Levene's test) for three of the nutrient variables: protein, NMES and sodium,
191 which can lead to an increase in type 1 error rate⁽⁵⁰⁾, the ANOVA and ANCOVA analyses for
192 these variables were carried out after equalizing the numbers in the three groups of children
193 by random sampling, to create three equal groups of 245 cases each (for large sample sizes
194 and modest levels of variance heterogeneity ANOVA is generally robust to inequality of
195 variances if group sizes are equal⁽⁵⁰⁾).

196 Within-person analysis was carried out on the children with an irregular breakfast habit (*n*
197 879), comparing their mean energy adjusted nutrient intakes for days on which they had
198 consumed breakfast with those for days on which they had not. It was assumed that any
199 degree of mis-reporting of dietary intakes by an individual would be similar across the diary
200 days and would therefore not have a significant effect on the within-person analysis. Paired
201 sample t-tests were conducted, followed by repeated measures ANOVA, the latter adjusted
202 for the covariates sex, age, BMI, ethnicity and equivalised household income. These tests
203 were also carried out on the sample split into two age groups: 4-10 years (*n* 384) and 11-18
204 years (*n* 495) and on the subset of 4-18-year-olds with an irregular breakfast habit after
205 removal of weekend diary days. Due to collinearity of nutrient variables, for all the tests
206 carried out a p value of <0.01 was considered as statistically significant to allow for multiple
207 testing, rather than using the potentially overly conservative Bonferroni method of
208 adjustment⁽⁵¹⁾. We did not apply the weighting as provided with the NDNS 2008-12 RP
209 dataset, as the calibration weights may not reflect the characteristics of the subsample of
210 children that we analysed, which may not be representative of the UK population.

211

212 **Results**

213 *Description of population*

214 The characteristics of the children, split by breakfast habit are set out in **Table 1**. There was a
215 significant variation in breakfast habit between girls and boys ($P=0.001$), with 19.9% of girls
216 skipping breakfast every diary day compared to 14.5% of boys. At 6.5%, the proportion of
217 the 4-10-year-olds skipping breakfast every diary day was less than a quarter of the
218 proportion of 11-18-year-olds (26.8%), and the proportion in the younger age group
219 consuming breakfast every day was 45.6%, more than twice that of the older age group
220 ($P<0.001$). Mean equivalised household income varied significantly with breakfast habit
221 ($P=0.001$), with a mean household income of £28,194 (SD £18,349) for those children
222 consuming breakfast every diary day compared to a mean household income of £23,587 (SD

223 £16,374) for those children skipping breakfast every diary day. There was no significant
224 difference in breakfast habits based on ethnicity or weight status. Data on whether or not the
225 children were dieting during the diary period was only available for 11-18-year-olds in years
226 3 and 4 of the NDNS RP. A significantly higher percentage of girls were dieting than boys
227 (10.7% v. 3.4%, $P=0.003$, n 430), but no significant variation in breakfast habit with dieting
228 behaviour was observed for the children as a whole ($P=0.456$), or for the girls alone
229 ($P=0.419$, n 224) (the chi-square analysis was invalid for the boys as more than 20% of the
230 expected cell values were less than 5). There were no significant differences found in sex
231 ($P=0.457$), age group ($P=0.470$) or breakfast habit ($P=0.844$) for children with missing BMI
232 and/or income data compared to those with complete data.

233

234 *Breakfast habits and micronutrient DRVs*

235 Significant increases ($P<0.001$) were noted in the proportion of children meeting their RNI
236 for each of the micronutrients folate, vitamin C, calcium, iron and iodine as breakfast
237 frequency increased (**Table 2**). The same trend was observed in the proportion of children
238 meeting the LRNI for folate, calcium, iron and iodine ($P<0.001$). No children who consumed
239 breakfast daily had a folate intake below their LRNI, compared to 7.3% of those who did not
240 eat breakfast on any diary day. The proportions of children not meeting their LRNI of
241 calcium, iron and iodine in the daily breakfast group were 2.9%, 4.4% and 3.3% respectively,
242 compared to 19.0%, 31.5% and 21.5% respectively in the breakfast skipping group. Similar
243 results were obtained for the sensitivity analysis including only plausible reporters of energy
244 intake (n 1505) (invalid LRNI chi-square results for folate and vitamin C as more than 20%
245 of cells had expected count of less than 5).

246

247 *Breakfast habits and nutrient intakes*

248 After adjustment for covariates, we observed significant increases in mean intakes of energy
249 (as a percentage of EAR) ($P=0.009$) and carbohydrates (as a percentage of energy) ($P=0.01$)
250 and decreases in mean intakes of fat (as a percentage of energy) ($P=0.005$), with increasing
251 number of breakfast days (**Table 3**). However, after eliminating implausible reporters of
252 energy intakes (energy intake as a percentage of EAR more than two SDs from 100% (i.e.
253 outside the range 54-146%)), the increases in energy intakes became non-significant
254 ($P=0.088$, n 1271). The percentage of plausible reporters in the sample was 89%, with a
255 significantly higher proportion of plausible reporters within the 4-10-year-olds than 11-18-
256 year-olds (97% v. 83%, $P<0.001$).

257 In the ANCOVA analysis we observed no statistically significant variations in intakes of
258 protein, NMES, vitamin C or sodium, but energy adjusted intakes of fibre, folate, calcium,

259 iron and iodine all increased significantly with increasing frequency of breakfast
260 consumption ($P<0.001$).

261

262 *Within-person comparison of nutrient intakes on breakfast days v. non-breakfast days*

263 For 4-18-year-olds, after adjustment for covariates, we observed significantly higher mean
264 intakes of energy, carbohydrate, folate, calcium and iodine and significantly lower mean
265 intakes of protein and sodium for days on which breakfast was consumed compared to days
266 on which it was not eaten (Table 4(a)). The mean energy intake for days on which breakfast
267 was consumed was 87kcal higher than for days on which it was not consumed (95%CI
268 52,121kcal, $P<0.001$). No significant differences were observed when comparing breakfast
269 with non-breakfast days for mean intakes of fat, NMES, fibre, vitamin C or iron.

270 For 4-10-year-olds there was no significant difference between mean energy intake on days
271 on which breakfast was eaten and mean intake on non-breakfast days (Table 4(b)). In
272 comparison, in the older age group (11-18-year-olds), after adjustment for covariates, energy
273 intake was significantly higher on breakfast days (118kcal (95% CI 66,169 kcal), $P<0.001$)
274 (Table 4(c)). For both age groups carbohydrate intake was significantly higher on breakfast
275 days compared to non-breakfast days. There was a significant decrease in protein intake on
276 breakfast days compared to non-breakfast days for the older but not the younger age group.
277 For 4-10-year-olds mean energy adjusted intakes of fibre, folate, vitamin C, calcium and
278 iodine were significantly higher on days on which breakfast was eaten, but for 11-18-year-
279 olds out of these five nutrients a significant increase was only observed for calcium intakes.
280 Whereas for the 4-10 years age group there was no significant difference in mean sodium
281 intakes comparing breakfast with non-breakfast days, sodium intake was significantly lower
282 for the 11-18 years age group on breakfast days.

283 For 4-18-year-olds with an irregular breakfast habit after removal of weekend diary days,
284 there were significantly higher intakes of energy (158kcal (95%CI 93,223kcal), folate,
285 calcium and iodine for days on which breakfast was consumed compared to days on which it
286 was not (Table 5). In contrast to the analysis of all diary days, no significant differences in
287 intakes of any of the macronutrients (protein, fat or carbohydrate) were observed and there
288 was no significant difference in sodium intake.

289

290 **Discussion**

291 Our findings suggest that the overall nutritional profile of the children in terms of fibre and
292 micronutrient intake was superior in regular breakfast consumers, for whom higher energy-
293 adjusted intakes of fibre, folate, calcium, iron and iodine were observed compared to
294 breakfast skippers. These findings are supported by our observation that significantly higher

295 proportions of breakfast-consuming children met their RNI of folate, vitamin C, calcium, iron
296 and iodine compared to breakfast skippers.

297 In accordance with previous studies^(21,23,25), for both the between and within-person analyses
298 and both age groups a lack of breakfast was associated with lower calcium intakes. A recent
299 Mexican study of breakfast dietary patterns among 4-13 year old children (*n* 3760) found that
300 the differences in overall daily nutrient intake profile between breakfast consumers and
301 skippers varied with the type of breakfast consumed, but all breakfast types were associated
302 with a higher daily intake of calcium than breakfast skipping⁽²²⁾. This is a particularly
303 important finding given that calcium is a vital nutrient for bone growth and needs in
304 childhood are high due to rapid growth and bone mass accretion⁽⁵²⁾. In the within-person
305 analysis, higher intakes of fibre, folate, vitamin C and iodine were observed on the days that
306 the 4-10-year-olds ate breakfast, but no significant difference in intakes of these
307 micronutrients was noted for the 11-18-year olds. This suggests that more foods rich in these
308 micronutrients (for example, fortified cereals, milk and fruit juice) are consumed for
309 breakfast in the younger compared to the older age group, possibly due to greater parental
310 supervision of the younger children's meals. A Spanish study of 8-17-year-old children (*n*
311 4332) evaluating the utility of a breakfast quality index, in which points were awarded for the
312 consumption of cereals, fruits, vegetables and dairy products at breakfast, found that the
313 score decreased with age ($P=0.001$)⁽⁵³⁾. For the within-person analyses including all diary
314 days, intakes of sodium are significantly higher on no breakfast days for 4-18-year olds and
315 11-18-year-olds, but not 4-10-year-olds, suggesting that the older age group may compensate
316 for the lack of breakfast by the consumption of salty snacks later in the day (protein intakes
317 are also higher on non-breakfast days for the older but not the younger group). However this
318 association was no longer significant when weekend diary days were removed from the
319 within-person analysis (and is not present in the between-person analysis), so may simply be
320 a reflection of a different dietary pattern at weekends (for example, late or no breakfast with
321 greater daily amounts of protein rich and salty foods) in the older age group.

322 We did not find any evidence to support the oft-quoted hypothesis that breakfast skipping
323 leads to increased overall daily energy intake due to compensatory overeating later in the
324 day^(9,54). On the contrary, in our analyses we observed that breakfast skipping was associated
325 with either no difference or a significantly lower daily energy intake. These results are in
326 accordance with those of most (but not all) recent observational studies of children^(9,18,20-22).

327 Similarly, small cross-over trials in children have reported no significant differences in
328 overall daily energy intakes on breakfast and non-breakfast days^(55,56). The findings on
329 energy intake in this and other studies might suggest caution when recommending breakfast
330 consumption as a weight management tool in children, despite the large body of

331 epidemiological evidence linking breakfast skipping and excess weight in children. Also,
332 they undermine the argument for a causal link between breakfast consumption and
333 overweight and obesity based on excess energy intake. Interestingly, a large, longitudinal US
334 study of breakfast habits and weight gain in 9-14 year olds found that, although breakfast
335 skipping was associated with higher BMI overall, overweight breakfast skippers tended to
336 lose weight over the study period compared to overweight breakfast eaters, whereas the
337 reverse was true for normal weight breakfast skippers compared to normal weight breakfast
338 eaters⁽⁵⁷⁾. Alternative theories for the link observed between adiposity and breakfast skipping
339 include the presence of confounding factors such as sleep duration and circadian rhythms⁽¹²⁾,
340 and lower physical activity levels in breakfast skippers^(4,58,59).

341 The results of previous studies of children investigating the effect on overall macronutrient
342 profile of breakfast habit vary⁽²³⁾. In this study small but statistically significant variations in
343 macronutrient profile were noted, with the proportion of carbohydrate consumed generally
344 higher for breakfast consumers (for the between-person analysis) and on breakfast days (for
345 the within-person analysis), at the expense of either protein or fat intake. However, no
346 significant variations in macronutrient profile when comparing intakes on breakfast days with
347 those on non-breakfast days were found for the within-person analysis after weekend diary
348 days had been removed from the analysis. This may be due to a different dietary pattern for
349 the children at the weekend. In studies that have analysed breakfast habit by type of
350 breakfast consumed, different overall daily nutrient intake profiles have been associated with
351 different breakfast types^(20,22). For example, in a large US cross-sectional study of breakfast
352 consumption in 9-18 year olds (*n* 9659), consumers of ready-to-eat cereals had higher intakes
353 of carbohydrate and lower intakes of fat than breakfast skippers, but for consumers of other
354 types of breakfast there was no significant difference in intakes of these macronutrients
355 compared with breakfast skippers⁽¹⁸⁾.

356 No link was observed between overall NMES intake (as a percentage of energy) and
357 breakfast consumption. In the within-person analysis we observed no significant difference in
358 the proportion of fat or NMES consumed between breakfast and non-breakfast days,
359 suggesting that intermittent breakfast skipping did not lead to an increase in consumption of
360 poor quality, high sugar and high fat foods and beverages on non-breakfast days in this study
361 population, as has been postulated elsewhere⁽⁵⁴⁾.

362 In our analysis of the characteristics of the children in the sample we found no significant
363 differences in the proportion of normal weight, overweight and obese children in each
364 breakfast habit category, which is at odds with the findings of some but not all cross-sectional
365 studies⁽⁴⁻⁷⁾ in children. In line with other studies in children^(9,23,60,61), we observed higher
366 levels of breakfast skipping in girls and older children and lower mean household incomes

367 for breakfast skippers. There is evidence that frequency of breakfast skipping in teenagers is
368 related to dieting and other weight-control behaviours⁽⁹⁾, which may explain its greater
369 incidence in girls⁽⁶²⁾. A cross-sectional study of UK 11-16 year olds (*n* 1019) found that
370 almost twice as many girls reported dieting and those girls that were dieting were 3 times
371 more likely to skip breakfast than non-dieters⁽⁶³⁾. In our study a significantly higher
372 proportion of girls in the 11-18 year old age group stated that they were currently dieting than
373 boys. However, reported dieting behaviour did not vary significantly with breakfast habit for
374 the girls in our sample, but this could be due to the smaller sample size (*n* 224) resulting in
375 lower statistical power (dieting data was only available for last 2 years of the NDNS RP
376 2008-2012).

377 The greater level of breakfast skipping among older children may be influenced by a
378 reduction in parental control enforcing a “healthy” breakfast habit. It may also reflect the
379 shift in circadian rhythms in adolescence to a later wake/sleep cycle⁽⁶⁴⁾. During puberty an
380 individual’s chronotype, that is their preference for an early or late wake/sleep cycle, shifts
381 from early to late, with sleep schedules moving progressively later between the ages of 10
382 and 20 years⁽⁶⁵⁾. During the school week children’s wake/sleep cycle is dictated by the school
383 routine but at the weekend the wake/sleep cycle is generally less restricted. A study of food
384 logs of German adolescents (*n* 152, mean age 13.23 years) found that wake times at the
385 weekend were on average 2:40 hours later, which translated to later breakfast times: the
386 average breakfast time on weekdays was 6:36am, compared to 9:15am at weekends⁽⁶⁶⁾. Wake
387 time data was not available for the study sample, so in our analysis we have defined breakfast
388 in relation to a specific, fixed time period, namely 6:00-8:59am, rather than relate it to intake
389 within a certain time of waking. However, because of this weekend shift in breakfast times a
390 meal eaten shortly after waking may not fall within this fixed time period. To address this
391 issue, a separate within-person analysis was carried out after removing weekend diary days.

392 Chronotype not only varies with age but it also depends on genetic and environmental
393 factors^(65,67). Chronotype may be an important confounding factor in the between-person
394 analysis in this and other studies. There is evidence that not only are adolescents with later
395 chronotypes more likely to skip breakfast, they are also more likely to have poorer overall
396 diets⁽⁶⁶⁾ and lower levels of physical activity⁽⁶⁸⁾. The impact of chronotype and other possible
397 residual confounding factors, which are always an issue in cross-sectional studies, should be
398 less of a factor in the within- compared to the between-person analysis. Adjustments have
399 been made in the ANOVA for age, sex, BMI, ethnicity and equivalised household income,
400 however we were not able to adjust for physical activity level due to lack of complete and
401 consistent data across all ages⁽³⁵⁾.

402 We relied on data from 4-d estimated food diaries and an objective definition of breakfast to
403 categorise children by their breakfast habit, rather than on responses to an eating habits
404 questionnaire. This avoids issues resulting from inconsistent personal definitions of breakfast.
405 Nevertheless, it is recognised that the 100 kcal threshold chosen for our definition of
406 breakfast is, to some extent, arbitrary. Also, the fixed time frame we used to define breakfast
407 may result in a late weekend breakfast not being captured by the definition, however this was
408 allowed for in the within-person analysis by removing weekend diary days.

409 In common with other dietary surveys, there is a possibility of mis-reporting of dietary
410 intakes. Doubly labelled water techniques used in the NDNS RP to validate energy intakes
411 for a sample of survey participants suggest that under-reporting may have been more
412 prevalent for the children in the 11-18 years age group, who completed their own food
413 diaries, in contrast to children in the younger age group, whose diaries were completed by
414 their parents⁽⁶⁹⁾. The results of our sensitivity analysis suggest that there were higher levels of
415 mis-reporting in the older age group. Where appropriate, to assess the impact of mis-
416 reporting we reran analyses omitting implausible reporters.

417 Many different methods of dietary data collection have been used in previous studies of
418 breakfast habits in children⁽²³⁾, with varying degree of reliability. Strengths of our study
419 include the large sample size and the method of dietary assessment used, namely a 4-d
420 estimated food diary. In other recent cross-sectional surveys of children's breakfast habits
421 and nutrient intakes⁽¹⁸⁻²²⁾, the 24-hour recall method was used, which relies heavily on the
422 accuracy of the child's or their parent's memory. We are aware of two small crossover
423 studies involving US children which report on the impact of breakfast skipping on overall
424 daily energy intake^(55,56) but, to our knowledge, there has not yet been a randomized,
425 controlled trial assessing the impact of breakfast consumption in children on intakes of
426 individual macro- and micronutrients. The approach we adopted of approximating a
427 crossover study design (thereby reducing residual confounding) in free living individuals by
428 conducting a within-person analysis of subjects with an irregular breakfast habit has been
429 carried out previously in adults⁽⁷⁰⁾, but not children.

430 In our study we examined how macro- and micronutrient intakes, plus energy intake, varied
431 with children's breakfast habit. A possible topic for future research would be to look at
432 associations in UK children between breakfast habit and daily intakes of specific foods,
433 incorporating a diet quality index⁽⁷¹⁾ to further investigate associations between breakfast
434 habit and overall diet quality. It would also be interesting to investigate how the quality of the
435 breakfast foods consumed by UK children varies with age, to ascertain if the differences we
436 noted between the two age groups in our within-person analysis might be attributed to lower
437 quality breakfasts in the older age group. Work has already been carried out on developing

438 breakfast quality indices for use in relation to children and adolescents in a Mediterranean
439 setting^(16,53,71); further work is required to adapt these for use in a UK population.
440 In conclusion, the connection between the consumption of breakfast and good health appears
441 to involve many different factors, and is still some way from being fully elucidated. A causal
442 link with obesity is, as yet, unsupported by the available evidence⁽⁷²⁾. However, this study
443 adds to the existing body of data linking breakfast consumption with higher quality dietary
444 intake in school-age children, particularly the 4-10 years age group, supporting the promotion
445 of breakfast as an important element of a healthy dietary pattern in children.

446 **Acknowledgements**

447 The authors would like to thank everyone involved in the National Diet and Nutrition Survey
448 Rolling Programme (2008-2012), the data from which formed the basis of this study.

449

450 **Financial Support**

451 This research received no specific grant from any funding agency, commercial or not-for-
452 profit sectors.

453

454 **Conflict of interest**

455 None.

456

457 **Authorship:** JDC formulated the research question, prepared the data for analysis, analysed
458 and interpreted the data and wrote the manuscript, all under the supervision of GKP and LP.
459 All authors approved the final draft before publication.

460 **References**

- 461 1. NHS Change4Life Get off to a great start with a healthy breakfast!
462 www.nhs.uk/Change4Life/Breakfast-for-life.aspx (accessed November 2016).
- 463 2. Rampersaud GC (2009) Benefits of breakfast for children and adolescents: update and
464 recommendations for practitioners. *Am. J. Lifestyle Med.* **3**, 86–103.
- 465 3. Moreno LA, Rodriguez G, Fleta J, et al. (2010) Trends of dietary habits in adolescents.
466 *Crit. Rev. Food Sci. Nutr.* **50**, 106–112.
- 467 4. Szajewska H & Rusczyński M (2010) Systematic review demonstrating that breakfast
468 consumption influences body weight outcomes in children and adolescents in Europe.
469 *Crit. Rev. Food Sci. Nutr.* **50**, 113–119.
- 470 5. De La Hunty A, Gibson S & Ashwell M (2013) Does regular breakfast cereal
471 consumption help children and adolescents stay slimmer? A systematic review and
472 meta-analysis. *Obes. Facts* **6**, 70–85.
- 473 6. Zakrzewski JK, Gillison FB, Cumming S, et al. (2015) Associations between breakfast
474 frequency and adiposity indicators in children from 12 countries. *Int. J. Obes. Suppl.* **5**,
475 S80–S88.
- 476 7. Kelly Y, Patalay P, Montgomery S, et al. (2016) BMI development and early
477 adolescent psychosocial well-being: UK Millennium Cohort Study. *Pediatrics* **138**,
478 e20160967.
- 479 8. Haug E, Rasmussen M, Samdal O, et al. (2010) Overweight in school-aged children
480 and its relationship with demographic and lifestyle factors: results from the WHO-
481 collaborative Health Behaviour in School-aged Children (HBSC) Study. *Int. J. Public*
482 *Health* **54**, 167–179.
- 483 9. Timlin MT, Pereira MA, Story M, et al. (2008) Breakfast eating and weight change in
484 a 5-year prospective analysis of breakfast eating and weight change in a 5-year
485 prospective analysis of adolescents: Project EAT (Eating Among Teens). *Pediatrics*
486 **121**, 638–645.
- 487 10. Dhurandhar EJ, Dawson J, Alcorn A, et al. (2014) The effectiveness of breakfast
488 recommendations on weight loss: a randomized controlled trial. *Am. J. Clin. Nutr.* **100**,
489 507–513.
- 490 11. Casazza K, Brown A, Astrup A, et al. (2015) Weighing the evidence of common
491 beliefs in obesity research. *Crit. Rev. Food Sci. Nutr.* **55**, 2014–2053.
- 492 12. Dhurandhar EJ (2016) True, true, unrelated? A review of recent evidence for a causal

- 493 influence of breakfast on obesity. *Curr. Opin. Endocrinol. Diabetes Obes.* **23**, 384–
494 388.
- 495 13. van Nassau F, Singh AS, Cerin E, et al. (2014) The Dutch Obesity Intervention in
496 Teenagers (DOiT) cluster controlled implementation trial: intervention effects and
497 mediators and moderators of adiposity and energy balance-related behaviours. *Int. J.*
498 *Behav. Nutr. Phys. Act.* **11**, 158–168.
- 499 14. Albertson AM, Anderson GH, Crockett SJ, et al. (2003) Ready-to-eat cereal
500 consumption: its relationship with BMI and nutrient intake of children aged 4 to 12
501 years. *J. Am. Diet. Assoc.* **103**, 1613–1619.
- 502 15. Albertson AM, Affenito SG, Bauserman R, et al. (2009) The relationship of ready-to-
503 eat cereal consumption to nutrient intake, blood lipids, and body mass index of
504 children as they age through adolescence. *J. Am. Diet. Assoc.* **109**, 1557–1565.
- 505 16. van den Boom A, Serra-Majem L, Ribas L, et al. (2006) The contribution of ready-to-
506 eat cereals to daily nutrient intake and breakfast quality in a Mediterranean setting. *J.*
507 *Am. Coll. Nutr.* **25**, 135–143.
- 508 17. Matthys C, De Henauw S, Bellemans M, et al. (2007) Breakfast habits affect overall
509 nutrient profiles in adolescents. *Public Health Nutr.* **10**, 413–421.
- 510 18. Deshmukh-Taskar PR, Nicklas TA, O’Neil CE, et al. (2010) The relationship of
511 breakfast skipping and type of breakfast consumption with nutrient intake and weight
512 status in children and adolescents: The National Health and Nutrition Examination
513 Survey 1999-2006. *J. Am. Diet. Assoc.* **110**, 869–878.
- 514 19. Grieger JA & Cobiac L (2012) Comparison of dietary intakes according to breakfast
515 choice in Australian boys. *Eur. J. Clin. Nutr.* **66**, 667–672.
- 516 20. Barr SI, DiFrancesco L & Fulgoni III VL (2014) Breakfast consumption is positively
517 associated with nutrient adequacy in Canadian children and adolescents. *Br J Nutr*
518 **112**, 1373–1383.
- 519 21. Fayet-Moore F, Kim J, Sritharan N, et al. (2016) Impact of breakfast skipping and
520 breakfast choice on the nutrient intake and body mass index of Australian children.
521 *Nutrients* **8**, 487–498.
- 522 22. Afeiche MC, Taillie LS, Hopkins S, et al. (2017) Breakfast dietary patterns among
523 Mexican children are related to total-day diet quality. *J. Nutr.* **147**, 404–412.
- 524 23. Rampersaud GC, Pereira MA, Girard BL, et al. (2005) Breakfast habits, nutritional
525 status, body weight, and academic performance in children and adolescents. *J. Am.*

- 526 *Diet. Assoc.* **105**, 743–760.
- 527 24. Barton BA, Eldridge AL, Thompson D, et al. (2005) The relationship of breakfast and
528 cereal consumption to nutrient intake and body mass index: The National Heart, Lung,
529 and Blood Institute Growth and Health Study. *J. Am. Diet. Assoc.* **105**, 1383–1389.
- 530 25. Affenito SG, Thompson DR, Barton BA, et al. (2005) Breakfast consumption by
531 African-American and white adolescent girls correlates positively with calcium and
532 fiber intake and negatively with body mass index. *J. Am. Diet. Assoc.* **105**, 938–945.
- 533 26. Sjöberg A, Hallberg L, Höglund D, et al. (2003) Meal pattern, food choice, nutrient
534 intake and lifestyle factors in The Göteborg Adolescence Study. *Eur. J. Clin. Nutr.* **57**,
535 1569–1578.
- 536 27. Williams P (2007) Breakfast and the diets of Australian children and adolescents: an
537 analysis of data from the 1995 National Nutrition Survey. *Int. J. Food Sci. Nutr.* **58**,
538 201–216.
- 539 28. Nicklas TA, Reger C, Myers L, et al. (2000) Breakfast consumption with and without
540 vitamin-mineral supplement use favorably impacts daily nutrient intake of ninth-grade
541 students. *J. Adolesc. Heal.* **27**, 314–321.
- 542 29. Nicklas TA, O’Neil C & Myers L (2004) The importance of breakfast consumption to
543 nutrition of children, adolescents, and young adults. *Nutr. Today* **39**, 30–39.
- 544 30. Sampson A, Dixit S, Meyers A, et al. (1995) The nutritional impact of breakfast
545 consumption on the diets of inner-city African-American elementary school children.
546 *J. Natl. Med. Assoc.* **87**, 195–202.
- 547 31. Antonogeorgos G, Panagiotakos DB, Papadimitriou A, et al. (2012) Breakfast
548 consumption and meal frequency interaction with childhood obesity. *Pediatr. Obes.* **7**,
549 65–72.
- 550 32. Berkey C, Rockett H, Gillman M, et al. (2003) Longitudinal study of skipping
551 breakfast and weight change in adolescents. *Int. J. Obes.* **27**, 1258–1266.
- 552 33. O’Neil CE, Byrd-Bredbenner C, Hayes D, et al. (2014) The role of breakfast in health:
553 definition and criteria for a quality breakfast. *J. Acad. Nutr. Diet.* **114**, S8–S26.
- 554 34. Betts JA, Chowdhury EA, Gonzalez JT, et al. (2016) Is breakfast the most important
555 meal of the day? *Proc. Nutr. Soc.*, 1–11.
- 556 35. Public Health England (2014) National Diet and Nutrition Survey Results from Years
557 1 , 2 , 3 and 4 (combined) of the Rolling Programme (2008/2009-2011/2012). [Bates
558 B, Lennox A, Prentice A, et al., editors].

- 559 [https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/594361/](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/594361/NDNS_Y1_to_4_UK_report_full_text_revised_February_2017.pdf)
560 [NDNS_Y1_to_4_UK_report_full_text_revised_February_2017.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/594361/NDNS_Y1_to_4_UK_report_full_text_revised_February_2017.pdf) (accessed May
561 2017).
- 562 36. NatCen Social Research, MRC Human Nutrition Research & University College
563 London (2014) *National Diet and Nutrition Survey Years 1-4, 2008/09-2011/12*
564 *[computer file]. 6th Edition*. Colchester, Essex: UK Data Archive [distributor]; SN:
565 6533, <http://dx.doi.org/10.5255/UKDA-SN-6533-5>.
- 566 37. Lennox A, Fitt E, Whitton C, et al. (2014) *Appendix A to National Diet and Nutrition*
567 *Survey. Results from Years 1-4 (combined) of the Rolling Programme (2008/2009 -*
568 *2011/12). Dietary data collection and editing*. London: Public Health England.
- 569 38. Bates B, Lennox A, Prentice A, et al. (2014) *P2753 National Diet and Nutrition*
570 *Survey (NDNS) Year 4 Interviewer Project Instructions*. Colchester, Essex: UK Data
571 Archive [distributor]; SN: 6533, <http://dx.doi.org/10.5255/UKDA-SN-6533-5>.
- 572 39. Public Health England (2014) *National Diet and Nutrition Survey Rolling Programme*
573 *Years 1-4 2008/09-2011/12 User Guide for UK Core Sample Data*. Colchester, Essex:
574 UK Data Archive [distributor]; SN: 6533, [http://dx.doi.org/10.5255/UKDA-SN-6533-](http://dx.doi.org/10.5255/UKDA-SN-6533-5)
575 [5](http://dx.doi.org/10.5255/UKDA-SN-6533-5).
- 576 40. Food Standards Agency (2002) *McCance and Widdowson's The Composition of*
577 *Foods*. 6th ed. Cambridge: Royal Society of Chemistry.
- 578 41. SACN (2015) Carbohydrates and Health.
579 <https://www.gov.uk/government/publications/sacn-carbohydrates-and-health-report>
580 (accessed May 2017).
- 581 42. Englyst HN, Quigley ME & Hudson GJ (1994) Determination of dietary fibre as non-
582 starch polysaccharides with gas-liquid chromatographic, high-performance liquid
583 chromatographic or spectrophotometric measurement of constituent sugars. *Analyst*
584 **119**, 1497–1509.
- 585 43. SACN (2011) Dietary Reference Values for Energy 2011.
586 [https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/339317/](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/339317/SACN_Dietary_Reference_Values_for_Energy.pdf)
587 [SACN_Dietary_Reference_Values_for_Energy.pdf](https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/339317/SACN_Dietary_Reference_Values_for_Energy.pdf) (accessed May 2017).
- 588 44. COMA (1991) *Dietary Reference Values for Food Energy and Nutrients for the*
589 *United Kingdom*. London: The Stationery Office.
- 590 45. Pot GK, Richards M, Prynne CJ, et al. (2014) Development of the Eating Choices
591 Index (ECI): a four-item index to measure healthiness of diet. *Public Health Nutr.* **17**,

- 592 1–7.
- 593 46. Vanderpump MPJ, Lazarus JH, Smyth PP, et al. (2011) Iodine status of UK
594 schoolgirls: a cross-sectional survey. *Lancet* **377**, 2007–2012.
- 595 47. Cole T, Freeman J & Preece M (1995) Body mass index reference curves for the UK,
596 1990. *Arch. Dis. Child.* **73**, 25–29.
- 597 48. Public Health England (2014) *P2752 National Diet and Nutrition Survey Year 4 CAPI*
598 *Interviewer Schedule*. Colchester, Essex: UK Data Archive [distributor]; SN: 6533,
599 <http://dx.doi.org/10.5255/UKDA-SN-6533-5>.
- 600 49. McCrory MA, Hajduk CL & Roberts SB (2002) Procedures for screening out
601 inaccurate reports of dietary energy intake. *Public Health Nutr.* **5**, 873–882.
- 602 50. Rogan JC & Keselman HJ (1977) Is the ANOVA F-test robust to variance
603 heterogeneity when sample sizes are equal?: An investigation via a coefficient of
604 variation. *Am. Educ. Res. J.* **14**, 493–498.
- 605 51. Feise RJ (2002) Do multiple outcome measures require p-value adjustment? *BMC*
606 *Med. Res. Methodol.* **2**, 8–11.
- 607 52. Cashman KD (2002) Calcium intake, calcium bioavailability and bone health. *Br. J.*
608 *Nutr.* **87 Suppl 2**, S169–S177.
- 609 53. Monteagudo C, Palacin-Arce A, del Mar Bibiloni M, et al. (2013) Proposal for a
610 Breakfast Quality Index (BQI) for children and adolescents. *Public Health Nutr.* **16**,
611 639–644.
- 612 54. Miech RA, Kumanyika SK, Stettler N, et al. (2006) Trends in the association of
613 poverty with overweight among US adolescents, 1971–2004. *J. Am. Med. Assoc.* **295**,
614 2385–2393.
- 615 55. Leidy HJ & Racki EM (2010) The addition of a protein-rich breakfast and its effects
616 on acute appetite control and food intake in ‘breakfast-skipping’ adolescents. *Int. J.*
617 *Obes.* **34**, 1125–1133.
- 618 56. Kral TVE, Whiteford LM, Heo M, et al. (2011) Effects of eating breakfast compared
619 with skipping breakfast on ratings of appetite and intake at subsequent meals in 8- to
620 10-y-old children. *Am. J. Clin. Nutr.* **93**, 284–291.
- 621 57. Berkey CS, Rockett HRH, Gillman MW, et al. (2003) Longitudinal study of skipping
622 breakfast and weight change in adolescents. *Int. J. Obes.* **27**, 1258–1266.
- 623 58. Schembre SM, Wen CK, Davis JN, et al. (2013) Eating breakfast more frequently is
624 cross-sectionally associated with greater physical activity and lower levels of adiposity

- 625 in overweight Latina and African American girls. *Am. J. Clin. Nutr.* **98**, 275–281.
- 626 59. Betts JA, Richardson JD, Chowdhury EA, et al. (2014) The causal role of breakfast in
627 energy balance and health : a randomized controlled trial in lean adults. *Am. J. Clin.*
628 *Nutr.* **100**, 539–547.
- 629 60. Hoyland A, McWilliams K, Duff R, et al. (2012) Breakfast consumption in UK
630 schoolchildren and provision of school breakfast clubs. *Nutr. Bull.* **37**, 232–240.
- 631 61. Adolphus K, Lawton CL & Dye L (2013) The effects of breakfast on behavior and
632 academic performance in children and adolescents. *Front. Hum. Neurosci.* **7**, 425.
- 633 62. Cohen B, Evers S, Manske S, et al. (2003) Smoking, physical activity and breakfast
634 consumption among secondary school students in a southwestern Ontario community.
635 *Can. J. Public Heal.* **94**, 41–44.
- 636 63. Lattimore PJ & Halford JCG (2003) Adolescence and the diet-dieting disparity:
637 healthy food choice or risky health behaviour? *Br. J. Health Psychol.* **8**, 451–463.
- 638 64. Crowley SJ, Acebo C & Carskadon MA (2007) Sleep, circadian rhythms, and delayed
639 phase in adolescence. *Sleep Med.* **8**, 602–612.
- 640 65. Roenneberg T, Kuehnle T, Pramstaller PP, et al. (2004) A marker for the end of
641 adolescence. *Curr. Biol.* **14**, R1038–R1039.
- 642 66. Fleig D & Randler C (2009) Association between chronotype and diet in adolescents
643 based on food logs. *Eat. Behav.* **10**, 115–118. Elsevier Ltd.
- 644 67. Roenneberg T, Kuehnle T, Juda M, et al. (2007) Epidemiology of the human circadian
645 clock. *Sleep Med. Rev.* **11**, 429–438.
- 646 68. Schaal S, Peter M & Randler C (2010) Morningness-eveningness and physical activity
647 in adolescents. *Int. J. Sport Exerc. Psychol.* **8**, 147–159.
- 648 69. Coulthard JD & Pot GK (2016) The timing of the evening meal: how is this associated
649 with weight status in UK children? *Br. J. Nutr.* **115**, 1616–1622.
- 650 70. Kant AK & Graubard BI (2015) Within-person comparison of eating behaviors, time
651 of eating, and dietary intake on days with and without breakfast : NHANES 2005-
652 2010. *Am. J. Clin. Nutr.* **102**, 661–670.
- 653 71. Gorgulho BM, Pot GK, Sarti FM, et al. (2016) Indices for the assessment of nutritional
654 quality of meals: a systematic review. *Br. J. Nutr.* **115**, 2017–2024.
- 655 72. Brown AW, Brown MMB & Allison DB (2013) Belief beyond the evidence: using the
656 proposed effect of breakfast on obesity to show 2 practices that distort scientific

657 evidence. *Am. J. Clin. Nutr.* **98**, 1298–1308.

658

659 **Tables and Figures**

660 **Table 1.** Characteristics of children by breakfast eating habit showing sex, age, ethnicity,
661 weight status and household income (*n* 1686) (Numbers and percentages; mean values and
662 standard deviations)

663

664 **Table 2.** Comparison of number and percentage of children below RNI and LRNI for folate,
665 vitamin C, calcium, iron and iodine depending on breakfast habit* (*n* 1686) (Numbers and
666 percentages)

667

668 **Table 3.** Comparison of daily nutrient intakes for children aged 4-18 years with different
669 breakfast habits (Mean values and standard deviations)

670

671 **Table 4. (a)** Within-person difference in daily nutrient intakes for days on which breakfast
672 consumed compared to days on which breakfast not consumed, 4-18 year olds (*n* 879) (Mean
673 values and standard deviations; mean differences and 95% confidence intervals)

674

675 **Table 4. (b)** Within-person difference in daily nutrient intakes for days on which breakfast
676 consumed compared to days on which breakfast not consumed, 4-10 year olds (*n* 384) (Mean
677 values and standard deviations; mean differences and 95% confidence intervals)

678

679 **Table 4. (c)** Within-person difference in daily nutrient intakes for days on which breakfast
680 consumed compared to days on which breakfast not consumed, 11-18 year olds (*n* 495)
681 (Mean values and standard deviations; mean differences and 95% confidence intervals)

682

683 **Table 5.** Within-person difference in daily nutrient intakes for days on which breakfast
684 consumed compared to days on which breakfast not consumed, no Saturdays or Sundays, 4-
685 18 year olds (*n* 365) (Mean values and standard deviations; mean differences and 95%
686 confidence intervals)

587
588
589**Table 1.** Characteristics of children by breakfast eating habit showing sex, age, ethnicity, weight status and household income (*n* 1686)
(Numbers and percentages; mean values and standard deviations)

		Breakfast no diary days		Breakfast on at least one but not all diary days		Breakfast every diary day		P*
		<i>n</i>	row %	<i>n</i>	row %	<i>n</i>	row %	
Total		289	17.1	879	52.1	518	30.7	
Sex								
	Male	124	14.5	443	51.6	291	33.9	0.001
	Female	165	19.9	436	52.7	227	27.4	
Age								
	4-10 years	52	6.5	384	47.9	366	45.6	<0.001
	11-18 years	237	26.8	495	56.0	152	17.2	
Ethnicity								
	White	240	16.5	757	52.1	455	31.3	0.168
	Non-white	49	20.9	122	52.1	63	26.9	
Weight Statust								
	Normal	195	17.7	553	50.2	353	32.1	0.258
	Overweight	33	14.9	120	54.3	68	30.8	
	Obese	52	17.5	167	56.2	78	26.3	
		Mean	SD	Mean	SD	Mean	SD	P†
Household income (£)§		23 587	16 374	25 108	16 998	28 194	18 349	0.001

*Differences between children with different breakfast habits analysed using Pearson chi-square test. P values of ≤ 0.01 considered significant, shown in bold.

†Differences between children with different breakfast habits analysed using ANOVA. P value of ≤ 0.01 considered significant, shown in bold.

‡The BMI measurements for the children were compared with British 1990 growth reference (UK90) charts to assess whether children were normal weight, overweight (85th centile cut-off) or obese (95th centile cut-off) *n* 1619 (67 missing values).

§Equivalised household income (using McClements equivalence scale), *n* 1470 (216 missing values).

590

591
592
593**Table 2.** Comparison of number and percentage of children below RNI and LRNI for folate, vitamin C, calcium, iron and iodine depending on breakfast habit* (*n* 1686)
(Numbers and percentages)

		Meeting RNI/LRNI	Breakfast no diary days		Breakfast on at least one but not all diary days		Breakfast every diary day	
			<i>n</i>	%	<i>n</i>	%	<i>n</i>	%
Folate	RNI	Yes	125	43.3	588	66.9	426	82.2
		No	164	56.7	291	33.1	92	17.8
	LRNI	Yes	268	92.7	858	97.6	518	100.0
		No	21	7.3	21	2.4	0	0.0
Vitamin C	RNI	Yes	213	73.7	762	86.7	470	90.7
		No	76	26.3	117	13.3	48	9.3
	LRNI	Yes	284	98.3	876	99.7	517	99.8
		No	5	1.7	3	0.3	1	0.2
Calcium	RNI	Yes	92	31.8	470	53.5	392	75.7
		No	197	68.2	409	46.5	126	24.3
	LRNI	Yes	234	81	819	93.2	503	97.1
		No	55	19.0	60	6.8	15	2.9
Iron	RNI	Yes	56	19.4	353	40.2	303	58.5
		No	233	80.6	526	59.8	215	41.5
	LRNI	Yes	198	68.5	751	85.4	495	95.6
		No	91	31.5	128	14.6	23	4.4
Iodine	RNI	Yes	90	31.1	400	45.5	332	64.1
		No	199	68.9	479	54.5	186	35.9
	LRNI	Yes	227	78.5	793	90.2	501	96.7
		No	62	21.5	86	9.8	17	3.3

RNI, reference nutrient intake (as set by Committee on Medical Aspects of Food Policy (COMA)), LRNI, lower reference nutrient intake (as set by COMA).

*P-values for chi-square analysis all <0.001, except for Vitamin C LRNI where results invalid as 50% of cells had expected count <5.

594

595
596
597
598**Table 3.** Comparison of daily nutrient intakes for children aged 4-18 years with different breakfast habits
(Mean values and standard deviations)

	Breakfast no diary days (n 289)		Breakfast at least one but not all diary days (n 879)		Breakfast every diary day (n 518)		ANOVA (n 1686)	ANCOVA§ (n 1416)
	Mean	SD	Mean	SD	Mean	SD	P†	P†
Energy (%EAR)	71.0	22.1	82.2	22.5	92.4	21.3	<0.001	0.009
Protein (%Energy)	14.9	3.5	14.6	2.6	14.8	2.5	0.309	0.041
Fat (%Energy)	34.2	5.3	33.8	4.8	33.1	4.6	0.005	0.005
CHO (%Energy)	49.7	6.1	51.3	5.3	52.1	4.8	<0.001	0.010
NMES (%Energy)	15.5	7.1	15.3	5.8	14.5	5.4	0.030	0.034
Fibre (g/1000kcalEnergy)	6.5	1.9	7.0	1.8	7.5	1.9	<0.001	<0.001
Folate (mcg/1000kcalEnergy)†	114	39	124	39	134	41	<0.001	<0.001
Vitamin C (mg/1000kcalEnergy)†	44.6	29.4	51.5	34.8	56.7	32.6	<0.001	0.472
Sodium (mg/1000kcalEnergy)*	1262	314	1255	284	1198	220	0.027	0.127
Calcium (mg/1000kcalEnergy)*	426	132	471	140	532	143	<0.001	<0.001
Iron (mg/1000kcalEnergy)	5.2	1.5	5.5	1.2	5.8	1.3	<0.001	<0.001
Iodine (mcg/1000kcalEnergy)†	68.0	30.8	77.3	37.2	89.8	35.8	<0.001	<0.001

EAR, Estimated Average Requirement of energy as set by the Scientific Advisory Committee on Nutrition; CHO, carbohydrate; NMES, non-milk extrinsic sugars.. *Square root transformation applied to calculate significance; means and standard deviations shown for untransformed variables.

† Natural logarithm transformation applied to calculate significance; means and standard deviations shown for untransformed variables.

†P value of ≤ 0.01 considered significant, shown in bold.

§ANOVA adjusted for the covariates: age, sex, body mass index, ethnicity (white/non white) and equivalised household income (using McClements equivalence scale); sample size reduced by 270 cases due to missing values for BMI and equivalised income.

||Due to inequality of variances for these variables ANOVA and ANCOVA performed after random sample selection to form three equal groups of 245 cases (n 735).

599

700
701
702**Table 4. (a)** Within-person difference in daily nutrient intakes for days on which breakfast consumed compared to days on which breakfast not consumed, 4-18 year olds (*n* 879) (Mean values and standard deviations; mean differences and 95% confidence intervals)

	Non-breakfast days		Breakfast days		T-test paired difference			Adjusted paired difference§		
	Mean	SD	Mean	SD	Mean difference	95% CI	P†	Mean difference	95% CI	P†
Energy (kcal)	1629	550	1721	535	92	60, 124	<0.001	87	52, 121	<0.001
Protein (%Energy)	15.1	3.8	14.5	3.2	-0.57	-0.85, -0.30	<0.001	-0.66	-0.96, -0.35	<0.001
Fat (%Energy)	33.9	6.6	33.3	5.8	-0.58	-1.07, -0.10	0.018	-0.49	-1.02, 0.05	0.075
CHO (%Energy)	50.7	7.6	52.0	6.3	1.29	0.76, 1.82	<0.001	1.28	0.71, 1.86	<0.001
NMES (%Energy)	15.3	7.7	15.1	7.0	-0.26	-0.79, 0.27	0.339	-0.21	-0.78, 0.37	0.483
Fibre (g/1000kcalEnergy)	7.0	2.4	7.1	2.1	0.11	-0.06, 0.27	0.215	0.10	-0.08, 0.27	0.272
Folate (mcg/1000kcalEnergy)†	123	50	129	50	5.9	0.03, 0.09	<0.001	5.2	0.03, 0.09	<0.001
Vitamin C (mg/1000kcalEnergy)†	52.1	43.2	52.7	39.5	0.6	-0.002, 0.114	0.057	0.3	-0.03, 0.10	0.257
Sodium (mg/1000kcalEnergy)*	1283	371	1237	348	-47	-1.03, -0.22	0.002	-55	-1.19, -0.30	0.001
Calcium (mg/1000kcalEnergy)*	455	173	491	165	36	0.64, 1.15	<0.001	36	0.60, 1.16	<0.001
Iron (mg/1000kcalEnergy)	5.5	1.6	5.7	1.6	0.18	0.05, 0.30	0.004	0.17	0.03, 0.30	0.016
Iodine (mcg/1000kcalEnergy)†	77.2	56.3	79.2	41.5	1.9	0.02, 0.09	0.001	1.3	0.01, 0.09	0.007

CHO, carbohydrate; NMES, non-milk extrinsic sugars.

*Square root transformation applied to calculate significance; means, standard deviations and mean differences shown for untransformed variables.

† Natural logarithm transformation applied to calculate significance; means, standard deviations and mean differences shown for untransformed variables.

†P value of ≤ 0.01 considered significant, shown in bold.§Repeated measures ANOVA adjusted for the covariates: age, sex, BMI, ethnicity (white/non white) and equivalised household income (using McClements equivalence scale). Sample size reduced by 145 cases (*n* 734) due to missing values for BMI and equivalised income.703
704

705
706
707**Table 4. (b)** Within-person difference in daily nutrient intakes for days on which breakfast consumed compared to days on which breakfast not consumed, 4-10 year olds (*n* 384) (Mean values and standard deviations; mean differences and 95% confidence intervals)

	Non-breakfast days		Breakfast days		T-test paired difference			Adjusted paired difference§		
	Mean	SD	Mean	SD	Mean difference	95% CI	P†	Mean difference	95% CI	P†
Energy (kcal)	1479	443	1525	355	46	6, 85	0.023	47	4, 90	0.032
Protein (%Energy)	14.8	3.4	14.6	2.8	-0.26	-0.63, 0.10	0.159	-0.36	-0.76, 0.03	0.073
Fat (%Energy)	33.9	6.5	33.1	5.3	-0.86	-1.54, -0.18	0.014	-0.77	-1.52, -0.01	0.048
CHO (%Energy)	51.2	7.1	52.3	5.8	1.14	0.39, 1.89	0.003	1.15	0.35, 1.94	0.005
NMES (%Energy)	15.1	7.3	14.7	5.7	-0.40	-1.08, 0.29	0.255	-0.56	-1.32, 0.19	0.142
Fibre (g/1000kcalEnergy)	7.0	2.3	7.3	2.0	0.37	0.14, 0.61	0.002	0.42	0.17, 0.67	0.001
Folate (mcg/1000kcalEnergy)†	125	46	133	45	8.5	0.04, 0.12	<0.001	8.4	0.04, 0.12	<0.001
Vitamin C (mg/1000kcalEnergy)†	56.0	43.5	59.4	37.6	3.4	0.54, 0.21	0.001	3.5	0.05, 0.23	0.002
Sodium (mg/1000kcalEnergy)*	1259	374	1213	299	-46	-1.09, 0.04	0.069	-46	-1.15, 0.08	0.085
Calcium (mg/1000kcalEnergy)*	494	190	524	176	29	0.34, 1.12	<0.001	33	0.38, 1.24	<0.001
Iron (mg/1000kcalEnergy)	5.5	1.6	5.7	1.5	0.26	0.08, 0.44	0.005	0.22	0.03, 0.42	0.027
Iodine (mcg/1000kcalEnergy)†	85.8	50.1	90.2	47.2	4.4	0.02, 0.13	0.005	5.1	0.03, 0.14	0.005

CHO, carbohydrate; NMES, non-milk extrinsic sugars.

*Square root transformation applied to calculate significance; means, standard deviations and mean differences shown for untransformed variables.

† Natural logarithm transformation applied to calculate significance; means, standard deviations and mean differences shown for untransformed variables.

‡P value of ≤0.01 considered significant, shown in bold.

§Repeated measures ANOVA adjusted for the covariates: age, sex, BMI, ethnicity (white/non white) and equivalised household income (using McClements equivalence scale). Sample size reduced by 62 cases (*n* 322) due to missing values for BMI and equivalised income.708
709
710

711
712
713**Table 4. (c)** Within-person difference in daily nutrient intakes for days on which breakfast consumed compared to days on which breakfast not consumed, 11-18 year olds (*n* 495) (Mean values and standard deviations; mean differences and 95% confidence intervals)

	Non-breakfast days		Breakfast days		T-test paired difference			Adjusted paired difference§		
	Mean	SD	Mean	SD	Mean difference	95% CI	P†	Mean difference	95% CI	P†
Energy (kcal)	1746	595	1874	598	128	80, 175	<0.001	118	66, 169	<0.001
Protein (%Energy)	15.3	4.1	14.5	3.4	-0.82	-1.22, -0.42	<0.001	-0.88	-1.33, -0.44	<0.001
Fat (%Energy)	33.8	6.7	33.4	6.1	-0.37	-1.04, 0.30	0.281	-0.27	-1.00, 0.47	0.478
CHO (%Energy)	50.3	7.9	51.7	6.6	1.41	0.67, 2.15	<0.001	1.39	0.58, 2.20	0.001
NMES (%Energy)	15.5	8.0	15.4	7.8	-0.15	-0.93, 0.63	0.704	0.08	-0.76, 0.91	0.859
Fibre (g/1000kcalEnergy)	7.0	2.5	6.9	2.2	-0.10	-0.33, 0.13	0.383	-0.16	-0.40, 0.09	0.214
Folate (mcg/1000kcalEnergy)†	122	52	126	54	4.0	0.01, 0.09	0.023	2.7	-0.002, 0.085	0.062
Vitamin C (mg/1000kcalEnergy)†	49.1	42.7	47.4	40.1	-1.7	-0.09, 0.08	0.917	-2.1	-0.13, 0.04	0.289
Sodium (mg/1000kcalEnergy*)	1302	368	1255	380	-47	-1.28, -0.14	0.015	-61	-1.53, -0.28	0.005
Calcium (mg/1000kcalEnergy*)	424	152	466	152	42	0.68, 1.36	<0.001	38	0.57, 1.31	<0.001
Iron (mg/1000kcalEnergy)	5.5	1.6	5.6	1.6	0.11	-0.05, 0.28	0.178	0.12	-0.07, 0.30	0.204
Iodine (mcg/1000kcalEnergy)†	70.6	59.9	70.5	34.1	0.0	-0.003, 0.092	0.063	-1.8	-0.02, 0.08	0.258

CHO, carbohydrate; NMES, non-milk extrinsic sugars.

*Square root transformation applied to calculate significance; means, standard deviations and mean differences shown for untransformed variables.

† Natural logarithm transformation applied to calculate significance; means, standard deviations and mean differences shown for untransformed variables.

†P value of ≤0.01 considered significant, shown in bold.

§Repeated measures ANOVA adjusted for the covariates: age, sex, BMI, ethnicity (white/non white) and equivalised household income (using McClements equivalence scale). Sample size reduced by 83 cases (*n* 412) due to missing values for BMI and equivalised income.714
715

716
717
718
719

Table 5. Within-person difference in daily nutrient intakes for days on which breakfast consumed compared to days on which breakfast not consumed, no Saturdays or Sundays, 4-18 year olds (*n* 365)
(Mean values and standard deviations; mean differences and 95% confidence intervals)

	Non-breakfast days		Breakfast days		T-test paired difference			Adjusted paired difference§		
	Mean	SD	Mean	SD	Mean difference	95% CI	P†	Mean difference	95% CI	P†
Energy (kcal)	1540	619	1688	571	148	90, 207	<0.001	158	93, 223	<0.001
Protein (%Energy)	15.2	4.4	14.6	4.0	-0.52	-1.06, 0.02	0.061	-0.61	-1.22, 0.01	0.052
Fat (%Energy)	33.4	7.8	33.5	7.1	0.11	-0.77, 0.99	0.807	0.13	-0.88, 1.13	0.802
CHO (%Energy)	51.1	8.2	51.7	7.7	0.56	-0.38, 1.49	0.240	0.65	-0.40, 1.71	0.225
NMES (%Energy)	15.2	8.6	15.3	8.0	0.13	-0.82, 1.09	0.782	0.38	-0.69, 1.46	0.484
Fibre (g/1000kcalEnergy)	7.2	2.8	7.0	2.5	-0.21	-0.53, 0.10	0.186	-0.29	-0.62, 0.04	0.080
Folate (mcg/1000kcalEnergy)†	120	53	127	52	7.7	0.03, 0.13	0.003	9.0	0.03, 0.14	0.004
Vitamin C (mg/1000kcalEnergy)*	51.3	48.4	53.7	45.2	2.4	-0.05, 0.58	0.100	1.4	-0.19, 0.52	0.369
Sodium (mg/1000kcalEnergy)*	1270	412	1236	411	-35	-1.21, 0.26	0.206	-41	-1.39, 0.24	0.165
Calcium (mg/1000kcalEnergy)*	434	181	475	172	41	0.59, 1.52	<0.001	43	0.60, 1.64	<0.001
Iron (mg/1000kcalEnergy)	5.4	1.9	5.6	1.7	0.16	-0.06, 0.37	0.149	0.15	-0.08, 0.38	0.208
Iodine (mcg/1000kcalEnergy)†	70.6	44.3	75.1	44.4	4.5	0.02, 0.15	0.007	5.5	0.03, 0.17	0.008

CHO, carbohydrate; NMES, non-milk extrinsic sugars.

*Square root transformation applied to calculate significance; means, standard deviations and mean differences shown for untransformed variables.

†Natural logarithm transformation applied to calculate significance; means, standard deviations and mean differences shown for untransformed variables.

‡P value of ≤ 0.01 considered significant, shown in bold.

§Repeated measures ANOVA adjusted for the covariates: age, sex, BMI, ethnicity (white/non white) and equivalised household income (using McClements equivalence scale). Sample size reduced by 67 cases (*n* 298) due to missing values for BMI and equivalised income.

720
721