



Heikkil, K; Rissanen, H; Helivaara, M; Knekt, P; Mki, M; Kaukinen, K (2017) Associations of tissue transglutaminase antibody seropositivity with coronary heart disease: Findings from a prospective cohort study. *Nutrition, metabolism, and cardiovascular diseases*. ISSN 0939-4753 DOI: <https://doi.org/10.1016/j.numecd.2017.06.005>

Downloaded from: <http://researchonline.lshtm.ac.uk/4258861/>

DOI: [10.1016/j.numecd.2017.06.005](https://doi.org/10.1016/j.numecd.2017.06.005)

#### Usage Guidelines


Please refer to usage guidelines at <http://researchonline.lshtm.ac.uk/policies.html> or alternatively contact [researchonline@lshtm.ac.uk](mailto:researchonline@lshtm.ac.uk).

Available under license: <http://creativecommons.org/licenses/by-nc-nd/2.5/>

# Accepted Manuscript

Associations of Tissue Transglutaminase Antibody Seropositivity with Coronary Heart Disease: Findings from a Prospective Cohort Study

Katriina Heikkila, Harri Rissanen, Markku Heliövaara, Paul Knekt, Markku Mäki, Katri Kaukinen



PII: S0939-4753(17)30126-6

DOI: [10.1016/j.numecd.2017.06.005](https://doi.org/10.1016/j.numecd.2017.06.005)

Reference: NUMECD 1736

To appear in: *Nutrition, Metabolism and Cardiovascular Diseases*

Received Date: 5 January 2017

Revised Date: 18 April 2017

Accepted Date: 8 June 2017

Please cite this article as: Heikkila K, Rissanen H, Heliövaara M, Knekt P, Mäki M, Kaukinen K, Associations of Tissue Transglutaminase Antibody Seropositivity with Coronary Heart Disease: Findings from a Prospective Cohort Study, *Nutrition, Metabolism and Cardiovascular Diseases* (2017), doi: 10.1016/j.numecd.2017.06.005.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Associations of Tissue Transglutaminase Antibody Seropositivity with Coronary Heart Disease: Findings from a Prospective Cohort Study**

Katriina Heikkilä<sup>1,2</sup>, Harri Rissanen<sup>2</sup>, Markku Heliövaara<sup>2</sup>, Paul Knekt<sup>2</sup>, Markku Mäki<sup>3</sup>,  
Katri Kaukinen<sup>3,4</sup>

<sup>1</sup> Department of Health Services Research and Policy, London School of Hygiene and Tropical Medicine, London, UK

<sup>2</sup> National Institute for Health and Welfare, Helsinki, Finland

<sup>3</sup> Tampere Centre for Child Health Research, University of Tampere and Tampere University Hospital, Finland

<sup>4</sup> Department of Internal Medicine, Tampere University Hospital, Tampere, Finland

**Corresponding author:**

Dr Katriina Heikkilä

Department of Health Services Research and Policy, Faculty of Public Health and Policy,

London School of Hygiene and Tropical Medicine

15-17 Tavistock Place, London WC1H 9SH, United Kingdom

E: [katriina.heikkila@lshtm.ac.uk](mailto:katriina.heikkila@lshtm.ac.uk), T: +44 (0)20 7958 8299, F: +44 (0)20 7927 2701

**Abstract**

**Background and aims** Clinical experience and observational studies suggest that individuals with coeliac disease are at increased risk of coronary heart disease (CHD), but the precise mechanism for this is unclear. Laboratory studies suggest that it may relate to tissue transglutaminase antibodies (tTGAs). Our aim was to examine whether seropositivity for tTGA and endomysial antibodies (EMAs) are associated with incident CHD in humans.

**Methods** We used data from Mini-Finland Health Survey, a prospective cohort study of Finnish men and women aged 35-80 at study baseline 1978-80. TTGA and EMA seropositivities were ascertained from baseline blood samples and incident CHD events were identified from national hospitalisation and death registers. Poisson regression was used to examine the associations between antibody seropositivity and incident CHD.

**Results** Of 6,887 men and women 562 were seropositive for tTGAs and 72 for EMAs. During a median follow-up of 26 years 2,367 individuals experienced a CHD event. We found no clear evidence for an association between tTGA positivity (hazard ratio, HR: 1.04, 95% confidence interval, CI: 0.83, 1.30) or EMA positivity (HR: 1.16, 95% CI: 0.77, 1.74) and incident CHD, once pre-existing CVD and known CHD risk factors had been adjusted for.

**Conclusion** We found no clear evidence for an association of tTGA or EMA seropositivity with incident CHD outcomes, suggesting that tTG autoimmunity is unlikely to be the biological link between coeliac disease and CHD.

**Keywords:** coronary heart disease, CHD, tissue transglutaminase antibody, tTGA, endomysial antibody, EMA, cohort study

## Introduction

Coeliac disease is an autoimmune-mediated condition triggered by the ingestion of gluten, a storage protein in wheat, rye and barley[1] Prolamine peptides from these cereals are deamidated by tissue transglutaminase (tTG) in the small intestine and in genetically susceptible individuals the deamidated peptides activate a T-cell driven immune response and production of immunoglobulin A class autoantibodies against tTG [2]. When this autoimmune reaction become chronic, it leads to small intestinal mucosal villous atrophy and crypt hyperplasia, which are currently the diagnostic gold standard for coeliac disease [3].

Coeliac disease can be effectively managed by excluding gluten from the patients' diet, thus stopping the autoimmune response and reversing the mucosal damage [4]. Individuals with coeliac disease who consume gluten, either because they have not been diagnosed or because their adherence to a gluten-free diet is poor, have gluten-induced elevated circulating concentrations of tissue transglutaminase antibodies (tTGAs) and endomysial antibodies (EMAs) [5, 6]. These autoantibodies have been used as serological markers for unrecognised coeliac disease in general population-based screening studies, the findings of which suggest that clinically unrecognised coeliac disease is more common than was previously thought [7-10]: it has been estimated that in addition to the 0.1-0.6% of the population with diagnosed coeliac disease, unrecognised coeliac disease may be present in up to 2% of the population in high income countries [11-13] .

Clinically diagnosed coeliac disease co-occurs with various comorbidities, including neurological and liver abnormalities [14], certain cancers [15] and bone fractures [16], and appears to be associated with an increased risk of cardiovascular outcomes, including coronary heart disease (CHD) [17]. However, the exact mechanism for these associations is

unclear. Laboratory studies have pointed to tissue transglutaminase autoimmunity as one potential link between coeliac and cardiovascular diseases. Findings from *in-vivo* and *in-vitro* studies suggest that tTG has the ability to enhance angiogenesis and that, by contrast, tTGAs can inhibit blood vessel formation. [18-20]. It is not known, however, whether these findings translate into humans. To explore this, we have examined the associations of tTGA and EMA seropositivity with incident coronary heart disease (CHD) using data from Mini-Finland Health Survey, a prospective population-based cohort study of over 6,000 Finnish men and women.

## **Materials and methods**

### *Participants*

Details of the design and data collection in Mini-Finland Health Survey have been reported previously and are provided in the Online Supplement [21]. Briefly, 7,217 men and women (90% of those recruited) aged 35-80 attended a baseline examination in 1978-80. The analyses presented here are based on 6,887 participants with complete data on serology, CHD outcomes and covariates (Online Supplement, Figure S1. Participant flow chart).

### *Exposures and outcomes*

Seropositivity for tTGAs and EMAs were determined from baseline serum samples. Details of the assays have been reported previously and are provided in the Online Supplement [22]. All serum samples were tested for tTGA (Eu-tTG umana IGA, Eurospital S.p.A., Trieste, Italy; abbreviated as Eu-tTG). Positive samples were re-tested using another tTGA test (Celikey Tissue Transglutaminase IgA Antibody Assay, Phadia, Freiburg, Germany; abbreviated as Celikey-tTG) and an EMA test (indirect immunofluorescence, in-house test,

Coeliac Disease Study Group, Tampere University and Tampere University Hospital; abbreviated as EMA).

Incident CHD events were ascertained from national hospitalisation and death registers; baseline cardiovascular disease (CVD) was ascertained from hospitalisation records and baseline health examination. Two outcome measures were used: any CHD (CHD death, non-fatal MI or angina) and hard endpoint CHD (coronary death or non-fatal MI, excluding angina). Full details of the assessment of outcomes and covariates are provided in the Online Supplement.

#### *Statistical analyses*

We estimated the incidence of CHD outcomes per 1000 person years, and used Cox proportional hazards regression to model the associations between seropositivity and incident CHD. Full details of the statistical methods are provided in the Online Supplement. Multivariable-adjusted Cox models were adjusted for age (as the timescale in the model), sex (binary), alcohol intake (grams of ethanol per week: 0, 1-49, 50-249,  $\geq 250$ ), years of education (0-8, 9-12,  $\geq 12$ ), BMI ( $\text{kg/m}^2$ :  $< 18.5$ , 18.5-24.9, 25-29.9, 30-34.9,  $\geq 35$ ), smoking (never, quit, pipe or cigars or 1-19 cigarettes per day,  $\geq 20$  cigarettes per day), systolic and diastolic blood pressure (mmHg), total and HDL cholesterol, baseline diabetes and homoeostasis model assessment-estimated insulin resistance (HOMA-IR). Blood pressure, cholesterol and HOMA-IR were natural log-transformed for the analyses. All analyses were conducted using Stata SE 14 (Stata Corporation, College Station, Texas, United States).

## **Results**

Our analyses were based on 6,887 men and women, of whom 562 were seropositive for Eu-tTGA, 197 for Celikey-tTGA and 72 for EMA. In all, 937 participants had CVD at baseline. CVD was more common among seropositive than seronegative participants, but after adjustment for age, the odds of baseline CVD did not significantly differ between these groups (Table 1).

During a median follow-up of 26 years (range: <1 to 32 years), 2,367 individuals experienced a CHD event. Age-adjusted Cox models provided no evidence for an association between antibody seropositivity and any CHD or hard endpoint CHD, and additional adjustment for pre-existing CVD or known CHD risk factors did not markedly alter the results (Table 2 and Online Supplement, Tables S1 and S1).

To investigate whether tTGA positivity is related to CHD incidence independently of EMA, we post-hoc investigated CHD risk in participants who were positive for Eu-tTGA and Celikey-tTGA but negative for EMA (n=130). Again, multivariable-adjusted analyses provided no clear evidence for an association independently of age, sex, pre-existing CVD and other covariates (Table 2).

## **Discussion**

### *Summary of main findings*

Our prospective investigation of over 6,800 Finnish men and women, who were followed up for up to three decades, provided no clear evidence for an association of tTGA or EMA seropositivity with incident CHD. Our findings support those of a previous register-based study conducted in Sweden, in which seropositivity was not associated with the risk of myocardial infarction or angina pectoris (HR: 1.14, 95% confidence interval, CI: 0.87, 1.50).



[23] In the latter investigation, however, the seropositive group included individuals who were positive for gliadin antibodies [23], which can be detected not only in coeliac disease patients but also in healthy people as well as those with other autoimmune diseases [24-26]. It is thus possible that the seropositive group in this study represented a population heterogeneous in terms of the origins of their serological status. In the present study similar may be true of Eu-tTGA positive participants. In Mini-Finland Health Survey the seropositivity for Eu-tTGA (8.2%) was higher than the estimated prevalence of coeliac disease in the Finnish population overall (2.4%) [27]. This suggests that this group includes individuals with a mix of tTGA-associated conditions, such as ulcerative colitis or various forms of arthritis [28]. However, EMAs are pathognomonic for coeliac disease and the prevalence for seropositivity for the other two tests, Celikey-tTGA (2.9% of our study population) and EMA (1.0% of our participants) are closer to the estimated prevalence of coeliac disease, suggesting that the individuals positive for these tests are more likely to have elevated antibody levels related to gluten-induced autoimmunity.

Taken together, our observations and the findings reported before suggest that seropositivity for tTGAs or EMAs is unlikely to explain the previously observed elevated CHD risk in individuals with coeliac disease. One possible explanation for the apparent discrepancy between laboratory-based investigations and human studies is that the angiogenesis-regulating properties of tTGAs reported in cells and animal models [18-20] do not translate to humans. It is also possible that the association is weak and that considerably larger studies than ours (with 235 incident CHD cases in the seropositive group) or Ludvigsson and colleagues' (with 62 incident CHD cases) would be needed to ascertain sufficient power to detect it. However, the clinical significance of such a weak association is uncertain. Either way, although early diagnosis and initiation of gluten-free diet will undoubtedly improve the

prognosis and well-being of individuals with coeliac disease, they are unlikely to have a significant impact on CHD burden in this patient population.

An important strength of our investigation is that we used prospectively collected data and register-based CHD outcomes, which are generally not prone to recall or other biases. Indeed, validation studies suggest that the Finnish hospitalisation and death registers have good coverage and reasonable diagnostic accuracy for CHD outcomes [29]. As Mini-Finland Health Survey had a 90% participation rate, with negligible amounts of missing data or attrition, it is unlikely that the latter have unduly biased our findings. Antibody measurements were missing from 3.1% of participants due to serum samples having been used up in other assays, and such data, likely to be missing completely at random, may have slightly biased our estimates towards the null. We examined the possibility of exposure misclassification in our data by testing a randomly selected sub-sample of 128 Eu-tTGA negative individuals for Celikey-tTG and EMA. No-one tested positive, suggesting that such exposure misclassification is unlikely. A further strength of our investigation is that we were able to adjust our estimates for a large number of covariates, which tend not to be available in studies based on electronic hospitalisation records only. We cannot, of course, exclude the possibility that our observations occurred due to chance.

#### *Implications for future research*

The mechanisms for the apparent association between coeliac disease and cardiovascular disease are not fully understood, but currently available evidence suggests several possibilities. Coeliac disease patients often suffer from chronic low level inflammation, which can drive atherosclerosis and vascular damage. [30] A further possibility is that the malabsorption of nutrients, often observed in untreated disease or patients whose adherence

to gluten-free diet is poor, can lead to low circulating concentrations of folate or elevated concentrations of homocysteine, [31] which are implicated in the cardiovascular disease pathology. [32, 33] It is also possible that coeliac disease patients are more likely to receive diagnoses of other diseases and conditions due to their regular contact with healthcare professionals. These mechanisms would merit further research.

### **Conclusion**

In a prospective study of over 6,800 Finnish men and women we found no clear evidence for an association of tTGA or EMA seropositivity with incident CHD outcomes, suggesting that tTG autoimmunity is unlikely to be the biological link between coeliac disease and CHD. Thus, although early diagnosis and initiation of gluten-free diet will undoubtedly improve the prognosis and well-being of individuals with coeliac disease, they are unlikely to have a significant impact on CHD burden in this patient population.

Table 1. Participant characteristics

Baseline characteristic	Eu-tTGA negative (n=6 325)	Eu-tTGA positive (n=562)	p vs. Eu-tTGA negative	Celikey tTGA positive (n=197)	p vs. Eu-tTGA negative	EMA positive (n=72)	p vs. Eu-tTGA negative
CVD (N, %)	812 (12.8)	125 (22.2)	<0.0001	44 (22.3)	<0.0001	6 (8.3)	0.3
Female (N, %)	3 439 (54.4)	280 (49.8)	0.038	125 (63.5)	0.012	51 (70.8)	0.005
Age (years) (mean, SD)	50.4 (13.9)	57.9 (14.2)	<0.0001	59.6 (14.0)	<0.0001	50.2 (11.4)	0.9
Years of education (N, %)							
0-8	4 241 (67.1)	427 (76.0)	<0.0001	151 (76.7)	0.013	45 (62.5)	0.6
9-12	1 349 (21.3)	98 (17.4)		33 (16.8)		19 (26.4)	
≥12	735 (11.6)	37 (6.6)		13 (6.6)		8 (11.1)	
Alcohol intake (N, %)							
None	2 826 (44.7)	292 (52.0)	0.008	119 (60.4)	<0.0001	40 (55.6)	0.07
1-49 gr ethanol/week	2 749 (43.5)	215 (38.3)		61 (31.0)		30 (41.7)	
50-249 gr ethanol/week	257 (4.10)	16 (2.9)		5 (2.5)		1 (1.4)	
250+ gr ethanol/week	493 (7.8)	39 (6.9)		12 (6.1)		1 (1.4)	
BMI (kg/m <sup>2</sup> ) (N, %)							
<18.5	308 (4.9)	31 (5.5)	0.3	15 (7.6)	0.1	6 (8.3)	0.3
18.5-24.9	2 611 (41.3)	206 (36.7)		65 (33.0)		30 (41.7)	
25-29.9	2 461 (38.9)	236 (42.0)		87 (44.2)		30 (41.7)	
30-34.9	779 (12.3)	73 (13.0)		24 (12.2)		4 (5.6)	
≥35	166 (2.6)	16 (2.9)		6 (3.1)		2 (2.8)	
Smoking (N, %)							
Never smoked	3 499 (55.3)	333 (59.3)	0.048	134 (68.0)	0.003	51 (70.8)	0.06
Ex-smoker	1 302 (20.6)	123 (21.9)		35 (17.8)		8 (11.1)	
Pipe/cigars/1-19 cigarettes/day	952 (15.1)	68 (12.1)		18 (9.1)		8 (11.1)	
20+ cigarettes/day	572 (9.0)	38 (6.8)		10 (5.1)		5 (6.9)	
Diabetes (N, %)	329 (5.2)	52 (9.3)	<0.0001	28 (14.2)	<0.0001	2 (2.8)	0.4
Systolic blood pressure (mmHg) (mean, SD)	145.5 (23.6)	152.1 (26.0)	<0.0001	153.8 (26.7)	<0.0001	143.9 (25.7)	0.6
Diastolic blood pressure (mmHg) (mean, SD)	87.1 (11.7)	88.4 (13.1)	0.0109	87.1 (12.8)	0.9	85.4 (10.9)	0.2
Total cholesterol (mmol/L) (mean, SD)	6.9 (1.4)	6.9 (1.4)	0.6	6.6 (1.4)	0.0002	6.3 (1.2)	0.0002
HDL cholesterol (mm/L) (mean, SD)	1.7 (0.4)	1.6 (0.4)	<0.0001	1.6 (0.4)	<0.0001	1.5 (0.3)	0.0008
HOMA-IR <sup>3</sup> (mean, SD)	2.5 (7.5)	2.8 (4.1)	0.4	3.6 (6.2)	0.0471	2.1 (1.5)	0.7

<sup>1</sup> CVD: cardiovascular disease.<sup>2</sup> HDL: high-density lipoprotein cholesterol<sup>3</sup> HOMA-IR: homeostasis model assessment; insulin resistance

Table 2. Associations of antibody seropositivity with incident CHD

<b>Antibody test status</b>	<b>Person-years at risk</b>	<b>N CHD<sup>1</sup> (incidence per 1000 person- years)</b>	<b>HR<sup>2</sup> (95% CI) Adjusted for age</b>	<b>HR<sup>2</sup> (95% CI) Adjusted for age, sex and baseline CVD<sup>3</sup></b>	<b>HR<sup>2</sup> (95% CI) Adjusted for age, sex, baseline CHD and covariates<sup>3</sup></b>
Eu-tTGA negative (n=6 325)	141 907	2 132 (15.0)	1 (ref.cat.)	1 (ref.cat.)	1 (ref.cat.)
Eu-tTGA positive (n=562)	10 094	235 (23.3)	1.16 (1.05, 1.33)	1.06 (0.93, 1.22)	1.05 (0.92, 1.21)
Celikey-tTGA positive (n=197)	3 470	80 (23.5)	1.04 (0.83, 1.30)	1.04 (0.82, 1.30)	1.04 (0.83, 1.30)
EMA positive (n=72)	1 789	24 (13.4)	0.80 (0.54, 1.21)	1.00 (0.67, 1.51)	1.16 (0.77, 1.74)
Eu-tTGA and Celikey-tTGA positive, EMA negative (n=130)	1 815	57 (31.4)	1.18 (0.90, 1.54)	1.06 (0.81, 1.38)	1.01 (0.77, 1.31)

<sup>1</sup>CHD: coronary heart disease

<sup>2</sup>HR: hazard ratio

<sup>3</sup>Covariates: level of education, BMI, smoking, alcohol intake, blood pressure, total cholesterol, HDL cholesterol, diabetes and HOMA-IR.

**Acknowledgements**

We thank all the men and women in Mini-Finland Health Survey for taking part in the study.

**Supporting Information**

Online Supplement 1.

**Acknowledgement of grant support**

This work was supported by grants to the Coeliac Disease Study Group from the Academy of Finland, the Sigrid Juselius Foundation and the Competitive State Research Financing of the Expert Responsibility Area of the Tampere University Hospital (grant no. 9S041). The funding bodies had no role in design or conduct of the study, analysing or interpreting the data, preparing the manuscript or decision to submit the manuscript for publication.

**Ethics statement**

Mini-Finland Health Survey precedes the current legislation on ethics in medical research, so no formal ethics committee approval was required at the time of the study baseline. All study participants received written information about the aims and purpose of the study, about the use and storage of the data collected, and that participation was voluntary. Agreeing to participate in the baseline examination was taken to indicate informed consent.

**Author contributions**

KH and KK designed the study, with input from MH and MM. KH and MH planned the statistical analyses, with input from HR and PK. KH conducted the statistical analyses and wrote the first draft of the paper. KH, MH, HR, PK, MM and KK critically reviewed the manuscript and participated in writing the final version.

**Disclosure/ Conflict of interest**

The authors declare no conflict of interest.

ACCEPTED MANUSCRIPT

**References**

1. Green PH, Cellier C. Celiac disease. *N Engl J Med* 2007;**357**(17):1731-43
2. Klock C, Diraimondo TR, Khosla C. Role of transglutaminase 2 in celiac disease pathogenesis. *Semin Immunopathol* 2012;**34**(4):513-22
3. CG86: Coeliac disease: Recognition and assessment of coeliac disease. Secondary CG86: Coeliac disease: Recognition and assessment of coeliac disease 2009. <http://publications.nice.org.uk/coeliac-disease-cg86/guidance>.
4. See J, Murray JA. Gluten-free diet: the medical and nutrition management of celiac disease. *Nutr Clin Pract* 2006;**21**(1):1-15
5. Kurppa K, Collin P, Viljamaa M, et al. Diagnosing mild enteropathy celiac disease: a randomized, controlled clinical study. *Gastroenterology* 2009;**136**(3):816-23
6. Vilppula A, Kaukinen K, Luostarinen L, et al. Clinical benefit of gluten-free diet in screen-detected older celiac disease patients. *BMC Gastroenterol* 2011;**11**:136
7. Dickey W, Hughes DF, McMillan SA. Patients with serum IgA endomysial antibodies and intact duodenal villi: clinical characteristics and management options. *Scand J Gastroenterol* 2005;**40**(10):1240-3
8. Godfrey JD, Brantner TL, Brinjikji W, et al. Morbidity and mortality among older individuals with undiagnosed celiac disease. *Gastroenterology* 2010;**139**(3):763-9
9. Lewis NR, Scott BB. Meta-analysis: deamidated gliadin peptide antibody and tissue transglutaminase antibody compared as screening tests for coeliac disease. *Aliment Pharmacol Ther* 2010;**31**(1):73-81
10. Rubio-Tapia A, Kyle RA, Kaplan EL, et al. Increased prevalence and mortality in undiagnosed celiac disease. *Gastroenterology* 2009;**137**(1):88-93
11. Lohi S, Mustalahti K, Kaukinen K, et al. Increasing prevalence of coeliac disease over time. *Aliment Pharmacol Ther* 2007;**26**(9):1217-25
12. Rubio-Tapia A, Ludvigsson JF, Brantner TL, et al. The prevalence of celiac disease in the United States. *Am J Gastroenterol* 2012;**107**(10):1538-44; quiz 37, 45
13. West J, Logan RF, Hill PG, et al. The iceberg of celiac disease: what is below the waterline? *Clin Gastroenterol Hepatol* 2007;**5**(1):59-62
14. Lewis NR, Holmes GK. Risk of morbidity in contemporary celiac disease. *Expert Rev Gastroenterol Hepatol* 2010;**4**(6):767-80
15. Ludvigsson JF. Mortality and malignancy in celiac disease. *Gastrointest Endosc Clin N Am* 2012;**22**(4):705-22
16. Heikkila K, Pearce J, Maki M, et al. Coeliac disease and bone fractures: a systematic review and meta-analysis. *J Clin Endocr Metab* 2015;**100**(1):25-34
17. Heikkila K, Koskinen OA, Agarwal A, et al. Associations of coeliac disease with coronary heart disease and cerebrovascular disease: A systematic review and meta-analysis. *Nutr Metab Cardiovasc Dis* 2015;**25**(9):816-31
18. Kalliokoski S, Sulic AM, Korponay-Szabo IR, et al. Celiac Disease-Specific TG2-Targeted Autoantibodies Inhibit Angiogenesis and in Mice by Interfering with Endothelial Cell Dynamics. *PLoS One* 2013;**8**(6):e65887
19. Nadalutti CA, Korponay-Szabo IR, Kaukinen K, et al. Celiac disease patient IgA antibodies induce endothelial adhesion and cell polarization defects via extracellular transglutaminase 2. *Cell Mol Life Sci* 2013
20. Sane DC, Kontos JL, Greenberg CS. Roles of transglutaminases in cardiac and vascular diseases. *Front Biosci* 2007;**12**:2530-45
21. Heliovaara M, Aromaa A, Klaukka T, et al. Reliability and validity of interview data on chronic diseases. The Mini-Finland Health Survey. *J Clin Epidemiol* 1993;**46**(2):181-91
22. Lohi S, Maki M, Montonen J, et al. Malignancies in cases with screening-identified evidence of coeliac disease: a long-term population-based cohort study. *Gut* 2009;**58**(5):643-7
23. Ludvigsson JF, James S, Askling J, et al. Nationwide cohort study of risk of ischemic heart disease in patients with celiac disease. *Circulation* 2011;**123**(5):483-90



24. Husby S, Koletzko S, Korponay-Szabo IR, et al. European Society for Pediatric Gastroenterology, Hepatology, and Nutrition guidelines for the diagnosis of coeliac disease. *J Pediatr Gastroenterol Nutr* 2012;**54**(1):136-60
25. Rostom A, Dube C, Cranney A, et al. The diagnostic accuracy of serologic tests for celiac disease: a systematic review. *Gastroenterology* 2005;**128**(4 Suppl 1):S38-46
26. Maki M. The humoral immune system in coeliac disease. *Baillieres Clin Gastroenterol* 1995;**9**(2):231-49
27. Mustalahti K, Catassi C, Reunanen A, et al. The prevalence of celiac disease in Europe: results of a centralized, international mass screening project. *Ann Med* 2010;**42**(8):587-95
28. Lerner A, Matthias T. Rheumatoid arthritis-celiac disease relationship: Joints get that gut feeling. *Autoimmunity Reviews* 2015;**14**(11):1038-47
29. Pajunen P, Koukkunen H, Ketonen M, et al. The validity of the Finnish Hospital Discharge Register and Causes of Death Register data on coronary heart disease. *Eur J Cardiovasc Prev Rehabil* 2005;**12**(2):132-7
30. Libby P, Ridker PM, Hansson GK. Inflammation in atherosclerosis: from pathophysiology to practice. *J Am Coll Cardiol* 2009;**54**(23):2129-38
31. Dickey W, Ward M, Whittle CR, et al. Homocysteine and related B-vitamin status in coeliac disease: Effects of gluten exclusion and histological recovery. *Scand J Gastroenterol* 2008;**43**(6):682-8
32. Homocysteine and risk of ischemic heart disease and stroke: a meta-analysis. *JAMA* 2002;**288**(16):2015-22
33. Huo Y, Qin X, Wang J, et al. Efficacy of folic acid supplementation in stroke prevention: new insight from a meta-analysis. *Int J Clin Pract* 2012;**66**(6):544-51

**Study Highlights****Associations of Tissue Transglutaminase Antibody Seropositivity with Coronary Heart Disease: Findings from a Prospective Cohort Study**

Katriina Heikkilä, Harri Rissanen, Markku Heliövaara, Paul Knekt, Markku Mäki, Katri Kaukinen

Clinical experience and observational studies suggest that individuals with coeliac disease are at increased risk of coronary heart disease (CHD), but the mechanism for this is unclear.

Laboratory studies suggest that it may relate to tissue transglutaminase antibodies (tTGAs).

In our prospective, population-based study of over 6,800 Finnish men and women, we found no clear evidence for an association of seropositivity for tTGA and endomysial antibodies (EMAs) with incident CHD in humans.

These findings suggest that though early diagnosis and initiation of gluten-free diet will undoubtedly improve the well-being of individuals with coeliac disease, they are unlikely to have a significant impact on CHD burden in this patient population.