



**QUEEN'S  
UNIVERSITY  
BELFAST**

## **Alcoholic beverage preference and diabetes incidence across Europe the Consortium on Health and Ageing Network of Cohorts in Europe and the United States (CHANCES) project**

Sluik, D., Jankovic, N., O'Doherty, M. G., Schöttker, B., Drygas, W., Rolandsson, O., ... Feskens, E. J. M. (2017). Alcoholic beverage preference and diabetes incidence across Europe the Consortium on Health and Ageing Network of Cohorts in Europe and the United States (CHANCES) project. *European Journal of Clinical Nutrition*, 71(5), 659-668. DOI: 10.1038/ejcn.2017.4

### **Published in:**

European Journal of Clinical Nutrition

### **Document Version:**

Peer reviewed version

### **Queen's University Belfast - Research Portal:**

[Link to publication record in Queen's University Belfast Research Portal](#)

### **Publisher rights**

© 2017 Macmillan Publishers Limited

This work is made available online in accordance with the publisher's policies. Please refer to any applicable terms of use of the publisher.

### **General rights**

Copyright for the publications made accessible via the Queen's University Belfast Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### **Take down policy**

The Research Portal is Queen's institutional repository that provides access to Queen's research output. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [openaccess@qub.ac.uk](mailto:openaccess@qub.ac.uk).

1 **Alcoholic beverage preference and diabetes incidence across Europe: the Consortium on**  
2 **Health and Ageing Network of Cohorts in Europe and the United States (CHANCES)**  
3 **project**

4  
5 Diewertje Sluik<sup>1</sup>, Nicole Jankovic<sup>1,2</sup>, Maria Hughes<sup>3</sup>, Mark G. O’Doherty<sup>3</sup>, Ben Schöttker<sup>4,5</sup>,  
6 Wojciech Drygas<sup>6</sup>, Olov Rolandsson<sup>7</sup>, Satu Männistö<sup>8</sup>, José M. Ordóñez-Mena<sup>4,9,10</sup>, Jean  
7 Ferrieres<sup>11</sup>, Christina Bamia<sup>12,13</sup>, Giovanni de Gaetano<sup>14</sup>, Jessica C. Kieft-De Jong<sup>15,16</sup>, Oscar  
8 H. Franco<sup>16</sup>, Ivonne Sluijs<sup>17</sup>, Annemieke M.W. Spijkerman<sup>18</sup>, Susana Sans<sup>19</sup>, Sture Eriksson<sup>7</sup>,  
9 Daan Kromhout<sup>1</sup>, Antonia Trichopoulou<sup>12,20</sup>, Tom Wilsgaard<sup>21</sup>, Herman Brenner<sup>4,9</sup>, Kari  
10 Kuulasmaa<sup>8</sup>, Tiina Laatikainen<sup>8,22,23</sup>, Stefan Söderberg<sup>24</sup>, Licia Iacoviello<sup>14</sup>, Paolo  
11 Boffetta<sup>12,25</sup>, Frank Kee<sup>3</sup>, Edith J.M. Feskens<sup>1</sup>

12  
13 <sup>1</sup> Division of Human Nutrition, Wageningen University, Wageningen, the Netherlands;  
14 <sup>2</sup> Centre of Clinical Epidemiology, Institute for Medical Informatics, Biometry and  
15 Epidemiology, Faculty of Medicine, University Duisburg-Essen, Germany; <sup>3</sup> UKCRC Centre  
16 of Excellence for Public Health, School of Medicine, Dentistry, and Biomedical Sciences,  
17 Queen’s University Belfast, Belfast, Northern Ireland; <sup>4</sup> Division of Clinical Epidemiology  
18 and Aging Research, German Cancer Research Center (DKFZ), Heidelberg, Germany;  
19 <sup>5</sup> Institute of Health Care and Social Sciences, FOM University, Essen, Germany;  
20 <sup>6</sup> Department of Epidemiology, CVD Prevention and Health Promotion, Institute of  
21 Cardiology , Warsaw, Poland; <sup>7</sup> Department of Public Health and Clinical Medicine, Family  
22 Medicine, Umeå University, Umeå, Sweden; <sup>8</sup> Department of Health, National Institute for  
23 Health and Welfare (THL), Helsinki, Finland; <sup>9</sup> Network of Aging Research (NAR),  
24 Heidelberg University, Heidelberg, Germany; <sup>10</sup> Nuffield Department of Primary Care and  
25 Health Sciences, University of Oxford, Oxford, United Kingdom; <sup>11</sup> Department of

26 Cardiology, Toulouse University School of Medicine, Toulouse, France; <sup>12</sup> Hellenic Health  
27 Foundation, Athens, Greece; <sup>13</sup> Department of Hygiene, Epidemiology and Medical Statistics,  
28 University of Athens Medical School, Athens, Greece; <sup>14</sup> Department of Epidemiology and  
29 Prevention. IRCCS Istituto Neurologico Mediterraneo Neuromed, Pozzilli, Italy; <sup>15</sup> Global  
30 Public Health, Leiden University College, the Hague, the Netherlands; <sup>16</sup> Department of  
31 Epidemiology, Erasmus Medical Center, Rotterdam, the Netherlands; <sup>17</sup> Julius Center for  
32 Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht, the  
33 Netherlands; <sup>18</sup> National Institute for Public Health and the Environment (RIVM), Bilthoven,  
34 the Netherlands; <sup>19</sup> Catalan Department of Health, Barcelona, Spain; <sup>20</sup> WHO Collaborating  
35 Center for Nutrition and Health, Unit of Nutritional Epidemiology and Nutrition in Public  
36 Health, Department of Hygiene, Epidemiology and Medical Statistics, University of Athens  
37 Medical School, Athens, Greece; <sup>21</sup> Department of Community Medicine, UiT the Arctic  
38 University of Norway, Tromsø, Norway; <sup>22</sup> Institute of Public Health and Clinical Nutrition,  
39 University of Eastern Finland, Kuopio, Finland; <sup>23</sup> Hospital District of North Karelia, Finland;  
40 <sup>24</sup> Department of Public Health and Clinical Medicine, Cardiology and Heart Centre, Umeå  
41 University, Umeå, Sweden; <sup>25</sup> Icahn School of Medicine, Mount Sinai School of Medicine,  
42 New York, USA.

43

44 **Correspondence and offprint requests to:**

45 Diewertje Sluik, DrPH

46 E-mail: [Diewertje.Sluik@wur.nl](mailto:Diewertje.Sluik@wur.nl)

47 P.O. Box 17

48 6700 AA Wageningen

49 The Netherlands

50 Phone: +31 317 485953 / Fax: +31 317 482782

51 **Funding:** The research of DS was supported by the Dutch Beer Institute and the European  
52 Foundation for Alcohol Research (ERAB). The sponsor did not have any role in the design  
53 and conduct of the study; collection, management, analysis and interpretation of the data; and  
54 preparation, review or approval of the manuscript. All authors declare no conflicts of interest.

55 **Running head:** Alcoholic beverage preference and diabetes in Europe

56 **Abbreviations:** Body Mass Index (BMI); coronary heart disease (CHD); confidence Interval  
57 (CI); the Consortium on Health and Ageing: Network of Cohorts in Europe and the United  
58 States (CHANCES); European Prospective Investigation into Cancer and Nutrition (EPIC);  
59 the Epidemiological Study on Chances for Prevention, Early Detection, and Optimized  
60 THERapy of Chronic Diseases at Old Age (ESTHER); Hazard Ratio (HR); MOnica Risk,  
61 Genetics, Archiving and Monograph (MORGAM)

62 **Abstract**

63 *Background/Objectives:* It is unknown if wine, beer, and spirit intake lead to a similar  
64 association with diabetes. We studied the association between alcoholic beverage preference  
65 and type 2 diabetes incidence in persons who reported to consume alcohol.

66 *Subjects/Methods:* Ten European cohort studies from the Consortium on Health and Ageing:  
67 Network of Cohorts in Europe and the United States (CHANCES) were included, comprising  
68 participant data of 62 458 adults who reported alcohol consumption at baseline. Diabetes  
69 incidence was based on documented and/or self-reported diagnosis during follow-up.

70 Preference was defined as  $\geq 70\%$  of total alcohol consumed was either beer, wine or spirits.

71 Adjusted hazard ratios (HRs) were computed using Cox proportional hazard regression.

72 Single cohort HRs were pooled by random-effects meta-analysis.

73 *Results:* Beer, wine, or spirit preference was not related to diabetes risk compared with having  
74 no preference. The pooled HRs were HR 1.06 (95%CI 0.93, 1.20) for beer, HR 0.99 (95%CI  
75 0.88, 1.11) for wine, and HR 1.19 (95%CI 0.97, 1.46) for spirit preference. Absolute wine  
76 intake, adjusted for total alcohol, was associated with a lower diabetes risk: pooled HR per 6  
77 grams/day was 0.96 (95% CI 0.93, 0.99). A spirit preference was related to a higher diabetes  
78 risk in those with a higher BMI, in men and women separately, but not after excluding  
79 persons with prevalent diseases

80 *Conclusions:* This large individual-level meta-analysis among persons who reported alcohol  
81 consumption revealed that the preference for beer, wine, and spirits was similarly associated  
82 with diabetes incidence compared with having no preference.

## 83 **Introduction**

84 Diabetes mellitus is the fourth to fifth leading cause of death in most high-income countries<sup>1</sup>.

85 In 2014, the International Diabetes Federation estimated the prevalence at 7.9% in Europe<sup>1</sup>.

86 Two systematic reviews and meta-analyses, including 20 and 26 cohort studies each, revealed  
87 a non-linear U-shaped relationship between alcohol consumption and type 2 diabetes

88 incidence in both men and women<sup>2, 3</sup>. The protective effect of alcohol consumption was

89 largest with light to moderate consumption. Higher levels of ethanol consumption were not

90 associated with diabetes or were associated with a higher risk<sup>2, 3</sup>. On the other hand, a more

91 recent meta-analysis of 38 studies concluded these risk reductions might have been

92 overestimated by including less healthy former consumers in the reference group<sup>4</sup>. Moreover,

93 the protective association might be confined to women and non-Asian populations only<sup>4</sup>.

94

95 Further research has indicated that the associations between alcohol and diabetes might be

96 beverage-specific. A recent systematic review and meta-analysis of 13 prospective studies

97 showed a strong protective association for wine consumption and type 2 diabetes, while for

98 beer or spirits only a slight trend of a protective association was observed<sup>5</sup>. Within the

99 European Prospective Investigation into Cancer and Nutrition (EPIC)-InterAct Study,

100 moderate alcohol consumption was also related to a lower diabetes risk, in particular the

101 consumption of red wine<sup>6</sup>. It was suggested that the association between alcohol and diabetes

102 was likely to be explained by ethanol itself. Indeed, intervention studies have shown that

103 alcohol increases levels of HDL-cholesterol, apolipoprotein A1, and adiponectin, and reduces

104 fibrinogen, fasting insulin and HbA<sub>1c</sub> concentrations<sup>7, 8</sup>. Hence, the observed differences in

105 association between wine, beer, and spirits and health outcomes might be due to socio-

106 demographic and lifestyle factors associated with the preference and consumption of these

107 beverages<sup>9, 10</sup>. However, differential effects of beer and wine on the glycemic response, as

108 expressed by their glycemic index, have also been observed. Beer induces a higher glucose  
109 response than wine, which may be related to the development of diabetes<sup>11, 12</sup>. Furthermore,  
110 due to its higher polyphenol content, red wine may exert additional benefits including  
111 reduction of blood pressure and inflammation and improving endothelial function<sup>13</sup>.

112

113 Alcohol consumption is a complex exposure that can be characterized in different ways: the  
114 absolute amount, the drinking frequency, and the beverage type. It is statistically difficult to  
115 distinguish between the overall alcohol effect and the specific effects of beer, wine, and spirits  
116 in observational studies<sup>14</sup>. We aimed to disentangle beverage-specific effects, independent of  
117 those from the absolute ethanol consumption, by studying the association between alcoholic  
118 beverage consumption and preference and type 2 diabetes incidence. This was done by  
119 performing a meta-analysis of harmonized individual participant data from several European  
120 cohorts including a large proportion of elderly participants. Because this study focused on the  
121 type of alcoholic beverage, the analyses were restricted to persons who reported alcohol  
122 consumption. Moreover, because the consumption of wine, beer, or spirits is mainly  
123 determined by factors including age, sex, socio-economic status, country, and lifestyle, these  
124 variables will be taken as much as possible into account in the analyses to strengthen  
125 potentially causal inference.

126

## 127 **Subjects and methods**

### 128 *Study design and population*

129 The Consortium on Health and Ageing: Network of Cohorts in Europe and the United States  
130 (CHANCES) project is a coordinated multi-country study which aims to harmonize data from  
131 ongoing prospective cohort studies in Europe and the USA in order to produce evidence on

132 ageing-related health characteristics and on determinants of healthy ageing among the elderly  
133 in these countries<sup>15</sup>.

134

135 The CHANCES project includes cohorts from 14 studies across Europe and the USA. In most  
136 CHANCES cohorts, elderly are defined as those who were 60 years or older at recruitment.

137 The CHANCES project as a whole has received ethical approval by the Hellenic Health

138 Foundation Committee on Bioethics (HHFCB). In the individual cohorts, all participants

139 signed informed consent for the original studies. The authors of this study did not have any

140 access to personal information regarding the participants included in this paper. All data that

141 have been analyzed are based on the CHANCES harmonized variables and are completely

142 anonymized. For the present study, the following ten European cohorts were eligible for

143 analysis: the Zutphen Elderly Study (the Netherlands)<sup>16</sup>, Rotterdam Study (the Netherlands)<sup>17</sup>,

144 the study centers in the Netherlands, Greece, and Sweden from the European Prospective

145 Investigation into Cancer and Nutrition (EPIC) – Elderly study<sup>18</sup>, the Tromsø Study

146 (Norway)<sup>19</sup>, the Epidemiological Study on Chances for Prevention, Early Detection, and

147 Optimized THERapy of Chronic Diseases at Old Age (ESTHER) study (Germany)<sup>20</sup>, and

148 from the MONica Risk, Genetics, Archiving and Monograph (MORGAM) study, the cohorts

149 of FINRISK (Finland), Northern Sweden (Sweden), and MOLI-SANI (Italy)<sup>21, 22</sup>. An

150 extensive overview of the cohorts included in the CHANCES project and data assessment has

151 been published elsewhere<sup>23</sup>. **Table 1** displays the main characteristics of the included ten

152 cohorts and participants.

153

154 Within the cohorts that were eligible for the present study, analyses were conducted upon all

155 subjects who reported to consume alcohol, without any missing data on alcohol and followed

156 up for diabetes incidence. Subjects with self-reported or independently ascertained prevalent



157 diabetes at baseline or with missing information on prevalent diabetes at baseline were  
158 excluded from analysis. **Supplemental Figure 1** shows the participant flow-charts of the ten  
159 included cohorts, comprising a total sample size of 63 458.

160

#### 161 *Data assessment and harmonization*

162 Data in the CHANCES project have been collected within the framework of independent  
163 cohort studies, with different protocols for data collection and distinct original research foci.  
164 Data harmonization was a major task of the project and the data harmonization and  
165 conversion rules of the CHANCES project have been described elsewhere<sup>23</sup>.

166

#### 167 *Alcoholic consumption and beverage preference*

168 Baseline alcohol data were recorded either by self-administered or interview-based  
169 questionnaires. The EPIC-Elderly and Rotterdam Study applied a validated food frequency  
170 questionnaire (FFQ)<sup>17, 24, 25</sup> to assess alcohol intake. The Zutphen Elderly Study used a  
171 validated dietary history method to assess diet including alcohol<sup>16</sup>. The Tromsø Study, the  
172 ESTHER study, and MORGAM cohorts derived alcohol consumption from a general  
173 questionnaire. In the FINRISK Study, alcohol consumption during the previous week was  
174 assessed. If not already defined, average daily alcohol consumption in grams was estimated  
175 by adding the amounts of ethanol found in each standard drink or cohort specific size for beer,  
176 wine, and spirits. To ensure comparability across cohorts, a conversion rule was applied using  
177 standardized portion sizes (330 ml for a bottle of beer, 175 ml for a glass of wine, and 25 ml  
178 for a shot of spirit) and alcohol percentages in beer (4.5%), wine (12%), and spirits (37.5%).

179

180 As defined in previous studies, a person was classified as having a preference for beer, wine,  
181 or spirits, when the alcohol consumption from the respective drink comprised 70% or more of

182 the total alcohol consumption in grams per day. When the average alcohol consumption from  
183 beer, wine, or spirits did not add up to 70% of the total alcohol consumption, a person was  
184 classified as having no preference<sup>26, 27</sup>. To assess robustness of this definition, a sensitivity  
185 analysis was performed using a cut-off of 50%. Associations between the preference for beer,  
186 wine, or spirits and diabetes incidence compared to having no specific preference was  
187 assessed. Non-consumers, comprising never and former consumers, were not included in the  
188 analyses.

189

190 Next, the association between average daily intake from beer, wine, and spirits and diabetes  
191 incidence was studied. The absolute intakes of beer, wine, and spirits were adjusted for total  
192 alcohol consumption by the residual method<sup>27</sup>. In this procedure, intakes of the respective  
193 beverage were regressed upon their total alcohol consumption and the residuals from the  
194 regression were used in the analysis. These residuals represent the differences between each  
195 individual's actual intake and the intake predicted by their total alcohol consumption. Because  
196 residuals, by definition, have a mean of zero, a constant representing the mean intake in each  
197 population was added to every value to reflect actual consumption values<sup>28</sup>. The beer, wine,  
198 and spirit residuals are uncorrelated with total alcohol consumption and this allows variation  
199 due to the intake of beer, wine, and spirits to be evaluated directly. The beer, wine, and spirit  
200 residuals were analyzed in tertiles and per 6 g/day.

201

202 Information on drinking patterns, i.e. consumption frequency, was not available for all  
203 cohorts. Sensitivity analyses were performed adjusting the associations additionally for  
204 frequency of consumption (less than once a week, 1-2 days/week, 3-5 days/week, or 6-7  
205 days/week) in the Tromsø Study, ESTHER, and MORGAM.

206 *Diabetes ascertainment*

207 Diabetes incidence was based on documented or self-reported type 2 diabetes during follow-  
208 up or based on fasting glucose measures, depending on the available options within the  
209 cohorts shown in Table 1.

210

211 *Covariate assessment*

212 Socio-demographic, lifestyle, and disease history data were assessed by self-administered  
213 questionnaires or in interviews. Weight and height were either measured or self-reported, and  
214 blood samples were drawn to determine total and HDL cholesterol. Diet quality was assessed  
215 with the Healthy Diet Indicator (HDI) as developed by Jankovic *et al.*<sup>29</sup>. The HDI score  
216 reflects adherence to the 2003 WHO dietary guidelines. The score ranges from 0 to 70 points  
217 and includes 6 nutrients (saturated fatty acids, polyunsaturated fatty acids, mono- and  
218 disaccharides, protein, cholesterol, dietary fiber) and 1 food group (fruit and vegetables) of  
219 the 14 WHO guideline goals, which were available for the cohorts providing nutrition data<sup>29</sup>.  
220 Dietary intake data to calculate the HDI score were available for the Zutphen Elderly Study,  
221 Rotterdam Study, and EPIC-Elderly. Self-reported physical activity was assessed by  
222 questionnaires in the Zutphen Elderly Study, Rotterdam Study, EPIC-Elderly the Netherlands  
223 and Greece, and ESTHER.

224

225 *Statistical analysis*

226 The statistical analyses were performed using SAS, version 9.3, software (SAS Institute, Inc.,  
227 Cary, North Carolina). Hazard Ratios (HRs) and 95% confidence intervals (CIs) for diabetes  
228 were calculated using Cox proportional hazard regression. The proportional hazard  
229 assumption was tested and not violated. Missing values for any of the covariates were  
230 imputed using the multiple imputation method, in which all variables included in the

231 statistical models were included in the procedure. For each cohort, five duplicate datasets  
232 were produced and after statistical inference on the duplicate datasets, pooled estimates were  
233 calculated with PROC MIANALYZE<sup>30</sup>. In Model 1, HRs were adjusted for socio-  
234 demographic factors: age (continuous; years), gender (not applicable for the Zutphen Elderly  
235 Study, which is composed only of men), education (categorical: primary or less (low), more  
236 than primary but less than college or university (middle), college or university (high)),  
237 employment status (categorical: full-time or part-time employment and not of pensionable  
238 age, self-employed, housewife and not of pensionable age, pensionable age and still working,  
239 pensionable age and not working, stopped work before retirement age due to poor health,  
240 unemployed and not of pensionable age; not applicable for SENECA and the Zutphen Elderly  
241 Study, where only retired subjects are included), and prevalent coronary heart disease (CHD;  
242 yes/no) or cancer (yes/no). Model 2 was additionally adjusted for the lifestyle factors:  
243 smoking status (categorical: never, former, current), sports activity (continuous: hours per  
244 week; physical activity data were not available for EPIC-Elderly Sweden, the Tromsø Study,  
245 and MORGAM; total physical activity was used in the Rotterdam Study), and HDI-score  
246 (continuous; dietary intake data to generate the HDI score was not available for ESTHER, the  
247 Tromsø Study, and MORGAM).

248

249 Because the definition of alcoholic beverage preference is not based upon absolute alcohol  
250 consumption, persons with a beer preference might, for instance, have a higher absolute  
251 alcohol intake than persons with a wine preference. Thus, total alcohol consumption might be  
252 a confounding factor. Due to the U-shaped relationship between total alcohol and diabetes<sup>2,3</sup>,  
253 additional adjustment for absolute alcohol consumption (gram/day) was evaluated using  
254 fractional polynomials where the best fit regression model was selected with the SAS Macro  
255 “Multivariable Fractional Polynomials”<sup>31</sup>. This macro uses an algorithm to determines the

256 inclusion and transformation of continuous covariates while taking into account their non-  
257 linearity. In a stepwise approach, the algorithm constructs a fractional polynomial  
258 transformation for the continuous covariate. Backward elimination selects the best  
259 transformation of the covariate, e.g. linear, first degree or second degree. Depending on the P-  
260 values associated with the best transformations, covariates may be eliminated from the model.  
261 In all cohorts, absolute alcohol consumption was omitted from the best fit model. Because the  
262 residuals of beer, wine, and spirit consumption are uncorrelated with total alcohol intake,  
263 these HR were not adjusted for total alcohol.

264

265 Adjustment model 3 was additionally adjusted for BMI (linearly or second degree;  $\text{kg/m}^2$ );  
266 this adjustment for BMI was also evaluated using fractional polynomials. BMI was omitted  
267 from the best fit model in the Zutphen Elderly Study, included as a second degree variable in  
268 ESTHER and FINRISK, and included linearly in the remaining cohorts. BMI is one of the  
269 most important risk factors for diabetes, but is also on a possible causal pathway between  
270 alcohol consumption and diabetes. Therefore, crude and adjusted BMI across alcoholic  
271 beverage preference categories was estimated with multiple linear regression. To investigate  
272 effect modification by BMI, stratified analyses were performed on persons with a BMI  $<25$   
273 and  $\geq 25 \text{ kg/m}^2$  and the P-value for interaction was checked after including a product term in  
274 the regression models. Furthermore, stratified analyses were performed for men and women  
275 separately to check for potential effect modification. Finally, subjects with prevalent CHD or  
276 cancer at baseline or a follow-up less than 2 years were excluded in a sensitivity analysis.

277

278 Cohort-specific HR estimates and 95% CIs for diabetes incidence from having a beer, wine,  
279 or spirit preference compared with no preference and for a beer, wine, or spirit consumption  
280 (per 6 gram/day) were pooled in meta-analyses, using adjustment model 3. Inverse variance

281 weighting was applied to give the largest weight to the study with the lowest variance. The  
282 random-effects model takes into account the between-study variance and the within-study  
283 variance. Heterogeneity between studies was assessed by the Q statistic and the I<sup>2</sup> index. I<sup>2</sup>  
284 was calculated as  $I^2 = ((Q - df)/Q)*100$ , where “df” stands for degrees of freedom, i.e. total  
285 number of studies (k) minus 1. Random-effects meta-analyses with inverse variance  
286 weighting were performed using the R package “meta” (R version 3.3.1). Statistical tests were  
287 two-sided and P-values <0.05 were considered statistically significant.

288

## 289 **Results**

290 In most cohorts, persons with a wine preference constituted the largest group, ranging from  
291 44% in ESTHER (Germany) to 79% in MOLI-SANI (Italy) (**Supplemental Table 1**). In  
292 EPIC-Elderly Sweden, the Tromsø Study, FINRISK, and Northern-Sweden persons with no  
293 preference formed the largest group and in the Zutphen Elderly Study (the Netherlands),  
294 persons with a spirit preference comprised the largest group, i.e. 62%. Across all cohorts,  
295 those who preferred wine were relatively more highly educated and were more likely to be a  
296 never smoker, and female. Furthermore, those with a beer or spirit preference were more  
297 likely to be male and current smoker. Persons with no specific preference generally had the  
298 highest absolute alcohol consumption. After adjustment for age, sex, education, employment,  
299 prevalent diseases, smoking, alcohol, sports activity, diet, BMI was lowest among those with  
300 a beer or wine preference and BMI was highest among persons with a spirit preference  
301 (**Supplemental Table 1**).

302

303 The pooled HRs from the random-effects meta-analyses showed no significant association  
304 between having a preference for beer, wine, or spirits and diabetes incidence compared with  
305 having no specific preference after adjustment for age, sex, education, prevalent diseases,

306 lifestyle factors, and BMI (**Figure 1-3**). Pooled HR was 1.06 (95%CI 0.93, 1.20) for a beer  
307 preference, HR 0.99 (95%CI 0.88, 1.11) for a wine preference, and HR 1.19 (95%CI 0.97,  
308 1.46) for having a spirit preference. Based on the  $I^2$  index and the Q-statistic, between-study  
309 heterogeneity was observed for the effect estimates of having a spirit preference.

310

311 Separate HRs and 95% CIs for the associations between a beer, wine, or spirits and diabetes  
312 incidence according to the different levels of adjustment are shown in Supplemental Table 1.  
313 Compared with persons with no preference, a preference for beer, wine, or spirits was in most  
314 cohorts not significantly associated with diabetes incidence. In the Rotterdam Study, beer or  
315 spirit preference had a significant association with a higher diabetes incidence. In EPIC-  
316 Elderly Greece, having a wine preference tended to be associated with a lower diabetes  
317 incidence. Within the cohorts, additional adjustment for BMI (Model 3) had mixed, but small  
318 effects on the observed associations.

319

320 The pooled HR for the association between alcohol preference and incident diabetes among  
321 sub-groups and with additional adjustments are shown in **Table 2**. Diabetes risk among  
322 persons with a spirit preference was higher in those with a higher BMI, in men and in women,  
323 but not after excluding persons with prevalent diseases. Excluding persons with prevalent  
324 diseases yielded similar results to the findings including those persons. P-values for  
325 interaction by BMI were not significant for all cohorts and did not therefore give indication  
326 for effect modification. Furthermore, additional adjustment for consumption frequency and  
327 alternative analysis using 50% as a cut-off in the definition of preference showed similar  
328 associations.

329 Additionally, the association between the residuals of beer, wine, and spirit intake per 6  
330 gram/day and diabetes incidence was assessed (**Figure 4-6**). Pooled HR was 1.03 (95%CI  
331 0.99, 1.06) per 6 grams of beer intake, HR 0.96 (95%CI 0.93, 0.99) per 6 grams of wine  
332 intake, and HR 1.02 (95%CI 0.98, 1.06) per 6 grams of spirit intake. Cohort-specific HR  
333 according to tertiles and per 6 grams/day generally showed similar associations  
334 (**Supplemental Table 2**).

335

## 336 **Discussion**

337 This meta-analysis of individual participant data from ten prospective European cohorts  
338 comprising ~60,000 adults who reported at least some alcohol consumption showed that a  
339 preference for beer, wine or spirits was not associated with a lower or higher diabetes risk  
340 compared with having no specific preference, taking into account several socio-demographic  
341 and lifestyle variables.

342

343 To our knowledge, no other studies have investigated the association between alcoholic  
344 beverage *preference* and diabetes risk. However, a number of observational studies have  
345 assessed associations of absolute beverage-specific *consumption* and diabetes, showing  
346 inconsistent results. Among 36,527 Australian adults, Hodge *et al.* observed an inverse  
347 association between wine consumption and risk of type 2 diabetes, but not for beer or spirits<sup>32</sup>.  
348 In the EPIC-InterAct Study, a prospective case-cohort study of 16,154 participants and 12,403  
349 incident diabetes cases, consumption of wine and fortified wine were most strongly related  
350 with a reduced diabetes risk<sup>6</sup>. Moreover, compared with a light consumption, men who did  
351 not consume beer had a reduced risk of diabetes in the EPIC-InterAct Study: HR 0.84 (95%CI  
352 0.74, 0.95) and in women higher spirit consumption was associated with an higher diabetes  
353 risk (P-trend 0.044). Fagherazzi *et al.* observed an inverse association between wine



354 consumption and diabetes risk when compared to other types of alcoholic beverage among  
355 66,485 women from the French E3N-EPIC cohort<sup>33</sup>. In contrast, Conigrave *et al.*, did not find  
356 a protective effect of red wine on diabetes risk among 46,892 U.S. male health professionals,  
357 whereas inverse associations for beer, spirits and white wine were similar and independent<sup>34</sup>.  
358 Moreover, two other studies in large U.S. cohorts also did not observe a specific protective  
359 effect of wine consumption on diabetes risk compared with beer or spirit consumption<sup>35,36</sup>. In  
360 their meta-analysis of 13 prospective studies, Huang, Wang, and Zhang presented a pooled  
361 RR of 0.85 (95% CI 0.80-0.89) for wine consumption, and RR 0.96 (95% CI 0.92, 1.00) for  
362 beer consumption and RR 0.95 (95%CI 0.89-1.03) for spirit consumption and type 2 diabetes  
363 risk compared to no or rare alcohol consumption<sup>5</sup>. In the present study, the pooled HR for  
364 residuals of beer, wine, and spirit intake showed similar results: a higher wine intake was  
365 related to a lower diabetes risk, even after fully taking into account total alcohol consumption.  
366 This further confirms the consistent finding that constituents other than ethanol in red wine  
367 may exert additional health benefits<sup>13</sup>.

368

369 Several other studies have found differential effects for the type of alcoholic beverage and  
370 diabetes risk, with a stronger beneficial association for wine consumption compared to  
371 abstinence. These observations could either be explained by a true beneficial effect of wine  
372 compared with beer and spirits, or by an artefact arising from residual confounding. Firstly, a  
373 true differential effect for beer, wine, and spirits and diabetes incidence might be caused by  
374 beneficial compounds other than ethanol in particular those found in wine. For example, a  
375 randomized controlled cross-over trial in 67 men at high cardiovascular risk showed that red  
376 wine rich in polyphenols with or without alcohol improved glucose metabolism<sup>37</sup>. This was  
377 not confirmed in our study, where we have found no additional beneficial association for  
378 having a wine preference compared to having no preference. Secondly, the observation might

379 be an artefact caused by confounding factors associated with the type of alcoholic beverage  
380 consumed. Indeed, the choice of alcoholic beverage is associated with a wide range of  
381 cultural, socio-demographic and lifestyle factors<sup>9,38</sup>, which may confound the association  
382 between alcohol and diabetes risk. Moreover, other important determinants of diabetes risk  
383 including age, gender, smoking status and overall drinking patterns differ across alcoholic  
384 beverage preference and study populations<sup>10</sup>. Therefore, we have adjusted the associations for  
385 age, gender, socio-economic status, and lifestyle factors including absolute alcohol  
386 consumption and BMI. However, we cannot exclude any residual confounding as a result of  
387 unmeasured or imprecisely measured confounders. Lastly, in previous studies there is a  
388 tendency to find an association for the alcoholic beverage that is most consumed. In the above  
389 mentioned studies into beverage type and diabetes risk, most of the alcohol was consumed as  
390 wine<sup>6,32,33</sup>. In our study, ten cohorts from seven European countries were included with  
391 varying preferences, suggesting this might have less influence on our findings and provided a  
392 wider insight into alcohol preference across Europe.

393

394 We observed a tendency toward a higher diabetes risk among persons with a spirit preference  
395 compared to those having no specific preference among men and women when analyzed  
396 separately and those with a higher BMI. This was also seen within EPIC-InterAct<sup>6</sup>. Cross-  
397 sectional studies have shown that a spirit preference is associated with an unhealthier lifestyle:  
398 persons who preferred spirits have been shown to have a higher BMI, are more likely to be  
399 smokers, and display unhealthier diet. Furthermore, spirits may be more often used for heavy  
400 binge drinking compared to wine<sup>9,10</sup>. In the present study, we could only take consumption  
401 frequency into account in a subset of five cohorts. Furthermore, the association was attenuated  
402 when excluding persons with prevalent diseases or a short follow-up, indicating that some

403 degree of reverse causation might be present. Finally, we were not able to take diet and  
404 physical activity into account in all cohorts.

405

406 To avoid any bias by the inclusion of former drinkers, the current analysis was restricted to  
407 persons who reported some alcohol consumption. Furthermore, persons with no specific  
408 preference, i.e. mixed drinkers, were used as a reference. Most other observational studies  
409 have used non-consumers as a reference; however, this has been contested. Non-consumers  
410 are in general a heterogeneous group comprising lifetime abstainers and former drinkers. In  
411 many high-income countries, lifetime abstinence of alcohol is not normative and this group  
412 differs from alcohol consumers in other health determinants<sup>39</sup>. Moreover, former drinkers may  
413 have quit because of ill health arising from their former (heavy) alcohol use. As a result, these  
414 individuals are more vulnerable for morbidity and mortality and their ill health may confound  
415 the association between alcohol consumption and health outcomes. Using non-consumers as a  
416 reference group may overestimate the beneficial effects of alcohol<sup>40</sup>. On the other hand, in the  
417 meta-analysis of Di Castelnuovo *et al.*, there was still a protective effect of alcohol  
418 consumption in the general population after exclusion of former drinkers<sup>41</sup>. We were unable  
419 to take into account former alcohol consumption in all cohorts, but by restricting the analyses  
420 to alcohol consumers, possible confounding by abstinence or former alcohol consumption  
421 could not influence our results.

422

423 The association between alcoholic beverage preference and diabetes may be partly driven by  
424 obesity: since adiposity is on one causal pathway between absolute alcohol consumption and  
425 diabetes, adjusting for BMI may lead to overadjustment bias<sup>42</sup>. However, after multiple  
426 adjustments, those with a spirit preference had the highest BMI and persons with a beer or  
427 wine preference had the lowest BMI. Because BMI is a strong risk factor for developing

428 diabetes, the effect of moderate alcohol consumption might be strongest, in absolute terms, in  
429 those at higher risk. In our analysis, the association between spirit preference and diabetes  
430 incidence was higher among those with overweight or obese (BMI>25 kg/m<sup>2</sup>). In contrast,  
431 Beulens *et al.* found that moderate alcohol consumption was more strongly related to a  
432 reduced diabetes risk in overweight men and women than in their normal weight  
433 counterparts<sup>6</sup>. Moreover, in the French E3N-EPIC cohort, overweight women consuming two  
434 or more glasses of wine per day had a lower diabetes risk, whereas in normal weight women  
435 consuming the same amount, no association was observed<sup>33</sup>.

436

437 We aimed to disentangle beverage-specific effects, independent of those from the absolute  
438 ethanol consumption, by studying the association between alcoholic beverage preference and  
439 type 2 diabetes incidence. Beverage preference was used to classify the study population  
440 according to their alcohol intake. This approach of studying preference rather than absolute  
441 intake can thus be considered as a qualitative approach. Independent from the biological  
442 mechanisms associated with the chemical composition of the beverages, beverage preference  
443 *per se* may not be directly associated with diabetes incidence. Therefore, we have additionally  
444 studied the residuals of beer, wine and spirit intake, fully adjusted for total alcohol  
445 consumption by the residual method<sup>28</sup>; these analyses yielded similar findings. The number of  
446 cases distributed by beverage preference differed across the cohorts and were in some cohorts  
447 quite low, which may have affected the statistical power of the analyses. Non-consumers were  
448 excluded from the analysis to prevent non-consumers and former (heavy) consumers to affect  
449 the results. As a result, these findings only apply to alcohol consumers.

450

451 The CHANCES project is a large-scale multi-national collaboration of cohort studies  
452 including a large number of elderly persons. Pooled analyses of the individual participant data

453 from the different cohorts is a cost-efficient analytical approach and increases statistical  
454 power considerably. However, we had to rely on secondary data collected according to  
455 different study objectives and protocols, which may be a weakness<sup>15</sup>. With respect to type 2  
456 diabetes incidence, the identification and verification of diabetes cases varied across the  
457 cohorts. The oral glucose tolerance test is considered the gold standard of diabetes  
458 ascertainment, but recently fasting glucose has been shown to be the most accurate method of  
459 diabetes diagnosis<sup>43</sup>. Most cohort studies relied on self-reports, linkage with registries, HbA<sub>1c</sub>,  
460 or fasting blood glucose measures. As a result, misclassification could have been present and  
461 we might have underestimated the number of diabetes cases. However, only if this  
462 misclassification is differential and related to alcohol preference, would it have influenced the  
463 direction of the effect estimates, and yet our observed associations across cohorts were  
464 broadly consistent. Furthermore, it is difficult to distinguish between type 1 and type 2  
465 diabetes; therefore, some cohorts may not have been able to appropriately distinguish between  
466 the types. Moreover, this issue is not restricted to this study only. Diagnosing diabetes can be  
467 equivocal: the clinical diagnosis is based on a pre-specified cut-off point on a continuous  
468 scale of declining glycemic control, but clinical practice will dictate how assiduously the  
469 necessary tests are applied. Furthermore, the diagnosis is often based on the occurrence of  
470 complications of the disease and the disease can remain asymptomatic for years. Hence, it has  
471 been estimated that up to 50% of all type 2 diabetes patients are undiagnosed<sup>44</sup>. As a result,  
472 the true association may have been underestimated.

473

474 This meta-analysis of individual participant data from ten cohorts among Europeans who  
475 reported at least some alcohol consumption showed that beer, wine, and spirits were similarly  
476 associated with diabetes incidence. The recommendations of the American Diabetes  
477 Association for the prevention of diabetes suggest that if adults choose to drink alcohol, daily

478 intake should be limited to a moderate amount, i.e. no more than one drink per day for women  
479 and two drinks per day for men<sup>45</sup>. Our analysis offers little support for making beverage  
480 specific recommendation for diabetes prevention.

481

## 482 **Acknowledgements**

483 The research of DS was supported by the Dutch Beer Institute and the European Foundation  
484 for Alcohol Research (ERAB). The sponsor did not have any role in the design and conduct of  
485 the study; collection, management, analysis and interpretation of the data; and preparation,  
486 review or approval of the manuscript. No other authors declare conflicts of interest.

487 This analysis was part of the Consortium on Health and Ageing (CHANCES) project funded  
488 in the FP7 framework programme of the Directorate-General for Research & Innovation in  
489 the European Commission (grant 242244). The CHANCES project is coordinated by the  
490 Hellenic Health Foundation, Greece. Harmonization of the data from the MORGAM cohorts  
491 was also supported by European Union FP 7 project BiomarCaRE (278913).

492

493 DS, FK, EJMF designed the study and formulated the research question; OHF, DK, AT, TW,  
494 HB, KK, TL, SS, LI, and PB acquired the data and contributed reagents/materials/analysis  
495 tools; DS carried out the study, analyzed the data, and drafted the manuscript; All authors  
496 critically revised the manuscript for important intellectual content and approved of the final  
497 version to be published.

498

## 499 **References**

500 1. International Diabetes Federation. IDF Diabetes Atlas, Sixth Edition [Retrieved 12  
501 March 2015]. Available from: <http://www.idf.org/diabetesatlas> 2014.

- 502 2. Li XH, Yu FF, Zhou YH, He J. Association between alcohol consumption and the risk  
503 of incident type 2 diabetes: a systematic review and dose-response meta-analysis. *The*  
504 *American journal of clinical nutrition* 2016; **103**(3): 818-829. e-pub ahead of print  
505 2016/02/05; doi: 10.3945/ajcn.115.114389
- 506 3. Baliunas DO, Taylor BJ, Irving H, Roerecke M, Patra J, Mohapatra S *et al.* Alcohol as  
507 a risk factor for type 2 diabetes: A systematic review and meta-analysis. *Diabetes care*  
508 2009; **32**(11): 2123-2132. e-pub ahead of print 2009/10/31; doi: 10.2337/dc09-0227
- 509 4. Knott C, Bell S, Britton A. Alcohol Consumption and the Risk of Type 2 Diabetes: A  
510 Systematic Review and Dose-Response Meta-analysis of More Than 1.9 Million  
511 Individuals From 38 Observational Studies. *Diabetes care* 2015; **38**(9): 1804-1812. e-  
512 pub ahead of print 2015/08/22; doi: 10.2337/dc15-0710
- 513 5. Huang J, Wang X, Zhang Y. Specific Types of Alcoholic Beverage Consumption and  
514 Risk of Type 2 Diabetes: A Systematic Review and Meta-analysis. *Journal of diabetes*  
515 *investigation* 2016. e-pub ahead of print 2016/05/18; doi: 10.1111/jdi.12537
- 516 6. Beulens JW, van der Schouw YT, Bergmann MM, Rohrmann S, Schulze MB, Buijsse  
517 B *et al.* Alcohol consumption and risk of type 2 diabetes in European men and  
518 women: influence of beverage type and body size The EPIC-InterAct study. *Journal of*  
519 *internal medicine* 2012; **272**(4): 358-370. e-pub ahead of print 2012/02/23; doi:  
520 10.1111/j.1365-2796.2012.02532.x
- 521 7. Brien SE, Ronksley PE, Turner BJ, Mukamal KJ, Ghali WA. Effect of alcohol  
522 consumption on biological markers associated with risk of coronary heart disease:  
523 systematic review and meta-analysis of interventional studies. *Bmj* 2011; **342**: d636. e-  
524 pub ahead of print 2011/02/24; doi: 10.1136/bmj.d636
- 525 8. Schrieks IC, Heil AL, Hendriks HF, Mukamal KJ, Beulens JW. The effect of alcohol  
526 consumption on insulin sensitivity and glycemic status: a systematic review and meta-

- 527 analysis of intervention studies. *Diabetes care* 2015; **38**(4): 723-732. e-pub ahead of  
528 print 2015/03/26; doi: 10.2337/dc14-1556
- 529 9. Gronbaek M. Confounders of the relation between type of alcohol and cardiovascular  
530 disease. *Annals of Epidemiology* 2007; **17**(5): S13-S15. doi:  
531 10.1016/j.annepidem.2007.01.004
- 532 10. Sluik D, Bezemer R, Sierksma A, Feskens E. Alcoholic Beverage Preference and  
533 Dietary Habits: A Systematic Literature Review. *Critical reviews in food science and*  
534 *nutrition* 2015; 0. e-pub ahead of print 2015/02/13; doi:  
535 10.1080/10408398.2013.841118
- 536 11. Brand-Miller JC, Fatema K, Middlemiss C, Bare M, Liu V, Atkinson F *et al.* Effect of  
537 alcoholic beverages on postprandial glycemia and insulinemia in lean, young, healthy  
538 adults. *The American journal of clinical nutrition* 2007; **85**(6): 1545-1551.
- 539 12. Hatonen KA, Virtamo J, Eriksson JG, Perala MM, Sinkko HK, Leiviska J *et al.*  
540 Modifying effects of alcohol on the postprandial glucose and insulin responses in  
541 healthy subjects. *The American journal of clinical nutrition* 2012; **96**(1): 44-49. e-pub  
542 ahead of print 2012/06/01; doi: 10.3945/ajcn.111.031682
- 543 13. Arranz S, Chiva-Blanch G, Valderas-Martinez P, Medina-Remon A, Lamuela-  
544 Raventos RM, Estruch R. Wine, beer, alcohol and polyphenols on cardiovascular  
545 disease and cancer. *Nutrients* 2012; **4**(7): 759-781. e-pub ahead of print 2012/08/02;  
546 doi: 10.3390/nu4070759
- 547 14. Wacholder S, Schatzkin A, Freedman LS, Kipnis V, Hartman A, Brown CC. Can  
548 energy adjustment separate the effects of energy from those of specific  
549 macronutrients? *American journal of epidemiology* 1994; **140**(9): 848-855.
- 550 15. Boffetta P, Bobak M, Borsch-Supan A, Brenner H, Eriksson S, Grodstein F *et al.* The  
551 Consortium on Health and Ageing: Network of Cohorts in Europe and the United



- 552 States (CHANCES) project--design, population and data harmonization of a large-  
553 scale, international study. *European journal of epidemiology* 2014; **29**(12): 929-936.  
554 e-pub ahead of print 2014/12/17; doi: 10.1007/s10654-014-9977-1
- 555 16. Buijsse B, Feskens EJ, Kok FJ, Kromhout D. Cocoa intake, blood pressure, and  
556 cardiovascular mortality: the Zutphen Elderly Study. *Archives of internal medicine*  
557 2006; **166**(4): 411-417. e-pub ahead of print 2006/03/01; doi:  
558 10.1001/archinte.166.4.411
- 559 17. Hofman A, Brusselle GG, Darwish Murad S, van Duijn CM, Franco OH,  
560 Goedegebure A *et al.* The Rotterdam Study: 2016 objectives and design update.  
561 *European journal of epidemiology* 2015; **30**(8): 661-708. doi: 10.1007/s10654-015-  
562 0082-x
- 563 18. Trichopoulou A, Orfanos P, Norat T, Bueno-de-Mesquita B, Ocke MC, Peeters PH *et*  
564 *al.* Modified Mediterranean diet and survival: EPIC-elderly prospective cohort study.  
565 *Bmj* 2005; **330**(7498): 991. e-pub ahead of print 2005/04/12; doi:  
566 10.1136/bmj.38415.644155.8F
- 567 19. Jacobsen BK, Eggen AE, Mathiesen EB, Wilsgaard T, Njolstad I. Cohort profile: the  
568 Tromso Study. *Int J Epidemiol* 2012; **41**(4): 961-967. e-pub ahead of print  
569 2011/03/23; doi: 10.1093/ije/dyr049
- 570 20. Schottker B, Haug U, Schomburg L, Kohrle J, Perna L, Muller H *et al.* Strong  
571 associations of 25-hydroxyvitamin D concentrations with all-cause, cardiovascular,  
572 cancer, and respiratory disease mortality in a large cohort study. *The American journal*  
573 *of clinical nutrition* 2013; **97**(4): 782-793. e-pub ahead of print 2013/03/01; doi:  
574 10.3945/ajcn.112.047712

- 575 21. Evans A, Salomaa V, Kulathinal S, Asplund K, Cambien F, Ferrario M *et al.*  
576 MORGAM (an international pooling of cardiovascular cohorts). *Int J Epidemiol* 2005;  
577 **34(1)**: 21-27. e-pub ahead of print 2004/11/25; doi: 10.1093/ije/dyh327
- 578 22. Di Castelnuovo A, de Curtis A, Costanzo S, Persichillo M, Olivieri M, Zito F *et al.*  
579 Association of D-dimer levels with all-cause mortality in a healthy adult population:  
580 findings from the MOLI-SANI study. *Haematologica* 2013; **98(9)**: 1476-1480. e-pub  
581 ahead of print 2013/05/07; doi: 10.3324/haematol.2012.083410
- 582 23. Kuulasmaa K, Palosaari T, editors. Contributors from Partners of the Consortium on  
583 Health and Ageing: Network of Cohorts in Europe and the United States  
584 (CHANCES). CHANCES cohort descriptions, assessment of the availability and  
585 quality of data, and definitions of variables. MORGAM Project e-publications  
586 [Internet]. 2015; (6). URN:NBN:fi-fe201501151161. Available from  
587 URL:[http://www.thl.fi/publications/morgam/chances\\_d9/index.html](http://www.thl.fi/publications/morgam/chances_d9/index.html). Available from:  
588 URL (Accessed n Date Accessed Year)|.
- 589 24. Klipstein-Grobusch K, den Breeijen JH, Goldbohm RA, Geleijnse JM, Hofman A,  
590 Grobbee DE *et al.* Dietary assessment in the elderly: validation of a semiquantitative  
591 food frequency questionnaire. *European journal of clinical nutrition* 1998; **52(8)**: 588-  
592 596. e-pub ahead of print 1998/09/02;
- 593 25. Riboli E, Kaaks R. The EPIC Project: rationale and study design. European  
594 Prospective Investigation into Cancer and Nutrition. *Int J Epidemiol* 1997; **26 Suppl**  
595 **1**: S6-14. e-pub ahead of print 1997/01/01;
- 596 26. Sluik D, van Lee L, Geelen A, Feskens EJ. Alcoholic beverage preference and diet in  
597 a representative Dutch population: the Dutch national food consumption survey 2007-  
598 2010. *European journal of clinical nutrition* 2014; **68(3)**: 287-294. e-pub ahead of  
599 print 2014/01/09; doi: 10.1038/ejcn.2013.279

- 600 27. Sluik D, Jankovic N, O'Doherty MG, Geelen A, Schottker B, Rolandsson O *et al.*  
601 Alcoholic Beverage Preference and Dietary Habits in Elderly across Europe: Analyses  
602 within the Consortium on Health and Ageing: Network of Cohorts in Europe and the  
603 United States (CHANCES) Project. *PloS one* 2016; **11**(8): e0161603. e-pub ahead of  
604 print 2016/08/23; doi: 10.1371/journal.pone.0161603
- 605 28. Willett WC, Howe GR, Kushi LH. Adjustment for total energy intake in  
606 epidemiologic studies. *The American journal of clinical nutrition* 1997; **65**(4 Suppl):  
607 1220S-1228S; discussion 1229S-1231S. e-pub ahead of print 1997/04/01;
- 608 29. Jankovic N, Geelen A, Streppel MT, de Groot LC, Orfanos P, van den Hooven EH *et*  
609 *al.* Adherence to a healthy diet according to the World Health Organization guidelines  
610 and all-cause mortality in elderly adults from Europe and the United States. *American*  
611 *journal of epidemiology* 2014; **180**(10): 978-988. e-pub ahead of print 2014/10/17;  
612 doi: 10.1093/aje/kwu229
- 613 30. Sterne JA, White IR, Carlin JB, Spratt M, Royston P, Kenward MG *et al.* Multiple  
614 imputation for missing data in epidemiological and clinical research: potential and  
615 pitfalls. *Bmj* 2009; **338**: b2393. e-pub ahead of print 2009/07/01; doi:  
616 10.1136/bmj.b2393
- 617 31. Sauerbrei W, Meier-Hirmer C, Benner A, Royston P. Multivariable regression model  
618 building by using fractional polynomials: Description of SAS, STATA, and R  
619 programs. *Computational Statistics & Data Analysis* 2006; **50**: 3464-3485.
- 620 32. Hodge AM, English DR, O'Dea K, Giles GG. Alcohol intake, consumption pattern  
621 and beverage type, and the risk of Type 2 diabetes. *Diabetic medicine : a journal of*  
622 *the British Diabetic Association* 2006; **23**(6): 690-697. doi: 10.1111/j.1464-  
623 5491.2006.01864.x

- 624 33. Fagherazzi G, Vilier A, Lajous M, Boutron-Ruault MC, Balkau B, Clavel-Chapelon F  
625 *et al.* Wine consumption throughout life is inversely associated with type 2 diabetes  
626 risk, but only in overweight individuals: results from a large female French cohort  
627 study. *European journal of epidemiology* 2014; **29**(11): 831-839. doi:  
628 10.1007/s10654-014-9955-7
- 629 34. Conigrave KM, Hu BF, Camargo CA, Jr., Stampfer MJ, Willett WC, Rimm EB. A  
630 prospective study of drinking patterns in relation to risk of type 2 diabetes among men.  
631 *Diabetes* 2001; **50**(10): 2390-2395.
- 632 35. Kao WH, Puddey IB, Boland LL, Watson RL, Brancati FL. Alcohol consumption and  
633 the risk of type 2 diabetes mellitus: atherosclerosis risk in communities study.  
634 *American journal of epidemiology* 2001; **154**(8): 748-757. e-pub ahead of print  
635 2001/10/09;
- 636 36. Wannamethee SG, Camargo CA, Jr., Manson JE, Willett WC, Rimm EB. Alcohol  
637 drinking patterns and risk of type 2 diabetes mellitus among younger women. *Archives*  
638 *of internal medicine* 2003; **163**(11): 1329-1336. e-pub ahead of print 2003/06/11; doi:  
639 10.1001/archinte.163.11.1329
- 640 37. Chiva-Blanch G, Urpi-Sarda M, Ros E, Valderas-Martinez P, Casas R, Arranz S *et al.*  
641 Effects of red wine polyphenols and alcohol on glucose metabolism and the lipid  
642 profile: a randomized clinical trial. *Clinical nutrition* 2013; **32**(2): 200-206. e-pub  
643 ahead of print 2012/09/25; doi: 10.1016/j.clnu.2012.08.022
- 644 38. Gronbaek M. Factors influencing the relation between alcohol and mortality--with  
645 focus on wine. *Journal of internal medicine* 2001; **250**(4): 291-308. e-pub ahead of  
646 print 2001/09/29;
- 647 39. Rehm J, Irving H, Ye Y, Kerr WC, Bond J, Greenfield TK. Are lifetime abstainers the  
648 best control group in alcohol epidemiology? On the stability and validity of reported

- 649 lifetime abstention. *American journal of epidemiology* 2008; **168**(8): 866-871. e-pub  
650 ahead of print 2008/08/15; doi: 10.1093/aje/kwn093
- 651 40. Shaper AG, Wannamethee G, Walker M. Alcohol and mortality in British men:  
652 explaining the U-shaped curve. *Lancet* 1988; **2**(8623): 1267-1273. e-pub ahead of  
653 print 1988/12/03;
- 654 41. Di Castelnuovo A, Costanzo S, Bagnardi V, Donati MB, Iacoviello L, de Gaetano G.  
655 Alcohol dosing and total mortality in men and women: an updated meta-analysis of 34  
656 prospective studies. *Archives of internal medicine* 2006; **166**(22): 2437-2445. e-pub  
657 ahead of print 2006/12/13; doi: 10.1001/archinte.166.22.2437
- 658 42. Schisterman EF, Cole SR, Platt RW. Overadjustment bias and unnecessary adjustment  
659 in epidemiologic studies. *Epidemiology* 2009; **20**(4): 488-495. e-pub ahead of print  
660 2009/06/16; doi: 10.1097/EDE.0b013e3181a819a1
- 661 43. Collaboration NCDRF. Effects of diabetes definition on global surveillance of  
662 diabetes prevalence and diagnosis: a pooled analysis of 96 population-based studies  
663 with 331 288 participants. *The lancet. Diabetes & endocrinology* 2015; **3**(8): 624-637.  
664 e-pub ahead of print 2015/06/26; doi: 10.1016/S2213-8587(15)00129-1
- 665 44. Ryden L, Standl E, Bartnik M, Van den Berghe G, Betteridge J, de Boer MJ *et al.*  
666 Guidelines on diabetes, pre-diabetes, and cardiovascular diseases: executive summary.  
667 The Task Force on Diabetes and Cardiovascular Diseases of the European Society of  
668 Cardiology (ESC) and of the European Association for the Study of Diabetes (EASD).  
669 *European heart journal* 2007; **28**(1): 88-136. e-pub ahead of print 2007/01/16; doi:  
670 10.1093/eurheartj/ehl260
- 671 45. American Diabetes Association. Nutrition recommendations and interventions for  
672 diabetes: a position statement of the American Diabetes Association. *Diabetes care*  
673 2008; **31 Suppl 1**: S61-78. e-pub ahead of print 2008/01/10; doi: 10.2337/dc08-S061

- 674 46. Feskens EJ, Loeber JG, Kromhout D. Diet and physical activity as determinants of  
675 hyperinsulinemia: the Zutphen Elderly Study. *American journal of epidemiology*. Aug  
676 15 1994;**140**(4):350-360.
- 677 47. Beulens JW, Stolk RP, van der Schouw YT, Grobbee DE, Hendriks HF, Bots ML.  
678 Alcohol consumption and risk of type 2 diabetes among older women. *Diabetes Care*.  
679 Dec 2005;**28**(12):2933-2938.
- 680 48. Kulathinal S, Niemelä M, Niiranen T, Saarela O, Palosaari T, Tapanainen H, et al.  
681 Description of MORGAM Cohorts. MORGAM Project e-publications [Internet].  
682 2005-; (2). URN:NBN:fi-fe20051214. Available from URL:  
683 <http://www.thl.fi/publications/morgam/cohorts/index.html>.

684 **Figure legends:**

685

686 **Figure 1.** Forest plot with pooled Hazard Ratios (HRs) and 95% Confidence Intervals (CIs)  
687 for the association between beer preference and diabetes incidence compared to having no  
688 preference adjusted for age, sex, education, employment status, prevalent coronary heart  
689 disease and cancer, smoking status, physical activity (if available), Healthy Diet Indicator  
690 score (if available), and BMI.

691

692 **Figure 2.** Forest plot with pooled Hazard Ratios (HRs) and 95% Confidence Intervals (CIs)  
693 for the association between wine preference and diabetes incidence compared to having no  
694 preference adjusted for age, sex, education, employment status, prevalent coronary heart  
695 disease and cancer, smoking status, physical activity (if available), Healthy Diet Indicator  
696 score (if available), and BMI.

697

698 **Figure 3.** Forest plot with pooled Hazard Ratios (HRs) and 95% Confidence Intervals (CIs)  
699 for the association between spirit preference and diabetes incidence compared to having no  
700 preference adjusted for age, sex, education, employment status, prevalent coronary heart  
701 disease and cancer, smoking status, physical activity (if available), Healthy Diet Indicator  
702 score (if available), and BMI.

703


704 **Figure 4.** Forest plot with pooled Hazard Ratios (HRs) and 95% Confidence Intervals (CIs)  
705 for the association between residuals of beer consumption per 6 g/d and diabetes incidence  
706 adjusted for absolute alcohol intake, age, sex, education, employment status, prevalent  
707 coronary heart disease and cancer, smoking status, physical activity (if available), Healthy  
708 Diet Indicator score (if available), and BMI.

709 **Figure 5.** Forest plot with pooled Hazard Ratios (HRs) and 95% Confidence Intervals (CIs)  
710 for the association between residuals of wine consumption per 6 g/d and diabetes incidence  
711 adjusted for absolute alcohol intake, age, sex, education, employment status, prevalent  
712 coronary heart disease and cancer, smoking status, physical activity (if available), Healthy  
713 Diet Indicator score (if available), and BMI.


714 **Figure 6.** Forest plot with pooled Hazard Ratios (HRs) and 95% Confidence Intervals (CIs)  
715 for the association between residuals of spirit consumption per 6 g/d and diabetes incidence  
716 adjusted for absolute alcohol intake, age, sex, education, employment status, prevalent  
717 coronary heart disease and cancer, smoking status, physical activity (if available), Healthy  
718 Diet Indicator score (if available), and BMI.




**Figure 1.** Forest plot with pooled Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) for the association between beer preference and diabetes incidence compared to having no preference adjusted for age, sex, education, employment status, prevalent coronary heart disease and cancer, smoking status, physical activity (if available), Healthy Diet Indicator score (if available), and BMI.




**Figure 2.** Forest plot with pooled Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) for the association between wine preference and diabetes incidence compared to having no preference adjusted for age, sex, education, employment status, prevalent coronary heart disease and cancer, smoking status, physical activity (if available), Healthy Diet Indicator score (if available), and BMI.




**Figure 1.** Forest plot with pooled Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) for the association between beer preference and diabetes incidence compared to having no preference adjusted for age, sex, education, employment status, prevalent coronary heart disease and cancer, smoking status, physical activity (if available), Healthy Diet Indicator score (if available), and BMI.




**Figure 2.** Forest plot with pooled Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) for the association between wine preference and diabetes incidence compared to having no preference adjusted for age, sex, education, employment status, prevalent coronary heart disease and cancer, smoking status, physical activity (if available), Healthy Diet Indicator score (if available), and BMI.




**Figure 3.** Forest plot with pooled Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) for the association between spirit preference and diabetes incidence compared to having no preference adjusted for age, sex, education, employment status, prevalent coronary heart disease and cancer, smoking status, physical activity (if available), Healthy Diet Indicator score (if available), and BMI.




**Figure 4.** Forest plot with pooled Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) for the association between the residuals of beer consumption per 6 g/d and diabetes incidence adjusted for absolute alcohol intake, age, sex, education, employment status, prevalent coronary heart disease and cancer, smoking status, physical activity (if available), Healthy Diet Indicator score (if available), and BMI.



**Figure 5.** Forest plot with pooled Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) for the association between the residuals of wine consumption per 6 g/d and diabetes incidence adjusted for absolute alcohol intake, age, sex, education, employment status, prevalent coronary heart disease and cancer, smoking status, physical activity (if available), Healthy Diet Indicator score (if available), and BMI.




**Figure 6.** Forest plot with pooled Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) for the association between the residuals of spirit consumption per 6 g/d and diabetes incidence adjusted for absolute alcohol intake, age, sex, education, employment status, prevalent coronary heart disease and cancer, smoking status, physical activity (if available), Healthy Diet Indicator score (if available), and BMI.





**Supplemental Figure 1:** Participant flow-charts of the ten included European cohort studies from the Consortium on Health and Ageing: Network of Cohorts in Europe and the United States (CHANCES) project.



**Supplemental Table 1.** Selected general characteristics across categories of alcoholic beverage preference and Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) of the associations between having a beer, wine, or spirit preference and diabetes incidence compared to having no preference in 10 European cohorts based on multiple imputation results.

	Beer preference	Wine preference	Spirit preference	No preference
<b><u>Zutphen Elderly Study</u></b>				
N (%)	65 (12)	73 (13)	344 (62)	75 (13)
Age, years	70.5 (5.8)	72.1 (5.0)	72.2 (5.2)	71.8 (5.5)
Men, %	100	100	100	100
Ethanol consumption, g/day	13.5 [4.0-25.0]	3.8 [1.7-12.0]	18.8 [6.2-37.5]	20.3 [8.4-44.9]
University or college education, %	0	14	3	10
Current smokers, %	32	23	41	29
BMI, kg/m <sup>2</sup>				
- Crude	26.0 (0.4)	24.7 (0.4)*	25.5 (0.2)	25.3 (0.4)
- Adjusted <sup>a</sup>	25.9 (0.4)	25.0 (0.4)	25.5 (0.2)	25.3 (0.4)
Diabetes cases / Person Years	6 / 756	9 / 747	27 / 3529	6 / 848
HR (95% CI): Model 1	1.00 (0.32-3.18)	1.34 (0.46-3.89)	0.95 (0.39-2.32)	1.00 (ref)
HR (95% CI): Model 2	1.04 (0.32-4.15)	1.46 (0.52-4.15)	0.96 (0.38-2.42)	1.00 (ref)
HR (95% CI): Model 3	1.00 (0.31-3.27)	1.60 (0.55-4.63)	1.07 (0.42-2.77)	1.00 (ref)
<b><u>Rotterdam Study</u></b>				
N (%)	182 (7)	1292 (53)	582 (24)	372 (15)
Age, years	62.4 (5.8)	64.8 (6.8)	66.2 (6.5)	64.0 (6.1)

**Supplemental Table 1 (continued).** Selected general characteristics across categories of alcoholic beverage preference and Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) of the associations between having a beer, wine, or spirit preference and diabetes incidence compared to having no preference in 10 European cohorts based on multiple imputation results.

	<b>Beer preference</b>	<b>Wine preference</b>	<b>Spirit preference</b>	<b>No preference</b>
Men, %	88	17	72	71
Ethanol consumption, g/day	11.7 [3.0-25.7]	2.9 [0.6-9.1]	18.9 [7.5-31.8]	11.3 [3.6-24.9]
University or college education, %	13	8	11	18
Current smokers, %	31	17	28	22
BMI, kg/m <sup>2</sup>				
- Crude	25.4 (0.3)	26.1 (0.1)*	26.2 (0.1)*	25.7 (0.2)*****
- Adjusted <sup>a</sup>	25.6 (0.3)	26.0 (0.1)	26.4 (0.2)***	26.0 (0.2)***
Diabetes cases / Person Years	30 / 1756	155 / 13447	100 / 5321	32 / 3920
HR (95% CI): Model 1	2.10 (1.27-3.47)	1.43 (0.96-2.15)	2.30 (1.54-3.43)	1.00 (ref)
HR (95% CI): Model 2	2.15 (1.29-3.57)	1.48 (0.98-2.24)	2.38 (1.58-3.57)	1.00 (ref)
HR (95% CI): Model 3	2.21 (1.33-3.68)	1.49 (0.99-2.24)	2.28 (1.52-3.43)	1.00 (ref)
<b><u>EPIC-Elderly the Netherlands</u></b>				
N (%)	82 (2)	2802 (74)	384 (10)	525 (14)
Age, years	63.7 (2.6)	64.2 (2.7)	64.1 (2.5)	64.2 (2.7)
Men, %	39	2	13	12
Ethanol consumption, g/day	7.3 [1.3-20.1]	4.4 [1.3-13.1]	10.1 [1.8-25.7]	4.9 [1.5-12.8]
University or college education, %	7	16	8	12
Current smokers, %	40	15	29	17

**Supplemental Table 1 (continued).** Selected general characteristics across categories of alcoholic beverage preference and Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) of the associations between having a beer, wine, or spirit preference and diabetes incidence compared to having no preference in 10 European cohorts based on multiple imputation results.

	Beer preference	Wine preference	Spirit preference	No preference
BMI, kg/m <sup>2</sup>				
- Crude	25.2 (0.4)	25.5 (0.1)	26.4 (0.2)*, **	25.6 (0.2)***
- Adjusted <sup>a</sup>	25.0 (0.4)	25.7 (0.1)	26.3 (0.2)*, **	25.6 (0.2)***
Diabetes cases / Person Years	4 / 416	167 / 13601	26 / 1927	22 / 2559
HR (95% CI): Model 1	0.75 (0.25-2.22)	1.28 (0.81-2.02)	1.22 (0.69-2.17)	1.00 (ref)
HR (95% CI): Model 2	0.88 (0.29-2.57)	1.23 (0.78-1.94)	1.20 (0.67-2.13)	1.00 (ref)
HR (95% CI): Model 3	0.91 (0.30-2.74)	1.14 (0.72-1.80)	1.11 (0.63-1.98)	1.00 (ref)
<b><u>EPIC-Elderly Greece</u></b>				
N (%)	510 (10)	2561 (48)	361 (7)	1884 (35)
Age, years	66.2 (4.5)	67.4 (4.5)	67.0 (4.3)	66.3 (4.3)
Men, %	47	46	74	52
Ethanol consumption, g/day	1.3 [0.6-8.5]	8.0 [1.2-16.0]	13.2 [4.4-21.3]	2.8 [1.3-12.5]
University or college education, %	4	2	3	7
Current smokers, %	12	13	29	16
BMI, kg/m <sup>2</sup>				
- Crude	29.2 (0.2)	28.9 (0.1)	28.9 (0.2)	28.8 (0.1)
- Adjusted <sup>a</sup>	29.0 (0.2)	28.7 (0.1)	29.4 (0.2)**	29.0 (0.1)
Diabetes cases / Person Years	72 / 5407	377 / 28107	58 / 3717	308 / 20618
HR (95% CI): Model 1	0.99 (0.77-1.28)	0.78 (0.67-0.91)	1.37 (1.03-1.82)	1.00 (ref)

**Supplemental Table 1 (continued).** Selected general characteristics across categories of alcoholic beverage preference and Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) of the associations between having a beer, wine, or spirit preference and diabetes incidence compared to having no preference in 10 European cohorts based on multiple imputation results.

	<b>Beer preference</b>	<b>Wine preference</b>	<b>Spirit preference</b>	<b>No preference</b>
HR (95% CI): Model 2	0.98 (0.76-1.27)	0.80 (0.69-0.94)	1.32 (1.00-1.76)	1.00 (ref)
HR (95% CI): Model 3	0.92 (0.71-1.19)	0.82 (0.70-0.95)	1.27 (0.95-1.68)	1.00 (ref)
<b><u>EPIC-Elderly Sweden</u></b>				
N (%)	958 (34)	413 (15)	137 (5)	1311 (47)
Age, years	60.4 (1.3)	60.3 (0.9)	60.4 (1.2)	60.3 (0.8)
Men, %	58	15	77	54
Ethanol consumption, g/day	0.9 [0.3-2.7]	1.6 [0.1-3.3]	2.0 [0.2-2.4]	2.9 [0.4-5.4]
University or college education, %	11	18	5	13
Current smokers, %	13	13	40	22
<b>BMI, kg/m<sup>2</sup></b>				
- Crude	25.6 (0.1)	25.5 (0.2)	26.7 (0.3)*,**	25.9 (0.1)***
- Adjusted <sup>a</sup>	25.5 (0.1)	25.5 (0.2)	26.8 (0.3)*,**	26.0 (0.1)*,***
Diabetes cases / Person Years	33 / 12680	14 / 5406	9 / 1821	53 / 17276
HR (95% CI): Model 1	0.81 (0.52-1.25)	1.00 (0.54-1.86)	1.34 (0.65-2.74)	1.00 (ref)
HR (95% CI): Model 2	0.80 (0.52-1.25)	1.00 (0.53-1.86)	1.31 (0.63-2.81)	1.00 (ref)
HR (95% CI): Model 3	0.81 (0.52-1.26)	1.00 (0.53-1.87)	0.92 (0.44-1.93)	1.00 (ref)
<b><u>The Tromsø Study</u></b>				
N (%)	722 (15)	1502 (30)	1042 (21)	1692 (34)

**Supplemental Table 1 (continued).** Selected general characteristics across categories of alcoholic beverage preference and Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) of the associations between having a beer, wine, or spirit preference and diabetes incidence compared to having no preference in 10 European cohorts based on multiple imputation results.

	<b>Beer preference</b>	<b>Wine preference</b>	<b>Spirit preference</b>	<b>No preference</b>
Age, years	59.6 (8.4)	59.8 (8.8)	62.1 (8.6)	58.7 (7.7)
Men, %	76	28	74	71
Ethanol consumption, g/day	3.3 [1.7-6.1]	4.7 [2.4-7.1]	1.1 [1.1-2.6]	4.7 [2.8-8.1]
University or college education, %	18	33	10	28
Current smokers, %	38	29	50	34
BMI, kg/m <sup>2</sup>				
- Crude	25.4 (0.1)	25.4 (0.1)	25.9 (0.1)*,**	25.9 (0.1)*,**
- Adjusted <sup>a</sup>	25.3 (0.1)	25.6 (0.1)	25.9 (0.1)*,**	25.8 (0.1)**
Diabetes cases / Person Years	45 / 9158	65 / 19853	72 / 12715	96 / 22094
HR (95% CI): Model 1	1.12 (0.78-1.60)	0.95 (0.68-1.34)	1.32 (0.96-1.81)	1.00 (ref)
HR (95% CI): Model 2	1.12 (0.78-1.60)	0.96 (0.68-1.34)	1.32 (0.96-1.81)	1.00 (ref)
HR (95% CI): Model 3	1.23 (0.86-1.77)	1.03 (0.73-1.44)	1.17 (0.85-1.61)	1.00 (ref)
<b><u>ESTHER</u></b>				
N (%)	1466 (28)	2305 (44)	107 (2)	1408 (27)
Age, years	61.5 (6.4)	61.5 (6.6)	63.2 (6.8)	62.0 (6.6)
Men, %	77	32	22	64
Ethanol consumption, g/day	6.6 [2.6-13.2]	5.6 [3.7-11.0]	0.8 [0.8-2.5]	9.1 [5.3-15.8]
Middle education, %	21	35	25	34
Current smokers, %	26	13	13	14

**Supplemental Table 1 (continued).** Selected general characteristics across categories of alcoholic beverage preference and Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) of the associations between having a beer, wine, or spirit preference and diabetes incidence compared to having no preference in 10 European cohorts based on multiple imputation results.

	Beer preference	Wine preference	Spirit preference	No preference
BMI, kg/m <sup>2</sup>				
- Crude	27.6 (0.1)	26.8 (0.1)*	28.0 (0.4)**	27.1 (0.1)*, **, ***
- Adjusted <sup>a</sup>	27.4 (0.1)	27.0 (0.1)*	28.2 (0.4)*, **	27.1 (0.1)*, ***
Diabetes cases / Person Years	165 / 9671	207 / 16394	14 / 736	140 / 9792
HR (95% CI): Model 1	1.07 (0.85-1.35)	0.98 (0.79-1.23)	1.42 (0.81-2.47)	1.00 (ref)
HR (95% CI): Model 2	1.05 (0.83-1.32)	0.97 (0.78-1.22)	1.39 (0.80-2.43)	1.00 (ref)
HR (95% CI): Model 3	0.98 (0.78-1.24)	0.96 (0.76-1.20)	1.25 (0.71-2.18)	1.00 (ref)
<b><u>MORGAM: FINRISK</u></b>				
N (%)	6410 (32)	3200 (16)	3410 (17)	6739 (34)
Age, years	41.5 (11.5)	47.3 (12.3)	46.8 (11.8)	44.6 (11.7)
Men, %	63	26	66	62
Ethanol consumption, g/day	7.0 [4.0-15.0]	3.0 [2.0-9.0]	8.0 [3.0-14.0]	13.0 [7.0-21.0]
University or college education, %	9	19	6	15
Current smokers, %	38	17	34	29
BMI, kg/m <sup>2</sup>				
- Crude	25.7 (0.1)	25.9 (0.1)*	25.9 (0.1)*, **	26.2 (0.1)*, **, ***
- Adjusted <sup>aw</sup>	25.9 (0.1)	26.1 (0.1)	25.5 (0.1)*, **	26.2 (0.1)*, ***
Diabetes cases / Person Years	487 / 101570	229 / 50342	437 / 59893	557 / 105820
HR (95% CI): Model 1	0.98 (0.87-1.11)	0.89 (0.76-1.05)	1.08 (0.95-1.22)	1.00 (ref)

**Supplemental Table 1 (continued).** Selected general characteristics across categories of alcoholic beverage preference and Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) of the associations between having a beer, wine, or spirit preference and diabetes incidence compared to having no preference in 10 European cohorts based on multiple imputation results.

	<b>Beer preference</b>	<b>Wine preference</b>	<b>Spirit preference</b>	<b>No preference</b>
HR (95% CI): Model 2	0.97 (0.86-1.10)	0.91 (0.78-1.07)	1.07 (0.94-1.21)	1.00 (ref)
HR (95% CI): Model 3	1.05 (0.93-1.18)	0.94 (0.80-1.10)	1.00 (0.88-1.13)	1.00 (ref)
<b><u>MORGAM: Northern-Sweden</u></b>				
N (%)	794 (26)	511 (17)	146 (5)	1581 (52)
Age, years	45.5 (13.4)	46.6 (11.6)	49.7 (11.6)	44.3 (11.7)
Men, %	72	10	81	65
Ethanol consumption, g/day	2.0 [1.0-6.0]	2.0 [1.0-2.0]	2.0 [1.0-2.0]	4.0 [3.0-6.0]
University or college education, %	15	26	2	19
Current smokers, %	23	27	40	30
<b>BMI, kg/m<sup>2</sup></b>				
- Crude	25.4 (0.1)	24.8 (0.2)*	26.7 (0.3)*, **, ***	25.2 (0.1)** , ***
- Adjusted <sup>a</sup>	25.2 (0.2)	25.2 (0.2)	26.1 (0.3)*, **	25.2 (0.1)***
Diabetes cases / Person Years	40 / 15215	25 / 10413	7 / 2728	77 / 31392
HR (95% CI): Model 1	0.99 (0.67-1.46)	1.05 (0.63-1.73)	0.88 (0.40-1.93)	1.00 (ref)
HR (95% CI): Model 2	1.04 (0.71-1.54)	1.10 (0.66-1.83)	0.87 (0.40-1.90)	1.00 (ref)
HR (95% CI): Model 3	1.06 (0.72-1.56)	1.15 (0.69-1.93)	0.68 (0.31-1.50)	1.00 (ref)
<b><u>MORGAM: MOLI-SANI</u></b>				
N (%)	618 (4)	11522 (79)	102 (1)	2268 (16)



**Supplemental Table 1 (continued).** Selected general characteristics across categories of alcoholic beverage preference and Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) of the associations between having a beer, wine, or spirit preference and diabetes incidence compared to having no preference in 10 European cohorts based on multiple imputation results.

	<b>Beer preference</b>	<b>Wine preference</b>	<b>Spirit preference</b>	<b>No preference</b>
Age, years	49.6 (9.9)	56.7 (11.7)	50.7 (9.3)	49.6 (9.7)
Men, %	62	60	34	64
Ethanol consumption, g/day	4.0 [1.0-17.0]	18.0 [10.0-34.0]	2.0 [2.0-7.0]	8.0 [3.0-20.0]
University or college education, %	13	12	14	17
Current smokers, %	35	22	25	27
BMI, kg/m <sup>2</sup>				
- Crude	27.3 (0.2)	27.8 (0.0)*	26.9 (0.4)**	27.6 (0.1)**
- Adjusted <sup>a</sup>	27.7 (0.2)	27.7 (0.0)	27.5 (0.4)	28.0 (0.1)**
Diabetes cases / Person Years	10 / 2758	255 / 20198	0 / 478	30 / 10319
HR (95% CI): Model 1	1.21 (0.59-2.49)	1.17 (0.80-1.73)	no cases	1.00 (ref)
HR (95% CI): Model 2	1.19 (0.58-2.44)	1.17 (0.80-1.73)	no cases	1.00 (ref)
HR (95% CI): Model 3	1.29 (0.63-2.65)	1.22 (0.89-1.80)	no cases	1.00 (ref)

\* P-value <0.05 versus beer preference; \*\* P-value <0.05 versus wine preference; \*\*\* P-value <0.05 versus spirit preference.

<sup>a</sup> BMI adjusted for age, sex, education, employment, prevalent coronary heart disease or cancer, smoking status, sports activity (if available), and Healthy Diet Indicator (if available).

Model 1: Adjusted for age, sex, education, employment, and prevalent coronary heart disease or cancer;

Model 2: Model 1 additionally adjusted for smoking status, sports activity (if available), and Healthy Diet Indicator score (if available).

Model 3: Model 2 additionally adjusted for BMI.

**Supplemental Table 2.** Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) of the associations between residuals of absolute consumption of beer, wine, and spirits and diabetes incidence in 10 European cohorts.

<b>Zutphen Elderly Study</b>	<b>Q1</b>	<b>Q2</b>	<b>Q3</b>	<b>Per 6 g/d</b>
<b>Beer consumption (residuals)</b>				
N	185	186	186	557
Cases / Person Years	14 / 1968	18 / 1903	16 / 2009	48 / 5880
HR (95% CI): Model 1	1.00 (ref)	1.47 (0.71-3.02)	1.25 (0.60-2.64)	0.98 (0.80-1.20)
HR (95% CI): Model 2	1.00 (ref)	1.47 (0.71-3.05)	1.30 (0.61-2.77)	0.98 (0.79-1.21)
HR (95% CI): Model 3	1.00 (ref)	1.47 (0.70-3.07)	1.23 (0.58-2.62)	0.97 (0.77-1.21)
<b>Wine consumption (residuals)</b>				
N	190	181	186	557
Cases / Person Years	15 / 1915	14 / 1859	19 / 2106	48 / 5880
HR (95% CI): Model 1	1.00 (ref)	0.92 (0.44-1.91)	1.25 (0.62-2.52)	0.89 (0.64-1.23)
HR (95% CI): Model 2	1.00 (ref)	0.93 (0.45-1.95)	1.19 (0.58-2.43)	0.85 (0.60-1.20)
HR (95% CI): Model 3	1.00 (ref)	0.90 (0.43-1.90)	1.12 (0.55-2.30)	0.84 (0.59-1.21)
<b>Spirit consumption (residuals)</b>				
N	189	184	184	557
Cases / Person Years	20 / 2095	13 / 1952	15 / 1833	48 / 5880
HR (95% CI): Model 1	1.00 (ref)	0.69 (0.34-1.41)	0.77 (0.39-1.54)	1.06 (0.88-1.26)
HR (95% CI): Model 2	1.00 (ref)	0.73 (0.35-1.50)	0.81 (0.41-1.63)	1.07 (0.89-1.28)
HR (95% CI): Model 3	1.00 (ref)	0.81 (0.39-1.68)	0.94 (0.47-1.90)	1.08 (0.89-1.31)

**Supplemental Table 2 (continued).** Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) of the associations between residuals of absolute consumption of beer, wine, and spirits and diabetes incidence in 10 European cohorts.

<b>Rotterdam Study</b>	<b>Q1</b>	<b>Q2</b>	<b>Q3</b>	<b>Per 6 g/d</b>
<b>Beer consumption (residuals)</b>				
N (%)	809	815	804	2428
Cases / Person Years	109 / 8118	112 / 8283	96 / 8042	317 / 24444
HR (95% CI): Model 1	1.00 (ref)	1.07 (0.82-1.41)	0.88 (0.67-1.17)	0.98 (0.87-1.10)
HR (95% CI): Model 2	1.00 (ref)	1.08 (0.82-1.42)	0.88 (0.66-1.17)	0.97 (0.86-1.10)
HR (95% CI): Model 3	1.00 (ref)	1.06 (0.80-1.39)	0.87 (0.62-1.15)	0.97 (0.87-1.09)
<b>Wine consumption (residuals)</b>				
N (%)	810	819	799	2428
Cases / Person Years	125 / 7627	99 / 8250	93 / 8567	317 / 24444
HR (95% CI): Model 1	1.00 (ref)	0.74 (0.55-1.00)	0.67 (0.50-0.90)	0.89 (0.79-1.01)
HR (95% CI): Model 2	1.00 (ref)	0.74 (0.55-1.00)	0.68 (0.50-0.91)	0.90 (0.79-1.01)
HR (95% CI): Model 3	1.00 (ref)	0.79 (0.54-0.98)	0.72 (0.54-0.98)	0.92 (0.82-1.04)
<b>Spirit consumption (residuals)</b>				
N (%)	810	819	799	2428
Cases / Person Years	89 / 8471	110 / 8359	118 / 7613	317 / 24444
HR (95% CI): Model 1	1.00 (ref)	1.28 (0.96-1.70)	1.43 (1.08-1.91)	1.09 (0.99-1.19)
HR (95% CI): Model 2	1.00 (ref)	1.30 (0.97-1.72)	1.41 (1.06-1.87)	1.09 (0.99-1.19)
HR (95% CI): Model 3	1.00 (ref)	1.26 (0.95-1.68)	1.33 (1.00-1.77)	1.07 (0.98-1.17)

**Supplemental Table 2 (continued).** Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) of the associations between residuals of absolute consumption of beer, wine, and spirits and diabetes incidence in 10 European cohorts.

<b>EPIC-ELDERLY: The Netherlands</b>	<b>Q1</b>	<b>Q2</b>	<b>Q3</b>	<b>Per 6 g/d</b>
<b>Beer consumption (residuals)</b>				
N	1264	1265	1264	3793
Cases / Person Years	67 / 6097	71 / 6120	81 / 6286	219 / 18503
HR (95% CI): Model 1	1.00 (ref)	1.01 (0.72-1.41)	0.97 (0.69-1.35)	1.07 (0.85-1.36)
HR (95% CI): Model 2	1.00 (ref)	0.95 (0.67-1.35)	0.96 (0.68-1.35)	1.10 (0.87-1.40)
HR (95% CI): Model 3	1.00 (ref)	0.79 (0.55-1.14)	0.83 (0.59-1.17)	1.13 (0.89-1.44)
<b>Wine consumption (residuals)</b>				
N	1265	1264	1264	3793
Cases / Person Years	78 / 6291	87 / 6172	54 / 6040	219 / 18503
HR (95% CI): Model 1	1.00 (ref)	1.20 (0.88-1.64)	0.87 (0.61-1.24)	0.90 (0.79-1.03)
HR (95% CI): Model 2	1.00 (ref)	1.16 (0.84-1.59)	0.85 (0.59-1.22)	0.88 (0.76-1.01)
HR (95% CI): Model 3	1.00 (ref)	1.03 (0.66-1.37)	0.95 (0.66-1.37)	0.90 (0.78-1.04)
<b>Spirit consumption (residuals)</b>				
N	1265	1266	1262	3793
Cases / Person Years	54 / 6068	87 / 6159	78 / 6276	219 / 18503
HR (95% CI): Model 1	1.00 (ref)	1.43 (1.01-2.02)	1.18 (0.83-1.69)	1.09 (0.95-1.26)
HR (95% CI): Model 2	1.00 (ref)	1.41 (0.99-2.00)	1.19 (0.83-1.70)	1.12 (0.97-1.29)
HR (95% CI): Model 3	1.00 (ref)	1.14 (0.79-1.63)	1.04 (0.72-1.49)	1.08 (0.93-1.26)

**Supplemental Table 2 (continued).** Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) of the associations between residuals of absolute consumption of beer, wine, and spirits and diabetes incidence in 10 European cohorts.

<b>EPIC-ELDERLY: Greece</b>	<b>Q1</b>	<b>Q2</b>	<b>Q3</b>	<b>Per 6 g/d</b>
<b>Beer consumption (residuals)</b>				
N	1784	1775	1757	5316
Cases / Person Years	251 / 19614	296 / 19127	268 / 19109	815 / 57849
HR (95% CI): Model 1	1.00 (ref)	1.48 (1.24-1.76)	1.29 (1.09-1.54)	1.04 (0.96-1.12)
HR (95% CI): Model 2	1.00 (ref)	1.46 (1.23-1.74)	1.26 (1.06-1.50)	1.02 (0.94-1.10)
HR (95% CI): Model 3	1.00 (ref)	1.39 (1.17-1.66)	1.20 (1.00-1.43)	1.03 (0.95-1.12)
<b>Wine consumption (residuals)</b>				
N	1769	1765	1782	5316
Cases / Person Years	273 / 18828	280 / 19263	262 / 19759	815 / 57849
HR (95% CI): Model 1	1.00 (ref)	0.84 (0.71-1.00)	0.66 (0.55-0.78)	0.93 (0.88-0.98)
HR (95% CI): Model 2	1.00 (ref)	0.88 (0.73-1.04)	0.69 (0.58-0.82)	0.94 (0.90-0.99)
HR (95% CI): Model 3	1.00 (ref)	0.89 (0.63-0.89)	0.75 (0.63-0.89)	0.95 (0.90-1.01)
<b>Spirit consumption (residuals)</b>				
N	1770	1820	1726	5316
Cases / Person Years	260 / 19497	284 / 20037	271 / 18315	815 / 57849
HR (95% CI): Model 1	1.00 (ref)	1.23 (1.03-.46)	1.49 (1.26-1.78)	1.08 (1.02-1.14)
HR (95% CI): Model 2	1.00 (ref)	1.21 (1.02-1.45)	1.45 (1.22-1.72)	1.06 (1.01-1.12)
HR (95% CI): Model 3	1.00 (ref)	1.12 (0.94-1.34)	1.35 (1.13-1.61)	1.05 (0.99-1.11)

**Supplemental Table 2 (continued).** Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) of the associations between residuals of absolute consumption of beer, wine, and spirits and diabetes incidence in 10 European cohorts.

<b>EPIC-ELDERLY: Sweden</b>	<b>Q1</b>	<b>Q2</b>	<b>Q3</b>	<b>Per 6 g/d</b>
<b>Beer consumption (residuals)</b>				
N	932	949	936	2819
Cases / Person Years	27 / 12276	48 / 12507	34 / 12400	109 / 37183
HR (95% CI): Model 1	1.00 (ref)	1.99 (1.22-3.25)	1.14 (0.68-1.91)	0.99 (0.48-2.03)
HR (95% CI): Model 2	1.00 (ref)	2.02 (1.24-3.31)	1.16 (0.69-1.95)	1.00 (0.48-2.06)
HR (95% CI): Model 3	1.00 (ref)	1.99 (1.22-3.77)	1.31 (0.78-2.22)	1.20 (0.56-2.56)
<b>Wine consumption (residuals)</b>				
N	941	921	957	2819
Cases / Person Years	39 / 12435	43 / 12173	27 / 12576	109 / 37183
HR (95% CI): Model 1	1.00 (ref)	1.31 (0.83-2.08)	0.85 (0.50-1.45)	0.54 (0.23-1.29)
HR (95% CI): Model 2	1.00 (ref)	1.32 (0.83-2.10)	0.85 (0.50-1.45)	0.54 (0.23-1.29)
HR (95% CI): Model 3	1.00 (ref)	1.21 (0.46-1.35)	0.79 (0.46-1.35)	0.44 (0.17-1.10)
<b>Spirit consumption (residuals)</b>				
N	939	956	924	2819
Cases / Person Years	21 / 12366	43 / 12604	45 / 12213	109 / 37183
HR (95% CI): Model 1	1.00 (ref)	2.14 (1.26-3.62)	2.04 (1.21-3.44)	1.76 (0.86-3.61)
HR (95% CI): Model 2	1.00 (ref)	2.13 (1.26-3.63)	2.02 (1.19-3.43)	1.76 (0.85-3.64)
HR (95% CI): Model 3	1.00 (ref)	2.05 (1.20-3.49)	1.86 (1.10-3.15)	1.65 (0.74-3.68)

**Supplemental Table 2 (continued).** Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) of the associations between residuals of absolute consumption of beer, wine, and spirits and diabetes incidence in 10 European cohorts.

<b>Tromsø</b>	<b>Q1</b>	<b>Q2</b>	<b>Q3</b>	<b>Per 6 g/d</b>
<b>Beer consumption (residuals)</b>				
N	1562	1752	1644	4958
Cases / Person Years	79 / 20158	97 / 22559	102 / 21103	278 / 63819
HR (95% CI): Model 1	1.00 (ref)	1.11 (0.82-1.50)	1.07 (0.79-1.45)	1.11 (0.91-1.36)
HR (95% CI): Model 2	1.00 (ref)	1.11 (0.82-1.50)	1.07 (0.79-1.45)	1.11 (0.91-1.36)
HR (95% CI): Model 3	1.00 (ref)	1.03 (0.76-1.39)	1.16 (0.85-1.57)	1.16 (0.95-1.41)
<b>Wine consumption (residuals)</b>				
N	1662	1733	1563	4958
Cases / Person Years	114 / 20596	100 / 22534	64 / 20689	278 / 63819
HR (95% CI): Model 1	1.00 (ref)	0.92 (0.70-1.21)	0.70 (0.50-0.97)	0.88 (0.71-1.09)
HR (95% CI): Model 2	1.00 (ref)	0.92 (0.70-1.22)	0.70 (0.50-0.98)	0.88 (0.71-1.09)
HR (95% CI): Model 3	1.00 (ref)	0.95 (0.56-1.10)	0.79 (0.56-1.10)	0.91 (0.73-1.15)
<b>Spirit consumption (residuals)</b>				
N	1708	1617	1633	4958
Cases / Person Years	80 / 22496	80 / 21103	118 / 20220	278 / 63819
HR (95% CI): Model 1	1.00 (ref)	1.02 (0.75-1.40)	1.41 (1.05-1.91)	1.07 (0.74-1.54)
HR (95% CI): Model 2	1.00 (ref)	1.02 (0.75-1.40)	1.41 (1.04-1.92)	1.06 (0.73-1.53)
HR (95% CI): Model 3	1.00 (ref)	0.85 (0.62-1.16)	1.08 (0.79-1.46)	0.84 (0.57-1.25)

**Supplemental Table 2 (continued).** Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) of the associations between residuals of absolute consumption of beer, wine, and spirits and diabetes incidence in 10 European cohorts.

<b>ESTHER</b>	<b>Q1</b>	<b>Q2</b>	<b>Q3</b>	<b>Per 6 g/d</b>
<b>Beer consumption (residuals)</b>				
N	1589	1932	1765	5286
Cases / Person Years	143 / 11176	186 / 13686	197 / 11731	526 / 36593
HR (95% CI): Model 1	1.00 (ref)	1.08 (0.86-1.34)	1.12 (0.89-1.40)	1.06 (0.95-1.18)
HR (95% CI): Model 2	1.00 (ref)	1.10 (0.88-1.37)	1.11 (0.89-1.39)	1.05 (0.95-1.17)
HR (95% CI): Model 3	1.00 (ref)	1.07 (0.86-1.34)	1.05 (0.84-1.32)	1.05 (0.94-1.16)
<b>Wine consumption (residuals)</b>				
N	1667	1792	1827	5286
Cases / Person Years	191 / 11018	175 / 12681	160 / 12894	526 / 36593
HR (95% CI): Model 1	1.00 (ref)	0.88 (0.71-1.10)	0.83 (0.67-1.04)	0.94 (0.85-1.05)
HR (95% CI): Model 2	1.00 (ref)	0.90 (0.72-1.13)	0.84 (0.67-1.06)	0.95 (0.85-1.05)
HR (95% CI): Model 3	1.00 (ref)	0.92 (0.72-1.13)	0.90 (0.72-1.13)	0.96 (0.87-1.07)
<b>Spirit consumption (residuals)</b>				
N	1865	1834	1687	5286
Cases / Person Years	166 / 12075	184 / 12834	176 / 11683	526 / 36593
HR (95% CI): Model 1	1.00 (ref)	1.18 (0.94-1.47)	1.11 (0.89-1.37)	1.06 (0.74-1.53)
HR (95% CI): Model 2	1.00 (ref)	1.20 (0.96-1.49)	1.12 (0.91-1.39)	1.05 (0.73-1.51)
HR (95% CI): Model 3	1.00 (ref)	1.12 (0.90-1.39)	1.03 (0.83-1.27)	0.93 (0.64-1.35)



**Supplemental Table 2 (continued).** Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) of the associations between residuals of absolute consumption of beer, wine, and spirits and diabetes incidence in 10 European cohorts.

<b><u>MORGAM: FINRISK</u></b>	<b>Q1</b>	<b>Q2</b>	<b>Q3</b>	<b>Per 6 g/d</b>
<b>Beer consumption (residuals)</b>				
N	6455	6069	7235	19759
Cases / Person Years	645 / 103702	493 / 99754	572 / 114169	1710 / 317624
HR (95% CI): Model 1	1.00 (ref)	0.82 (0.73-0.92)	0.96 (0.86-1.08)	0.97 (0.93-1.02)
HR (95% CI): Model 2	1.00 (ref)	0.85 (0.75-0.96)	0.96 (0.86-1.07)	0.97 (0.93-1.02)
HR (95% CI): Model 3	1.00 (ref)	0.95 (0.84-1.07)	1.09 (0.97-1.22)	1.02 (0.98-1.07)
<b>Wine consumption (residuals)</b>				
N	6643	6477	6639	19759
Cases / Person Years	705 / 108162	548 / 108167	457 / 101296	1710 / 317624
HR (95% CI): Model 1	1.00 (ref)	0.81 (0.72-0.91)	0.79 (0.69-0.89)	0.95 (0.88-1.02)
HR (95% CI): Model 2	1.00 (ref)	0.84 (0.75-0.95)	0.82 (0.72-0.93)	0.96 (0.89-1.02)
HR (95% CI): Model 3	1.00 (ref)	0.95 (0.78-1.01)	0.89 (0.78-1.01)	0.98 (0.92-1.04)
<b>Spirit consumption (residuals)</b>				
N	7069	6188	6502	19759
Cases / Person Years	511 / 105964	445 / 1005545	754 / 111116	1710 / 317624
HR (95% CI): Model 1	1.00 (ref)	0.81 (0.71-0.92)	1.03 (0.92-1.16)	1.06 (1.01-1.11)
HR (95% CI): Model 2	1.00 (ref)	0.85 (0.75-0.97)	1.05 (0.94-1.18)	1.05 (1.01-1.10)
HR (95% CI): Model 3	1.00 (ref)	0.83 (0.73-0.95)	0.92 (0.82-1.03)	0.99 (0.94-1.03)

**Supplemental Table 2 (continued).** Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) of the associations between residuals of absolute consumption of beer, wine, and spirits and diabetes incidence in 10 European cohorts.

<b><u>MORGAM: MOLI-SANI</u></b>	<b>Q1</b>	<b>Q2</b>	<b>Q3</b>	<b>Per 6 g/d</b>
<b>Beer consumption (residuals)</b>				
N	4811	5065	4634	14510
Cases / Person Years	110 / 20994	117 / 22226	68 / 20533	295 / 63753
HR (95% CI): Model 1	1.00 (ref)	1.32 (1.00-1.74)	1.04 (0.76-1.42)	1.04 (0.92-1.18)
HR (95% CI): Model 2	1.00 (ref)	1.33 (1.00-1.75)	1.04 (0.76-1.42)	1.04 (0.92-1.17)
HR (95% CI): Model 3	1.00 (ref)	1.26 (0.95-1.66)	1.03 (0.75-1.41)	1.01 (0.90-1.14)
<b>Wine consumption (residuals)</b>				
N	4797	5271	4442	14510
Cases / Person Years	75 / 21702	115 / 23140	105 / 18911	295 / 63753
HR (95% CI): Model 1	1.00 (ref)	1.15 (0.85-1.56)	0.97 (0.71-1.33)	0.98 (0.87-1.10)
HR (95% CI): Model 2	1.00 (ref)	1.16 (0.85-1.57)	0.97 (0.71-1.33)	0.98 (0.87-1.10)
HR (95% CI): Model 3	1.00 (ref)	1.14 (0.73-1.36)	1.00 (0.73-1.36)	1.00 (0.89-1.12)
<b>Spirit consumption (residuals)</b>				
N	4750	5075	4685	14510
Cases / Person Years	104 / 19857	106 / 22123	85 / 21772	295 / 63753
HR (95% CI): Model 1	1.00 (ref)	1.23 (0.92-1.64)	1.08 (0.80-1.45)	0.93 (0.68-1.27)
HR (95% CI): Model 2	1.00 (ref)	1.23 (0.92-1.65)	1.09 (0.81-1.46)	0.93 (0.68-1.27)
HR (95% CI): Model 3	1.00 (ref)	1.21 (0.91-1.62)	1.02 (0.76-1.37)	0.93 (0.69-1.26)

**Supplemental Table 2 (continued).** Hazard Ratios (HRs) and 95% Confidence Intervals (CIs) of the associations between residuals of absolute consumption of beer, wine, and spirits and diabetes incidence in 10 European cohorts.

<b><u>MORGAM: North-Sweden</u></b>	<b>Q1</b>	<b>Q2</b>	<b>Q3</b>	<b>Per 6 g/d</b>
<b>Beer consumption (residuals)</b>				
N	1101	911	1020	3032
Cases / Person Years	55 / 21790	44 / 18329	50 / 19628	149 / 59748
HR (95% CI): Model 1	1.00 (ref)	1.11 (0.74-1.65)	1.04 (0.70-1.55)	1.44 (0.84-2.46)
HR (95% CI): Model 2	1.00 (ref)	1.15 (0.77-1.73)	1.11 (0.74-1.65)	1.46 (0.87-2.47)
HR (95% CI): Model 3	1.00 (ref)	1.15 (0.77-1.72)	1.11 (0.75-1.66)	1.49 (0.86-2.60)
<b>Wine consumption (residuals)</b>				
N	784	1224	1024	3032
Cases / Person Years	40 / 14853	63 / 24261	46 / 20633	149 / 59748
HR (95% CI): Model 1	1.00 (ref)	0.96 (0.64-1.44)	0.93 (0.58-1.48)	0.87 (0.44-1.69)
HR (95% CI): Model 2	1.00 (ref)	0.98 (0.65-1.48)	0.92 (0.57-1.48)	0.87 (0.45-1.67)
HR (95% CI): Model 3	1.00 (ref)	0.99 (0.66-1.68)	1.05 (0.66-1.68)	0.98 (0.49-1.96)
<b>Spirit consumption (residuals)</b>				
N	1042	1012	978	3032
Cases / Person Years	46 / 20310	47 / 20377	56 / 19060	149 / 59748
HR (95% CI): Model 1	1.00 (ref)	0.98 (0.65-1.48)	1.09 (0.72-1.65)	0.93 (0.68-1.27)
HR (95% CI): Model 2	1.00 (ref)	0.99 (0.66-1.50)	1.05 (0.70-1.60)	0.67 (0.29-1.54)
HR (95% CI): Model 3	1.00 (ref)	0.89 (0.59-1.34)	0.88 (0.58-1.33)	0.51 (0.21-1.26)

Model 1: Adjusted for age, sex, education, employment, and prevalent coronary heart disease or cancer;

Model 2: Model 1 additionally adjusted for smoking status, sports activity (if available), and Healthy Diet Indicator score (if available).

Model 3: Model 2 additionally adjusted for BMI.

**Table 1.** Cohort characteristics of the ten included European cohorts from the Consortium on Health and Ageing: Network of Cohorts in Europe and the United States (CHANCES) project and number of included persons who reported to consume alcohol.

<b>Cohort</b>	<b>Sub-cohort or country</b>	<b>Baseline period</b>	<b>Follow-up period</b>	<b>Follow-up length, median [P25-P75]</b>	<b>Ascertainment of incident diabetes type 2</b>	<b>N</b>	<b>Age category</b>	<b>Males, %</b>
<b>Zutphen Elderly Study</b>	The Netherlands	1985	1985-2010	9.7 [5.0-15.0]	Self-report and current treatment and non-fasting glucose measures using the WHO definition <sup>46</sup> .	557	≥60 y	100
<b>Rotterdam Study</b>	The Netherlands	1990	1997-2013	12.2 [7.0-13.1]	Followed-up using information from general practitioners, pharmacies' databases, and follow-up examinations. Defined as being registered by a general practitioner as having type 2 diabetes and meeting at least one of the following four criteria: fasting plasma glucose concentration ≥7.0 mmol/L, random plasma glucose concentration ≥11.1 mmol/L, use of anti-diabetic medication, and/or following dietary guidelines for type 2 diabetes.	2428	≥55 y	44
<b>EPIC-Elderly</b>	The Netherlands	1993-1997	1993-2005	4.9 [4.1-5.0]	Self-reported diagnosis in the follow-up questionnaires and/or a urinary glucose strip test for detection of glucosuria, and/or linkage with the Dutch register of hospital discharge diagnoses <sup>47</sup> .	3793	≥60 y	6

**Table 1 (continued).** Cohort characteristics of the ten included European cohorts from the Consortium on Health and Ageing: Network of Cohorts in Europe and the United States (CHANCES) project and number of included persons who reported to consume alcohol.

Cohort	Sub-cohort or country	Baseline period	Follow-up period	Follow-up length, median [P25-P75]	Ascertainment of incident diabetes type 2	N	Age category	Males, %
	Greece	1994-1999	1994-2011	11.4 [9.9-12.5]	Collected during follow-up through self-report and current treatment; cases were not validated.	5316	≥60 y	50
	Sweden	1992-1996	1992-2011	13.2 [12.1-14.2]	Followed up through linkage with the Swedish diabetes register and verified by biomarker measurements of impaired glucose tolerance and impaired fasting glucose in a few cases.	2819	≥60 y	51
<b>The Tromsø Study</b>	Norway	1994-1995	1994-2010	15.6 [10.0-16.0]	Linkage with diabetes-related or cardiovascular diseases discharge diagnosis at the only hospital serving the Tromsø population or verified by self-report or observed HbA <sub>1c</sub> -values >6.5% during follow-up. Some of the cases were validated using medical records or a non-fasting glucose measurement.	4958	≥45 y	59

**Table 1 (continued).** Cohort characteristics of the ten included European cohorts from the Consortium on Health and Ageing: Network of Cohorts in Europe and the United States (CHANCES) project and number of included persons who reported to consume alcohol.

Cohort	Sub-cohort or country	Baseline period	Follow-up period	Follow-up length, median [P25-P75]	Ascertainment of incident diabetes type 2	N	Age category	Males, %
<b>ESTHER</b>	Germany	2000-2003	2000-2007	7.9 [5.3-8.1]	The cohort was systematically searched for diabetes events and incident cases were validated with medical records during follow-up. In addition, subjects with HbA <sub>1c</sub> $\geq$ 6.5% at 8-year follow-up were classified as subjects with incident type 2 diabetes in order to identify undiagnosed cases.	5286	48-75 y	53
<b>MORGAM</b>	FINRISK (Finland)	1982-2002	1982-2010	14.0 [8.9-23.8]	Through linkage to the national Hospital Discharge Register, Causes of Death Register, and drug reimbursement registers <sup>48</sup> .	19759	24-74 y	57
	MOLI-SANI (Italy)	2005-2010	2005-2011	4.3 [3.3-5.4]	Cases were identified and validated through linkage to the National Medication Register and to the Local Diagnosis Registers <sup>48</sup> .	14510	35-99 y	60
	Northern-Sweden (Sweden)	1986-1994	1986-2011	20.8 [17.8-24.5]	Based on self-reported diagnosis in a phone interview and/or linkage with Hospital Discharge Records <sup>48</sup> .	3032	24-74 y	58

**Table 2.** Pooled Hazard Ratios (95% CI) from random-effects meta-analyses for the association between having a beer, wine, or spirit preference compared to having no preference according to sub-groups and additional analyses.

	Beer preference	Wine preference	Spirit preference	No preference
<b>BMI</b>				
- <25 kg/m <sup>2</sup>	1.24 (0.95-1.61)	1.07 (0.85-1.36)	0.89 (0.44-1.82)	1.00 (ref)
- ≥25 kg/m <sup>2</sup>	1.05 (0.95-1.15)	0.95 (0.85-1.06)	1.26 (1.06-1.51)	1.00 (ref)
<b>Sex</b>				
- Men	1.26 (0.93-1.71)	1.17 (0.92-1.50)	1.27 (1.01-1.59)	1.00 (ref)
- Women	1.23 (0.89-1.70)	1.08 (0.85-1.37)	1.33 (1.11-1.60)	1.00 (ref)
<b>Excluding persons with prevalent diseases*</b>	1.04 (0.95-1.15)	0.99 (0.87-1.13)	1.16 (0.95-1.42)	1.00 (ref)
<b>Additional adjustment for frequency pattern</b>	1.05 (0.95-1.16)	0.98 (0.87-1.10)	1.01 (0.90-1.14)	1.00 (ref)
<b>Definition alcoholic beverage preference with cut-off 50%</b>	1.06 (0.93-1.21)	0.93 (0.77-1.12)	1.10 (0.93-1.31)	1.00 (ref)

\* Defined as persons with prevalent heart disease or cancer or a follow-up ≤2 years.

Models adjusted for age, sex, education, employment, prevalent coronary heart disease or cancer, smoking status, sports activity (if available), Healthy Diet Indicator score (if available), and BMI.