

1 **Spinal CX3CL1/CX3CR1 May Not Directly Participate in the Development of**
2 **Morphine Tolerance in Rats**

3 Yawen Peng¹, Genhua Guo², Bin Shu¹, Daiqiang Liu¹, Peng Su¹, Xuming Zhang³,
4 Feng Gao^{1*}

5 1 Department of Anesthesiology, Tongji Hospital, Tongji Medical College, Huazhong
6 University of Science & Technology, Wuhan, Hubei, PR China

7 2 Department of Anesthesiology, The Central People's Hospital of Ji'an City, 106
8 Jinggangshan Road, Ji'an 343000, PR China

9 3 School of Life & Health Sciences, Aston University, Aston Triangle, Birmingham
10 B4 7ET, United Kingdom

11 * Corresponding author: Feng Gao

12 E-mail: 13971587381@163.com

13 **Running title:** CX3CL1/CX3CR1 and Morphine Tolerance

14 **Keywords**

15 Morphine tolerance; Chemokine; CX3CL1; CX3CR1

16 **Abbreviations**

17 CCL2, C-C motif ligand 2; CCL5, C-C motif ligand 5; CCR1, C-C motif chemokine
18 receptor 1; CXCL12, C-X-C motif ligand 12; CX3CL1, C-X3-C motif chemokine 1;

19 CX3CR1, C-X3-C motif chemokine receptor 1; CXCR4, C-X-C motif chemokine

20 receptor 4; rrCX3CL1, recombinant rat CX3CL1 protein; DAMGO, [D-Ala², N-

21 MePhe⁴, Gly-ol]-enkephalin; DOR, delta opioid receptor; ERK, extracellular

22 regulated protein kinases; GAPDH, glyceraldehyde-3-phosphate dehydrogenase;

23 GFAP, glial fibrillary acidic protein; GLAST, glutamate-aspartate transporters; GLT-1,

1 glutamate transporter-1; Iba-1, ionized calcium-binding adapter molecule 1; IL-1 β ,
2 interleukin-1 β ; MPE, maximal possible antinociceptive effect; MOR, mu opioid
3 receptor; NeuN, neuronal nuclei; PAG, periaqueductal gray; p38MAPK, p38 mitogen-
4 activated protein kinase; PBS, phosphate buffer saline; TNF- α , tumor necrosis factor
5 α .

6 **Acknowledgement**

7 No.

8 **Funding sources:** Funding was provided by the National Natural Science Foundation
9 of China (No. 81471143; 81171259).

10 **Abstract**

11 CX3CL1 (fractalkine), the sole member of chemokine CX3C family, is implicated in
12 inflammatory and neuropathic pain *via* activating its receptor CX3CR1 on neural cells
13 in spinal cord. However, it has not been fully elucidated whether CX3CL1 or CX3CR1
14 contributes to the development of morphine tolerance. In this study, we found that
15 chronic morphine exposure did not alter the expressions of CX3CL1 and CX3CR1 in
16 spinal cord. And neither exogenous CX3CL1 nor CX3CR1 inhibitor could affect the
17 development of morphine tolerance. The cellular localizations of spinal CX3CL1 and
18 CX3CR1 changed from neuron and microglia, respectively, to all the neural cells during
19 the development of morphine tolerance. A microarray profiling revealed that 15
20 members of chemokine family excluding CX3CL1 and CX3CR1 were up-regulated in
21 morphine-treated rats. Our study provides evidence that spinal CX3CL1 and CX3CR1
22 may not be involved in the development of morphine tolerance directly.

1 **Introduction**

2 Morphine is the most important and frequently used opioid for acute and chronic pain
3 in clinical practice. However, repeated usage of morphine can induce drug tolerance, a
4 consequence requiring higher doses of morphine to maintain the same analgesic effect.
5 Dose escalation of morphine could potentially cause serious side effects including
6 respiratory depression [1], hypotension, nausea, constipation, dizziness and addiction
7 [2]. Morphine tolerance could hinder the clinical utilization of morphine and impair the
8 quality of life in patients. Thereby, understanding the mechanism of morphine tolerance
9 is critical for improving pain management.

10 Chemokines play the pivotal roles in neuroinflammation, nerve injury-induced pain [3-
11 5] and morphine analgesia [6-8]. CX3CL1 (fractalkine) is the only member of
12 chemokine CX3C family [9] and activates its sole receptor CX3CR1. Previous studies
13 have demonstrated that intrathecal injection of CX3CR1 neutralizing antibody can
14 effectively delay the development of mechanical allodynia and thermal hyperalgesia in
15 neuropathic pain, inflammatory pain and cancer pain [3,10,11]. In addition, CX3CL1
16 has been reported to be involved in diminishing the analgesic effect of opioids in
17 periaqueductal grey [12]. Thus, CX3CL1 plays an important role in the mechanisms of
18 chronic pain.

19 In central nervous system, microglia are generally regarded as the source and target of
20 chemokines [13]. Microglia can regulate and receive chemokine signal between
21 astrocyte and neuron through autocrine and paracrine communications, thus
22 contributing to the development of neuropathic pain [14] and traumatic brain injury
23 [15]. Although it is clear that microglia activation is involved in morphine tolerance

1 [16,17], the potential signals that causing glial activation have not been well understood.
2 Morphine tolerance and chronic pain may share the similar cellular mechanisms.
3 Previous study reported that CX3CL1 is mainly released by neurons and its receptor
4 CX3CR1 is primarily expressed on microglia [18]. As CX3CL1 plays an important role
5 in chronic pain, these results indicate a potential involvement of CX3CL1/CX3CR1
6 signaling axis in the activation of microglia, and thus the mechanism of morphine
7 tolerance. Thereby, the present study was designed to investigate the possible roles of
8 spinal CX3CL1/CX3CR1 in the development of morphine tolerance in rats.

9 **Materials and methods**

10 **Animals**

11 Specific pathogen free adult male Sprague-Dawley rats, weighing 220-240 g, were
12 purchased from Laboratory Animal Center, Tongji Medical College, Huazhong
13 University of Science & Technology. Animals were housed under controlled conditions
14 (22 ± 0.5 °C, relative humidity 40-60 %, alternate light-dark cycles, food and water *ad*
15 *libitum*). To keep the integrity of catheter, rats were housed individually after surgery.
16 All experimental procedures and protocols were reviewed and approved by
17 Experimental Animal Care and Use Committee of Tongji Medical College, Huazhong
18 University of Science & Technology, and carried out in accordance with the National
19 Institutes of Health Guidelines for the Care and Use of Laboratory Animals.

20 **Intrathecal catheterization**

21 Lumbosacral indwelling catheters were constructed and planted using a lumbar
22 approach, as described previously [19]. Briefly, rats were deeply anesthetized with 1 %

1 pentobarbital sodium [60 mg/kg, intraperitoneal injection (i.p.)]. The lumbar region of
2 rat was shaved, and intrathecal catheterization was performed by implanting a sterile
3 PE-0503 catheter (outer diameter 0.5 mm, inner diameter 0.3 mm. Anilab Software &
4 Instruments, Ningbo, China) into subarachnoid cavity between L4 and L5 vertebrae.
5 The catheter was subcutaneously tunneled, externalized and fixed to the back of neck.
6 Wounds were sutured after disinfection with 75 % (v/v) ethanol. Proper location of
7 catheter was confirmed by a temporary motor block of both hind limbs after intrathecal
8 injection of 10 μ L of 2 % lidocaine. The rats were allowed a 7-day recovery period
9 before the following experiments. Rats with hind limb paralysis or paresis after surgery
10 were excluded and euthanized with overdose of pentobarbital sodium.

11 **Drugs administration**

12 The drugs used in this study were prepared as follows. Morphine hydrochloride was
13 diluted by saline (North-East Pharmaceutical Group, China). Recombinant rat CX3CL1
14 protein (rrCX3CL1, R&D, Minneapolis, MN, USA) was dissolved in saline and
15 injected intrathecally (100 ng or 500 ng). Rabbit anti-rat CX3CR1 (Torrey Pine Biolabs,
16 East Orange, NJ, USA) and placebo rabbit IgG (Sigma, St. Louis, MO, USA) were
17 diluted by saline and administered intrathecally (5 μ g or 10 μ g). All drugs or vehicle
18 solutions were injected 30 min before morphine administration in a volume of 5 μ L
19 followed by 10 μ L of saline to flush the catheter. All the doses of each drug used in this
20 study were determined according to the previous experiments [3,12,20].

21 **Morphine tolerance**

22 To induce chronic tolerance to morphine, rats were intrathecally injected with morphine

1 (10 µg) twice daily for 7 days. Rats in the control group received an equivalent volume
2 of saline at the same time points. The development of morphine tolerance was assessed
3 by behavioral tests on day 1, 3, 5 and 7 [21].

4 **Behavioral assessment**

5 Thermal pain thresholds in rats were measured by a tail-flick latency test [22] before
6 drug administration and at 30 min after morphine administration on day 1, 3, 5, and 7
7 [21]. Briefly, rats were placed in plastic containers to hold the body without restraining
8 the head and tail, and one-third to the tip of tail was immersed into water. The
9 temperature of water was adjusted to 50 ± 0.2 °C because this was the proper
10 temperature to record an average tail-flick latency of 2 - 4 s in naïve rats. A cutoff time
11 of 15 seconds was determined to prevent tail damage. The test was repeated three times
12 and the mean of three trials was considered as the final latency. The percentage of
13 maximal possible antinociceptive effect (%MPE) was calculated by comparing the test
14 latency before (baseline, BL) and after drug injection (TL) using following
15 equation: %MPE= [(TL-BL)/ (cutoff time-BL)] ×100. All behavioral tests were carried
16 out under blind conditions.

17 **Quantitative real-time polymerase chain reaction (qRT-PCR)**

18 Under deep anesthesia with 1 % pentobarbital sodium, L1-L5 spinal cord segments of
19 rats were quickly removed on day 7. Total RNA was extracted from tissue sample and
20 the reverse transcription procedure was performed by using RNAiso Plus (Takara,
21 Shiga, Japan) according to the manufacturer's instructions. One microgram of total
22 RNA from each sample was added into 20 µL reactive solution of reverse transcription,

1 respectively. Specific primers for rat CX3CL1, CX3CR1, mu opioid receptor (MOR)
2 and endogenous control rat GAPDH were obtained from GeneCopoeia Company
3 (USA). The catalogs of each primer were as following: CX3CL1 (RQP052632),
4 CX3CR1 (RQP052471), MOR (RQP048316), GAPDH (RQP049537). StepOne Real-
5 Time PCR System (Applied Biosystems, USA) was used to conduct the qRT-PCR.
6 Relative quantification of mRNA was performed by $2^{-\Delta\Delta C_t}$ method.

7 **Western blots**

8 L1-L5 spinal cord segments of all animals were quickly removed and dissected. Total
9 protein of spinal cord tissue from each group was extracted by using radio
10 immunoprecipitation assay lysis buffer according to the manufacturer's instructions
11 (Beyotime, Wuhan, China). The protein concentration of supernatants was measured
12 by using bicinchoninic acid assay. 50 μ g protein from each sample was loaded on 10 %
13 SDS-PAGE gel after boiling in the sample buffer. Electrophoresis was conducted at 60
14 V constant voltage for stacking gel and 100 V for separating gel. The proteins were
15 subsequently electro-transferred (200 mA, 60-90 min) to a PVDF membrane (Millipore,
16 Bellerica, MA, USA). The membrane was blocked with 5 % BSA (Bovine Serum
17 Albumin) for 1 h at room temperature followed by incubating with the following
18 primary antibodies: mouse anti-ionized calcium-binding adapter molecule 1 (Iba1)
19 antibody (1: 200, Santa Cruz, Dallas, TX, USA), goat anti-CX3CL1 antibody (1: 50,
20 R&D, Minneapolis, MN, USA), rabbit anti-CX3CR1 antibody (1: 1000, Abcam,
21 Cambridge, MA, USA), mouse anti-MOR antibody (1: 500, R&D, Minneapolis, MN,
22 USA), or rabbit anti-GAPDH (1: 2500, Aspen, Wuhan, China) overnight at 4 °C. After
23 being thoroughly washed, the membrane was incubated with HRP-conjugated rabbit

1 anti-goat IgG (1: 5000, EarthOx, Millbrae, CA, USA), HRP-conjugated goat anti-rabbit
2 IgG (1: 4000, Aspen, Wuhan, China), or HRP-conjugated goat anti-mouse IgG (1: 1000,
3 Bioyartech, Wuhan, China) for 2 h at room temperature. Finally, proteins were
4 detected by ECL reagents (Beyotime, Wuhan, China) and visualized by exposing to X-
5 ray film. The ImageJ analysis system (NIH, Bethesda, MD) was used for the
6 quantification of specific bands. The levels of Iba-1, CX3CL1, CX3CR1 and MOR
7 were exhibited as density relative to the density of GAPDH.

8 **Double immunofluorescent staining**

9 After being treated with morphine or saline for 7 days, rats were perfused with saline,
10 followed by 4 % ice-cold paraformaldehyde (PFA) in 0.1 M phosphate buffer saline
11 (PBS) under deep anesthesia with pentobarbital sodium. The L1-L5 spinal cord
12 segments were removed and post-fixed for 24 h at 4 °C, then dehydrated in 30 %
13 sucrose solution. After being treated with 0.3 % Triton X-100 and blocked with 10 %
14 donkey serum for 40 min at room temperature, 25 µm-thick sections were incubated
15 overnight at 4 °C with mixtures of the following primary antibodies: goat anti-CX3CL1
16 antibody (1: 25, R&D, Minneapolis, MN, USA) or rabbit anti-CX3CR1 antibody (1:
17 200, Abcam, Cambridge, MA, USA) and mouse anti-NeuN (1: 200, Millipore,
18 Bellerica, MA, USA), mouse anti-GFAP (1: 200, CST, Beverly, MA, USA), goat anti-
19 Iba1 (1: 100, Abcam, Cambridge, MA, USA) or rabbit anti-Iba1 (1: 200, Wako, Osaka,
20 Japan). Then sections were incubated with mixtures of the following secondary
21 antibodies: Cy3-conjugated donkey anti-goat IgG (1: 300, Proteintech, Wuhan, China)
22 and FITC-conjugated donkey anti-mouse IgG (1: 100, Proteintech, Wuhan, China) or
23 FITC-conjugated donkey anti-rabbit IgG (1: 100, Proteintech, Wuhan, China), or

1 IFKine Red labeled donkey anti-rabbit IgG (1: 500, Abbkine, Redlands, CA, USA) and
2 FITC-conjugated donkey anti-mouse IgG (1: 100, Proteintech, Wuhan, China) or FITC-
3 conjugated donkey anti-goat IgG (1: 100, Proteintech, Wuhan, China) for 2 h at room
4 temperature and stained with 4, 6-diamidino-2-phenylindole (DAPI, Boster, Wuhan,
5 China) for 10 min. The stained sections were examined by using Fluorescence
6 Microscope (DM2500, Leica, German) to capture the fluorescent images. Five spinal
7 sections were selected randomly for each rat and the immunoreactivities of CX3CL1
8 and CX3CR1 were counted in a blinded fashion [23]. The stained sections were
9 analyzed by Image Pro Plus 4 software (Media Cybernetics, Maryland, MD, USA).

10 **Microarray mRNA profiling**

11 Gene expression profile of spinal cord tissues was established by using Affymetrix Rat
12 Genome 230 2.0 Arrays. L1-L5 spinal cord segments of morphine-treated or saline-
13 treated rats were isolated on day 7 and RNAlater RNA Stabilization Reagent (Qiagen,
14 Germany) was used for stabilization of RNA in tissue samples. Total RNA isolation
15 was performed with TRIzol reagent (Invitrogen, USA) and NucleoSpin® RNA Clean-
16 up (MACHEREY-NAGEL, Germany). The cRNA was generated and labeled by one-
17 cycle target labeling method. Affymetrix Rat Genome 230 2.0 microarray (CapitalBio
18 Corporation, Beijing, China) which contains 31,000 probe sets including 65 probe sets
19 of chemokine family was used to screen the differential expressions of chemokines.
20 The acceptance criteria for RNA quality were 260/280 ratio ≥ 1.80 and RNA integrity
21 number ≥ 8.0 . The cRNA generated from each sample was hybridized to a single array
22 according to standard Affymetrix protocols. Initial image analysis of microarray chips
23 was performed using the Genechip® Command Console® Software. Data were

1 exported to Significance Analysis of Microarrays software for screening differentially
2 expressed genes. The screening criterion was set as fold change ≥ 2 or fold change \leq
3 0.5 with false discovery rate (FDR) q -value ≤ 0.05 .

4 **Statistical analysis**

5 Animal sample size for behavioral experiment was decided by power analysis using
6 SSize2021 software (National University of Singapore, Singapore) (version 2). With
7 anticipated population proportion $P_1 = 0.95$, $P_2 = 0.05$, significance level 0.05 and
8 power of test 0.09, the sample size was estimated to be four per group. All data were
9 presented as mean \pm SEM. Behavioral test was analyzed by two-way repeated measure
10 ANOVA (treatment group \times time) to detect overall differences among treatment groups
11 followed by Bonferroni's test to detect the changes to %MPE after drug injection over
12 time. The results of qRT-PCR and western blots were analyzed by one-way ANOVA.
13 Individual comparisons were conducted with unpaired t -test. Statistical analyses were
14 performed with GraphPad Prism 5 (GraphPad Software Inc.) with statistical
15 significance set at $P < 0.05$.

16 **Results**

17 *Chronic morphine treatment induced drug tolerance and activated microglia*

18 Rats were intrathecally administered with morphine (10 $\mu\text{g}/5 \mu\text{L}$) or saline (5 μL) twice
19 daily for consecutive 7 days. Behavioral tests were conducted before drug
20 administration and at 30 min after the last drug administration on day 1, 3, 5, and 7. As
21 shown in Fig. 1A, rats received morphine exhibited significantly higher %MPE
22 compared with saline-treated rats on day 1 ($P < 0.001$) and day 3 ($P < 0.01$). On day 5,

1 there was no significant difference of %MPE levels between morphine-treated and
2 saline-treated rats ($P > 0.05$), suggesting that chronic morphine tolerance had been
3 successfully established. Iba-1 is the marker of activated microglia. On day 7, the
4 increased expression of Iba-1 was detected in spinal cord (Fig. 1B), indicating the
5 activation of microglia induced by chronic morphine exposure. Previous study has
6 demonstrated that chronic morphine treatment significantly decreases the expression of
7 MOR in hypothalamus but not in locus ceruleus and nucleus accumbens [24],
8 suggesting that the cellular adaptation for morphine is tissue-specific. In this study, we
9 did not detect any changes in MOR expression in spinal cord of morphine-treated rats
10 compared to that in saline-treated rats on day 7 (Fig. 1C and 1D).

11 *Chronic morphine treatment did not affect the expressions of CX3CL1 and CX3CR1*

12 Previous studies have demonstrated that CX3CL1 and its receptor could play the
13 important roles in antinociceptive effects of opioid agonists in periaqueductal grey
14 [12,25]. To determinate whether CX3CL1 and CX3CR1 participate in the development
15 of morphine tolerance, the expressions of CX3CL1 and CX3CR1 in spinal cord of rats
16 were examined on day 7 of morphine administration. As shown in Fig. 2, neither the
17 levels of CX3CL1 mRNA ($F_{2,15} = 0.901$, $P = 0.427$) and CX3CR1 mRNA ($F_{2,15} = 1.314$,
18 $P = 0.298$), nor expressions of CX3CL1 protein ($F_{2,12} = 0.999$, $P = 0.397$) and CX3CR1
19 protein ($F_{2,21} = 0.833$, $P = 0.449$) was affected by intrathecal administration of morphine
20 when compared to saline-treated rats.

21 *Exogenous CX3CL1 or CX3CR1 inhibitor had no effect on behavioral responses during* 22 *the development of morphine tolerance*

23 There was no significant difference in baseline levels of tail-flick latency measured

1 prior to drug administration among all groups, indicating that intrathecal catheterization
2 did not affect behavioral responses of rats (Fig. 3A). To further investigate the roles of
3 CX3CL1 and CX3CR1 in the development of morphine tolerance, recombinant rat
4 CX3CL1 (100 ng or 500 ng), anti-CX3CR1 neutralizing antibody (5 μ g or 10 μ g) or
5 control IgG (100 ng or 5 μ g) was intrathecally injected 30 min before morphine
6 administration, respectively. As shown in Fig. 3B, neither 100 ng nor 500 ng rrCX3CL1
7 exhibited statistically significant effect on %MPE in rats treated with morphine
8 compared with that in IgG-morphine treated rats ($F_{2, 45} = 1.498$, $P = 0.255$ for 100 ng
9 rrCX3CL1; $F_{2, 45} = 0.903$, $P = 0.426$ for 500 ng rrCX3CL1). Moreover, interaction
10 between rrCX3CL1 treatment and time was not considered significantly ($F_{9, 60} = 0.527$,
11 $P = 0.849$). As shown in Fig. 3C, there was no significant effect of 5 μ g or 10 μ g anti-
12 CX3CR1 neutralizing antibody on %MPE in rats treated with morphine when
13 compared with that in IgG-morphine treated rats ($F_{2, 45} = 0.905$, $P = 0.426$ for 5 μ g; $F_{2,$
14 $45 = 1.107$, $P = 0.356$ for 10 μ g). There was no significant interaction between anti-
15 CX3CR1 neutralizing antibody treatment and time ($F_{9, 60} = 1.770$, $P = 0.093$). These
16 results suggest that both exogenous CX3CL1 stimulation and CX3CR1 inhibition could
17 not markedly affect the development of morphine tolerance that assessed by tail flick
18 test.

19 *Exogenous CX3CL1 or CX3CR1 inhibitor had no influence on the activation of*
20 *microglia induced by morphine*

21 In order to further clarify the roles of CX3CL1 and CX3CR1, exogenous CX3CL1 (100
22 ng or 500 ng) or CX3CR1 inhibitor (5 μ g or 10 μ g) was intrathecally administered
23 respectively and their effects on the activation of spinal microglia in morphine tolerant
24 rats were assessed. As shown in Fig. 4, the expressions of Iba-1 were significantly

1 increased in all morphine-treated rats. Neither rrCX3CL1 nor anti-CX3CR1
2 neutralizing antibody had statistically significant effect on morphine-induced
3 expressions of Iba-1 ($F_{5, 12} = 0.138, P = 0.980$).

4 *Cellular localizations of spinal CX3CL1 and CX3CR1 in morphine tolerant rats*

5 The communication between neurons and glia mediated by CX3CL1 and CX3CR1
6 contributes to the mechanisms of inflammatory and neuropathic pain [3,26,27]. The
7 changes of cellular distribution of CX3CL1 and CX3CR1 in spinal cord have also been
8 reported to be associated with pain conditions [18]. In this study, we examined the
9 cellular localizations of CX3CL1 and CX3CR1 in rat spinal dorsal horn. The results
10 showed that CX3CL1 expression was extensively distributed to all layers of spinal
11 dorsal horn, and CX3CR1 was mainly expressed in lamina I to lamina III of spinal cord
12 (Fig. 5A). There was no significant difference in the expression of CX3CL1 or
13 CX3CR1 between morphine-treated rats and saline-treated rats (Fig. 5B). In saline-
14 treated rats, the immunoreactivity of CX3CL1 was co-localized with neuronal marker
15 NeuN, while CX3CR1 was co-localized with microglia marker Iba-1 (Fig. 5C).
16 However, both CX3CL1 and CX3CR1 were found to be co-localized with NeuN, GFAP
17 and Iba-1 in morphine-treated rats (Fig. 5D and 5E). These indicate the shift of
18 CX3CL1/CX3CR1 expressions occurred during the development of morphine
19 tolerance.

20 *The mRNA expression profiling screened spinal chemokines related to morphine* 21 *tolerance*

22 To screen the possible chemokines which might be involved in the development of
23 morphine tolerance in spinal cord, mRNA of L1-L5 lumbar spinal cord of respective

1 animal was analyzed using microarrays which contains 65 probe sets of chemokines.
2 As shown in Fig. 6, expressions of 15 chemokines were identified to be upregulated in
3 morphine-treated rats when compared with saline-treated rats. All the upregulated
4 genes of chemokine were listed in Table. 1. However, the changes of CX3CL1 and
5 CX3CR1 expressions were not detected by microarray analysis.

6 **Discussion**

7 The results of our study showed that chronic morphine treatment can induce
8 antinociceptive tolerance, but did not affect the expressions of CX3CL1 and its receptor
9 CX3CR1 in spinal cord. Neither intrathecal administration of exogenous CX3CL1 nor
10 CX3CR1 inhibitor affected the development of morphine tolerance. However,
11 morphine treatment could influence the cellular localization of CX3CL1 and CX3CR1
12 in spinal dorsal horn in rats.

13

14 Various signaling pathways have been found to be involved in the mechanism of opioid
15 tolerance. Opioid tolerance could be prevented, attenuated or reversed by inhibiting
16 proinflammatory cytokines interleukin-1 β (IL-1 β) and tumor necrosis factor α (TNF- α)
17 [28,29]; blocking the activations of extracellular regulated protein kinases (ERK) and
18 p38 protein [30-32]; increasing glial glutamate transporter-1 (GLT-1) and glutamate-
19 aspartate transporters (GLAST) [33,34]. Recently, toll-like receptor 4 (TLR4) has been
20 reported to participate in the development of opioid tolerance *via* increasing tumor
21 necrosis factor and IL-1 β expressions and downregulating the expressions of GLT-1
22 and GLAST [35, 36]. Opioid tolerance could be considered as a drug-specific side
23 effect. The mechanisms of drug tolerance induced by different opioids may be distinct.

1 The internalization of MOR, which is the typical feature of opioid tolerance, highly
2 depends on the type of agonist [37]. Endogenous opioids as well as synthetic peptide
3 DAMGO ([D-Ala², N-MePhe⁴, Gly-ol]-enkephalin) promote the rapid endocytosis of
4 MOR, but the highly addictive opioid, such as morphine, fails to induce detectable
5 endocytosis [38,39]. Changes in MOR expression in response to chronic opioid
6 treatment have long been speculated to directly contribute to the development of opioid
7 tolerance. Following chronic treatment with various agonists, the expression of opioid
8 receptor in brain tissue is either increased, decreased or unchanged, indicating that the
9 regulation of opioid receptor expression depends on the type of opioid [40], agonist
10 [41,42], and the region of brain [26,43]. In addition, opioid tolerance is not only due to
11 the rapid decrease of receptor activity but also the compensatory mechanism
12 counteracting the function of opioid receptor [44-46]. Therefore, it is reasonable to
13 comprehend the expression of MOR in spinal cord was unchanged in our study during
14 the development of morphine tolerance.

15

16 Previous studies have shown that CX3CL1/CX3CR1 signaling axis participate in
17 numerous physiological and pathological processes, including neuropathic pain [26],
18 maturation of synaptic connection [47,48], neuronal survival [49], insulin secretion [50]
19 and atherosclerosis [51]. However, our results revealed the unchanged expressions of
20 CX3CL1 and CX3CR1 in spinal cord of morphine tolerant rats. Recently study in
21 opioid tolerant patients also showed that the concentration of CX3CL1 in cerebrospinal
22 fluid is not significantly different from that in naïve-control patients [6]. Although
23 previous study reported that intrathecal administration of 30 ng exogenous CX3CL1
24 could induce the behavioral effects such as mechanical allodynia and thermal

1 hyperalgesia [4], the development of morphine tolerance was not affected by much
2 higher doses of exogenous CX3CL1 in our study. Previous study showed that
3 intrathecal administration of 3 µg anti-CX3CR1 neutralizing antibody could effectively
4 inhibit monoarthritis-induced mechanical allodynia and thermal hyperalgesia [3],
5 which illustrates that the doses of anti-CX3CR1 neutralizing antibody used in our study
6 (5 µg and 10 µg) should be sufficient to block the function of CX3CR1 in spinal cord.
7 However, in our study, intrathecal injection of CX3CR1 inhibitor did not affect either
8 the antinociceptive effect of morphine or the development of morphine tolerance. In
9 contrast, Johnston and colleagues reported that intrathecal injection of anti-CX3CR1
10 neutralizing antibody could attenuate the development of morphine tolerance [20]. This
11 discrepancy might be due to the different experimental protocols including the
12 evaluation of pain threshold. In Johnston's study, tail flick latencies were recorded
13 every 20 min for 2 hours after morphine infusions to calculate the average response
14 over this time on day 1 and day 5. Their behavioral assessment protocol is quite
15 different from ours which has been most commonly used in the previous studies
16 [21,22,37]. The analgesic effect of morphine occurs at 5 minutes after intrathecal
17 injection, lasts for about 60 minutes, and dissipates by 100 minutes [29]. Choosing the
18 average value of tail flick latencies over 2 hours as the pain threshold of morphine-
19 treated rats may fail to represent the maximum analgesic potency of morphine. Taken
20 together, CX3CL1 and its receptor CX3CR1 in spinal cord may not participate in the
21 mechanism of morphine tolerance directly. Although sufficient dosages of exogenous
22 CX3CL1 and anti-CX3CR1 neutralizing antibody were used in our study, we could not
23 definitively exclude the possibility that CX3CL1/CX3CR1 signaling play a positive
24 role in the mechanism of morphine tolerance yet. CX3CL1 or CX3CR1 knockout

1 animals might be the ideal option to verify the role of CX3CL1/CX3CR1 in morphine
2 tolerance.

3

4 Under physiological conditions, expression of CX3CL1 in spinal cord appears to be
5 restricted to neurons, whereas CX3CR1 in microglia [18]. However, CX3CL1 could be
6 detected not only in neurons but also in astrocytes after spinal nerve ligation [10],
7 suggesting that the distribution of CX3CL1 and CX3CR1 may depend on the diverse
8 pathological processes. Both CX3CR1 and opioid receptors are members of G protein
9 coupled receptor family. The formation of heterodimer among G protein coupled
10 receptors are common. Previous study has found the MOR-CX3CR1 co-localization on
11 neurons in several brain regions, including nucleus accumbens, ventral tegmental area
12 and periaqueductal gray and MOR-CX3CR1 heterologous desensitization has been
13 proved in periaqueductal gray [25]. We found that the cellular localizations of spinal
14 CX3CL1 and CX3CR1 changed from neuron and microglia, respectively, to all the
15 neural cells after chronic morphine administration. These findings support the
16 possibility that morphine treatment may stimulate the cleavage of CX3CL1 in neurons
17 [26] and promote the combination of CX3CL1 with its receptor on glia. It has been
18 reported that simultaneous activations of delta opioid receptor (DOR) and CXCR4 on
19 human peripheral blood mononuclear cells could promote the formation of non-
20 functional DOR-CXCR4 heterodimers which are unable to respond to the agonists [52].
21 Therefore, we speculate that the newly expressed CX3CR1 on neuron may bind to
22 MOR to form into the heterodimer, which at least partly contribute to morphine
23 analgesia or tolerance. Further studies are still needed to explore the potential
24 interaction between CX3CR1 and opioid receptor in the mechanism of morphine

1 tolerance.

2

3 Chemokines and opioids are effective regulators of immune, inflammatory and
4 neuronal responses in pain mechanism in central nervous system. Several chemokines
5 could increase the neuronal excitability and subsequently decrease opioid analgesic
6 efficacy, which may act as the key neuromodulators of pain pathways [53].

7 Administration of CCL5 or CXCL12 (binding to CCR1 or CXCR4, respectively) into
8 periaqueductal gray could attenuate acute opioid analgesia *via* heterologous
9 desensitization of opioid receptors [54,55]. Spinal glial CXCL12 has also been reported

10 to be associated with pain hypersensitivity process induced by bone cancer [56]. Other

11 evidences indicate that CXCL10 could serve as a negative regulator of morphine

12 analgesia [7]. Intrathecal administration of CCL2 neutralizing antibody could attenuate

13 the development of morphine tolerance [21]. Our microarrays analysis revealed the

14 expressions of 15 chemokines, which mainly belong to CXC and CC subfamilies, were

15 significantly increased due to chronic morphine exposure. These results further

16 excluded the direct involvement of CX3CL1 and CX3CR1 in the mechanism of

17 morphine tolerance and also indicated the potential contribution of chemokines to the

18 development of morphine tolerance.

19

20 In conclusion, our study reveals that chronic morphine exposure did not alter the

21 expressions of CX3CL1 and CX3CR1 in spinal cord and inhibiting CX3CL1 or

22 CX3CR1 could not affect the morphine analgesia and the development of drug

23 tolerance. But morphine could change the cellular localizations of spinal CX3CL1 and

24 CX3CR1 which indicates the complex interaction between neuron and glia during

1 morphine tolerance. We also found that 15 chemokines were upregulated significantly
2 during the development of morphine tolerant. These might provide the potential
3 research targets for the further studies in morphine tolerance in the future.

4 **Conflict of interest**

5 None declare.

6 **Reference**

- 7 1. Handley Carroll A, Ensberg Dorrence L (1945) A comparison of amphetamine
8 sulfate with other stimulants of the central nervous system in morphine respiratory
9 depression. *Anesthesiology* 6 (6):561-564
- 10 2. Benyamin R TA, Datta S, Buenaventura R, Adlaka R, Sehgal N, Glaser SE, Vallejo
11 R (2008) Opioid complications and side effects. *Pain Physician* 11 (2S):S105-120
- 12 3. Sun S, Cao H, Han M, Li TT, Pan HL, Zhao ZQ, Zhang YQ (2007) New evidence
13 for the involvement of spinal fractalkine receptor in pain facilitation and spinal glial
14 activation in rat model of monoarthritis. *Pain* 129 (1-2):64-75.
15 doi:10.1016/j.pain.2006.09.035
- 16 4. Milligan ED, Zapata V, Chacur M, Schoeniger D, Biedenkapp J, O'Connor KA, Verge
17 GM, Chapman G, Green P, Foster AC, Naeve GS, Maier SF, Watkins LR (2004)
18 Evidence that exogenous and endogenous fractalkine can induce spinal nociceptive
19 facilitation in rats. *Eur J Neurosci* 20 (9):2294-2302. doi:10.1111/j.1460-

1 9568.2004.03709.x

2 5. White FA, Bhangoo SK, Miller RJ (2005) Chemokines: integrators of pain and
3 inflammation. *Nat Rev Drug Discov* 4 (10):834-844. doi:10.1038/nrd1852

4 6. Lin CP, Kang KH, Lin TH, Wu MY, Liou HC, Chuang WJ, Sun WZ, Fu WM (2015)
5 Role of spinal CXCL1 (GROalpha) in opioid tolerance: a human-to-rodent translational
6 study. *Anesthesiology* 122 (3):666-676. doi:10.1097/ALN.0000000000000523

7 7. Ye D, Bu H, Guo G, Shu B, Wang W, Guan X, Yang H, Tian X, Xiang H, Gao F
8 (2014) Activation of CXCL10/CXCR3 signaling attenuates morphine analgesia:
9 involvement of Gi protein. *J Mol Neurosci* : MN 53 (4):571-579. doi:10.1007/s12031-
10 013-0223-1

11 8. Parsadaniantz SM, Rivat C, Rostene W, Goazigo AR-L (2015) Opioid and
12 chemokine receptor crosstalk: a promising target for pain therapy? *Nat Rev Neurosci*
13 16 (2):69-78. doi:10.1038/nrn3858

14 9. Bazan JF, Bacon KB, Hardiman G, Wang W, Soo K, Rossi D, Greaves DR, Zlotnik
15 A, Schall TJ (1997) A new class of membrane-bound chemokine with a CX3C motif.
16 *Nature* 385 (6617):640-644

17 10. Lindia JA, McGowan E, Jochnowitz N, Abbadie C (2005) Induction of CX3CL1
18 expression in astrocytes and CX3CR1 in microglia in the spinal cord of a rat model of

- 1 neuropathic pain. *J Pain* 6 (7):434-438. doi:10.1016/j.jpain.2005.02.001
- 2 11. Hu JH, Yang JP, Liu L, Li CF, Wang LN, Ji FH, Cheng H (2012) Involvement of
3 CX3CR1 in bone cancer pain through the activation of microglia p38 MAPK pathway
4 in the spinal cord. *Brain Res* 1465:1-9. doi:10.1016/j.brainres.2012.05.020
- 5 12. Chen X, Geller EB, Rogers TJ, Adler MW (2007) The chemokine
6 CX3CL1/fractalkine interferes with the antinociceptive effect induced by opioid
7 agonists in the periaqueductal grey of rats. *Brain Res* 1153:52-57.
8 doi:10.1016/j.brainres.2007.03.066
- 9 13. Hanisch U-K (2002) Microglia as a source and target of cytokines. *Glia* 40 (2):140-
10 155. doi:10.1002/glia.10161
- 11 14. Luo X, Tai WL, Sun L, Pan Z, Xia Z, Chung SK, Cheung CW (2016) Crosstalk
12 between astrocytic CXCL12 and microglial CXCR4 contributes to the development of
13 neuropathic pain. *Mol Pain* 12. doi:10.1177/1744806916636385
- 14 15. Morganti JM, Riparip L-K, Chou A, Liu S, Gupta N, Rosi S (2016) Age exacerbates
15 the CCR2/5-mediated neuroinflammatory response to traumatic brain injury. *J*
16 *Neuroinflammation* 13:80. doi:10.1186/s12974-016-0547-1
- 17 16. Holdridge SV, Armstrong SA, Taylor AMW, Cahill CM (2007) Behavioural and
18 morphological evidence for the involvement of glial cell activation in delta opioid

- 1 receptor function: implications for the development of opioid tolerance. *Mol Pain* 3:7.
2 doi:10.1186/1744-8069-3-7
- 3 17. Cui Y, Liao X-X, Liu W, Guo R-X, Wu Z-Z, Zhao C-M, Chen P-X, Feng J-Q (2008)
4 A novel role of minocycline: Attenuating morphine antinociceptive tolerance by
5 inhibition of p38 MAPK in the activated spinal microglia. *Brain Behav Immun* 22
6 (1):114-123. doi:10.1016/j.bbi.2007.07.014
- 7 18. Verge GM, Milligan ED, Maier SF, Watkins LR, Naeve GS, Foster AC (2004)
8 Fractalkine (CX3CL1) and fractalkine receptor (CX3CR1) distribution in spinal cord
9 and dorsal root ganglia under basal and neuropathic pain conditions. *Eur J Neurosci* 20
10 (5):1150-1160. doi:10.1111/j.1460-9568.2004.03593.x
- 11 19. McNally GP, WR (1998) Effects of systemic, intracerebral, or intrathecal
12 administration of an N-methyl-D-aspartate receptor antagonist on associative morphine
13 analgesic tolerance and hyperalgesia in rats. *Behav Neurosci* 112 (4):966-978
- 14 20. Johnston IN, Milligan ED, Wieseler-Frank J, Frank MG, Zapata V, Campisi J,
15 Langer S, Martin D, Green P, Fleshner M, Leinwand L, Maier SF, Watkins LR (2004)
16 A role for proinflammatory cytokines and fractalkine in analgesia, tolerance, and
17 subsequent pain facilitation induced by chronic intrathecal morphine. *J Neurosci* 24
18 (33):7353-7365. doi:10.1523/JNEUROSCI.1850-04.2004
- 19 21. Zhao CM, Guo RX, Hu F, Chen PX, Cui Y, Feng JQ, Meng JL, Mo LQ, Liao XX

- 1 (2012) Spinal MCP-1 Contributes to the Development of Morphine Antinociceptive
2 Tolerance in Rats. *Am J Med Sci* 344 (6):473-479.
3 doi:10.1097/MAJ.0b013e31826a82ce
- 4 22. Cui Y, Chen Y, Zhi JL, Guo RX, Feng JQ, Chen PX (2006) Activation of p38
5 mitogen-activated protein kinase in spinal microglia mediates morphine antinociceptive
6 tolerance. *Brain Res* 1069 (1):235-243. doi:10.1016/j.brainres.2005.11.066
- 7 23. Obata K, Yamanaka H, Kobayashi K, Dai Y, Mizushima T, Katsura H, Fukuoka T,
8 Tokunaga A, Noguchi K (2004) Role of Mitogen-Activated Protein Kinase Activation
9 in Injured and Intact Primary Afferent Neurons for Mechanical and Heat
10 Hypersensitivity after Spinal Nerve Ligation. *J Neurosci* 24 (45):10211
- 11 24. Zhu ZP, Badisa RB, Palm DE, Goodman CB (2012) Regulation of rat MOR-1 gene
12 expression after chronic intracerebroventricular administration of morphine. *Mol Med*
13 *Rep* 5 (2):513-516. doi:10.3892/mmr.2011.677
- 14 25. Heinisch S, Palma J, Kirby LG (2011) Interactions between chemokine and mu-
15 opioid receptors: anatomical findings and electrophysiological studies in the rat
16 periaqueductal grey. *Brain Behav Immun* 25 (2):360-372.
17 doi:10.1016/j.bbi.2010.10.020
- 18 26. Zhuang ZY, Kawasaki Y, Tan PH, Wen YR, Huang J, Ji RR (2007) Role of the
19 CX3CR1/p38 MAPK pathway in spinal microglia for the development of neuropathic

- 1 pain following nerve injury-induced cleavage of fractalkine. *Brain Behav Immun* 21
2 (5):642-651. doi:10.1016/j.bbi.2006.11.003
- 3 27. Milligan ED, Sloane EM, Watkins LR (2008) Glia in Pathological Pain: A Role for
4 Fractalkine. *J Neuroimmunol* 198 (1-2):113-120. doi:10.1016/j.jneuroim.2008.04.011
- 5 28. Raghavendra V, Rutkowski MD, DeLeo JA (2002) The role of spinal neuroimmune
6 activation in morphine tolerance/hyperalgesia in neuropathic and sham-operated rats. *J*
7 *Neurosci* 22(22):9980-9989
- 8 29. Hutchinson MR, Coats BD, Lewis SS, Zhang Y, Sprunger DB, Rezvani N, Baker
9 EM, Jekich BM, Wieseler JL, Somogyi AA, Martin D, Poole S, Judd CM, Maier SF,
10 Watkins LR (2008) Proinflammatory cytokines oppose opioid-induced acute and
11 chronic analgesia. *Brain Behav Immun* 22 (8):1178-1189.
12 doi:10.1016/j.bbi.2008.05.004
- 13 30. Wang Z, Ma W, Chabot JG, Quirion R (2009) Cell-type specific activation of p38
14 and ERK mediates calcitonin gene-related peptide involvement in tolerance to
15 morphine-induced analgesia. *FASEB J* 23(8):2576–2586
- 16 31. Wang Z, Ma W, Chabot JG, Quirion R (2010) Calcitonin gene-related peptide as a
17 regulator of neuronal CaMKII-CREB, microglial p38-NFκB and astroglial ERK-
18 Stat1/3 cascades mediating the development of tolerance to morphine-induced
19 analgesia. *Pain* 151(1):194–205

- 1 32. Wang Z, Ma W, Chabot JG, Quirion R (2010) Morphological evidence for the
2 involvement of microglial p38 activation in CGRP-associated development of
3 morphine antinociceptive tolerance. *Peptides* 31(12):2179–2184.
- 4 33. Mao J, Sung B, Ji RR, Lim G (2002) Chronic morphine induces downregulation of
5 spinal glutamate transporters: implications in morphine tolerance and abnormal pain
6 sensitivity. *J Neurosci* 22(18):8312–8323
- 7 34. Lin SL, Tsai RY, Shen CH, Lin FH, Wang JJ, Hsin ST, Wong CS (2010) Co-
8 administration of ultra-low dose naloxone attenuates morphine tolerance in rats via
9 attenuation of NMDA receptor neurotransmission and suppression of
10 neuroinflammation in the spinal cords. *Pharmacol Biochem Behav* 96(2):236–245
- 11 35. Hutchinson MR, Zhang Y, Shridhar M, Evans JH, Buchanan MM, Zhao TX, Slivka
12 PF, Coats BD, Rezvani N, Wieseler J, Hughes TS, Landgraf KE, Chan S, Fong S,
13 Phipps S, Falke JJ, Leinwand LA, Maier SF, Yin H, Rice KC, Watkins LR (2010)
14 Evidence that opioids may have toll-like receptor 4 and MD-2 effects. *Brain Behav*
15 *Immun* 24(1):83–95
- 16 36. Eidson LN, Inoue K, Young LJ, Tansey MG, Murphy AZ (2017) Toll-like Receptor
17 4 Mediates Morphine-Induced Neuroinflammation and Tolerance via Soluble Tumor
18 Necrosis Factor Signaling. *Neuropsychopharmacology* 42(3):661–670
- 19 37. Narita M, Suzuki M, Narita M, Niikura K, Nakamura A, Miyatake M, Yajima Y,

- 1 Suzuki T (2006) mu-Opioid receptor internalization-dependent and -independent
2 mechanisms of the development of tolerance to mu-opioid receptor agonists:
3 Comparison between etorphine and morphine. *Neuroscience* 138 (2):609-619.
4 doi:10.1016/j.neuroscience.2005.11.046
- 5 38. Arden JR, Segredo V, Wang Z, Lamah J, Sadée W (1995) Phosphorylation and
6 Agonist-Specific Intracellular Trafficking of an Epitope-Tagged μ -Opioid Receptor Ex
7 pressed in HEK 293 Cells. *J Neurochem* 65 (4):1636-1645. doi:10.1046/j.1471-
8 4159.1995.65041636.x
- 9 39. Koch T, Schulz S, Pfeiffer M, Klutzny M, Schroder H, Kahl E, Holtt V (2001) C-
10 terminal splice variants of the mouse mu-opioid receptor differ in morphine-induced
11 internalization and receptor resensitization. *J Biol Chem* 276 (33):31408-31414.
12 doi:10.1074/jbc.M100305200
- 13 40. Chakrabarti S, Madia PA, Gintzler AR (2015) Selective upregulation of functional
14 mu-opioid receptor splice variants by chronic opioids. *J Neurochem.*
15 doi:10.1111/jnc.13519
- 16 41. Caputi FF, Lattanzio F, Carretta D, Mercatelli D, Candeletti S, Romualdi P (2013)
17 Morphine and fentanyl differently affect MOP and NOP gene expression in human
18 neuroblastoma SH-SY5Y cells. *J Mol Neurosci : MN* 51 (2):532-538.
19 doi:10.1007/s12031-013-0019-3

- 1 42. Szentirmay AK, Kiraly KP, Lenkey N, Lacko E, Al-Khrasani M, Friedmann T,
2 Timar J, Gyarmati S, Toth G, Furst S, Riba P (2013) Spinal interaction between the
3 highly selective mu agonist DAMGO and several delta opioid receptor ligands in naive
4 and morphine-tolerant mice. *Brain Res Bull* 90:66-71.
5 doi:10.1016/j.brainresbull.2012.09.006
- 6 43. Xu J, Lu Z, Xu M, Rossi GC, Kest B, Waxman AR, Pasternak GW, Pan YX (2014)
7 Differential expressions of the alternatively spliced variant mRNAs of the micro opioid
8 receptor gene, OPRM1, in brain regions of four inbred mouse strains. *PloS one* 9
9 (10):e111267. doi:10.1371/journal.pone.0111267
- 10 44. Taylor DA, Fleming WW (2001) Unifying Perspectives of the Mechanisms
11 Underlying the Development of Tolerance and Physical Dependence to Opioids. *J*
12 *Pharmacol Exp Ther* 297 (1):11-18
- 13 45. Yan H, Yu LC (2013) Influences of calcitonin gene-related peptide on mu opioid
14 receptors in nucleus accumbens neurons of rats. *Neuropeptides* 47 (2):125-131.
15 doi:10.1016/j.npep.2012.10.008
- 16 46. Raehal KM, Bohn LM (2011) The role of beta-arrestin2 in the severity of
17 antinociceptive tolerance and physical dependence induced by different opioid pain
18 therapeutics. *Neuropharmacology* 60 (1):58-65.
19 doi:10.1016/j.neuropharm.2010.08.003

- 1 47. Paolicelli RC, Bolasco G, Pagani F, Maggi L, Scianni M, Panzanelli P, Giustetto M,
2 Ferreira TA, Guiducci E, Dumas L, Ragozzino D, Gross CT (2011) Synaptic Pruning
3 by Microglia Is Necessary for Normal Brain Development. *Science* 333 (6048):1456-
4 1458. doi:10.1126/science.1202529
- 5 48. Zhan Y, Paolicelli RC, Sforazzini F, Weinhard L, Bolasco G, Pagani F, Vyssotski
6 AL, Bifone A, Gozzi A, Ragozzino D, Gross CT (2014) Deficient neuron-microglia
7 signaling results in impaired functional brain connectivity and social behavior. *Nat*
8 *Neurosci* 17 (3):400-406. doi:10.1038/nn.3641
- 9 49. Limatola C, Ransohoff RM (2014) Modulating neurotoxicity through
10 CX3CL1/CX3CR1 signaling. *Front Cell Neurosci* 8:229.
11 doi:10.3389/fncel.2014.00229
- 12 50. Gregg B, Lumeng CN, Bernal-Mizrachi E (2014) Fractalkine signaling in regulation
13 of insulin secretion: Mechanisms and potential therapeutic implications? *Islets* 6
14 (1):e27861. doi:10.4161/isl.27861
- 15 51. Umehara H, Bloom ET, Okazaki T, Nagano Y, Yoshie O, Imai T (2004) Fractalkine
16 in vascular biology: from basic research to clinical disease. *Arterioscler Thromb Vasc*
17 *Biol* 24 (1):34-40. doi:10.1161/01.ATV.0000095360.62479.1F
- 18 52. Pello OM, Martínez-Muñoz L, Parrillas V, Serrano A, Rodríguez-Frade JM, Toro
19 MJ, Lucas P, Monterrubio M, Martínez-A C, Mellado M (2008) Ligand stabilization of

1 CXCR4/ δ -opioid receptor heterodimers reveals a mechanism for immune response
2 regulation. *Eur J Immunol* 38 (2):537-549. doi:10.1002/eji.200737630

3 53. Hutchinson MR, Shavit Y, Grace PM, Rice KC, Maier SF, Watkins LR (2011)
4 Exploring the Neuroimmunopharmacology of Opioids: An Integrative Review of
5 Mechanisms of Central Immune Signaling and Their Implications for Opioid Analgesia.
6 *Pharmacol Rev* 63 (3):772

7 54. Szabo I, Chen X-H, Xin L, Adler MW, Howard OMZ, Oppenheim JJ, Rogers TJ
8 (2002) Heterologous desensitization of opioid receptors by chemokines inhibits
9 chemotaxis and enhances the perception of pain. *Proc Natl Acad Sci* 99 (16):10276-
10 10281. doi:10.1073/pnas.102327699

11 55. Chen X, Geller EB, Rogers TJ, Adler MW (2007) Rapid heterologous
12 desensitization of antinociceptive activity between mu or delta opioid receptors and
13 chemokine receptors in rats. *Drug Alcohol Depend* 88 (1):36-41

14 56. Hu XM, Liu YN, Zhang HL, Cao SB, Zhang T, Chen LP, Shen W (2015)
15 CXCL12/CXCR4 chemokine signaling in spinal glia induces pain hypersensitivity
16 through MAPKs-mediated neuroinflammation in bone cancer rats. *J Neurochem* 132
17 (4):452-463. doi:10.1111/jnc.12985

18

19 **Figure 1. Expressions of MOR and Iba-1 in lumbar spinal cord of rats.**

20 **A.** Thermal pain threshold of rats was assessed using the percentage of maximal

1 possible antinociceptive effect (%MPE) according to the tail-flick latency of rats.
2 The %MPE in rats received morphine (10 µg, twice daily, intrathecally) on day 5 and 7
3 were dramatically decreased compared with the baseline on day 1. Values represent
4 mean ± SEM; two-way ANOVA, ** $P < 0.01$, *** $P < 0.001$, vs. naïve and sham, n =
5 6 in each group. **B.** The expression of Iba-1 protein was significantly increased in
6 morphine-treated rats measured by Western blots. Values represent mean ± SEM;
7 ANOVA, * $P < 0.05$ vs. naïve and sham, n = 6 in each group. **C and D.** The expressions
8 of MOR mRNA (C) and protein (D) were not affected by morphine treatment measured
9 by real-time PCR and Western blots, respectively. Values represent mean ± SEM;
10 ANOVA, * $P < 0.05$ vs. naïve and sham, n = 6 in each group.

11

12 **Figure 2. Expressions of CX3CL1 and CX3CR1 in lumbar spinal cord of rats.**

13 **A.** The expressions of CX3CL1 and CX3CR1 mRNA were not affected by morphine
14 treatment measured by real-time PCR. Values represent mean ± SEM; ANOVA, * $P <$
15 0.05 vs. naïve and sham, n = 6 in each group. **B and C.** The expressions of CX3CR1
16 (B) and CX3CL1 (C) protein were not affected by morphine treatment measured by
17 Western blots. Values represent mean ± SEM; ANOVA, * $P < 0.05$ vs. naïve and sham,
18 n = 8 in each group for CX3CR1, n = 5 in each group for CX3CL1.

19

20 **Figure 3. Effects of recombinant rat CX3CL1 and anti-CX3CR1 neutralizing** 21 **antibody on the development of morphine tolerance.**

22 **A.** Tail flick latency of rats in each group did not change after intrathecal catheterization.
23 **B and C.** Recombinant rat CX3CL1 (100 ng or 500 ng) (B), anti-CX3CR1 neutralizing
24 antibody (5 µg or 10 µg) (C) or normal IgG (100 ng for rrCX3CL1 group, 5 µg for anti-

1 CX3CR1 neutralizing antibody group as control dose.) was intrathecally administered
2 30 minutes before morphine treatment for 7 days. None of them significantly affected
3 morphine antinociception or alleviated the development of morphine tolerance. Values
4 represent mean \pm SEM; two-way ANOVA, **** $P < 0.001$, * $P < 0.05$ vs. naïve and
5 sham, n = 6 in each group.

6

7 **Figure 4. Effects of exogenous CX3CL1 and CX3CR1 inhibitor on the microglia**
8 **activation with repeated morphine administration for 7 consecutive days.**

9 Exogenous CX3CL1 (100 ng or 500 ng), CX3CR1 inhibitor (5 μ g or 10 μ g) and IgG (5
10 μ g) were administered 30 minutes before morphine treatment for 7day, respectively.
11 Microglia activity was increased significantly in morphine involved groups. However,
12 both exogenous CX3CL1 and CX3CR1 inhibitor showed no influence in quantification
13 of Iba-1 levels from Western blots compared to morphine treated rats. Values represent
14 mean \pm SEM; one-way ANOVA, * $P < 0.05$ vs. sham, n = 3 in each group.

15

16 **Figure 5. Changes in the cellular localizations of CX3CL1 and CX3CR1 during**
17 **the development of morphine tolerance.**

18 **A and B.** Expressions of CX3CL1 and CX3CR1 in the spinal cord. There were no
19 changes in the number of CX3CL1 (a, b) and CX3CR1 (c, d) immunoreactive cells in
20 spinal dorsal horn of morphine-tolerant rats compared with those in sham rats (n = 3,
21 scale bar 200 μ m). **C.** Double immunostaining of CX3CL1 (a, b, c) or CX3CR1 (d, e,
22 f) and cell-specific markers in the spinal cord in saline-treated rats. CX3CL1 was co-
23 localized with NeuN and CX3CR1 was co-localized with Iba-1. **a:** CX3CL1 and NeuN;
24 **b:** CX3CL1 and GFAP; **c:** CX3CL1 and Iba-1; **d:** CX3CR1 and NeuN; **e:** CX3CR1 and

1 GFAP; f: CX3CR1 and Iba-1. Scale bar: 200 μ m. **D.** Double immunostaining of
2 CX3CL1 and cell-specific markers in morphine-treated rats. CX3CL1 was co-localized
3 with NeuN, GFAP and Iba-1 (indicated by arrows). a, e and i: CX3CL1; b: NeuN; f:
4 GFAP; j: Iba-1; c and d: CX3CL1 merged with NeuN; g and h: CX3CL1 merged with
5 GFAP; k and l: CX3CL1 merged with Iba-1. Scale bars: 200 μ m (a, b, c, e, f, g, i, j and
6 k); scale bar: 100 μ m (d, h and l). **E.** Double immunostaining of CX3CR1 and cell-
7 specific markers in morphine-treated rats. CX3CR1 was co-localized with NeuN, GFAP
8 and Iba-1 (indicated by arrows). a, e and i: CX3CR1; b: NeuN; f: GFAP; j: Iba-1; c and
9 d: CX3CR1 merged with NeuN; g and h: CX3CR1 merged with GFAP; k and l:
10 CX3CR1 merged with Iba-1. Scale bar: 200 μ m (a, b, c, e, f, g, i, j and k); scale bar:
11 100 μ m (d, h and l).

12

13 **Figure 6. Heatmap of expression ratios of chemokines family mRNAs.**

14 The probe sets that expressions were changed in morphine tolerance rats were identified
15 by microarray analysis. Probe sets with similar expression profiles were clustered
16 together using a Pearson's correlation-based method with Cluster 3.0 and TreeView
17 software. The expression level of each chemokine probe set was displayed as a log₂
18 ratio of their expression values divided by their expression values in sham rats.