



Cassava brown streak disease: historical timeline, current knowledge and future prospects

Journal:	<i>Molecular Plant Pathology</i>
Manuscript ID	MPP-Rev-17-149.R2
Manuscript Type:	Review
Date Submitted by the Author:	07-Sep-2017
Complete List of Authors:	Tomlinson, Katie; School of Biological Sciences, University of Bristol Bailey, Andy; Univeristy of Bristol, School of Biological Sciences Seal, Sue; National Resources Institute Alicai, Titus; National Crops Resources Research Institute Foster, Gary; School of Biological Sciences, University of Bristol
Keywords:	CBSD, CBSV, UCBSV, CMD, Food Security, Virology

Title: Cassava brown streak disease: historical timeline, current knowledge and future prospects

Key words:

CBSD; CBSV; UCBSV; CMD; Food Security; Virology

Authors:

Katie R. Tomlinson (1); Andy M. Bailey (1); Titus Alicai (2); Sue Seal (3); Gary D. Foster (1)*

(1) School of Biological Sciences, University of Bristol, 24 Tyndall Avenue, Bristol, BS8 1TQ, UK

(2) National Crops Resources Research Institute, P.O. Box 7084, Kampala, Uganda

(3) Natural Resources Institute, University of Greenwich, Central Avenue, Chatham Maritime, Kent ME4 4TB, UK

*Corresponding author email: Gary.Foster@bristol.ac.uk, Tel: 0117 39 41178

Word count:

Abstract: 197

Review: 6716

Acknowledgements: 27

Figure Legends: 1 = 40; 2 = 9; 3 = 20; 4 = 6; 5 = 658

Suggested running title:

CBSD: timeline, current knowledge and future prospects

Abstract

Cassava is the second most important staple food crop in terms of *per capita* calories consumed in Africa and holds potential for climate change adaptation. Unfortunately, productivity in East and Central Africa is severely constrained by two viral diseases: cassava mosaic disease (CMD) and cassava brown streak disease (CBSD). CBSD was first reported in 1936 from northeast Tanzania. For approximately seventy years CBSD was restricted to coastal East Africa and so had a relatively low impact on food security compared to CMD. However, at the turn of the 21st century CBSD re-emerged further inland, in areas around Lake Victoria and it has since spread through many East and Central African countries, causing high yield losses and jeopardising the food security of subsistence farmers. This recent re-emergence has attracted intense scientific interest, with studies shedding light on CBSD viral epidemiology, sequence diversity, host interactions and potential sources of resistance within the cassava genome. This review reflects on 80 years of CBSD research history (1936 – 2016) with a timeline of key events. We provide insights into current CBSD knowledge, management efforts and future prospects for improved understanding needed to underpin effective control and mitigation of impacts on food security.

Introduction

Cassava (*Manihot esculenta* Crantz, family *Euphorbiaceae*) produces carbohydrate rich storage roots, which are a staple food crop for approximately 800 million people worldwide (FAO 2013). In Africa, cassava is the second most important food staple in terms of *per capita* calories consumed (Nweke 2004). Storage roots are used as a fresh carbohydrate source and can also be processed into flour, which may be consumed by the grower's family, sold in local markets or used to produce several industrial food products (Hillocks & Thresh 2002). Subsistence farmers rely on cassava for a vital energy source, as it can be planted and harvested throughout the year, tolerates periods of unpredictable droughts and grows on marginal soils (Hillocks & Thresh 2002). Recent modeling has suggested that cassava may be highly resilient to future climate change and could provide Africa with adaptation opportunities, which are not offered by other staple food crops (Jarvis *et al.* 2012).

Cassava was introduced into Africa from Brazil by Portuguese traders in the 16th century and subsequently integrated into local agriculture in countries across the continent (Jones 1959). Africa produces over half of global cassava (57%) (Bennett 2015), however the continent's average fresh yield (9.9 t ha⁻¹) lags behind potential yields (15 – 40 t ha⁻¹) achieved under experimental conditions (Fermont *et al.* 2009). There are many reasons behind reduced yields, including restricted access to labour, poor soil quality and premature harvesting (Fermont *et al.* 2009). Productivity in East and Central Africa is significantly constrained by two viral diseases: cassava mosaic disease (CMD) and cassava brown

1 streak disease (CBSD), which together are estimated to cause annual
2 losses worth US\$1billion (IITA 2014a) and adversely affect food security
3 in the entire region (Patil *et al.* 2015).
4
5

6
7 In this paper, we review CBSD research history, highlighting key events
8 in a timeline (Fig. 1) and provide future prospects for further
9 understanding and effective control. The review is split into two phases
10 according to the geographical distribution of CBSD. Phase one covers
11 the small number (n=65) of reports published between 1936 and the
12 early 1990s when CBSD was reported to be restricted to low altitude
13 areas (<1000 meters above sea level (masl)) along coastal East Africa
14 and lake shore districts of Malawi (Legg *et al.* 2011). Phase two
15 examines CBSD re-emergence after the mid 1990s, when CBSD spread
16 across East and Central Africa (Legg *et al.* 2011). We review the
17 corresponding increased number (n=277) of reports on CBSD
18 geographical expansion, viral molecular characterization, host
19 interactions, diagnostic techniques and control efforts.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35

36 We offer insights into what can be learnt from CBSD history, in particular
37 the need for application of knowledge to protect against and predict
38 multiple biotic threats to staple food crops through improved
39 understanding of CBSD epidemiology, diagnostics, surveillance and
40 predictive modeling. This calls for effective international scientific
41 collaborations across multiple areas of expertise and the rapid
42 application of research and technologies to solve problems affecting
43 farmers.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Initial emergence and symptom description (1930s – early 1990s)

The first report of CBSD from northeast Tanzania (then called Tanganyika) describe distinctive foliar symptoms on lower mature cassava leaves and rot of storage roots (Storey 1936). Nichols (1950) later reported that symptoms could be expressed on all parts of the plant and include storage root necrosis (Fig. 3A), radial root constrictions (Fig. 3B), foliar chlorosis (Fig. 3C) and occasionally brown streaks or lesions on stems (Fig. 3D).

It was noted that two main types of foliar symptoms exist: (1) feathery chlorosis along secondary vein margins, which eventually coalesce to form blotches, (2) chlorotic mottling with no veinal association (Nichols 1950). These distinctive symptoms lack the leaf distortion observed in CMD infected cassava plants. CBSD symptoms are variable in terms of severity, onset of symptom expression and parts of the plant affected, depending on the viral strain, cassava cultivar, environmental conditions and the age of the plant when infected (Nichols 1950). This variability makes diagnosis difficult for farmers (Nichols 1950) and can result in farmers being unaware that their crop is affected until they harvest storage roots (Legg & Kanju 2015). The difficulty in diagnosing CBSD has meant that infected stems have been transported to areas where CBSD has previously been absent and used for planting material. Symptom variability has also hampered epidemiology studies, as the disease can go unnoticed in an area for long periods.

The surveying and symptom scoring of infected plants across different geographical areas revealed that most plants with foliar symptoms usually also develop root necrosis (Hillocks *et al.* 2001). In the most

1 sensitive cultivars, CBSD reduces root weight by up to 70% with necrosis
2 developing at six months post planting (Hillocks *et al.* 2001). Whereas, it
3 was reported that the local Tanzanian cultivar "Nachinyaya" did not
4 develop root necrosis and so was relatively tolerant to CBSD (Hillocks *et*
5 *al.* 1996).
6
7
8
9
10

11 **Early geographical distribution**

12 Storey (1939) reported that CBSD was widespread in coastal Tanzania
13 and by 1950, the disease was endemic across coastal areas of East
14 Africa from northeast Kenya, Tanzania to northern Mozambique at
15 altitudes below 1000 masl (Nichols 1950). The disease was reported in
16 Uganda in 1945 and may have been introduced through infected cuttings
17 sent from the Amani research station in Tanzania (Nichols 1950;
18 Jameson 1964). Strict roguing of infected plants, replacement with non-
19 infected planting material and quarantine appear to have prevented
20 spread of CBSD in Uganda at this time (Nichols 1950). Significantly, a
21 lack of plant-to-plant vector transmission at higher altitudes was reported
22 (Nichols 1950; Jennings 1960).
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 **Causal agent characterization**

42 Storey (1936) suspected a viral causal agent, as CBSD was successfully
43 transmitted through grafting stem cuttings. Subsequently, Lister (1959)
44 mechanically transmitted CBSD to indicator hosts, including *Petunia*
45 *hybrida*, *Datura stramonium*, *Nicotiana tabacum*, *N. rustica* and *N.*
46 *glutinosa*, which produce a range of symptoms depending on the
47 sensitivity of the host and viral variant. In 1976, sap transmission of
48 CBSD from infected cassava material to *N. clevelandii* produced two
49
50
51
52
53
54
55
56
57
58
59
60

1 distinct symptom types, which suggested that two viral variants may be
2 responsible for CBSD (Bock & Guthrie 1976).
3

4
5 Virus particles were identified by electron microscopy analysis of CBSD
6 infected *N. debneyi* (Bock 1994). The infected samples contained 650nm
7 filamentous particles with a similar morphology to viruses within the
8 *Carlavirus* genus (Bock 1994). However, pinwheel inclusions, typical of
9 *Potyviridae* were identified in in CBSD infected *N. benthamiana* (Lennon
10 *et al.* 1985). Pinwheel inclusions were subsequently found through more
11 thorough electron microscopy of CBSD infected cassava samples, albeit
12 at low concentrations (Were *et al.* 2004).
13
14
15
16
17
18
19
20
21
22
23

24 The *Potyviridae* sequence identity was finally confirmed in 2001 through
25 reverse transcription PCR (RT-PCR) on CBSD infected *N. benthamiana*
26 samples (Monger *et al.* 2001a). When the RT-PCR product was
27 sequenced, it aligned most closely to the coat protein sequence of *Sweet*
28 *potato mild mottle virus* (SPMMV, genus *Ipomovirus*, family *Potyviridae*)
29 (Monger *et al.* 2001a). The same RT-PCR technique was used to detect
30 CBSV in symptomless cassava leaves, highlighting the sensitivity of the
31 RT-PCR technique (Monger *et al.* 2001a).
32
33
34
35
36
37
38
39
40
41
42

43 **Early control efforts**

44
45
46 In the 1930s, a cassava breeding programme was launched in Tanzania,
47 which included breeding for CBSD and CMD resistance at the Amani
48 research station (Jennings 1957; Nichols 1946). Early breeding to
49 develop virus resistant cultivars involved crossing cultivated cassava with
50 wild relatives, including *M. glaziovii*, *M. dichotoma*, *M. catingea*, *M.*
51 *saxicola* and *M. melanobasis*, which are believed to have higher levels of
52
53
54
55
56
57
58
59
60

1 CBSD resistance (Jennings 1957; Kawuki *et al.* 2016). The breeding
2 programme produced the *M. esculenta*-*M. glaziovii* hybrid known as
3 "Namikonga" in Tanzania or "Kaleso" in Kenya, which for many years
4 offered relatively high levels of CBSD tolerance (Hillocks & Jennings
5 2003; Kaweesi *et al.* 2014). However, "Namikonga" was not widely
6 distributed to farmers, which may be because of its susceptibility to CMD
7 (Hillocks & Jennings 2003; Kawuki *et al.* 2016).

17 **Initial vector transmission studies**

18
19
20 Until relatively recently, very little was known about vector transmission
21 of CBSVs. It had been noted that CBSD outbreaks tended to coincide
22 with increases in whitefly populations (Storey 1939; Hillocks & Jennings
23 2003). However initial attempts to transmit CBSV with whitefly (*B. tabaci*)
24 or aphid (*Myzus persicae*) were unsuccessful (Bock 1994).

32 **Geographical distribution in the early 1990s**

33
34
35 In the early 1990s, there were reports of high CBSD incidences in areas
36 of Tanzania, Mozambique and Malawi (Hillocks & Jennings 2003).
37 Surveys revealed CBSD incidences reaching 36% - 50% in cassava
38 fields along coastal areas of Tanzania (Legg & Raya 1998; Hillocks *et al.*
39 1999). Similarly, CBSD incidences in Malawi reached 75% in many fields
40 surrounding Lake Malawi and nearly all plants inspected in northern
41 coastal areas of Mozambique were expressing CBSD symptoms
42 (Hillocks *et al.* 2002; Hillocks & Jennings 2003). In a control effort, virus-
43 free CBSD tolerant cultivars were distributed to farmers in Mozambique
44 who depended heavily on CBSD sensitive cassava cultivars for food
45 security (Hillocks & Jennings 2003). CBSD was also re-discovered in
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Uganda in 1994 at a site near Entebbe (Thresh *et al.* 1994). This led
2
3 researchers to call for concerted efforts to understand CBSD through
4
5 improved surveillance (Hillocks & Jennings 2003).
6
7

8 **Reflections on initial emergence (1930s – early 1990s)**

9

10
11 Despite CBSD being endemic across coastal East Africa during this
12
13 period, relatively little work was done to understand and control CBSD.
14
15 This is reflected by the low number of scientific papers, reports or
16
17 reviews which feature CBSD published between 1936 and early 1990s
18
19 (n=65) (Fig. 2). The slight increase in references to CBSD in the 1970s is
20
21 due to a small number of reports (n= 27) on the threat posed by CBSD.
22
23 In hindsight, these reports should have served as a warning to take
24
25 control actions, which may have prevented the later expansion of CBSD
26
27 across the region.
28
29

30
31
32 There was a general lack of scientific interest in CBSD at this time due to
33
34 many factors, including the restricted occurrence of CBSD to low altitude
35
36 areas along coastal eastern Africa and the devastating impacts of CMD
37
38 on food security. During this period, CMD was a greater priority due to its
39
40 prevalence across all cassava-growing areas of Africa, resulting in
41
42 famines, higher economic losses and forcing many farmers to abandon
43
44 the crop (Thresh *et al.* 1994; Thresh & Cooter 2005; Alabi *et al.* 2011). To
45
46 help control the disease, CMD resistant cultivars were distributed to
47
48 areas severely affected (Legg & Thresh 2000). Unfortunately these
49
50 cultivars show varying levels of CBSD susceptibility (Legg *et al.* 2006). It
51
52 is not known whether the deployment of these cultivars has contributed
53
54 to the increased distribution of CBSD in the field.
55
56
57
58
59
60

Re-emergence and expansion across East and Central Africa (mid 1990s – 2016)

In 2004, the apparent restriction of CBSD to coastal lowlands changed with the re-emergence of CBSD at altitudes above 1000masl (Alicai *et al.* 2007). Infections of cassava plants showing CBSD symptoms at higher altitudes in Uganda were confirmed by RT-PCR. Coat protein sequences aligned to CBSV isolates from Mozambique and Tanzania with sequence identities from 77.0 to 82.9% (Alicai *et al.* 2007). It is not known whether CBSD had been re-introduced to Uganda through infected cuttings or whether the disease had existed at a low level since it was first introduced in the 1940s (Alicai *et al.* 2007). Shortly after this first report, overall incidence of CBSD in Uganda increased from 12% in 2008 to 27% in 2011 (T. Alicai personal communication) and similar increases were reported in Tanzania and Kenya (Mware *et al.* 2009; Ntawuruhunga & Legg 2007; Legg *et al.* 2011). There have since been CBSD reports from Burundi (Bigirimana *et al.* 2011), Rwanda (FAO 2011), eastern Democratic Republic of Congo (Mulimbi *et al.* 2012), South Sudan (T. Alicai personal communication) and Mayotte Island (Roux-Cuvelier *et al.* 2014).

It is difficult to obtain a truly accurate estimation of the economic damage caused by CBSD, however an overall loss of US\$750 million a year is estimated across Kenya, Tanzania, Uganda, and Malawi (Hillocks & Maruthi 2015). CBSD is now one of the leading causes of cassava losses in East Africa (Pennisi 2010) and its on-going spread threatens the major cassava growing areas of Central and West Africa (Legg *et al.* 2014).

1 The dramatic increase in the impact of CBSD on food security is reflected
2 in the increase of papers, reports and reviews, which refer to CBSD
3 published from the mid 1990s to 2016 (n= 277) (Fig. 2). The expansion of
4 the CBSD epidemic across the Great Lakes region of East and Central
5 Africa has necessitated the rapid development and implementation of
6 effective control strategies. Several important projects were initiated
7 following CBSD re-emergence, which aimed to develop research,
8 extension and policy capacity in the countries affected. Key targets have
9 been to breed or genetically engineer resistant cultivars, provide certified
10 virus-clean planting material and improve viral surveillance and diagnosis
11 (Legg *et al.* 2014).
12
13
14
15
16
17
18
19
20
21
22
23
24

25 **Recent local and regional CBSD epidemiology**

26 The reasons behind the sudden increase in CBSD incidence and
27 geographical range remain poorly understood. Studies have shown
28 CBSD spread and development is enhanced by high disease pressure,
29 use of susceptible genotypes and high whitefly numbers (Katono *et al.*
30 2015). CBSD is dispersed locally and over long distances through the
31 trade transportation of infected planting material, whereas whiteflies are
32 only able disperse and amplify CBSD locally (McQuaid *et al.* 2017).
33
34
35
36
37
38
39
40
41
42
43
44

45 The ability of *B. tabaci* to transmit CBSV from infected to healthy plants
46 was confirmed under quarantine insectary and glasshouse conditions in
47 Maruthi *et al.* (2005). It has since been shown that CBSD viruses are
48 transmitted semi-persistently, with whiteflies acquiring viruses in 5 – 10
49 minutes, retaining them for up to 48 hours, and transmitting them over
50 relatively short distances of less than 17 meters in a cropping season
51
52
53
54
55
56
57
58
59
60

1 (Maruthi *et al.* 2016). CBSD outbreaks occur from 3 to 12 years after
2 increases in whitefly numbers (Legg *et al.* 2011). Critically, one of the
3 primary cause for increases of both CMD and CBSD in the African Great
4 Lakes region appears to be super-abundant numbers of whiteflies (Fig.
5 4), which are able to thrive at altitudes above 1000 masl (Alicai *et al.*
6 2007; Jeremiah *et al.* 2015).
7
8
9
10
11
12

13
14 Survey data has revealed that the transportation of infected material to
15 areas where CBSD was previously absent has enabled the disease to
16 spread from independent hot spots (Legg *et al.* 2011). This is because
17 cassava stems used for vegetative planting material are exchanged by
18 farmers across localities and transported over long distances. One report
19 concluded that plants can also be infected through the use of
20 contaminated cutting tools, which could contribute to in-field spread
21 (Rwegasira & Chrissie 2015), however a similar study showed that such
22 practices did not result in transmission of CBSVs (Maruthi *et al.* 2016).
23
24
25
26
27
28
29
30
31
32
33
34
35

36 CBSD viruses are found only in Africa and so it appears that these
37 viruses have evolved within in East Africa on an unknown species and
38 subsequently jumped host into cassava in a new encounter situation
39 (Monger *et al.* 2010). Therefore there may be other hosts for CBSVs,
40 which could serve as a viral inoculum sources in the field (Monger *et al.*
41 2010). CBSV has been detected in the wild perennial species *M. glaziovii*
42 (Ogwok *et al.* 2014); the importance of this to CBSD epidemiology is not
43 currently known.
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Molecular characterisation of unusual CBSD viral genome features

CBSVs belong to the *Ipomovirus* genus of the *Potyviridae* family (Monger *et al.* 2001a). *Ipomoviruses* have positive sense, single-stranded genomes, which are translated as large polyproteins and autocatalytically cleaved by virus-encoded proteases into ten mature proteins with an additional P3N-PIPO protein produced through ribosomal frameshifting (Valli *et al.* 2015). The genome organisation of CBSVs is shown in Fig. 5.

Molecular characterization of coat protein sequences has revealed that there are at least two genetically distinct species: *Cassava brown streak virus* (CBSV) and *Ugandan Cassava brown streak virus* (UCBSV) (Monger *et al.* 2001b; Winter *et al.* 2010), which typically have 76–78% nucleotide and 87–90% amino acid identity (Mbanzibwa *et al.* 2009b).

Genome analysis has revealed that CBSVs share unusual features (Mbanzibwa *et al.* 2009a; Monger *et al.* 2010). Firstly, CBSVs lack the multi-functional helper-component proteinase protein (HCPro), which possesses silencing suppressor, vector transmission and long distance movement *in planta* activities in *Potyviridae* viruses (Valli *et al.* 2015). The HCPro protein is found in all other known *Potyviridae* viruses, except for *Squash vein yellowing virus* (SqVYV) and *Cucumber vein yellowing virus* (CVYV) (Mbanzibwa *et al.* 2009a). In CBSVs, HCPro appears to have been replaced by silencing suppressor activity of the P1 serine proteinase (Mbanzibwa *et al.* 2009a). CBSV and UCBSV P1 proteins are most closely related to P1 of SPMMV and P1b of SqVYV and CVYV, which are related to the tritomovirus P1 proteins (Mbanzibwa *et al.* 2009a). The CBSV and UCBSV P1 proteins both contain zinc finger and LXKA motifs (Mbanzibwa *et al.* 2009a). The zinc finger and LXKA motifs

1 in CVYV P1b are essential for silencing suppressor activity through the
2 binding and sequestering of siRNA required for silencing (Valli *et al.*
3 2008). It is therefore likely that the same motifs are responsible for the
4 silencing suppressor activities of CBSV and UCBSV P1 proteins
5 (Mbanzibwa *et al.* 2009a).
6
7
8
9

10 CBSV and UCBSV also encode novel Ham1 proteins with conserved
11 Maf/Ham1 motifs (Mbanzibwa *et al.* 2009a). Proteins with Maf/Ham1
12 domains are found across prokaryotic and eukaryotic organisms and
13 have nucleoside triphosphate pyrophosphatase activities, which reduce
14 mutation rates by preventing the incorporation of non-canonical
15 nucleotides into RNA and DNA (Galperin *et al.* 2006). The functions of
16 CBSV and UCBSV Ham1 proteins are yet to be elucidated but they are
17 likely to provide essential functions in the lifecycles of CBSVs. For
18 instance, Ham1 proteins may reduce mutation rates under oxidative
19 stress conditions in mature cassava leaves where CBSV viruses are
20 found at the highest concentrations within the plant (Ogwok *et al.* 2014).
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Euphorbia ring spot virus (EuRV, genus *Potyvirus*, family *Potyviridae*)
also encodes a Ham1 protein with an uncharacterized function (Knierim
et al. 2016). EuRV, CBSV and UCBSV are part of a small number of
viruses which are able to infect plants in the *Euphorbiaceae* family and
so perhaps Ham1 proteins are a euphorbia host adaptation (Monger *et al.* 2010).

Differences between CBSVs infections and genome sequences

In 2010, CBSV was found in infected cassava samples from Mozambique and Tanzania and UCBSV in Kenya, Uganda, Malawi and north-western Tanzania (Winter *et al.* 2010). However, recent

1 phylogenetic analysis of whole genome sequences has revealed that the
2 viral species are not limited to agro-ecological zones and that there may
3 be three separate species within the UCBSV clade (Ndunguru *et al.*
4 2015).
5
6
7
8
9

10 CBSV and UCBSV produce distinctly different symptoms on cassava and
11 indicator hosts. CBSV causes more severe root necrosis and feathery
12 chlorosis along vein margins, which develops into chlorotic blotches,
13 whereas UCBSV causes circular chlorotic blotches between veins in
14 cassava (Mohammed *et al.* 2012; Winter *et al.* 2010; Nichols 1950).
15 CBSV tends to accumulate to higher titers than UCBSV in cassava
16 (Kaweesi *et al.* 2014) and indicator plants (Mohammed *et al.* 2012;
17 Ogwok *et al.* 2014).
18
19
20
21
22
23
24
25
26
27

28 Sequence differences between CBSV and UCBSV genomes should
29 explain differences in symptom severities, viral loads and host-
30 interactions observed between the two viral species. Key areas of CBSV
31 and UCBSV genomes show relatively high levels of divergence, including
32 the P1 and Ham1 regions, with only 59% and 47% amino acid identities
33 respectively (Winter *et al.* 2010). One suggestion for the low level of
34 Ham1 sequence similarity is that Ham1 genes may have been acquired
35 separately by CBSV and UCBSV from a eukaryotic host (Monger *et al.*
36 2010). Alternatively CBSV and UCBSV Ham1 sequences may be derived
37 from a common ancestor, which have diverged due to differential
38 selection pressures on the genome sequences of the two species
39 (Monger *et al.* 2010).
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Evolution of CBSVs

1
2
3 Statistical analysis of CBSV and UCBSV genomes using the Empirical
4 Bayes approach has predicted amino acid sites in UCBSV and CBSV
5 coat protein and UCBSV Ham1 sequences, which appear to have been
6 under positive selection (Mbanzibwa *et al.* 2011a). It is possible that
7 positive selection at these different amino acid positions may be enabling
8 adaptive evolution of the two viral species (Mbanzibwa *et al.* 2011a).
9 Recent whole genome sequence analysis has revealed that there is a
10 higher diversity of CBSV isolates compared to UCBSV (Alicai *et al.*
11 2016). This diversity may be enabling CBSV to rapidly adapt to overcome
12 host resistance mechanisms, which breeders have been selecting for
13 (Alicai *et al.* 2016).
14
15

16
17 Whole genome analysis has also identified putative homologous
18 recombination sites within the genomes of CBSV and UCBSV isolates
19 (Ndunguru *et al.* 2015). To date there has been no evidence for
20 recombination between CBSV and UCBSV isolates (Mbanzibwa *et al.*
21 2011a; Ndunguru *et al.* 2015). However, analysis of more CBSV and
22 UCBSV genome sequences should provide insights into the importance
23 of recombination in CBSV viral evolution.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44

Potential for interactions between CBSVs

45
46 There is potential for CBSV and UCBSV isolates to interact as RT-PCR
47 has revealed that mixed infections are common, making up 34% - 50% of
48 tested infections in Kenya (Kathurima *et al.* 2016), Tanzania (Mbanzibwa
49 *et al.* 2011b) and Uganda (Ogwok *et al.* 2014). The potential interactions
50 between the two viral species are not currently understood. Two of the
51 CMD causal viruses: *African cassava mosaic virus* (ACMV) and *East*
52
53
54
55
56
57
58
59
60

1 *African cassava mosaic virus* (EACMV) were shown to interact
2 synergistically, leading to increased viral titers (Vanitharani *et al.* 2004). It
3 is therefore possible that similar synergistic interactions occur between
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Breeding for CBSD resistance

To date, there is no cassava cultivar with a high level of CBSD resistance available to farmers (Abaca *et al.* 2013). Breeding cassava is notoriously difficult due to high heterozygosity and a challenging cross-pollination process (Ceballos *et al.* 2012). Breeding is further complicated by cultivars showing variation in CBSD resistance across different environments, which necessitates the testing of cultivars in different agro-ecological zones to ensure their resistance is stable (Tumuhimbise *et al.* 2014).

Breeders and farmers across Tanzania, Kenya, Uganda and Malawi have been selecting cultivars which strongly express foliar symptoms but develop low levels of storage root necrosis (Hillocks *et al.* 2016). Twenty five best bet clones from five countries across East and Southern African were selected, virus-cleaned, shared and regionally evaluated across diverse environments for sources of CBSD and CMD resistance under the 5CP project (IITA 2014b). Breeding efforts also include a seven-year evaluation process of Tanzanian and Ugandan germplasm, whereby extensive intra-specific hybridizations have generated tolerant clones which develop relatively low levels of root necrosis of 12% compared to >80% in sensitive cultivars (Kawuki *et al.* 2016).

1 Although tolerant cultivars develop reduced symptoms, they remain
2 susceptible to CBSV viruses and thereby their adoption does not remove
3 viral inocula from the field. Therefore considerable efforts have been
4 made to screen and breed cassava cultivars for CBSV resistance, which
5 are able to restrict CBSV viral replication and/or movement. Promisingly,
6 protoplast studies have recently shown that the elite breeding line
7 KBH2006/18 line can inhibit CBSV viral replication, which offers exciting
8 opportunities to characterize resistance and resistance-breaking viral
9 virulence factors (Anjanappa *et al.* 2016).
10
11
12
13
14
15
16
17
18
19

20 21 **Responses of different cassava cultivars to CBSV**

22
23
24
25 Cassava cultivars respond very differently to infection by CBSVs; they
26 produce a range of symptoms and are associated with varying viral loads
27 at different time points of infection (Kaweesi *et al.* 2014). Sensitive
28 cultivars show severe shoot and root symptoms, whereas cultivars with
29 higher tolerance tend to express foliar symptoms but usually lack or
30 exhibit mild root necrosis (Hillocks & Jennings 2003). Cultivars such as
31 NASE 3 show high levels of resistance to UCBSV infection but remain
32 susceptible to CBSV (Ogwok *et al.* 2016). It has been shown that
33 cultivars, such as “Namikonga” support lower viral titers than susceptible
34 cultivars, such as Albert (Maruthi *et al.* 2014). However, symptom
35 severity is not always correlated with viral load, as the cultivar NASE 1
36 supports a relatively high viral load but produces no foliar or root necrosis
37 symptoms, whilst the cultivar NASE 14 supports a low viral load but
38 expresses severe root necrosis (Kaweesi *et al.* 2014).
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 This disparity between viral titers and symptom development has
2 necessitated the use of viral load quantification during breeding to
3 identify and select cultivars, which support low CBSD viral titers. Until
4 recently, quantification of CBSD viruses in cassava has been based on
5 quantitative RT-PCR, which measures the abundance of viral transcripts
6 relative to the abundance of plant reference gene transcripts (Kaweesi *et*
7 *al.* 2014; Moreno *et al.* 2011; Abarshi *et al.* 2012; Ogwok *et al.* 2014).
8 However, the expression of plant reference genes can vary in different
9 plant tissues, under varying developmental and environmental conditions
10 (Brunner & Yakovlev 2004) and during viral infection (Liu *et al.* 2012). To
11 overcome this, Shirima *et al.* (2017) have recently adapted the qRT-PCR
12 technique to enable absolute quantification of CBSV mRNA without
13 normalization to plant reference genes. The higher levels of accuracy
14 offered by this technique should be valuable in breeding efforts to
15 generate cassava cultivars, which support very low CBSD viral loads.
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34

35 **Identification of CBSD tolerance markers in cassava genomes**

36
37
38 Despite the importance of cassava in developing countries it has
39 received relatively little scientific attention when compared to maize, rice
40 and wheat (Varshney *et al.* 2012). Genomic studies of cassava are now
41 enabling the identification of genetic markers associated with tolerance
42 within the genomes of tolerant cultivars. In 2009 the first cassava
43 genome assembly and annotation was publicly released (Prochnik *et al.*
44 2012). Since then a large linkage map has been built using simple
45 sequence repeats (SSR) and single nucleotide polymorphisms (SNPs) to
46 identify quantitative trait markers associated with CBSD tolerance across
47 diverse African farmer-preferred cultivars (Prochnik *et al.* 2012; Patil *et*
48
49
50
51
52
53
54
55
56
57
58
59
60

1 *al.* 2015). This has revealed a number of putative CBSD tolerance alleles
2
3 across different genetic clusters in different cassava genotypes (Abaca *et*
4
5 *al.* 2013; Ferguson 2015). If validated these alleles will be useful as
6
7 markers in marker-assisted breeding and combined into cultivars for
8
9 effective and durable CBSD tolerance (Pariyo *et al.* 2013).
10

11 **Transcriptional responses to CBSD viruses in different cassava** 12 **cultivars**

13
14
15
16
17
18 To date very little is known about the function of these putative CBSD
19
20 tolerance alleles. RNA-sequencing analysis of transcripts, which are
21
22 overexpressed during CBSD infection of the tolerant cultivar
23
24 "Namikonga" has implicated NAC transcription factors, as well as genes
25
26 involved with jasmonic acid hormone signaling and the biosynthesis of
27
28 phenylpropanoid, terpenoid and steroid secondary metabolites (Maruthi
29
30 *et al.* 2014). In other plants, jasmonic acid and secondary metabolites are
31
32 linked to abiotic and biotic stress responses (Wasternack & Hause 2013;
33
34 Petrusa *et al.* 2013; Izbiańska *et al.* 2014).
35
36
37

38
39 Transcriptional studies are also helping to gain understanding into the
40
41 mechanisms behind these different interactions between cassava
42
43 cultivars and different CBSD viruses. Ogwok *et al.* (2016) recently
44
45 demonstrated that Dicer like proteins (DCL): 2 and 4 and
46
47 Argonaute (AGO) 2 are differentially expressed during CBSV and
48
49 UCBSV infections in different cassava cultivars. DCL and AGO proteins
50
51 are integral to the plant antiviral defense mechanism of silencing viral
52
53 RNA (Llave 2010). Further studies are required to gain a fuller
54
55 understanding how genes involved with host silencing of viral RNA are
56
57
58
59
60

1 differentially expressed in different cultivars in response to different
2 CBSVs.
3

4
5
6 Transcriptome analysis has also revealed that *beta-1,3-glucanase*, which
7 is involved with callose degradation at plasmodesmata is up-regulated
8 during CBSD infection of the susceptible cultivar 60444 but not in the
9 elite breeding line KBH2006/18, which shows relatively high levels of
10 CBSD resistance (Anjanappa *et al.* 2017). The degradation of callose at
11 plasmodesmata has previously been shown to promote viral movement
12 (Zavaliev *et al.* 2011). Anjanappa *et al.* (2017) suggest that enhanced
13 callose degradation at plasmodesmata during CBSD infection of 60444
14 may promote viral movement, whereas the higher amount of callose
15 present at plasmodesmata during KBH2006/18 infection is sufficient to
16 limit systemic viral movement and thereby restrict infection.
17
18
19
20
21
22
23
24
25
26
27
28
29
30

31 **CBSD resistance through genetic transformation**

32
33 There are promising attempts to introduce CBSD resistance into cassava
34 through genetic engineering. The mechanism utilized involves the
35 transgenic expression of inverted repeat CBSD viral sequences to trigger
36 post-transcriptional gene silencing (PTGS) of the corresponding
37 sequences during infection and hence confer viral resistance to the plant
38 (Patil *et al.* 2011). The approach was successful in *N. benthamiana*;
39 transgenic expression of UCBSV coat protein hairpin constructs resulted
40 in high levels of resistance to six diverse CBSV and UCBSV isolates
41 (Patil *et al.* 2011). The same construct was expressed in cassava and
42 conferred resistance to CBSV and UCBSV under field conditions with
43 high disease pressure (Yadav *et al.* 2011; Ogwok *et al.* 2012). Vegetative
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 stem cuttings taken from transgenic plants retained CBSD resistance,
2 enabling their use in vegetative propagation (Odipio *et al.* 2013).

3
4
5 To ensure that transgenic plants are resistant to both CBSV and UCBSV
6 viruses, the cultivar TME 204 was transformed with a construct (p5001)
7 containing fused tandem repeat coat protein sequences from both CBSV
8 and UCBSV to produce the transgenic line: TME 204 p5001 (Beyene *et al.*
9 2017). This transgenic line was resistant to CBSD when graft
10 challenged (Beyene *et al.* 2017) and grown within confined field trials
11 across different agro-ecological locations in Uganda and Kenya, where
12 plants were exposed to a range of both CBSV and UCBSV isolates over
13 multiple vegetative propagation cycles (Wagaba *et al.* 2017).

14
15
16 It is vitally important that improved cultivars are resistant to both CBSD
17 and CMD. Transgenic CBSD resistance was conferred to cultivars TME 7
18 and TME 204, which are naturally CMD resistant due to the presence of
19 the single dominant CMD2 resistance locus (Vanderschuren *et al.* 2012;
20 Beyene *et al.* 2016). Critically however, these TME cultivars lost their
21 CMD2 resistance through an unknown mechanism during somatic
22 embryogenesis (Beyene *et al.* 2016). Work is ongoing to cross the CBSD
23 resistant transgenic line TME 204 p5001 with a wild type CMD2 type
24 cultivar to combine durable CBSD and CMD resistance into a single
25 cultivar (Beyene *et al.* 2017).

26
27
28 Once biosafety issues have been addressed, the potential benefits of GM
29 cassava to smallholder famers are substantial. It was estimated that net
30 value for the release of CBSD resistant cultivars would be US\$436
31 million for western Kenya and US\$790 million for Uganda over 35 year
32 period starting in 2025 (Taylor *et al.* 2016). The Virus Resistant Cassava
33 for Africa (VIRCA Plus) project is working to deliver CMD and CBSD

1 resistant cassava cultivars to smallholder farmers in Uganda and Kenya
2
3 and so improve their livelihoods and food security (Taylor *et al.* 2016).
4
5
6

7 **Distribution of certified virus-clean planting material**

8
9 The lack of cultivars highly resistant to CBSD makes the existence of
10 clean seed system critical for effective management of CBSD. Clean
11 cassava seed systems are non-existent in most eastern Africa countries
12 where CBSD is a problem. The Great Lakes Cassava Initiative was
13 launched in 2008 with an overall goal to distribute certified virus-clean
14 CBSD tolerant cultivars to 1.15 million farmers across six East and
15 Central African countries over a four year period (Catholic Relief Services
16 2010). As tolerant cultivars still retain viruses within their stems, planting
17 material must be subjected to a cleaning process and highly sensitive
18 diagnostic testing before it can be multiplied and supplied to farmers.
19 This should reduce disease pressure in affected areas, as at least initially
20 the majority of crops will be disease-free (Mwangangi 2014). The
21 production of certified virus-clean cassava germplasm is particularly
22 important during the transportation of vegetative planting material due to
23 the risks CBSD poses to cassava growing areas which are currently
24 unaffected (Legg *et al.* 2011). The cleaning process involves culturing
25 meristem tissue *in vitro*, and subjecting it to thermo and/or
26 chemotherapy, which inactivates viruses and prevents viral replication or
27 movement within tissues.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51 Mathematical modeling has shown that in order for the clean seed
52 system to be sustainable, multiplication sites should only be set up in
53 areas with low-disease pressure and low vector population density
54 (McQuaid *et al.* 2015). Modeling has also shown that to reduce CBSD
55
56
57
58
59
60

1 dispersal and increase cassava yields, virus-free planting material should
2 be distributed to a number of different growers across a widespread area
3 with restricted trade (McQuaid *et al.* 2017). Once certified virus-clean
4 material has been distributed, farmers must also be thoroughly trained in
5 the identification of disease symptoms to enable sufficient roguing to
6 further reduce CBSD spread (McQuaid *et al.* 2015; Legg *et al.* 2017).
7 Cassava clean seed system projects have recently been piloted in
8 Uganda, Tanzania and Nigeria. It is hoped that similar systems will be
9 established and effective across other African countries, including Kenya
10 (CSS Report 2016).
11
12
13
14
15
16
17
18
19
20
21
22
23
24

25 **CBSVs diagnostics**

26
27 As many CBSD infected plants remain symptomless, highly sensitive
28 diagnostic techniques are required in the production and transportation of
29 material (Abarshi *et al.* 2010). There have been several important
30 advancements in cassava disease diagnostic techniques, including the
31 optimization of RT-PCR to enable reliable simultaneous detection of
32 CBSV and UCBSV (Mbanzibwa *et al.* 2011b), as well as cassava mosaic
33 begomoviruses in a single multiplex RT-PCR reaction (Abarshi *et al.*
34 2012b). Next generation high through-put sequencing (NGS) has been
35 used to screen large numbers of plants for the presence of CBSVs to
36 ensure it is virus-free before dissemination as planting material. Adams
37 *et al.* (2013) demonstrated that with NGS it was possible to detect 1%
38 infected plants out of a total of 300 plants with 95% probability. Although
39 useful tools, to date many of these techniques are relatively resource
40 intensive and so it is vitally important that affordable diagnostic tools are
41 available in African countries to enable sensitive CBSD detection locally,
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 even in cassava fields. One promising technique is reverse transcription
2 loop-mediated isothermal amplification (RT-LAMP), which is able to
3 detect and differentiate the presence of CBSV and UCBSV viruses with
4 lower consumables, resources and instrument costs than RT-PCR
5
6
7
8
9
10 (Tomlinson *et al.* 2013).

11 **Conclusions**

12
13
14
15
16
17 In the past 20 years, CBSD has become a major cause of food insecurity
18 across East and Central Africa and only since its recent geographic
19 expansion has the disease received the scientific interest it deserves.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Once the CBSD pandemic unraveled, it was largely too late to restrict the
disease to limited outbreak areas. Lessons must be learnt from this to
prevent similar disease outbreaks in the future. Critically, that scientific
interest should be applied to predicting and preventing future outbreaks
before they are able to emerge and cause devastating yield losses
across large areas.

In terms of understanding CBSD, recent studies have begun to show that
CBSVs are diverse and that CBSV has a high evolutionary capacity
(Alicai *et al.* 2016). Many control efforts are being aided by advancing
molecular techniques, including marker-assisted breeding, development
of genetically modified resistant lines, provision of certified virus-clean
planting material and the use of sensitive diagnostics. Despite this
progress, there are still many areas of CBSD biology and epidemiology,
which remain poorly understood and offer opportunities to further
understand and control the disease.

Future prospects

Understand key drivers in CBSD epidemiology

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Relatively little is known about the complex interactions between viral variants, vectors, cassava cultivars and environmental conditions and how they may be influencing the spread of CBSD. Therefore CBSD incidence, prevalence and whitefly populations in farmers' fields need to be regularly monitored in major cassava producing areas to track periodic changes in the general status of the disease in affected countries and those at risk. Where control interventions are deployed they should be evaluated for their impact in controlling CBSD. Availability of the above information is required in development of predictive models that will provide an evidence base for disease control decisions and resource allocation. The effectiveness of CBSD control strategies also heavily depends on the level of farmer engagement and awareness. In Uganda, extension work includes efforts to raise farmer awareness of CBSD and deliver information on its management (Kumakech *et al.* 2013).

Gain insights into viral populations

We currently know very little about viral populations within wild hosts, which may serve as important sources of viral inoculum and enable the evolution of CBSD and other emerging viral diseases. Next generation deep sequencing can be used to detect viral populations of which very little sequence information is known (Prabha *et al.* 2013). It would be fascinating to apply this to cassava and characterize viral populations within CBSD infected cassava and wild hosts surrounding cassava crops.

1 This could shed light on viral evolution and the contribution of wild hosts
2 in epidemiology. It may also help to identify potential unknown viral
3 diseases, against which preemptive control could be taken in anticipation
4 of emerging diseases (Newbery *et al.* 2016).
5
6
7
8

9 10 11 **Measures to restrict CBSD spread into unaffected areas**

12
13
14 To date, CBSD viruses are only found in East and Central Africa.
15 However, CBSD distribution could increase should infected material be
16 transported to other cassava growing areas of Africa, Latin America and
17 Asia, which would result in huge economic losses and food insecurity (J.
18 Legg *et al.* 2014b). Therefore movement of cassava material from CBSD
19 affected countries should be subject to strict quarantine measures to
20 ensure that planting material is virus-free before transportation. Such
21 measures will facilitate movement of superior cultivars for production or
22 breeding purposes.
23
24
25
26
27
28
29
30
31
32

33 34 35 36 **Utilize diverse cultivars for genomic resources**

37
38 It is important to continue to maintain and investigate diverse cassava
39 germplasm from across Africa and Latin America and their wild relatives
40 for potential sources of disease resistance and other beneficial
41 agronomic traits (Turyagyenda *et al.* 2012). This will enable farmers to
42 adapt to changing environmental, socio-cultural and market conditions
43 (Pautasso *et al.* 2013).
44
45
46
47
48
49
50

51 52 53 54 **Surveillance of viral diseases**

55
56 To target control efforts, it is vitally important to accurately survey viral
57 disease distribution. The IITA has recently launched the Cassava
58
59
60

1 Disease Surveillance platform in Nigeria, which offers opportunities for
2 cassava breeders and extension workers to upload images of plants
3 suspected to be infected with CBSD and other diseases. The images are
4 analysed by a team of experts to enable rapid diagnosis and coordination
5 of emergency control responses (IITA 2016). Similarly the West African
6 Virus Epidemiology project launched in 2015 aims to use field surveys to
7 gain a clear understanding of the viruses which affect cassava in West
8 Africa to predict viral emergence and inform policy decisions. Structured
9 surveys under the Cassava Virus Diagnostics project in eastern and
10 southern Africa are tracking area-wide changes in cassava viral diseases
11 over time. This will provide the basis for disease control intervention
12 decision-making and impact assessment.
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

29 **Predicted affects of climate change on cassava production**

30 Cassava demonstrates relatively high levels of resilience to temperature
31 and rainfall fluctuations predicted in climate change models (El-Sharkawy
32 2004). A model based on temperature and rainfall projections across
33 Africa has predicted that compared to other staple food crops, overall
34 cassava is the least likely to be adversely affected by climate change
35 (Jarvis *et al.* 2012). This makes cassava an attractive food security crop
36 for climate change adaptation in Africa. However, climate change is also
37 predicted to affect the distribution and abundance of cassava pests and
38 diseases, including *B. tabaci* (Jarvis *et al.* 2012). Recent ecological niche
39 modeling has predicted that with climate change, the potential distribution
40 of CBSD and CMD carrying *B. tabaci* will spread over West, Central and
41 the south-western coast of Africa where cassava production is high and
42 CBSD is currently absent (Herrera Campo *et al.* 2011). Therefore
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 monitoring and controlling *B. tabaci* populations is a major priority. The
2 deep sequencing technique could be extended to *B. tabaci*, enabling the
3 mapping of the most active and abundant viral species carried by *B.*
4 *tabaci* populations across different agricultural regions (Ng *et al.* 2011).
5
6
7
8
9

10 **Understanding CBSVs infection mechanisms and virulence** 11 **determinants**

12 Despite the increasing number of sequenced, CBSVs genomes, there is
13 very little is known about virulence determinants within CBSV and
14 UCBSV genomes responsible for key functions during infection and their
15 effect on disease symptomatology. To date, only the silencing
16 suppression activity of the UCBSV P1 protein has been characterized
17 (Mbanzibwa *et al.* 2009a). The construction of infectious clones will
18 enable the targeted mutagenesis of key viral sequences to identify the
19 functions of viral proteins and the host proteins they interact with, which
20 should serve as potential targets to restrict viral infection. Current work to
21 develop and manipulate CBSVs infectious clones is ongoing at various
22 institutions.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42

43 **Collaborative sharing of information and resources**

44 There are many opportunities to exploit recent progress made in
45 understanding CBSD through progress in cassava, viral and vector
46 research. There is a need for this research to be integrated into a central,
47 easily accessible platform (Ayling *et al.* 2012). This will require experts
48 across diverse backgrounds and countries openly communicate, engage,
49 share data and collaborate through networks such the Global Cassava
50 Partnership for the 21st Century. Such partnerships should help to
51
52
53
54
55
56
57
58
59
60

1 generate solutions to controlling CBSD and enable cassava to fulfill its
2
3 potential of feeding billions of people by 2050 (Legg *et al.* 2014).
4
5
6

7 **Acknowledgements**

8
9 This work was made possible by the South West Training Partnership,
10
11 funded by Biotechnology and Biological Sciences Research Council, UK.
12

13
14 All photos were taken by Katie Tomlinson.
15
16
17

18 **Conflict of interest**

19
20 The authors declare that they have no conflicts of interest with the
21
22 contents of this article.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- 1
2
3 Abaca, A. *et al.*, 2013. Genetic relationships of cassava genotypes that are
4 susceptible or tolerant to Cassava brown streak disease in Uganda.
5 *Journal of Agricultural Science*, 5(7).
6
- 7 Abarshi, M.M. *et al.*, 2012. Multiplex RT-PCR assays for the simultaneous
8 detection of both RNA and DNA viruses infecting cassava and the
9 common occurrence of mixed infections by two cassava viruses in East
10 Africa. *Journal of Virological Methods*, 179(1), pp.176–184.
11
- 12 Abarshi, M.M. *et al.*, 2010. Optimization of diagnostic RT-PCR protocols and
13 sampling procedures for the reliable and cost-effective detection of
14 *Cassava brown streak virus*. *Journal of Virological Methods*, 163(2),
15 pp.353–9.
16
- 17 Adams, I.P. *et al.*, 2013. High throughput real-time RT-PCR assays for
18 specific detection of Cassava brown streak disease causal viruses, and
19 their application to testing of planting material. *Plant Pathology*, 62(1),
20 pp.233–242.
21
- 22 Alabi, O., Kumar, P. & Naidu, R., 2011. Cassava mosaic disease: a curse to
23 food security in sub-Saharan Africa. *APSnet Features*,
24 doi:10.1094/APSnetFeature-2011-0701.
25
- 26 Alicai, T. *et al.*, 2016. Characterization by next generation sequencing reveals
27 the molecular mechanisms driving the faster evolutionary rate of
28 *Cassava brown streak virus* compared with *Ugandan cassava brown*
29 *streak virus*. *Scientific Reports*, 6, p.36164.
30
- 31 Alicai, T. *et al.*, 2007. Re-emergence of Cassava brown streak disease in
32 Uganda. *Plant Disease*, 91(1), pp.24–29.
33
- 34 Anjanappa, R.B. *et al.*, 2017. Early transcriptome analysis of the brown streak
35 virus-cassava pathosystem provides molecular insights into virus
36 susceptibility and resistance. *BioRxiv*, doi: 10.1101/100552.
37
- 38 Anjanappa, R.B. *et al.*, 2016. Characterization of virus-resistant cassava.
39 *Molecular Plant-Microbe Interactions*, 29(7), pp.527–534.
40
- 41 Ayling, S. *et al.*, 2012. Information resources for cassava research and
42 breeding. *Tropical Plant Biology*, 5(1), pp.140–151.
43
- 44 Bennett, B., 2015. Guest editorial: smallholder cassava production and the
45 cassava processing sector in Africa. *Food Chain*, 5(1–2), pp.1–3.
46
- 47 Beyene, G. *et al.*, 2017. A virus-derived stacked RNAi construct confers
48 robust resistance to Cassava brown streak disease. *Frontiers in Plant*
49 *Science*, 7, p.2052.
50
- 51 Beyene, G. *et al.*, 2016. Loss of CMD2-mediated resistance to Cassava
52 mosaic disease in plants regenerated through somatic embryogenesis.
53 *Molecular Plant Pathology*, 17(7), pp.1095–110.
54
- 55 Bigirimana, S., Barumbanze, P. & Ndayihanzamaso, P., 2011. First report of
56 Cassava brown streak disease and associated *Ugandan cassava brown*
57 *streak virus* in Burundi. *New Disease Reports*, 24(26).
58
- 59 Bock, K., 1994. Studies on *Cassava brown streak virus* disease in Kenya.
60 *Tropical Science*, 34(1), pp.134–145.

- 1 Bock, K.R. & Guthrie, E.J., 1976. Recent advances in research on cassava
2 viruses in East Africa. *East African Agricultural And Forestry Journal*, pp.
3 11-16.
- 4 Brunner, A. & Yakovlev, I., 2004. Validating internal controls for quantitative
5 plant gene expression studies. *BMC plant*, 4:14.
- 6
7 Catholic Relief Services, 2010. Final Report on the Great Lakes Cassava
8 Initiative.
- 9
10 Ceballos, H., Kulakow, P. & Hershey, C., 2012. Cassava breeding: current
11 status, bottlenecks and the potential of biotechnology tools. *Tropical*
12 *Plant Biology*, 5(1), pp.73–87.
- 13
14 Clean Seed System Report, 2016. Commercializing clean cassava planting
15 material delivery system in Uganda. The second annual review and
16 planning meeting held at the National Crops Resources Research
17 Insitute: 28th - 30th June 2016.
- 18
19 El-Sharkawy, M.A., 2004. Cassava biology and physiology. *Plant Molecular*
20 *Biology*, 56(4), pp.481–501.
- 21
22 FAO, 2013. *Save and Grow Cassava: a guide to sustainable production*
23 *intensification*. Rome: Food and Agriculture Organization of the United
24 Nations.
- 25
26 FAO, 2011. *Cassava virus on verge of epidemic in East Africa: experts urge*
27 *funding, swift action to protect staple food crop*. Rome: Food and
28 Agricultural Organisation.
- 29
30 Ferguson, M., 2015. QTL Mapping for Resistance to Cassava Viruses in
31 Tanzania. *Conference Abstract: Plant and Animal Genome XXIII*.
- 32
33 Fermont, A.M. *et al.*, 2009. Closing the cassava yield gap: An analysis from
34 smallholder farms in East Africa. *Field Crops Research*, 112(1), pp.24–
35 36.
- 36
37 Galperin, M.Y. *et al.*, 2006. House cleaning, a part of good housekeeping.
38 *Molecular Microbiology*, 59(1), pp.5–19.
- 39
40 Herrera Campo, B.V., Hyman, G. & Bellotti, A., 2011. Threats to cassava
41 production: known and potential geographic distribution of four key biotic
42 constraints. *Food Security*, 3(3), pp.329–345.
- 43
44 Hillocks, R. *et al.*, 2016. Disparity between leaf and root symptoms and crop
45 losses associated with Cassava brown streak disease in four countries in
46 eastern Africa. *Journal of Phytopathology*, 164(2), pp.86–93.
- 47
48 Hillocks, R.J. & Maruthi, M.N., 2015. Post-harvest impact of Cassava brown
49 streak disease in four countries in eastern Africa. *Food Chain*, 5(1–2),
50 pp.116–122.
- 51
52 Hillocks, R. & Jennings, D., 2003. Cassava brown streak disease: A review of
53 present knowledge and research needs. *International Journal of Pest*
54 *Management*, 49(3), pp.225–234.
- 55
56 Hillocks, R.J. *et al.*, 2002. Cassava brown streak disease in northern
57 Mozambique. *International Journal of Pest Management*, 48(3), pp.178–
58 181.
- 59
60 Hillocks, R.J. & Thresh, J.M. eds., 2002. *Cassava: biology, production and*

- 1 *utilization*, Wallingford: CABI.
- 2 Hillocks, R.J. *et al.*, 2001. Effects of brown streak virus disease on yield and
3 quality of cassava in Tanzania. *Journal of Phytopathology*, 149(7–8),
4 pp.389–394.
- 5
6 Hillocks, R., J., Raya, M. & Thresh, J., M., 1999. Factors affecting the
7 distribution, spread and symptom expression of Cassava brown streak
8 disease in Tanzania. *International Journal of Pest Management*, 42,
9 pp.285–289.
- 10
11 Hillocks, R.J., Raya, M. & Thresh, J.M., 1996. The association between root
12 necrosis and above-ground symptoms of brown streak virus infection of
13 cassava in southern Tanzania. *International Journal of Pest*
14 *Management*, 42(4), pp.285–289.
- 15
16 IITA, 2016. IITA Bulletin, Issue 2324. IITA and NAQS pilot digital Cassava
17 Disease Surveillance platform.
- 18
19 IITA, 2014a. IITA Bulletin, Issue 2215. Nigeria: International Institute of
20 Tropical Agriculture.
- 21
22 IITA, 2014b. IITA Report. 5CP Updates: Highlights of the progress of the new
23 cassava varieties and clean seed to combat CBSD and CMD project
24 (5CP).
- 25
26 Izbiańska, K., Arasimowicz-Jelonek, M. & Deckert, J., 2014. Phenylpropanoid
27 pathway metabolites promote tolerance response of lupine roots to lead
28 stress. *Ecotoxicology and Environmental Safety*, 110, pp.61–67.
- 29
30 Jameson, J.D., 1964. Cassava mosaic disease in Uganda. *East African*
31 *Agricultural and Forestry Journal*, 29(3).
- 32
33 Jarvis, A. *et al.*, 2012. Is cassava the answer to African climate change
34 adaptation? *Tropical Plant Biology*, 5(1), pp.9–29.
- 35
36 Jennings, D.L., 1960. Observations on virus diseases of cassava in resistant
37 and susceptible varieties. II. Brown streak disease. *Empire Journal of*
38 *Experimental Agriculture*, 28(111), pp.261–270.
- 39
40 Jennings, D.L., 1957. Further studies in breeding cassava for virus
41 resistance. *The East African Agricultural Journal*, 22(4), pp.213–219.
- 42
43 Jeremiah, S.C. *et al.*, 2015. The dynamics and environmental influence on
44 interactions between Cassava brown streak disease and the whitefly,
45 *Bemisia tabaci*. *Phytopathology*, 105(5), pp.646–655.
- 46
47 Jones, W., 1959. *Manioc in Africa*, Standford University Press.
- 48
49 Kathurima, T. *et al.*, 2016. Genetic diversity and distribution of *Cassava*
50 *brown streak virus* and *Ugandan cassava brown streak virus* in major
51 cassava-growing regions in Kenya. *Annual Research & Review in*
52 *Biology*, 10(5), pp.1–9.
- 53
54 Katono, K. *et al.*, 2015. Influence of host plant resistance and disease
55 pressure on spread of Cassava brown streak disease in Uganda.
56 *American Journal of Experimental Agriculture*, 7(5), pp.284–293.
- 57
58 Kaweesi, T. *et al.*, 2014. Field evaluation of selected cassava genotypes for
59 Cassava brown streak disease based on symptom expression and virus
60 load. *Virology Journal*, 11(1), p.216.
- 61
62 Kawuki, R.S. *et al.*, 2016. Eleven years of breeding efforts to combat Cassava
63 brown streak disease. *Breeding Science*, 66(4), pp.560–571.

- 1 Knierim, D., Menzel, W. & Winter, S., 2016. Analysis of the complete genome
2 sequence of *Euphorbia ringspot virus*, an atypical member of the genus
3 *Potyvirus*. *Archives of Virology*, pp.1–3.
- 4 Kumakech, A. *et al.*, 2013. The case of cassava brown streak disease
5 awareness in Northern Uganda. *African Journal of Plant Science*
6 *Knowledge on cassava disease management*, 7(12), pp.597–601.
- 7
8 Llave, C. 2010. Virus-derived small interfering RNAs at the core of plant–virus
9 interactions. *Trends in Plant Science*, 15(12), 701–707.
- 10
11 Legg, J. *et al.*, 2017. Community phytosanitation to manage cassava brown
12 streak disease. *Virus Research*.
- 13
14 Legg, J. & Kanju, E., 2015. *Virus Diseases of Tropical and Subtropical Crops*,
15 CABI Plant Protection Series.
- 16
17 Legg, J. *et al.*, 2014. A global alliance declaring war on cassava viruses in
18 Africa. *Food Security*, 6(2), pp.231–248.
- 19
20 Legg, J.P. *et al.*, 2011. Comparing the regional epidemiology of the Cassava
21 mosaic and *Cassava brown streak virus* pandemics in Africa. *Virus*
22 *Research*, 159(2), pp.161–70.
- 23
24 Legg, J.P. *et al.*, 2006. Cassava Mosaic Virus Disease in East and Central
25 Africa: Epidemiology and Management of A Regional Pandemic.
26 *Advances in Virus Research*, 67, pp. 355–418.
- 27
28 Legg, J. P., & Thresh, J. M., 2000. Cassava mosaic virus disease in East
29 Africa: a dynamic disease in a changing environment. *Virus Research*,
30 71(1), 135–149.
- 31
32 Legg, J.P. & Raya, M.D., 1998. Survey of cassava virus diseases in
33 Tanzania. *International Journal of Pest Management*, 44(1), pp.17–23.
- 34
35 Lennon, A., Aiton, M. & Harrison, B., 1985. Cassava viruses from Africa: third
36 country quarantine. *Report for 1985. Scottish Crop Research Institute*,
37 *Invergowrie, UK, pp 168–169.*
- 38
39 Lister, R.M., 1959. Mechanical transmission of *Cassava brown streak virus*.
40 *Nature*, 183(4675), pp.1588–1589.
- 41
42 Liu, D. *et al.*, 2012. Validation of reference genes for gene expression studies
43 in virus-infected *Nicotiana benthamiana* using quantitative real-time PCR.
44 *PloS one*, 7(9), p.e46451.
- 45
46 Maruthi, M.N. *et al.*, 2016. Virus-vector relationships and the role of whiteflies,
47 *Bemisia tabaci*, and farmer practices in the spread of *Cassava brown*
48 *streak viruses*. *Phytopathology*, ISSN 0031.
- 49
50 Maruthi, M.N. *et al.*, 2014. Transcriptional response of virus-infected cassava
51 and identification of putative sources of resistance for Cassava brown
52 streak disease. *PLOS ONE*, 9(5), p.e96642.
- 53
54 Maruthi, M.N. *et al.*, 2005. Transmission of *Cassava brown streak virus* by
55 *Bemisia tabaci* (*Gennadius*). *Journal of Phytopathology*, 153(5), pp.307–
56 312.
- 57
58 Mbanzibwa, D.R. *et al.*, 2011a. Evolution of Cassava brown streak disease-
59 associated viruses. *The Journal of General Virology*, 92(Pt 4), pp.974–
60 87.
- 61
62 Mbanzibwa, D.R. *et al.*, 2011b. Simultaneous virus-specific detection of the
63 two Cassava brown streak-associated viruses by RT-PCR reveals wide

- 1 distribution in East Africa, mixed infections, and infections in *Manihot*
2 *glaziovii*. *Journal of Virological Methods*, 171(2), pp.394–400.
- 3 Mbanzibwa, D.R. *et al.*, 2009a. *Cassava brown streak virus (Potyviridae)*
4 encodes a putative Maf/HAM1 pyrophosphatase implicated in reduction
5 of mutations and a P1 proteinase that suppresses RNA silencing but
6 contains no HC-Pro. *Journal of Virology*, 83(13), pp.6934–40.
- 7
8 Mbanzibwa, D.R. *et al.*, 2009b. Genetically distinct strains of *Cassava brown*
9 *streak virus* in the Lake Victoria basin and the Indian Ocean coastal area
10 of East Africa. *Archives of Virology*, 154(2), pp.353–359.
- 11 McQuaid, C.F. *et al.*, 2017. Spatial dynamics and control of a crop pathogen
12 with mixed-mode transmission. *PLOS Computational Biology*, 13(7),
13 p.e1005654.
- 14
15
16 McQuaid, C.F. *et al.*, 2015. Cassava brown streak disease and the
17 sustainability of a clean seed system. *Plant Pathology*, 65(2), pp.299–
18 309.
- 19
20 Mohammed, I.U. *et al.*, 2012. The symptom and genetic diversity of *Cassava*
21 *brown streak viruses* infecting cassava in East Africa. *Advances in*
22 *Virology*, 2012, p.795697.
- 23
24 Monger, W.A. *et al.*, 2010. The complete genome sequence of the Tanzanian
25 strain of *Cassava brown streak virus* and comparison with the Ugandan
26 strain sequence. *Archives of Virology*, 155(3), pp.429–433.
- 27
28 Monger, W.A. *et al.*, 2001a. Molecular characterization of the *Cassava brown*
29 *streak virus* coat protein. *Plant Pathology*, 50(4), pp.527–534.
- 30
31 Monger, *et al.*, 2001b. Identification of different isolates of *Cassava brown*
32 *streak virus* and development of a diagnostic test. *Plant Pathology*, 50(6),
33 pp.768–775.
- 34
35 Moreno, I., Gruissem, W. & Vanderschuren, H., 2011. Reference genes for
36 reliable potyvirus quantitation in cassava and analysis of *Cassava brown*
37 *streak virus* load in host varieties. *Journal of virological methods*, 177(1),
38 pp.49–54.
- 39
40 Mulimbi, W. *et al.*, 2012. First report of *Ugandan cassava brown streak virus*
41 on cassava in Democratic Republic of Congo. *New Disease Reports*, 26,
42 p.11.
- 43
44 Mware, B. O. *et al.*, 2009. Transmission and distribution of cassava
45 brown streak virus disease in cassava growing areas of Kenya. *Journal*
46 *of Applied Biosciences*, 16: 864 - 870.
- 47
48 Mwangangi, M.M., 2014. Evaluation of meristem tip culture, chemotherapy
49 and thermochemistry on the reduction of *Cassava brown streak virus* in
50 infected cassava (Guzo variety) cuttings. *Master of Science in*
51 *Biotechnolog thesis submitted to Tomo Kenyatta University of Agriculture*
52 *and Technology*.
- 53
54 Ndunguru, J. *et al.*, 2015. Analyses of twelve new whole genome sequences
55 of *Cassava brown streak viruses* and *Ugandan cassava brown streak*
56 *viruses* from East Africa: diversity, supercomputing and evidence for
57 further speciation. *PLOS ONE*, 10(10), p.e0139321.
- 58
59 Newbery, F., Qi, A. & Fitt, B.D., 2016. Modelling impacts of climate change on
60 arable crop diseases: progress, challenges and applications. *Current*

- 1 *Opinion in Plant Biology*, 32, pp.101–109.
- 2 Ng, T.F.F. *et al.*, 2011. Exploring the diversity of plant DNA viruses and their
3 satellites using vector-enabled metagenomics on whiteflies. *PLOS ONE*,
4 6(4), p.e19050.
- 5
6 Nichols, R.F.W., 1950. The brown streak disease of cassava. *The East
7 African Agricultural Journal*, 15(3), pp.154–160.
- 8
9 Nichols, R.F.W., 1946. Breeding cassava for virus resistance. *The East
10 African Agricultural Journal*, 12(3), pp.184-194.
- 11 Ntawuruhunga, P. & Legg, J., 2007. New spread of *Cassava brown streak
12 virus* disease and its implications for the movement of cassava
13 germplasm in the East and Central African region. *Crop Crisis Control
14 C3P Project Report*.
- 15
16 Nweke, F., 2004. New challenges in the cassava transformation in Nigeria
17 and Ghana. *International Food Policy Research Institute*, 118.
- 18
19 Odipio, J. *et al.*, 2013. RNAi-derived field resistance to Cassava brown streak
20 disease persists across the vegetative cropping cycle. *GM Crops & Food*,
21 5(1), pp.16-19.
- 22
23 Ogwok, E. *et al.*, 2016. Comparative analysis of virus-derived small RNAs
24 within cassava (*Manihot esculenta* Crantz) infected with *Cassava brown
25 streak viruses*. *Virus Research*, 215, pp.1–11.
- 26
27 Ogwok, E. *et al.*, 2014. Distribution and accumulation of *Cassava brown
28 streak viruses* within infected cassava (*Manihot esculenta* Crantz) plants.
29 *Plant Pathology*, 64(5), p.12343.
- 30
31 Ogwok, E. *et al.*, 2012. Transgenic RNA interference (RNAi)-derived field
32 resistance to Cassava brown streak disease. *Molecular Plant Pathology*,
33 13(9), pp.1019–31.
- 34
35 Pariyo, A. *et al.*, 2013. Simple sequence repeat (SSR) diversity of cassava in
36 South, East and Central Africa in relation to resistance to Cassava brown
37 streak disease. *African Journal of Biotechnology*, 12(28), pp.4453–4464.
- 38
39 Patil, B.L. *et al.*, 2015. Cassava brown streak disease: a threat to food
40 security in Africa. *Journal of General Virology*, 96(Pt5), pp.956–968.
- 41
42 Patil, B.L. *et al.*, 2011. RNAi-mediated resistance to diverse isolates
43 belonging to two virus species involved in Cassava brown streak
44 disease. *Molecular Plant Pathology*, 12(1), pp.31–41.
- 45
46 Pautasso, M. *et al.*, 2013. Seed exchange networks for agrobiodiversity
47 conservation. A review. *Agronomy for Sustainable Development*, 33(1),
48 pp.151–175.
- 49
50 Pennisi, E., 2010. Armed and dangerous. *Science (New York, N.Y.)*,
51 327(5967), pp.804–5.
- 52
53 Petrusa, E. *et al.*, 2013. Plant flavonoids: biosynthesis, transport and
54 involvement in stress responses. *International journal of molecular
55 sciences*, 14(7), pp.14950–73
- 56
57 Prabha, K., Baranwal, V.K. & Jain, R.K., 2013. Applications of next generation
58 high throughput sequencing technologies in characterization, discovery
59 and molecular interaction of plant viruses. *Indian Journal of Virology*,
60 24(2), pp.157–165.

- 1 Prochnik, S.E. *et al.*, 2012. The Cassava Genome: Current Progress, Future
2 Directions. *Tropical Plant Biology*, 5(1), pp.88–94.
- 3 Roux-Cuvelier, M., Teyssedre, D. & Chesneau, T., 2014. First report of
4 Cassava brown streak disease and associated *Ugandan cassava brown*
5 *streak virus* in Mayotte Island. *New Disease*, 30(28).
- 6
7 Rwegasira, G. & Chrissie, M., 2015. Efficiency of non-vector methods of
8 *Cassava brown streak virus* transmission to susceptible cassava plants.
9 *African Journal of Food, Agriculture, Nutrition and Development*, 15(4),
10 pp.10335–10351.
- 11
12 Shirima, R.R. *et al.*, 2017. Absolute quantification of *Cassava brown streak*
13 *virus* mRNA by real-time qPCR. *Journal of Virological Methods*, 245,
14 pp.5–13.
- 15
16 Storey, H., 1939. Report of the Plant Pathologist. *Annual Report East African*
17 *Agricultural Research*.
- 18
19 Storey, H.H., 1936. Virus Diseases of East African Plants. *The East African*
20 *Agricultural Journal*, 1(4), pp.333–337.
- 21
22 Taylor, N.J. *et al.*, 2016. *Disease-resistant GM cassava in Uganda and Kenya*
23 *during a pandemic*. B4FA.
- 24
25 Thresh, J. M., & Cooter, R. J., 2005. Strategies for controlling Cassava
26 mosaic virus disease in Africa. *Plant Pathology*, 54(5), 587–614.
- 27
28 Thresh, J.M., Fargette, D. & Otim-Nape, G.W., 1994. The viruses and virus
29 diseases of cassava in Africa. *African Crop Science Journal*, 2(4),
30 pp.459–478.
- 31
32 Tomlinson, J.A. *et al.*, 2013. Loop-mediated isothermal amplification for rapid
33 detection of the causal agents of Cassava brown streak disease. *Journal*
34 *of Virological Methods*, 191(2), pp.148–54.
- 35
36 Tumuhimbise, R., Shanahan, P., Melis, R., & Kawuki, R., 2014. Combining
37 ability analysis of storage root yield and related traits in cassava at the
38 seedling evaluation stage of breeding. *Journal of Crop Improvement*,
39 28(4), 530–546.
- 40
41 Turyagyenda, L. *et al.*, 2012. Genetic diversity among farmer-preferred
42 cassava landraces in Uganda. *African Crop Science Journal*, 20(1), Issue
43 Supplement s1, pp. 15 - 30.
- 44
45 Valli, A. *et al.*, 2015. *Potyviridae*. In eLS. Chichester, UK: John Wiley & Sons,
46 Ltd, pp. 1–10.
- 47
48 Valli, A., Dujovny, G. & García, J.A., 2008. Protease activity, self interaction,
49 and small interfering RNA binding of the silencing suppressor p1b from
50 *Cucumber vein yellowing ipomovirus*. *Journal of Virology*, 82(2), pp.974–
51 86.
- 52
53 Vanderschuren, H. *et al.*, 2012. Exploiting the combination of natural and
54 genetically engineered resistance to Cassava mosaic and Cassava
55 brown streak viruses impacting cassava production in Africa. *PLOS ONE*,
56 7(9), p.e45277.
- 57
58 Vanitharani, R. *et al.*, 2004. Differential Roles of AC2 and AC4 of cassava
59 geminiviruses in mediating synergism and suppression of
60 posttranscriptional gene silencing. *Journal of Virology*, 78(17), pp.9487–

9498.

- Varshney, R.K. *et al.*, 2012. Can genomics boost productivity of orphan crops? *Nature Biotechnology*, 30(12), pp.1172–1176.
- Wagaba, H. *et al.*, 2017. Field level RNAi-mediated resistance to Cassava brown streak disease across multiple cropping cycles and diverse East African agro-ecological locations. *Frontiers in Plant Science*, 7, p.2060.
- Wasternack, C. & Hause, B., 2013. Jasmonates: biosynthesis, perception, signal transduction and action in plant stress response, growth and development. *Annals of Botany*, 6, pp.1021–1058.
- Were, H.K., Winter, S. & Maiss, E., 2004. Viruses Infecting cassava in Kenya. *Plant Disease*, 88(1), pp.17–22.
- Winter, S. *et al.*, 2010. Analysis of *Cassava brown streak viruses* reveals the presence of distinct virus species causing Cassava brown streak disease in East Africa. *The Journal of General Virology*, 91(5), pp.1365–72.
- Yadav, J.S. *et al.*, 2011. RNAi-mediated resistance to *Cassava brown streak Uganda virus* in transgenic cassava. *Molecular Plant Pathology*, 12(7), pp.677–687.
- Zavaliev, R. *et al.*, 2011. Biology of callose (β -1,3-glucan) turnover at plasmodesmata. *Protoplasma*, 248(1), pp.117–130.

Figure 1: Key events in CBSD geographical distribution and research history (1936 – 2016).

Figure 2: Cumulative number of scientific papers, reports or reviews which refer to CBSD, published in each decade between 1936 and 2016; dramatic increase in publications from the mid 1990s following the re-emergence of CBSD (accessed on Google scholar in December 2016).

Figure 3: CBSD storage root necrosis (A), radial root constrictions (B), foliar chlorosis (C) and brown streaks or lesions on stems (D).

Figure 4: Super-abundant whiteflies on cassava in Uganda.

Figure 5: CBSVs genomes encode a large polyprotein which is auto-catalytically cleaved at specific cleavage sequences by virus-encoded proteases into ten mature proteins and an additional P3N-PIPO protein is produced through a +2 ribosomal frameshift in the P3 region (Valli *et al.* 2015). CBSVs proteins are: P1

1 = serine protease/silencing suppressor, P3 = third protein, PIPO = pretty
2
3 interesting Potyviridae ORF, 6K1 and 6K2 = 6-kDa proteins, CI = cylindrical
4
5 inclusion protein, Vpg = viral genome-linked protein, NIa-Pro = main viral
6
7 protease, NIb = viral RNA dependent RNA polymerase, Ham1 = putative
8
9 pyrophosphatase, CP = coat protein. Note unusual features: presence of single
10
11 P1 protein, absence of HC-Pro and presence of novel Ham1 protein
12
13
14 (Mbanzibwa et al. 2009a).
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Proof




Figure 1: Key events in CBSV geographical distribution and research history (1936 – 2016).

254x179mm (72 x 72 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2: Cumulative number of scientific papers, reports or reviews which refer to CBDSD, published in each decade between 1936 and 2016; dramatic increase in publications from the mid 1990s following the re-emergence of CBDSD (accessed on Google scholar in December 2016).

220x116mm (72 x 72 DPI)

ooof




Figure 3: CBSD storage root necrosis (A), radial root constrictions (B), foliar chlorosis (C) and brown streaks or lesions on stems (D).

167x129mm (72 x 72 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Figure 4: Super-abundant whiteflies on cassava in Uganda.

114x88mm (72 x 72 DPI)

proof




Figure 5: CBSVs genomes encode a large polyprotein which is auto-catalytically cleaved at specific cleavage sequences by virus-encoded proteases into ten mature proteins and an additional P3N-PIPO protein is produced through a +2 ribosomal frameshift in the P3 region (Valli et al. 2015). CBSVs proteins are: P1 = serine protease/silencing suppressor, P3 = third protein, PIPO = pretty interesting Potyviridae ORF, 6K1 and 6K2 = 6-kDa proteins, CI = cylindrical inclusion protein, VPg = viral genome-linked protein, NIa-Pro = main viral protease, NIB = viral RNA dependent RNA polymerase, Ham1 = putative pyrophosphatase, CP = coat protein. Note unusual features: presence of single P1 protein, absence of HC-Pro and presence of novel Ham1 protein (Mbanzibwa et al. 2009a).

226x37mm (72 x 72 DPI)

Proof