

Article

Infrared Emotions and Behaviours: Thermal Imaging in Psychology

Topalidou, Anastasia and Ali, Nazmin

Available at <http://clock.uclan.ac.uk/20352/>

Topalidou, Anastasia and Ali, Nazmin (2017) Infrared Emotions and Behaviours: Thermal Imaging in Psychology. International Journal of Prenatal & Life Sciences, 1 (01). pp. 65-70.

It is advisable to refer to the publisher's version if you intend to cite from the work.
10.24946/IJPLS.20.17.0101.110704

For more information about UCLan's research in this area go to <http://www.uclan.ac.uk/researchgroups/> and search for <name of research Group>.

For information about Research generally at UCLan please go to <http://www.uclan.ac.uk/research/>

All outputs in CLoK are protected by Intellectual Property Rights law, including Copyright law. Copyright, IPR and Moral Rights for the works on this site are retained by the individual authors and/or other copyright owners. Terms and conditions for use of this material are defined in the <http://clock.uclan.ac.uk/policies/>



DOI: 10.24946/IJPLS.20.17.0101.110704

Vol. 01 Issue 01

www.journalprenatalife.com

Infrared Emotions and Behaviours: Thermal Imaging in Psychology

Authors:

Anastasia Topalidou¹ and Nazmin Ali¹

¹ Research in Childbirth and Health Unit, School of Community Health and Midwifery, Faculty of Health and Wellbeing, University of Central Lancashire, UK

Corresponding author:

Anastasia Topalidou

Research Associate (Thermal Imaging)

University of Central Lancashire, Faculty of Health and Wellbeing, School of Community Health and Midwifery, Research in Childbirth and Health Unit (ReaCH)

Brook Building (BB224)

PR1 2HE, Preston, UK

Tel: +44 (0) 1772 89 5199

Email: atopalidou@uclan.ac.uk

COMMENTARY

Psychology is a social science that studies behavioural changes and mental processes. Usually, the theory and diagnosis of conditions has been based on a consultation process of working with a therapist through a series of questions, exercises and techniques. When we consider, for example, social anxiety – a fear of social situations and an avoidance of normal social interactions – a therapist traditionally talks through the issues with the client and relies purely on the descriptions of the client's feelings and emotions connected to these events. When working through a treatment program, the therapist would need to rely on the client reporting improvements in their approach to triggers, based on their changing emotional response to the situations that caused them anxiety. Although, there are demonstrative physiological responses, they are usually not measured as their recording requires a special process.

The physiological response is a change in the equilibrium in the body which can be felt as an increase in heart rate, blood pressure, breathing rate and changes in Gastrointestinal (GI) tract motility. This state of arousal does not quickly fade; in fact it can take between 20 to 60 minutes for the body to return to its pre-arousal levels (Changeux, 1985; Huitt, 1999; Faizi et al, 2014; Richardson et al, 2015). These physiological changes could be further investigated, analyzed and the results of these analyses provide additional knowledge and hence improved treatment or treatment techniques.

Within the last decades, several methods have been developed and used for the detection and recording of the physiological changes and activities. However, researchers studying the neurovascular elements behind social interactions and emotions are frequently confronted with the restrictions of methods commonly used in neuroscience and psychological research. Such methods involve applying electrodes, sensors or other contact instruments on the skin of participants to measure and monitor activity of the autonomic nervous system. However, using such methods can make it difficult to distinguish between spontaneous behavior and event-related behavior, and contact of the instrument with the participant is required (Bach et al, 2010). On the other hand, non-contact methods, for example functional magnetic resonance imaging (fMRI), require the participant to remain still to minimize the induction caused by motion inside the magnetic field (Wong et al, 2011).

A potentially novel solution used in recent years is thermal imaging (TI). TI is a contact free, non-invasive and free from any contra-indications method. TI records infrared radiation that is naturally emitted by the human body based on its temperature, therefore not submitting the participant to any form of radiation. All objects above the absolute zero temperature (0 K) emit infrared radiation. With the use of a thermal camera the temperature variations can be captured and displayed as an image, allowing participants to move freely and researcher to gather data in real time (Ring and Ammer, 2000; Modest, 2013; Ioannou et al, 2014; Topalidou and Downe, 2016).

A variety of emotional changes or conditions and/or responses to a trigger elicit thermal responses (changes in temperature) in several areas of human body, such as face and hands indicating that skin temperature variation can be suggestive to emotional or behavioral changes. It has been demonstrated that a range of emotional states, such as aggression (O'Kane et al, 2004) and emotional arousal (Zajonc et al, 1989; Shearn et al, 1990; Nozawa and Tacano, 2009; Nhan and Chau, 2010), stimulate thermal responses that are indicative of a certain state. Fear, for example, is associated with an increase in blood flow to the eyes (300 ms) and a decrease in temperature of the cheeks (Pavlidis et al, 2000); Levine et al, 2001). Whilst the responses associated with stress can differ depending on the activity and age. Stress in infants showed a decrease in forehead temperature due to lower arousal levels caused by separation from the mother (Mizukami et al, 1987; Mizukami et al, 1990). On the other hand, stress in adults correlated with an increase of blood flow to the frontal region (Puri et al, 2005; Merla and Romani, 2007) and the periorbital regions (Pavlidis and Levine, 2002); this was accompanied with a temperature decrease in the region of the jaw (Merla and Romani, 2007). Merla and Romani (2007) conducted a study to investigate the thermal response of stress, pain and sexual arousal in 10 healthy males. Pain and stress were

observed to cause an overall decrease in facial temperature, particularly in the perioral region, while sexual arousal lead to increased facial perfusion rates in specific regions: the forehead, lips and nose.

Thermographic measures of these conditions were correlated with other forms of arousal, for example galvanic skin response and penile turgidity. The results suggested that specific thermal signatures related to specific emotions possibly exist due to varying types of emotional arousal. Another study (Hahn et al, 2012) – more aimed at whether temperature changes can be stimulated by interpersonal social contact – saw similarities with the study of Merla and Romani (2007). Their results demonstrated that during interpersonal interactions, changes in skin temperature were a sensitive index of arousal.

Furthermore, another study which explored the temperature variations of the nasal tip and maxillary area averaged for a group of children and for a group of mothers, found a strong correlation in thermal variations between children and their mothers which were shown during the emotional phases of the experiment. During the experiment, children were invited by the researcher to play with a toy. The toy was designed to break during play, giving the impression that the child had accidentally broken it. Mothers were invited to observe their children in interaction with the researcher through a one-way mirror from a separated room. Results showed that during the experiment the emotional distress caused by the 'breaking' of the toy lead to thermal variations in the specified facial region of interest of the child. The facial thermal modulations observed in the mothers were surprisingly similar to those observed in the child. Thus, mother-child dyads demonstrated a significant and condition-related synchronicity between the autonomic reactions individually exhibited by each of them (Ebisch et al, 2012).

It has also been demonstrated that a specific physiological activity will have a distinct thermal signature; this is a result of the environmental demands the human body is faced with, such as the “fight or flight or freeze” response. Changes in the autonomic nervous system due to the “fight or flight or freeze” response will cause changes to cutaneous temperature, therefore affecting temperature control. However, temperature can vary depending on the level and type of arousal. The distinct thermal signature associated with this response can be clearly observed using an infrared camera (Ioannou et al, 2014). Most scientists have described TI as an emerging and favorable method for detecting stress as well as anxiety (Pollina, 2006). Yoshitomi et al, (1997) addressed the accuracy of TI method within their study with the aim of developing a TI process that would aid computer vision in recognition of facial expression. Along with thermal image processing, the Neural Network was used; data of temperature-difference was the input data for the Neural Network to distinguish between different facial expressions. The results showed that this method had 90% accuracy in detecting neutral, happy, surprised and sad facial expressions.

TI is a reliable method that offers new opportunities to those wanting to study psychological responses, emotional and behavioral changes. TI turns emotional and psychological changes that are invisible to the naked eye into a thermal infrared image with measurable data.

References

1. Bach DR, Friston KJ, Dolan RJ. Analytic measures for quantification of arousal from spontaneous skin conductance fluctuations. *Int J Psychophysiol*, 2010;76(1):52-55.
2. Changeux JP. *Neuronal Man: The Biology of Mind*. New York: Oxford University Press, 1985.
3. Ebisch SJ, Aureli T, Bafunno D, Cardone D, Romani GL, Merla A. Mother and child in synchrony: thermal facial imprints of autonomic contagion. *Biological Psychology*, 2012; 89(1):123-129.
4. Faizi F, Tavallaee A, Rahimi A, Saburi A, Saghafinia M. Quality assessment of randomized control trials applied psychotherapy for chronic pains in Iran: a systematic review of domestic trials. *Iran Red Crescent Med J*, 2014;16(9): e15312.
5. Hahn AC, Whitehead RD, Albrecht M, Lefevre CE Perrett DI. Hot or not? Thermal reactions to social contact. *Biol Lett* doi:10.1098/rsbl.2012.0338 Published online
6. Huitt W. *Reliability and validity. Educational Psychology Interactive*. Valdosta, GA: Valdosta State University, 1999.
7. Ioannou S, Gallese A, Merla A. Thermal infrared imaging in psychophysiology: Potentialities and limits. *Psychophysiology*, 2014;51(10):951-963.
8. Levine JA, Pavlidis I, Cooper M. The face offear. *Lancet*. 2001; 357:1757.
9. Merla A and Romani GL. Thermal signatures of emotional arousal: a functional infrared imaging study. Paper presented at the 29th annual international conference of the IEEE (EMBS 2007), Lyon, France.
10. Mizukami K, Kobayashi N, Iwata H, Ishii T. Telethermography in infant's emotional behavioural research. *Lancet*, 1987; 330:38-39.
11. Mizukami K, Kobayashi N, Ishii T, Iwata H. First selective attachment begins in early infancy: a study using telethermography. *Infant Behav*. 1990; 13: 257-271.
12. Modest, M.F. *Radiative Heat Transfer*. Academic Press: Waltham, MA, USA, 2013.

13. Nhan BR and Chau T. Classifying affective states using thermal infrared imaging of the human face. *Biomed Eng IEEE Trans.* 2010; 57:979–987.
14. Nozawa A and Tacano M. Correlation analysis on alpha attenuation and nasal skin temperature. *J Stat Mech Theory Exp.* 2009; 01:P01007.
15. O’Kane BL, Sandick P, Shaw T, Cook M. Dynamics of human thermal signatures. Paper presented at the *InfraMation Proceedings 2004*, Las Vegas, NV.
16. Pavlidis I, Levine J, Baukol P. Thermal imaging for anxiety detection. *Proceedings of the IEEE Workshop on Computer Vision Beyond the Visible Spectrum: Methods and Applications*, 16-6 June 2000, Washington, DC, pp. 104–109.
17. Pavlidis I, Levine J. Thermal image analysis for polygraph testing. *Eng Med Biol Mag IEEE.* 2002; 21:56–64.
18. Pollina D. Emerging Methods and Measures for Detecting Stress and Deception: Thermal Imaging. *The Journal of Credibility Assessment and Witness Psychology*, 2006; 7(2):108-115.
19. Puri C, Olson L, Pavlidis I, Levine J, Starren J. StressCam: non-contact measurement of users’ emotional states through thermal imaging. Paper presented at *CHI extended abstracts on human factors in computing systems*, April 2-7 2005, Portland, OR, USA, pp. 1725–1728.
20. Richardson R, Trépel D, Perry A, Ali S, Duffy S, Gabe R, Gilbody S, Glanville J, Hewitt C, Manea L, Palmer S, Wright B, McMillan D. Screening for psychological and mental health difficulties in young people who offend: a systematic review and decision model. *Health Technol Assess*, 2015;19(1):1-128.
21. Ring EFJ, Ammer K. The technique of infrared imaging in medicine. *Thermology Int*, 2000;10:7–14.
22. Shearn D, Bergman E, Hill K, Abel A. Facial coloration and temperature responses in blushing. *Psychophysiology*, 1990; 27: 687–693.
23. Topalidou A., Downe S. Investigation of the use of thermography for research and clinical applications in pregnant. *Infrared Physics and Technology*, 2016; 7(2016):59-64.

24. Wong SWH, Xue G, Bechara A. Integrating fMRI with psychophysiological measurements in the study of decision-making. *J Neurosci Psychol Econ*, 2011;4(2):85-94.
25. Yoshiton Y, Miyawaki N, Tomita S, Kimura S. Facial expression recognition using thermal image processing and neural network. *ACS'10 Proceedings of the 10th WSEAS international conference on applied computer science*. Pp 182-186.
26. Zajonc RB, Murphy ST, Inglehart M. Feeling and facial efference: implications of the vascular theory of emotion. *Psychol Rev*, 1989; 96:395-416.