


This document is the unedited Author's version of a Submitted Work that was subsequently accepted for publication in Organic Letters, copyright © American Chemical Society after peer review. To access the final edited and published work see <http://pubs.acs.org/doi/abs/10.1021/acs.orglett.5b01317>

One-Pot Hydrogen Peroxide and Hydrohalic acid Induced Ring Closure and Selective Aromatic Halogenation to give New Ring-Fused Benzimidazoles

Michael Gurry, Martin Sweeney, Patrick McArdle, and Fawaz Aldabbagh*

School of Chemistry, National University of Ireland Galway, University Road, Galway, Ireland

Supporting Information



ABSTRACT: A new series of selectively dichlorinated and dibrominated five to eight-membered ring [1,2-*a*] fused benzimidazoles and [1,4]oxazino[4,3-*a*]benzimidazoles are synthesized in mostly high yields of >80% using the reaction of hydrogen peroxide and hydrohalic acid with commercially available *o*-cyclic amine substituted anilines. Domestic bleach with HCl is also capable of a one-pot ring-closure and chlorination.


The combination of hydrogen peroxide and hydrohalic acid (HX, where X = Cl, Br) is a source of electrophilic chlorine and bromine that can be used for facile aromatic halogenations.¹⁻⁴ Moreover H₂O₂ and HCl has been reported to give 4,6-dichlorination of 5-hydroxybenzimidazole.⁵ 1-Chlorobenzimidazole can be formed by reacting benzimidazole with sodium hypochlorite (NaOCl) in CCl₄.⁶ The intermediate of the reaction between H₂O₂ and HCl is hypochlorous acid (HOCl), which is commonly used to disinfect water, and its salt is the active ingredient in domestic bleaches. The oxidizing solution is very cheap, low in molecular weight, and allows the *in situ* generation of elemental chlorine and bromine with the by-product being water. Therefore there are significant green and technical advantages to using H₂O₂-HX in organic synthesis.

More than fifty years ago, it was recognized that a combination of H₂O₂ and trifluoroacetic acid could be used to prepare ring-fused benzimidazoles in good yields from *o*-cyclic amine substituted anilines.⁷ The use of *o*-*tert*-aminoacetanilides with peroxide (including H₂O₂⁸ and Oxone⁹) in formic acid is recognized as a versatile method for preparing ring-fused benzimidazoles and imidazobenzimidazoles. The preparation of 2-aryl substituted benzimidazoles was reported from the condensation of aryl aldehydes with *o*-phenylenediamines in the presence of H₂O₂-HCl.¹⁰ Moderate yields of benzimidazoles and ring-fused benzimidazoles with tetrachlorination of the fused benzene part was reported using the reaction of sulfonyl chloride with *o*-aminodialkylanilines.¹¹

Halogenated benzimidazoles have anti-cancer,^{12, 13} anti-protozoal,¹³ anti-tuberculosis,¹⁴ and anti-hepatitis activity,¹⁵ as well as, dopamine-receptor binding.¹⁶ In addition to their biological activity, benzimidazoles chlorinated and brominated at specific sites provide promise as valuable synthetic intermediates. Halogenations of heterocycles are generally carried out in subsequent synthetic step(s) and require the use of difficult to handle Cl₂, Br₂ or organic reagents prepared from them. The halogenation of the heterocycle is often associated with low selectivity and when organic reagents are used waste by-products are generated. Our objective was thus to accomplish a one-pot reaction that combined the aromatic halogenation capacity of H₂O₂-HX with the oxidative cyclization of *o*-cyclic amine substituted anilines to form a new series of valuable halogenated ring-fused benzimidazoles. Herein, is reported the first preparation of specifically dichlorinat-

ed and dibrominated ring-fused benzimidazoles from commercially available anilines.

Table 1. Optimization of Reaction Conditions




entry	oxidant	solvent	temp	yield of 2a (%)
1	H ₂ O ₂	MeCN	rt	trace
2	H ₂ O ₂	MeCN	rt ^a	73
3	H ₂ O ₂	MeCN	50 °C	81
4	H ₂ O ₂	MeCN	reflux	90
5	H ₂ O ₂	THF	reflux	83
6	H ₂ O ₂	MeOH	reflux	50 ^b
7	NaOCl	MeCN	reflux	68 ^c
8	household bleach ^d	MeCN	reflux ^e	56

Conditions: Aniline **1a** (1.0 mmol), oxidant (10 mmol), HCl (12 mmol), solvent (10 mL); ^a4 h; ^brecovery of **1a** (38%); ^cPlus **3a** in 15% yield; ^d20 mL of Parazone[®] thin bleach; ^e1 h

Initially we attempted to establish the one-pot oxidative cyclization and chlorination on 5-bromo-2-piperidin-1-ylaniline **1a** in acetonitrile (Table 1). Excess molar amounts of concentrated HCl relative to H₂O₂ were found necessary, and the reaction times were reduced on heating. Monitoring of the reaction by TLC showed the total consumption of aniline **1a** within 20 minutes using the optimized conditions (entry 4). 6,8-Dichlorinated pyrido[1,2-*a*]benzimidazole **2a** was isolated in 90% yield after basic work-up without the requirement for chromatography. This transformation was found to also work in THF and methanol, however yields of **2a** were reduced, and a cleaner reaction occurred in acetonitrile. The sodium salt of hypochlorous acid formed *in situ* is commonly used in domestic bleaches. We thus decided to replace

H₂O₂ with NaOCl solution, and found that the desired **2a** could be isolated in 68% yield, although chromatography was required to separate some trichlorinated benzimidazole **3a** formed in 15% yield. The reaction was then carried out using a well-known brand of household bleach (containing an unspecified quantity of NaOCl) to exclusively give 6,8-dichloro adduct **2a** in 56% yield with the reduced yield attributed to the purification by chromatography, which separated a number of additives (presumably surfactants and perfumes) in the bleach.

Scheme 1. One-Pot Ring-Fused Benzimidazole Formation with Aromatic Trihalogenation




Using the optimized conditions (Table 1, entry 4) the scope and versatility was explored. Firstly we attempted to investigate *o*-cyclic amine substituted anilines devoid of other aromatic substituents (Scheme 1). 2-Piperidin-1-ylaniline (**1b**) and 2-morpholin-4-ylaniline (**1c**) gave the respective trichlorinated ring-fused benzimidazoles **2b** and **2c** in 71 and 89% yield. Tetrachlorination product **3b** in 12% yield was separated by column chromatography from **2b**. Replacing HCl by concentrated HBr transformed piperidine **1b** into dibrominated pyrido[1,2-*a*]benzimidazole **4b** in 94% yield (Table 3), although under the same conditions the morpholine analogue was tribrominated to give **4c** in 87% yield (Scheme 1). Attempts at isolating dibrominated benzimidazole from the reaction of **1c** at shorter reaction times were unsuccessful due to the isolation of mixtures of brominated benzimidazoles, and a clean transformation occurred by increasing the reaction time to 18 hours to give exclusively **4c**.

The preparation of selectively dichlorinated ring-fused benzimidazoles from various *o*-cyclic amine substituted anilines containing electron-donating and electron-withdrawing groups proved mostly facile with yields of 72-92% obtained of ring-fused benzimidazoles (Table 2). The substitution of the chlorine atoms was consistent, and did not vary with the nature of the substituent on the aniline, as confirmed by X-ray crystal structures of adducts **2d** and **2i** (Figure 1).¹⁷ In some cases chromatography was required to separate small amounts of fully chlorinated benzimidazoles **3b** and **3h**, although 5-fluoro-2-piperidin-1-ylaniline (**1f**) tended to prefer cyclization with monochlorination to give **3f** in 62% yield with the dichlorinated adduct **2f** given in smaller yield of 27%.


Beginning with the optimized conditions (Table 1, entry 4) and replacing HCl with HBr, the preparation of dibrominated ring-fused benzimidazoles was investigated (Table 3). In some cases the latter reaction conditions used successfully to yield dichlorinated ring-fused benzimidazoles (Table 2) did not require modification, although significantly longer reaction times were required for resonance activators (NHAc and OMe) on the aniline.

Table 2. One-Pot Ring-Fused Benzimidazole Formation with Aromatic Dichlorination




Conditions: Same as in Table 1, entry 4

Figure 1. Crystal Structures of 7,9-dichloro-3,4-dihydro-1H-[1,4]oxazino[4,3-*a*]benzimidazoles




1	R	Y	yield (%)
1a	Br	CH ₂	2a , 90
1d	Br	O	2d , 92
1e	Cl	CH ₂	2b , 73 + 3b , 15
1f	F	CH ₂	2f , 27 + 3f , 62
1g	Me	CH ₂	2g , 82
1h	CN	CH ₂	2h , 79 + 3h , 12
1i	CF ₃	O	2i , 72
1j	NHAc	CH ₂	2j , 88
1k	OMe	O	2k , 87



Nevertheless dibrominated ring-fused benzimidazoles were isolated in excellent yield of 74-94% and without the requirement for chromatography, save for three anilines containing electron-withdrawing groups (F, CN, CF₃). The substitution pattern was the same as for the chlorination, and the positions for the bromination did not vary with the nature of the substituent on the aniline, as confirmed by X-ray crystal structures of dibrominated adducts **4b**, **4f** and **4m** (Figure 2).¹⁷ Analogous to the results obtained with H₂O₂-HCl, 5-fluoro-2-piperidin-1-ylaniline (**1f**) gave significant monobromination at pyrido[1,2-*a*]benzimidazole C-8 with **5f** separated in 31% yield from the desired dibrominated adduct **4f** isolated in 56% yield. Surprisingly, the benzonitrile **1h**, which gave some polychlorination with H₂O₂-HCl after only 20 minutes, was found difficult to dibrominate with only the monobromide **5h** isolated in 62% yield. Further, no brominated benzimidazoles could be cleanly separated from the attempted reaction with 5-(trifluoromethyl)aniline **1i**, although the same substrate gave selectively 7,9-dichlorinated [1,4]oxazino[4,3-*a*]benzimidazole **2i** in 72% yield using H₂O₂-HCl (Table 2). Thus, it seems that the bromination is more strongly affected by substituents on the aniline than the chlorination with electron-withdrawing substituents making dibromination difficult.


Table 3. One-Pot Ring-Fused Benzimidazole Formation with Aromatic Dibromination



1	R	Y	time	yield (%)
1a	Br	CH ₂	20 min	4a, 89
1b	H	CH ₂	20 min	4b, 94
1d	Br	O	20 min	4c, 92
1e	Cl	CH ₂	40 min	4e, 93
1f	F	CH ₂	20 min	4f, 56 ^a + 5f, 31 ^a
1g	Me	CH ₂	20 min	4g, 84
1h	CN	CH ₂	20 min	5h, 62 ^{a,b}
1i	CF ₃	O	20 min	4i, 0 ^{a,c}
1j	NHAc	CH ₂	2 d	4j, 74 ^d
1k	OMe	O	8 h	4k, 92

Conditions: Aniline **1** (1.0 mmol), H₂O₂ (10 mmol), HBr (12 mmol) in MeCN (10 mL); ^aAdditional times and equivalents produced similar results; ^b4h observed but was not isolated; ^cIntractable mixture of mono and non-brominated ring-fused benzimidazole; ^dH₂O₂ (20 mmol), HBr (24 mmol) in MeCN (10 mL).


Figure 2. Crystal Structures of 6,8-dibromo-1,2,3,4-tetrahydropyrro[1,2-*a*]benzimidazoles



The synthesis of alternative [1,2-*a*] alicyclic ring-fused benzimidazoles was examined (Table 4). Selectively dichlorinated five to eight-membered [1,2-*a*] alicyclic ring fused benzimidazoles **2l-2o** were isolated in 78-93% yield after reaction times of 20 minutes. The brominations were found to be significantly slower than the analogous chlorinations, and double the number of equivalent of H₂O₂-HBr were required to obtain the desired products. This is in agreement with literature kinetic studies for the halogenation of *p*-xylene, where Br₂ was found to react more than 200 times slower than Cl₂.¹⁸ Attempts were made to reduce reac-

tion times by activating bromine towards electrophilic attack by adding a quaternary ammonium salt (TBAB),² but no effect on rate was observed. Our system is however different to the literature,² as there is an absence of a two-phase system due to the solubility of acetonitrile in water. It is conceivable that steric factors might be also influencing the rate of electrophilic aromatic bromination, since one would have expected greater polarizability in suspected bromination species, H₂O⁺-Br and Br₂ compared to the chlorine analogues.³ Nevertheless selectively dibrominated pyrro-, azepino- and azocino [1,2-*a*] ring-fused benzimidazoles **4m-4o** were isolated in high yields of 70-86%. Monitoring the reaction of **1l** with H₂O₂-HBr using ¹H NMR showed after 20 minutes the formation of a mixture of monobromides; 5- and 7-bromobenzimidazoles in an approximate 1:3 ratio and after 20 hours a mixture of di- and tribromides **4l** and **5l** remained in an approximate 2:1 ratio. It was thus not possible to cleanly isolate 5,7-dibromopyrro[1,2-*a*]benzimidazole **4l**, and it was decided to allow the tribromide **5l** to be regioselectively formed in 87% yield.

Table 4. One-Pot Five to Eight-Membered [1,2-*a*] Ring-Fused Benzimidazole Formation with Aromatic Dihalogenation



1	X	n	time	yield (%)
1l	Cl	1	20 min	2l, 93
1m	Cl	2	20 min	2m, 90
1n	Cl	3	20 min	2n, 89
1o	Cl	4	20 min	2o, 78
1l	Br	1	4 d	5l, 87
1m	Br	2	4 d	4m, 86
1n	Br	3	5 d	4n, 80
1o	Br	4	5 d	4o, 70

Conditions: Aniline **1** (1.0 mmol) in MeCN (10 mL) with H₂O₂ (10 mmol) and HCl (12 mmol) for X = Cl or with H₂O₂ (20 mmol) and HBr (24 mmol) for X = Br.

In conclusion we have successfully established a simple and inexpensive H₂O₂-HX preparation of halogenated ring-fused benzimidazoles with 31 new compounds now reported from commercial anilines. In most cases the one-pot transformation gives regioselectively the novel dihalogenated heterocycle in high yields.

■ ASSOCIATED CONTENT

Supporting Information

Experimental procedures and spectroscopic data for all compounds, including X-ray crystallographic data for compounds **2d**, **2i**, **4b**, **4f** and **4m** is available free of charge via the Internet at <http://pubs.acs.org>.

■ AUTHOR INFORMATION

Corresponding Author

*Email: fawaz.aldabbagh@nuigalway.ie; Tel.: +353 (0) 91 493120. Fax: +353 (0) 91 49557.

Notes

The Authors declare no competing financial interest.

■ ACKNOWLEDGMENT

The authors thank the College of Science, National University of Ireland Galway (NUI Galway) for a Postgraduate Scholarship for M.G. and the Irish Research Council (IRC) for a Government of Ireland Postgraduate Scholarship for M.S.

■ REFERENCES

1. (a) Ben-Daniel, R.; de Visser, S. P.; Shaik, S.; Neumann, R. *J. Am. Chem. Soc.* **2003**, *125*, 12116-12117; (b) Sivey, J. D.; Roberts, A. L. *Environ. Sci. Technol.* **2012**, *46*, 2141-2147.
2. Dakka, J.; Sasson, Y. *J. Chem. Soc. Chem. Commun.* **1987**, 1421-1422.
3. Sivey, J. D.; Arey, J. S.; Tentscher, P. R.; Roberts, A. L. *Environ. Sci. Technol.* **2013**, *47*, 1330-1338.
4. (a) Barhate, N. B.; Gajare, A. S.; Wakharkar, R. D.; Bedekar, A. V. *Tetrahedron Lett.* **1998**, *39*, 6349-6350. (b) Barhate, N. B.; Gajare, A. S.; Wakharkar, R. D.; Bedekar, A. V. *Tetrahedron* **1999**, *55*, 11127-11142. (c) Bogdal, D.; Lukasiewicz, M.; Pielichowski, J. *Green Chem.* **2004**, *6*, 110-113. (d) Koini, E. N.; Avlonitis, N.; Calogeropoulou, T. *Synlett* **2011**, 1537-1542.
5. Kuznetsov, Y. V.; Stolyarova, L. G.; Lezina, V. P.; Smirnov, L. D. *Bull. Acad. Sci. USSR. Div. Chem. Sci.* **1989**, *38*, 1494-1500.
6. De Rosa, M.; Canudas, N.; Arnold, D.; Yennawar, H. *J. Org. Chem.* **2013**, *78*, 7264-7267.
7. Nair, M. D.; Adams, R. *J. Am. Chem. Soc.* **1961**, *83*, 3518-3521.
8. Fahy, K.; Aldabbagh, F. *Tetrahedron Lett.* **2008**, *49*, 5235-5237.
9. Fagan, V.; Bonham, S.; McArdle, P.; Carty, M.P.; Aldabbagh, F. *Eur. J. Org. Chem.* **2012**, 1967-1975
10. Bahrami, K.; Khodaei, M. M.; Kavianiinia, I. *Synthesis* **2007**, 547-550.
11. Martin, J.; Meth-Cohn, O.; Suschitzky, H. *Tetrahedron Lett.* **1973**, *14*, 4495-4498.
12. Carpenter, R. D.; Natarajan, A.; Lau, E. Y.; Andrei, M.; Solano, D. M.; Lightstone, F. C.; DeNardo, S. J.; Lam, K. S.; Kurth, M. J. *Cancer Res.* **2010**, *70*, 5448-5456.
13. Andrzejewska, M.; Yépez-Mulia, L.; Cedillo-Rivera, R.; Tapia, A.; Vilpo, L.; Vilpo, J.; Kazimierczuk, Z. *Eur. J. Med. Chem.* **2002**, *37*, 973-978.
14. Gobis, K.; Foks, H.; Serocki, M.; Augustynowicz-Kopeć, E.; Napiórkowska, A. *Eur. J. Med. Chem.* **2015**, *89*, 13-20.
15. Borowski, P.; Deinert, J.; Schalinski, S.; Bretner, M.; Ginalski, K.; Kulikowski, T.; Shugar, D. *Eur. J. Biochem.* **2003**, *270*, 1645-1653.
16. Šukalović, V.; Andrić, D.; Roglić, G.; Kostić-Rajačić, S.; Šoškić, V. *Arch. Pharm.* **2004**, *337*, 376-382.
17. Crystallographic data (excluding structure factors) has been deposited with the Cambridge Crystallographic Data Centre as supplementary publication CCDC 1054365 for **(2d)** CCDC 1054369 for **(2i)**, CCDC 1054366 for **(4b)**, CCDC 1054368 for **(4f)**, and CCDC 1054367 for **(4m)**.
18. Voudrias, E. A.; Reinhard, M. *Environ. Sci. Technol.* **1988**, *22*, 1049-1056.