


RESEARCH ARTICLE

Chronic activation of PPAR α with fenofibrate reduces autophagic proteins in the liver of mice independent of FGF21

Eunjung Jo¹, Songpei Li¹, Qingning Liang², Xinmei Zhang¹, Hao Wang¹, Terence P. Herbert¹, Trisha A. Jenkins¹, Aimin Xu², Ji-Ming Ye^{1*}

1 School of Health and Biomedical Sciences, RMIT University, Melbourne, Victoria, Australia, **2** State Key Laboratory of Pharmaceutical Biotechnology, Department of Medicine, Hong Kong University, Hong Kong, China

* jiming.ye@rmit.edu.au


 OPEN ACCESS

Citation: Jo E, Li S, Liang Q, Zhang X, Wang H, Herbert TP, et al. (2017) Chronic activation of PPAR α with fenofibrate reduces autophagic proteins in the liver of mice independent of FGF21. PLoS ONE 12(4): e0173676. <https://doi.org/10.1371/journal.pone.0173676>

Editor: Hervé Guillou, INRA, FRANCE

Received: July 11, 2016

Accepted: February 25, 2017

Published: April 19, 2017

Copyright: © 2017 Jo et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper and its Supporting Information files.

Funding: This work was funded by a National Health and Medical Research Council of Australia Program Grant 535921 (allocation to JM Ye), Diabetes Australia Research Program GG12314 (allocation to JM Ye), and Block Grant Earmarked for Research (104) (201109176145) (allocation to A Xu).

Competing interests: The authors have declared that no competing interests exist.

Abstract

Autophagy is a catabolic mechanism to degrade cellular components to maintain cellular energy levels during starvation, a condition where PPAR α may be activated. Here we report a reduced autophagic capacity in the liver following chronic activation of PPAR α with fenofibrate (FB) in mice. Chronic administration of the PPAR α agonist FB substantially reduced the levels of multiple autophagy proteins in the liver (Atg3, Atg4B, Atg5, Atg7 and beclin 1) which were associated with a decrease in the light chain LC3II/LC3I ratio and the accumulation of p62. This was concomitant with an increase in the expression of lipogenic proteins mSREBP1c, ACC, FAS and SCD1. These effects of FB were completely abolished in PPAR α ^{-/-} mice but remained intact in mice with global deletion of FGF21, a key downstream mediator for PPAR α -induced effects. Further studies showed that decreased the content of autophagy proteins by FB was associated with a significant reduction in the level of FoxO1, a transcriptional regulator of autophagic proteins, which occurred independently of both mTOR and Akt. These findings suggest that chronic stimulation of PPAR α may suppress the autophagy capacity in the liver as a result of reduced content of a number of autophagy-associated proteins independent of FGF21.

Introduction

Autophagy is a process to degrade and recycle dysfunctional cellular components via the lysosome in order to maintain cellular homeostasis [1]. It is also important in maintaining energy during periods of starvation. Autophagy is regulated by the nutrient status of the cell via a number of nutrient-sensitive signalling pathways such as mammalian target of rapamycin (mTOR) and AMP-activated protein kinase (AMPK) pathways [2–4]. Forkhead box O (FoxO) family proteins also play an important role by controlling the expression of a number of autophagy related genes [3, 5, 6]. Another transcription factor that is critical for adaptive metabolism to starvation is peroxisome proliferator-activated receptor α (PPAR α). Under physiological

conditions, PPAR α is activated by mobilised fatty acids (FAs) but can also be activated pharmacologically by fibrates, a class of lipid-lowering drugs [7].

PPAR α is highly expressed in the liver and when activated it up-regulates genes for FA oxidation and gluconeogenesis to provide fuels for the body [7]. As expected from its role in promoting catabolism, recent studies have shown that hepatic autophagy is activated via PPAR α during fasting or after short-term treatment with PPAR α agonists both *in vivo* and *in vitro* in hepatocytes [8, 9]. Interestingly, it has been suggested that an increased autophagy activity may be gradually subsided or even reduced over time under certain conditions [10]. Therefore, the first aim of the present study was to examine the expression of autophagic proteins in the liver of both wild-type (PPAR $\alpha^{+/+}$) and PPAR $\alpha^{-/-}$ mice after chronic administration of the PPAR α activator fenofibrate (FB). As lipogenic proteins are up-regulated during PPAR α activation [11] or by inhibition of autophagy [12], our second aim was to investigate the relationship of changes in autophagic proteins with the expression of lipogenic proteins.

It has been suggested that fibroblast growth factor 21 (FGF21) is an important mediator for the physiological effects initiated by PPAR α activation [13–16] and this cytokine is up-regulated along with autophagy-related gene 5 (Atg5) [17]. Thus, our third aim was to determine whether FGF21 is required for PPAR α to exert its effects on the expression of autophagic proteins using FGF21 $^{-/-}$ mice. Finally, we examined the key signalling pathways that have been suggested to regulate autophagy during the chronic activation of PPAR α .

In this report we show that chronic activation of PPAR α by FB reduces the expression of autophagic proteins in the liver in a manner that is entirely independent of FGF21. PPAR α -induced suppression of autophagic proteins is possibly mediated by a decrease in FoxO1 expression rather than through changes in the activity of mTOR or Akt. These findings suggest a need to further investigate the dynamic changes of hepatic autophagy during PPAR α activation and associated implications for lipid metabolism.

Materials and methods

Animals

The studies were conducted in male mice starting at an age of 10–12 weeks, including wild-type (PPAR $\alpha^{+/+}$) and PPAR $\alpha^{-/-}$ on the background of C57BL/6N, and wild-type (FGF21 $^{+/+}$) and FGF21 $^{-/-}$ mice on the background of C57BL/6J originally obtained from Jackson Laboratories (Sacramento, CA, US). The mice were housed at 23±1°C in a 12-h light/dark cycle with free access to water and standard rodent diet consisting of 70% calories as starch, 10% calories as fat and 20% calories from proteins (Specialty Feeds, Australia). After 1–2 weeks of acclimatization, mice were fed the standard diet in the absence or presence of the PPAR α agonist FB for 3 weeks. FB (Sigma-Aldrich, Australia) was administered as an additive to diet at a lower dose (50 mg/kg/day) relative to our previous studies to minimize the possible influence of body weight reduction. Body weight and food intake were monitored daily. Blood samples were taken from the tail vein in week 3 after 5–7 hours of fasting and the mice were culled by cervical dislocation. Liver was removed quickly (<5 seconds), weighed on a balance and immediately freeze-clamped for storage at -80°C for subsequent analysis. All animal experiments were approved by the Animal Ethics Committee of the RMIT University or the University of Hong Kong, where animal studies were performed.

Determination of circulating levels of glucose and FGF21

Plasma glucose levels of PPAR $\alpha^{-/-}$ mice were determined by glucose assay according to the manufacturer's protocol (Sigma-Aldrich, Australia). The concentrations of plasma FGF21 in both FGF21 $^{+/+}$ and FGF21 $^{-/-}$ were measured by an ELISA kit (University of Hong Kong, Hong

Kong) during FB induced 24 hours of fasting according to the manufacturer's introductions [15].

Extraction of hepatic triglycerides contents

Liver triglycerides (TG) were extracted by the method of Folch and determined by a TG GPO-PAP kit (Roche Diagnostic, Australia) as previously described [18]. Briefly, 30–40mg of each liver sample was homogenized in 4 ml of chloroform/methanol (2:1) using a glass pestle tissue grinder. After the homogenization, the samples were rotated at room temperature overnight to ensure the complete solubilisation of the liver TG. The next day, 2 ml of 0.6% NaCl was added to each sample and followed by centrifugation to separate the aqueous from the organic phases. The lower chloroform layer contained liver TG were carefully transferred into a glass vial and dried completely under the nitrogen or air at 45°C. The extract was reconstituted in absolute ethanol for the determination TG using a POLARstar microplate reader (BMG Labtech, Germany).

Immunoblotting analysis

Immunoblotting analysis was performed as described in our recent reports [11, 18]. Briefly, liver tissues were homogenized in ice-cold RIPA lysis buffer supplemented with protease inhibitor cocktail and phosphatase inhibitor cocktail (Sigma Aldrich Pty Ltd, Australia) and DL-dithiothreitol. Protein samples were then denatured in a SDS sample buffer. Proteins of interest were analyzed by immunoblotting using specific antibodies from Cell Signaling (USA) unless indicated otherwise. Key autophagy proteins included Atg3, Atg4B, Atg5, Atg7, nucleoporin p62 (p62), light chain 3A/B (LC3A/B) and beclin-1, phosphor- (Ser2448) and total- mTOR, phosphor- (Thr389) and total- S6K, phosphor- (Thr37/46) and total- 4EBP. For investigating the mechanism pathway of FoxO1: phospho- and total- FoxO1, acetyl (D-19) FoxO1 (Santa Cruz, USA), sirtuin-silent mating type information regulation 2 homolog (SIRT1), phosphor- (Ser473) and total- Akt, phosphor- (Ser9) and total- glycogen synthase kinase 3 β (GSK3 β). Key lipogenic enzymes were examined using specific antibodies including matured form of mSREBP-1c, acetyl-CoA carboxylase (ACC), fatty acid synthase (FAS) and stearyl-CoA desaturase (SCD1) (Santa Cruz, USA). Acyl-coenzyme A oxidase 1 (ACOX1) (Santa Cruz, USA) was determined to indicate PPAR α action and peroxisomal FA oxidation. Proteins were analyzed and normalized against the housekeeper glyceraldehyde 3-phosphate dehydrogenase (GAPDH) and/or α -Tubulin, or its specific total form of protein. Immunolabeled proteins were visualized using a ChemiDoc densitometer and quantified by densitometry of Image Lab software (Bio-Rad Laboratories, USA) with inclusion of representative images.

Quantitative RT-PCR

RNA was extracted from the liver tissue using TRIzol Reagent (Invitrogen, Australia) and reverse transcribed using a high capacity cDNA reverse transcription kit (Applied Biosystems, Australia) according to the manufacturer's instructions. Primers (GeneWorks, Australia) and SYBER green supermix (Bio-Rad, USA) were used for quantitative real time PCR. The primer sequences for FoxO1 were: forward 5' -TTCAATTTCGCCACAATCTGTCC-3' and reverse 5' -GGGTGATTTTCCGCTCTTGC-3'. All reactions were performed on QIAGEN Rotor-Gene Q PCR system (Germany). 18s was used as the normalizing control gene.

Table 1. Metabolic responses to chronic activation of PPAR α with FB in PPAR α ^{-/-} mice.

	PPAR α ^{+/+}	PPAR α ^{+/+} -FB	PPAR α ^{-/-}	PPAR α ^{-/-} -FB
Body weight (g)				
Basal	21.5 ± 0.1	21.5 ± 0.2	26.4 ± 0.9 ^{††}	24.5 ± 1.1
After	24.2 ± 0.3	24.4 ± 0.4	28.7 ± 1.1 ^{††}	26.3 ± 1.8
Body weight gain (g)	2.7 ± 0.2	2.9 ± 0.2	2.3 ± 0.2	1.8 ± 0.7
Food intake (g/day/mouse)				
Basal	3.2 ± 0.1	3.2 ± 0.1	4.1 ± 0.2	5.0 ± 0.2
During treatment	3.3 ± 0.2	3.2 ± 0.1	4.8 ± 0.2	5.3 ± 0.2
Liver weight (g)	0.9 ± 0.0	1.6 ± 0.1 ^{**}	1.1 ± 0.0	1.2 ± 0.1
Liver weight/Body weight (%)	4.0 ± 0.0	7.0 ± 0.0 ^{**}	4.0 ± 0.0	5.0 ± 0.0
Plasma glucose level	10.0 ± 0.4	9.3 ± 0.3	6.9 ± 0.2 ^{††}	6.0 ± 0.2 ^{††}

PPAR α ^{+/+} and PPAR α ^{-/-} mice were administered with fenofibrate (FB, 50 mg/kg/day in diet) for 3 weeks. Data are means ± SEM (n = 5–6 mice/group).

* $p < 0.05$,

** $p < 0.01$ vs. vehicle control;

†† $p < 0.01$ vs. corresponding wild-type.

<https://doi.org/10.1371/journal.pone.0173676.t001>

Statistical analyses

Data are presented as means ± SEM. One-way analysis of variance (ANOVA) or the t-tests were used for comparison of relevant groups as needed. When significant differences were found from an ANOVA test, the Tukey-Kramer post-hoc multiple comparisons test was applied. Differences at $p < 0.05$ were considered to be statistically significant.

Results

Effects of FB on whole-body parameters in PPAR α ^{-/-} and FGF21^{-/-} mice

PPAR α ^{-/-} mice were approximately 16% heavier compared with PPAR α ^{+/+} mice at the start of the experiment (25.5 ± 0.8 vs. 21.5 ± 0.2, n = 12, $p < 0.01$). As shown in [Table 1](#), plasma levels of glucose were 40% lower in untreated PPAR α ^{-/-} mice ($p < 0.01$ vs. untreated PPAR α ^{+/+}). Chronic administration of FB did not influence the body weight gain or food intake in either PPAR α ^{+/+} or PPAR α ^{-/-} mice. Consistent with our previous reports [11, 18], PPAR α ^{+/+} mice treated with FB displayed a 70% increase in liver weight ($p < 0.01$ vs. untreated PPAR α ^{+/+} mice) but this effect were not detected in PPAR α ^{-/-} mice. In comparison, FGF21^{-/-} mice were 8% lighter compared with the age-matched FGF21^{+/+} mice at the start of the experiment (21.5 ± 0.2 vs. 25.5 ± 0.8, n = 12, $p < 0.01$, [Table 2](#)). In both of the FGF21^{+/+} and FGF21^{-/-} mice, chronic administration of FB increased liver weight by ~80% ($p < 0.01$ vs. corresponding untreated mice) but had no effect on body weight gain, food intake or plasma glucose.

Effects of FB on hepatic lipid metabolism and autophagic proteins in PPAR α ^{+/+} and PPAR α ^{-/-} mice

Chronic administration of FB had no significant effect on hepatic triglyceride (TG) content in either PPAR α ^{+/+} or PPAR α ^{-/-} mice ([Fig 1A](#)). In PPAR α ^{+/+} mice, FB up-regulated the protein expression of ACOX1 (~2.2-fold, [Fig 1B and 1C](#)), a PPAR α responsive enzyme catalysing FA oxidation in peroxisomes. The levels of lipogenic proteins mature SREPB-1c, ACC, FAS and SCD1 were increased by 2–3 fold ($p < 0.01$). However, these increases were abolished in PPAR α ^{-/-} mice, confirming the requirement of PPAR α for FB to up-regulate the lipogenic pathway.

Table 2. Metabolic responses to chronic activation of PPAR α with FB in FGF21 $^{-/-}$ mice.

	FGF21 $^{+/+}$	FGF21 $^{+/+}$ -FB	FGF21 $^{-/-}$	FGF21 $^{-/-}$ -FB
Body weight (g)				
Basal	25.9 \pm 1.4	25.1 \pm 0.5	21.1 \pm 1.1 [†]	22.0 \pm 1.0 [†]
After	29.5 \pm 1.3	29.5 \pm 0.6	24.3 \pm 1.2 [†]	25.2 \pm 1.0 [†]
Body weight gain (g)	3.6 \pm 0.1	4.4 \pm 0.1	3.2 \pm 0.1	3.2 \pm 0.0
Food intake (g/day/mouse)				
Basal	4.2 \pm 0.2	4.1 \pm 0.1	3.8 \pm 0.2	4.1 \pm 0.2
During treatment	4.4 \pm 0.2	4.2 \pm 0.1	3.9 \pm 0.1	4.0 \pm 0.2
Liver weight (g)	1.2 \pm 0.0	2.1 \pm 0.2**	1.1 \pm 0.1	2.0 \pm 0.1*
Liver weight/Body weight (%)	4.0 \pm 0.3	7.2 \pm 0.7*	4.4 \pm 0.2	8.0 \pm 0.2**
Blood glucose level	12.2 \pm 1.6	10.5 \pm 1.2	12.6 \pm 1.8	13.2 \pm 1.7 ^{††}

FGF21 $^{+/+}$ and FGF21 $^{-/-}$ mice were administered with fenofibrate (FB, 50 mg/kg/day in diet) for 3 weeks. Data are means \pm SEM (n = 5–6 mice/group).

* $p < 0.05$,

** $p < 0.01$ vs. vehicle control;

[†] $p < 0.05$,

^{††} $p < 0.01$ vs. corresponding wild-type.

<https://doi.org/10.1371/journal.pone.0173676.t002>

In PPAR α $^{+/+}$ mice, chronic administration of FB reduced the protein levels of Atg3, Atg4B, Atg5, Atg7 and beclin1 by more than 50% (all $p < 0.01$, Fig 2), suggesting a decreased autophagy capacity. The LC3II to LC3I ratio (indicative of autophagosome formation) was inhibited by ~50% (due to the accumulation of LC3I) whereas p62 protein (indicator of non-degraded aggregates) was accumulated by ~70%. In the liver of PPAR α $^{-/-}$ mice, there were increases in the expression of beclin 1 (~40%), Atg5 (~80%) and Atg7 (~60%) compared to untreated PPAR α $^{+/+}$ mice (all $p < 0.05$) but other measured autophagic proteins were similar. However, chronic administration of FB in PPAR α $^{-/-}$ mice had no effects on the level of autophagic proteins, indicating the requirement of PPAR α activation for FB to mediate the effect of FB to reduce the level of autophagic proteins.

Effects of FB on hepatic lipid metabolism and autophagic proteins in FGF21 $^{+/+}$ and FGF21 $^{-/-}$ mice

We next investigated whether or not FGF21 may be required for the effects of FB on those lipogenic and autophagic proteins. In FGF21 $^{-/-}$ mice, fasting-induced increase in plasma levels of FGF21 was completely diminished (Fig 3A), confirming the lack of FGF21 after its deletion. While liver TG content was not affected (Fig 3B) after chronic administration of FB, the effect of FB in increasing ACOX1 protein was significantly enhanced in FGF21 $^{-/-}$ mice ($p < 0.05$ vs. FB-treated FGF21 $^{+/+}$ mice, Fig 3C). Despite this, the effects of FB to up-regulate mSREPB-1c, ACC, FAS and SCD1 were similar in FGF21 $^{-/-}$ mice compared to FGF21 $^{+/+}$ mice. Atg3, Atg4B, Atg5, Atg7 and beclin1 were reduced (by more than 50%) along with an increase in p62 (2-fold) to the similar extent in both FGF21 $^{+/+}$ and FGF21 $^{-/-}$ mice after chronic administration of FB (Fig 4). These results suggest that FGF21 is not required for PPAR α -induced expression of proteins in the lipogenic and autophagic pathways in the liver.

Effects of FB on the mTOR and insulin signalling pathways in PPAR α $^{+/+}$ and PPAR α $^{-/-}$ mice

We next examined the effects of FB on mTOR and insulin signalling in both PPAR α $^{+/+}$ and PPAR α $^{-/-}$ mice as the activation of these pathways can inhibit autophagy activity. In PPAR α $^{+/+}$ mice, FB inhibited the phosphorylation of mTOR (by ~20%) and its downstream effectors S6K




Fig 1. Effects of FB on hepatic TG content and the lipogenic pathway in PPAR α ^{+/+} and PPAR α ^{-/-} mice. Fenofibrate (+ FB) was administered to PPAR α ^{+/+} and PPAR α ^{-/-} mice on the background of C57BL/6N at a dose (50 mg/kg/day) for 3 weeks. **(A)** Effects on triglyceride (TG) content. **(B)** Representative images and **(C)** Quantification of Western blots for ACOX1 (peroxisomal acyl-CoA oxidase 1, PPAR α activation marker) and lipogenic proteins: mSREBP1c (matured form of sterol regulatory element-binding protein 1 c), ACC (acetyl-CoA carboxylase), FAS (fatty acid synthase) and SCD1 (stearoyl-CoA desaturase 1). ** $p < 0.01$ vs. control (-FB); n.s. not statistically significant. $n = 5-6$ /group.

<https://doi.org/10.1371/journal.pone.0173676.g001>

(by ~65%) and 4EBP1 (~50%) (all $p < 0.05$) (Fig 5). The phosphorylation of Akt was significantly inhibited with a similar trend of change for GSK3 β . However, these effects by FB were all abolished in PPAR α ^{-/-} mice. These results suggest that the changes in the expression of autophagic proteins by PPAR α activation with FB are not due to the activation of the mTOR or insulin signalling pathway.

Effects of FB on FoxO1, SIRT1 and HDAC4 in PPAR α ^{+/+} and PPAR α ^{-/-} mice

We next investigated whether the reduced autophagic proteins after chronic administration of FB may be mediated by FoxO1, a critical transcription factor regulating autophagic proteins. In PPAR α ^{+/+} mice, the expression of FoxO1 was markedly reduced (by ~60%) by the activation of PPAR α with FB ($p < 0.01$, Fig 6A). This reduction was associated with increases in the ratios of both p-FoxO1/FoxO1 (~1.5 fold) and Ac-FoxO1/FoxO1 (~2 fold). However, the expression of FoxO1 mRNA was not altered by FB treatment (Fig 6B). Concomitant with the increased acetylation of FoxO1, the level of SIRT1 (a Class III deacetylase) was reduced




Fig 2. Effects of FB on hepatic autophagic proteins in PPAR α ^{+/+} and PPAR α ^{-/-} mice. Experiments were conducted as described in Fig 1. (A) Representative of images and (B) quantification of the Western blots for key autophagic proteins. Atg3 (autophagy-related gene protein 3), Atg4B (autophagy-related gene protein 4B), Atg5 (autophagy-related gene protein 5), Atg7 (autophagy-related gene protein 7), LC3 (microtubule-associated protein light chain 3) and p62 (polyubiquitin-binding protein p62). * $p < 0.05$, ** $p < 0.01$ vs control (-FB); † $p < 0.05$, †† $p < 0.01$ vs untreated PPAR α ^{+/+} mice; n.s. not statistically significant. $n = 5-6$ /group.

<https://doi.org/10.1371/journal.pone.0173676.g002>

(~60%) and the phosphorylation of histone deacetylase 4 (HDAC4, a Class I deacetylase) was decreased (~70%) (Fig 6C). In PPAR α ^{-/-} mice, all of these effects of FB were abolished, indicating that these observed changes were entirely dependent upon PPAR α .

Discussion

The present study investigated the effect of chronic activation of PPAR α on liver autophagy in mice and the results revealed several novel findings. Firstly, chronic administration of FB PPAR α -dependently down-regulate multiple autophagic proteins while up-regulating lipogenic proteins. Secondly, these PPAR α -dependent effects were entirely independent of FGF21, a key downstream mediator of PPAR α . Thirdly, PPAR α -induced the reductions in autophagic proteins was associated with a reduction in FoxO1 (master transcriptional factor for




Fig 3. Effects of FB on hepatic TG content and lipogenic proteins in FGF21^{+/+} and FGF21^{-/-} mice. (A) Plasma levels of FGF21 in response to fasting. For B and C, FB was administered to FGF21^{+/+} and FGF21^{-/-} mice on the background of C57BL/6J at a dose (50 mg/kg/day) for 3 weeks. (B) Liver TG content. (C) Representative images and quantification of the Western blots for ACOX1, mSREBP1c, ACC, FAS and SCD1. * $p < 0.05$, ** $p < 0.01$ vs control (-FB); n.s. not statistically significant, $n = 5-6$ /group.

<https://doi.org/10.1371/journal.pone.0173676.g003>

autophagic proteins). These findings suggest that chronic activation of PPAR α may reduce the capacity for autophagy in the liver by reducing autophagic proteins via FoxO1 to influence hepatic lipid metabolism.

PPAR α plays an important role in regulating a wide range of metabolic effects in the liver. Under physiological conditions, it is activated by elevated FAs to promote FA oxidation by up-regulating enzymes in mitochondria and peroxisomes including ACOX1 [11, 19]. It has been recently reported that short-term (24–48 hrs) activation of PPAR α in primary hepatocytes increase autophagic activity [9]. However, the effect of long-term activation of PPAR α on autophagy is not clear. The present study revealed that chronic administration of the PPAR α agonist FB down-regulates the expression of multiple autophagic proteins (i.e. Atg3, Atg4B, Atg5, Atg7 and beclin1). Along with this, autophagic activity may be compromised as suggested by the reduced LC3II/LC3I ratio and accumulation of p62, a polyubiquitin-binding protein p62 (SQSTM1) degraded by autophagy [1, 20]. These effects are specific to activation of PPAR α (indicated by increased ACOX1) because they are completely abolished in PPAR α ^{-/-} mice. These results lead us to suggest that persistent stimulation of PPAR α may result in a negative feedback mechanism to down-regulate autophagic proteins in the liver. Consistent with this notion, some autophagic proteins (beclin 1, Atg5 and Atg7) were found to be increased in PPAR α ^{-/-} in the absence of FB, suggesting a regulation of PPAR α on the expression of autophagic proteins under physiological conditions.




Fig 4. Effects of FB on the content of autophagic proteins in the liver of FGF21^{+/+} and FGF21^{-/-} mice. FGF21^{+/+} and FGF21^{-/-} mice were administrated with FB for 3 weeks as described in the methods. The liver samples were collected for Western blots for autophagic proteins including Atg3, Atg4B, Atg5, Atg7, Beclin-1, LC3 and p62. * $p < 0.05$; ** $p < 0.01$ vs group (- FB), $n = 5-6$ /group.

<https://doi.org/10.1371/journal.pone.0173676.g004>

Our previous work showed that chronic administration of FB promotes *de novo* lipogenesis in the liver by up-regulating lipogenic proteins in mice in the presence of PPAR α [18]. The results from the PPAR α ^{-/-} mice in the present study confirmed that the up-regulation of lipogenic proteins in response to FB is mediated specifically by PPAR α . It has been shown that an inhibition of autophagy can lead to increased *de novo* lipogenesis such as in high fructose feeding [12]. Similarly, there is an inverse relationship of autophagic activity with *de novo* lipogenesis in response to chronic activation of PPAR α . The results from this study suggest that the altered autophagy by chronic activation of PPAR α may also impact on *de novo* lipogenesis in the liver.

PPAR α is known to play a critical role in promoting hepatic gluconeogenesis and FA oxidation mediated by FGF21 during starvation [15, 21] where autophagy activity is also altered. However, the observed changes in autophagic proteins (Atg3, Atg4B, Atg5, Atg7, beclin1, LC3 and p62) and lipogenic proteins (mSREBP1, ACC, FAS and SCD1) by chronic activation PPAR α with FB remained intact in FGF21^{-/-} mice. These findings indicate that PPAR α -induced expressions of autophagic and lipogenic proteins are independent of its downstream mediator FGF21.

Autophagy activity can be regulated by the mTOR pathway [2-4]. However, chronic activation of PPAR α with FB actually inhibits the mTOR pathway (indicated by reduced pmTOR,




Fig 5. Effects of FB on the autophagy upstream pathways in PPAR α ^{+/+} and PPAR α ^{-/-} mice. Liver samples were collected after 3 weeks of FB administration for Western blotting. **(A)** Effects of FB on key proteins of the mTOR pathway: mTOR (mammalian target of rapamycin), S6K (P70S6 serine/threonine kinase), 4EBP1 (eukaryotic translation initiation factor 4E-binding protein). **(B)** Effects of FB on key proteins the insulin signaling pathway: Akt (protein kinase B) and GSK3 β (glycogen synthase kinase 3 β). * $p < 0.05$, ** $p < 0.01$ control (-FB); n.s. not statistically significant, $n = 5-6$ /group.

<https://doi.org/10.1371/journal.pone.0173676.g005>

pS6K and p4EBP1). Because mTOR can be activated by the stimulation of insulin signalling [22, 23], we examined insulin signalling and found that the activity of pAkt and pGSK3 β were decreased by chronic stimulation of PPAR α similarly to our recent reports [11, 18]. These results together indicate that PPAR α -mediated down-regulation of autophagic proteins cannot be attributed to the mTOR pathway. Such interpretation agrees with the notion that the constitutive activity of autophagy is insensitive to the mTOR pathway [10].

FoxO1 is a master transcription factor controlling the expression of autophagy proteins [3, 5, 6]. Interestingly our results showed the level of FoxO1 was markedly reduced following chronic activation of PPAR α as previously reported [24], suggesting that the down-regulation of autophagic proteins is possibly due to the inhibition of FoxO1. FoxO1 is degraded once translocated to the cytosol from the nucleus once phosphorylation and acetylation [5, 6]. Indeed, the reduction in FoxO1 protein level was associated with increases in its phosphorylation and acetylation but there was no change in the mRNA expression of FoxO1. Thus, we speculate that the reduced FoxO1 content is due to an increased degradation promoted by its phosphorylation and/or acetylation. It has been suggested that FoxO3 may also induce autophagy by controlling the transcription of LC3 and Bnip3 [25, 26]. Additional studies are warranted to investigate the role of FoxO3 in PPAR α -induced changes in the expression of autophagic proteins in the liver. However, we do not rule out the direct effect on autophagy proteins (including the degradation of autophagic proteins) from PPAR α because the




Fig 6. Effects of FB on FoxO1 and SIRT1 in PPAR α ^{+/+} and PPAR α ^{-/-} mice. Liver samples were collected after 3 weeks of FB administration. **(A)** Effects on the content, phosphorylation and acetylation (Ac) of FoxO1 the protein level. **(B)** Effects on the level of mRNA of FoxO1. **(C)** Effects on deacetylases, SIRT1 (silent mating type information regulation 2 homolog or NAD-dependent deacetylase sirtuin-1) and HDAC4 (histone deacetylases 4). * $p < 0.05$, ** $p < 0.01$ vs control (-FB); n.s. not statistically significant; n = 5–6/group.

<https://doi.org/10.1371/journal.pone.0173676.g006>

regulation of autophagy genes is multifactorial. The present study focused on the chronic effect of PPAR α activation (3 weeks) as opposed to acute effect reported by Lee et al [9] and Jiao et al. [8]. The different time points to capture the dynamic changes in autophagy during PPAR α activation may also contribute to the discrepancy.

Several recent studies suggest that an increase in the acetylation of FoxO1 may result from the inhibition of deacetylases [27, 28]. Our recent work showed that the acetylation of FoxO1 can be increased as a result of the suppression of the deacetylases SIRT1 and HDAC4 [29]. In the present study, level of SIRT1 and HDAC4 phosphorylation were reduced chronic activation of PPAR α . As deacetylase activity is a mechanism to retain FoxO1 in the nucleus of hepatocytes [30], our findings from this study suggest that the chronic activation of PPAR α may promote the acetylation FoxO1 via suppressing of SIRT1 and HDAC4 as a mechanism contributing to the degradation of FoxO1.

In summary, the present study suggests that chronic activation of PPAR α decreases the autophagic capacity in the liver by reducing multiple autophagic proteins while increasing

lipogenic proteins. These changes are independent of the PPAR α downstream mediator FGF21 and the mTOR pathway. The reduction in autophagic proteins may be due to a down-regulation due to reduction in FoxO1. The suppression of SIRT1 and HDAC4 by chronic activation of PPAR α may contribute to the acetylation of FoxO1. These findings may help explain why chronic activation of PPAR α fails to reduce hepatic steatosis in high fat fed mice despite increased FA oxidation [11, 31]. As the fibrate class drugs are chronically used in clinics, the observations from this study suggest a need to consider their potentially impact on level of autophagic proteins.

Supporting information

S1 File. Data file.
(XLSM)

Acknowledgments

The authors would like to thank Dr Xiu Zhou (RMIT University) for her assistance to the animal experiments.

Author Contributions

Conceptualization: JMY AX TJ TH.

Data curation: EJ SL QL XZ HW.

Formal analysis: EJ SL QL HW JMY.

Funding acquisition: JMY.

Investigation: EJ SL QL XZ HW.

Methodology: EJ SL QL.

Project administration: JMY AX.

Resources: JMY AX.

Supervision: JMY AX.

Validation: JMY.

Visualization: EJ HW TH JMY.

Writing – original draft: EJ TH JMY.

Writing – review & editing: XZ TH JMY.

References

1. Klionsky DJ, Abdelmohsen K, Abe A, Abedin MJ, Abeliovich H, Acevedo Arozena A, et al. Guidelines for the use and interpretation of assays for monitoring autophagy (3rd edition). *Autophagy*. 2016; 12(1):1–222. <https://doi.org/10.1080/15548627.2015.1100356> PMID: 26799652
2. Yin XM, Ding WX, Gao W. Autophagy in the liver. *Hepatology (Baltimore, Md)*. 2008; 47(5):1773–85. Epub 2008/04/09.
3. Rabinowitz JD, White E. Autophagy and metabolism. *Science (New York, NY)*. 2010; 330(6009):1344–8. Epub 2010/12/04.
4. Kim J, Kundu M, Viollet B, Guan KL. AMPK and mTOR regulate autophagy through direct phosphorylation of Ulk1. *Nature cell biology*. 2011; 13(2):132–41. Epub 2011/01/25. <https://doi.org/10.1038/ncb2152> PMID: 21258367

5. Sengupta A, Molkenin JD, Yutzey KE. FoxO transcription factors promote autophagy in cardiomyocytes. *The Journal of biological chemistry*. 2009; 284(41):28319–31. Epub 2009/08/22. <https://doi.org/10.1074/jbc.M109.024406> PMID: 19696026
6. Zhao Y, Yang J, Liao W, Liu X, Zhang H, Wang S, et al. Cytosolic FoxO1 is essential for the induction of autophagy and tumour suppressor activity. *Nature cell biology*. 2010; 12(7):665–75. Epub 2010/06/15. <https://doi.org/10.1038/ncb2069> PMID: 20543840
7. Kersten S. Integrated physiology and systems biology of PPAR α . *Molecular metabolism*. 2014; 3(4):354–71. Epub 2014/06/20. PMID: 24944896
8. Jiao M, Ren F, Zhou L, Zhang X, Zhang L, Wen T, et al. Peroxisome proliferator-activated receptor alpha activation attenuates the inflammatory response to protect the liver from acute failure by promoting the autophagy pathway. *Cell Death Dis*. 2014; 5:e1397. <https://doi.org/10.1038/cddis.2014.361> PMID: 25165883
9. Lee JM, Wagner M, Xiao R, Kim KH, Feng D, Lazar MA, et al. Nutrient-sensing nuclear receptors coordinate autophagy. *Nature*. 2014; 516(7529):112–5. <https://doi.org/10.1038/nature13961> PMID: 25383539
10. Schneider JL, Cuervo AM. Autophagy and human disease: emerging themes. *Current opinion in genetics & development*. 2014; 26:16–23. Epub 2014/06/08.
11. Chan SM, Zeng XY, Sun RQ, Jo E, Zhou X, Wang H, et al. Fenofibrate insulates diacylglycerol in lipid droplet/ER and preserves insulin signaling transduction in the liver of high fat fed mice. *Biochimica et biophysica acta*. 2015; 1852(7):1511–9. Epub 2015/04/25. <https://doi.org/10.1016/j.bbadis.2015.04.005> PMID: 25906681
12. Wang H, Sun RQ, Zeng XY, Zhou X, Li S, Jo E, et al. Restoration of autophagy alleviates hepatic ER stress and impaired insulin signalling transduction in high fructose-fed male mice. *Endocrinology*. 2015; 156(1):169–81. Epub 2014/10/25. <https://doi.org/10.1210/en.2014-1454> PMID: 25343276
13. Badman MK, Pissios P, Kennedy AR, Koukos G, Flier JS, Maratos-Flier E. Hepatic fibroblast growth factor 21 is regulated by PPAR α and is a key mediator of hepatic lipid metabolism in ketotic states. *Cell metabolism*. 2007; 5(6):426–37. Epub 2007/06/07. <https://doi.org/10.1016/j.cmet.2007.05.002> PMID: 17550778
14. Inagaki T, Dutchak P, Zhao G, Ding X, Gautron L, Parameswara V, et al. Endocrine regulation of the fasting response by PPAR α -mediated induction of fibroblast growth factor 21. *Cell metabolism*. 2007; 5(6):415–25. Epub 2007/06/07. <https://doi.org/10.1016/j.cmet.2007.05.003> PMID: 17550777
15. Liang Q, Zhong L, Zhang J, Wang Y, Bornstein SR, Triggie CR, et al. FGF21 maintains glucose homeostasis by mediating the cross talk between liver and brain during prolonged fasting. *Diabetes*. 2014; 63(12):4064–75. Epub 2014/07/16. <https://doi.org/10.2337/db14-0541> PMID: 25024372
16. Ye D, Wang Y, Li H, Jia W, Man K, Lo CM, et al. Fibroblast growth factor 21 protects against acetaminophen-induced hepatotoxicity by potentiating peroxisome proliferator-activated receptor coactivator protein-1 α -mediated antioxidant capacity in mice. *Hepatology (Baltimore, Md)*. 2014; 60(3):977–89. Epub 2014/03/05.
17. Sharara-Chami RI, Zhou Y, Ebert S, Pacak K, Ozcan U, Majzoub JA. Epinephrine deficiency results in intact glucose counter-regulation, severe hepatic steatosis and possible defective autophagy in fasting mice. *The international journal of biochemistry & cell biology*. 2012; 44(6):905–13. Epub 2012/03/13.
18. Chan SM, Sun RQ, Zeng XY, Choong ZH, Wang H, Watt MJ, et al. Activation of PPAR α ameliorates hepatic insulin resistance and steatosis in high fructose-fed mice despite increased endoplasmic reticulum stress. *Diabetes*. 2013; 62(6):2095–105. Epub 2013/01/26. <https://doi.org/10.2337/db12-1397> PMID: 23349482
19. Dreyer C, Krey G, Keller H, Givel F, Helftenbein G, Wahli W. Control of the peroxisomal beta-oxidation pathway by a novel family of nuclear hormone receptors. *Cell*. 1992; 68(5):879–87. Epub 1992/03/06. PMID: 1312391
20. Mizushima N, Yoshimori T. How to interpret LC3 immunoblotting. *Autophagy*. 2007; 3(6):542–5. Epub 2007/07/06. PMID: 17611390
21. Kim KH, Jeong YT, Oh H, Kim SH, Cho JM, Kim YN, et al. Autophagy deficiency leads to protection from obesity and insulin resistance by inducing Fgf21 as a mitokine. *Nature medicine*. 2013; 19(1):83–92. Epub 2012/12/04. <https://doi.org/10.1038/nm.3014> PMID: 23202295
22. Laplante M, Sabatini DM. mTOR signaling in growth control and disease. *Cell*. 2012; 149(2):274–93. Epub 2012/04/17. <https://doi.org/10.1016/j.cell.2012.03.017> PMID: 22500797
23. Vander Haar E, Lee SI, Bandhakavi S, Griffin TJ, Kim DH. Insulin signalling to mTOR mediated by the Akt/PKB substrate PRAS40. *Nature cell biology*. 2007; 9(3):316–23. Epub 2007/02/06. <https://doi.org/10.1038/ncb1547> PMID: 17277771

24. Qu S, Su D, Altomonte J, Kamagate A, He J, Perdomo G, et al. PPAR{alpha} mediates the hypolipidemic action of fibrates by antagonizing FoxO1. *American journal of physiology Endocrinology and metabolism*. 2007; 292(2):E421–34. Epub 2006/09/21. <https://doi.org/10.1152/ajpendo.00157.2006> PMID: 16985262
25. Mammucari C, Milan G, Romanello V, Masiere E, Rudolf R, Del Piccolo P, et al. FoxO3 controls autophagy in skeletal muscle in vivo. *Cell metabolism*. 2007; 6(6):458–71. Epub 2007/12/07. <https://doi.org/10.1016/j.cmet.2007.11.001> PMID: 18054315
26. Zhou J, Liao W, Yang J, Ma K, Li X, Wang Y, et al. FOXO3 induces FOXO1-dependent autophagy by activating the AKT1 signaling pathway. *Autophagy*. 2012; 8(12):1712–23. Epub 2012/08/31. <https://doi.org/10.4161/auto.21830> PMID: 22931788
27. Hariharan N, Maejima Y, Nakae J, Paik J, Depinho RA, Sadoshima J. Deacetylation of FoxO by Sirt1 Plays an Essential Role in Mediating Starvation-Induced Autophagy in Cardiac Myocytes. *Circulation research*. 2010; 107(12):1470–82. Epub 2010/10/16. <https://doi.org/10.1161/CIRCRESAHA.110.227371> PMID: 20947830
28. Motta MC, Divecha N, Lemieux M, Kamel C, Chen D, Gu W, et al. Mammalian SIRT1 represses forkhead transcription factors. *Cell*. 2004; 116(4):551–63. Epub 2004/02/26. PMID: 14980222
29. Zhou X, Zeng XY, Wang H, Li S, Jo E, Xue CC, et al. Hepatic FoxO1 Acetylation Is Involved in Oleanolic Acid-Induced Memory of Glycemic Control: Novel Findings from Study 2. *PloS one*. 2014; 9(9): e107231. Epub 2014/09/16. <https://doi.org/10.1371/journal.pone.0107231> PMID: 25222566
30. Frescas D, Valenti L, Accili D. Nuclear trapping of the forkhead transcription factor FoxO1 via Sirt-dependent deacetylation promotes expression of glucogenetic genes. *The Journal of biological chemistry*. 2005; 280(21):20589–95. Epub 2005/03/25. <https://doi.org/10.1074/jbc.M412357200> PMID: 15788402
31. Liu M, Montgomery MK, Fiveash CE, Osborne B, Cooney GJ, Bell-Anderson K, et al. PPARalpha-independent actions of omega-3 PUFAs contribute to their beneficial effects on adiposity and glucose homeostasis. *Sci Rep*. 2014; 4:5538. <https://doi.org/10.1038/srep05538> PMID: 24986106