



Article (refereed) - postprint

This is the peer reviewed version of the following article:

Staley, Joanna T.; Botham, Marc S.; Amy, Sam R.; Hulmes, Sarah; Pywell, Richard F. 2018. **Experimental evidence for optimal hedgerow cutting regimes for brown hairstreak butterflies.** *Insect Conservation and Diversity*, 11 (2). 213-218, which has been published in final form at <https://doi.org/10.1111/icad.12239>

This article may be used for non-commercial purposes in accordance with Wiley Terms and Conditions for Self-Archiving.

© 2017 The Royal Entomological Society

This version available <http://nora.nerc.ac.uk/517893/>

NERC has developed NORA to enable users to access research outputs wholly or partially funded by NERC. Copyright and other rights for material on this site are retained by the rights owners. Users should read the terms and conditions of use of this material at <http://nora.nerc.ac.uk/policies.html#access>

This document is the author's final manuscript version of the journal article, incorporating any revisions agreed during the peer review process. There may be differences between this and the publisher's version. You are advised to consult the publisher's version if you wish to cite from this article.

The definitive version is available at <http://onlinelibrary.wiley.com/>

Contact CEH NORA team at
noraceh@ceh.ac.uk

1

2

3

4

5 **Experimental evidence for optimal hedgerow cutting regimes for Brown hairstreak**
6 **butterflies**

7

8

9 Joanna T. Staley, Marc S. Botham, Sam R. Amy, Sarah Hulmes, Richard F. Pywell

10

11

12 NERC Centre for Ecology & Hydrology, Maclean Building, Benson Lane, Crowmarsh Gifford,
13 Wallingford, Oxfordshire OX10 8BB, UK

14

15 *Corresponding author: jnasta@ceh.ac.uk

16

17 **Running title:** Hedge management for Brown hairstreaks

18

19 **Keywords**

20 *Thecla betulae*; agri-environment scheme; blackthorn; Countryside Stewardship; Environmental
21 Stewardship; *Prunus spinosa*;

22 **Abstract**

23 1. The Brown hairstreak butterfly has declined in range and abundance over the past fifty
24 years, leading to designated conservation status in several European countries including England
25 and Wales. The Brown hairstreak's decline has been linked to changes in hedgerow
26 management, based on mortality of eggs over winter and female oviposition preferences.

27 2. We assessed Brown hairstreak egg abundance in late winter over four years in response
28 to hedgerow management treatments to manipulate the frequency, timing and the intensity of
29 trimming (reduced intensity resulting in an annual increase of approximately 10cm in height and
30 width), using a field experiment with a randomized block design.

31 3. Hedgerow plots cut every year to a standard height and width had the lowest Brown
32 hairstreak egg abundance; this is the most common hedgerow management outside agri-
33 environment schemes (AES). Cutting hedgerow plots at a reduced intensity nearly doubled the
34 number of surviving eggs in late winter. Plots cut at a reduced frequency in autumn (once every
35 three years), which forms part of current English AES, had 1.3 times more eggs than those cut
36 annually.

37 4. Current AES management prescriptions are likely to benefit the Brown hairstreak, but its
38 requirements need to be balanced with those of other taxa in relation to the timing of hedgerow
39 cutting. Cutting hedges at a reduced intensity has previously been shown to benefit the wider
40 Lepidoptera community as well as Brown hairstreak butterflies. Reduced intensity cutting does
41 not currently form part of AES hedgerow prescriptions, but could be considered for inclusion in
42 future schemes.

43 **Introduction**

44

45 The Brown hairstreak butterfly (*Thecla betula* L.) is a designated conservation priority species in
46 the UK and is on Red Lists in several northern European countries (de Vries et al., 2011), due to
47 reductions in its distribution and abundance over the past five decades. In the UK, there was a
48 49% decline in the number of 1km squares occupied by the Brown hairstreak between 1976 and
49 2014, and although there are some positive signs of locally increased occupancy over the last
50 decade, the abundance of Brown hairstreak continues to decrease (Fox et al., 2015). Brown
51 hairstreak females preferentially oviposit on young growth of *Prunus* species which protrudes
52 from hedges, and can result in eggs being cut off during hedge trimming (Thomas, 1974; Merckx
53 & Berwaerts, 2010). Conservation effort has largely focused on hedgerow management, as the
54 change towards annual cutting of hedges with mechanized flails since the middle of the twentieth
55 century is thought to be linked to the decline in Brown hairstreak range and abundance (Bourn &
56 Warren, 1998; Butterfly Conservation, 2013).

57

58 Hedgerow management prescriptions under agri-environment schemes (AES), which provide
59 payments to land managers in return for less frequent cutting of hedgerows, and trimming at
60 specific times of year, have considerable potential to support Brown hairstreak conservation
61 (Butterfly Conservation, 2013). AES hedgerow management prescriptions in England have had
62 high uptake over the last decade; in 2009 41% (163,712 kms) of the total length of English
63 hedgerows were managed under reduced frequency trimming regimes in the Environmental
64 Stewardship (ES) AES (Natural England, 2009). Links between hedgerow management and
65 Brown hairstreak conservation are based on its life cycle (Butterfly Conservation, 2013),

66 evidence of preference for young foliage for oviposition (Fartmann & Timmerman, 2006), and
67 high egg mortality measured following a single cutting event with a mechanised flail at a site in
68 Surrey (Thomas, 1974). Here, we measured Brown hairstreak egg abundance on hedgerows in
69 late winter over four years, in a replicated field experiment. The frequency, timing and intensity
70 of hedgerow cutting were manipulated, in the context of current and potentially future AES
71 management prescriptions. Specifically, we tested the following questions in relation to the
72 abundance of Brown hairstreak eggs: 1) Is abundance greater on hedges that are not cut
73 annually?; 2) Are there fewer eggs if hedges are cut in late winter rather than autumn?; 3) Does
74 reducing the intensity of hedge trimming increase egg abundance?

75 **Materials and methods**

76

77 *Study species*

78

79 Adult Brown hairstreak butterflies fly from late July through to early October (Bourn & Warren,
80 1998), and females lay eggs on the bark of Prunus species, predominantly blackthorn (*Prunus*
81 *spinosa* L.) (Merckx & Berwaerts, 2010). Adult Brown hairstreaks often aggregate in Ash tree
82 canopies, and make both short occasional flights and longer directional flights (e.g. across one or
83 several trees; Thomas, 1974). The majority of Brown hairstreak eggs are laid on young
84 blackthorn growth or on blackthorn suckers, in a fork or at the base of a bud, and between 50 and
85 170cm above ground level (Fartmann & Timmerman, 2006; Merckx & Berwaerts, 2010). Most
86 (88%) eggs are laid singly, though groups of two, three or more can be found occasionally
87 (Merckx & Berwaerts, 2010). Brown hairstreak overwinters in the egg stage, with larvae
88 hatching in late April or May, in synchrony with budburst of its host plant (de Vries et al., 2011).
89 The larvae continue to feed until late June or early July, when they pupate on the ground among
90 leaves or in grass tussocks at the base of the hedge (Thomas & Lewington, 2010; Butterfly
91 Conservation, 2013).

92

93 *Experimental design*

94

95 The field experiment was conducted at four hedgerows on a working farm in Yacombe, Devon
96 in south-west England (50°51'N, 3°03'W). The hedgerows consisted of traditional mixed
97 woody species growing on banks, and were planted 200 – 300 years ago (Staley et al., 2016).

98 The four dominant plant species in the hedgerows were: Field maple (*Acer campestre*, average
99 cover = 31.7%); Blackthorn (average cover = 25.9%); Hawthorn (*Crataegus monogyna*, average
100 cover = 11.2%) and Hazel (*Corylus avellana*, average cover = 11.2); all other woody species
101 were present at < 6% cover. The landscape around the experimental hedgerows consisted of a
102 patchwork of hedges, small (mainly grassland) fields and small woodlands, with small patches of
103 blackthorn in the hedges and woodlands.

104

105 Three experimental treatments were applied in full factorial combination: 1) frequency of cutting
106 (once every one vs. two vs. three years); 2) timing of cutting (early autumn, September vs. late
107 winter, January / February); and 3) intensity of cutting (standard cutting to the same hedgerow
108 height and width each time vs. incrementally raising the cutter bar by approximately 10 cm each
109 time the hedge is cut, resulting in a slightly wider and taller hedge).

110

111 Treatments were applied to 15 m long contiguous hedgerow plots, replicated in three randomised
112 blocks. Two blocks each consisted of a single hedge, and the third of two parallel hedges
113 bordering the same field. In addition, each block contained a control plot that was not cut during
114 the experiment, the position of which was not randomized but located at the end of each block
115 for practical reasons (Figure 1a). Hedge cutting treatments were applied to both sides and the top
116 of each plot using tractor mounted flails, operated by the farmer who regularly cut the hedges on
117 his farm, to ensure that the cutting was representative of typical hedgerow cutting. All
118 experimental plots including the controls were cut prior to the start of the experiment in late
119 winter (January / February 2010). Hedgerow cutting treatments were applied for five years from
120 September 2010.

121

122 *Brown hairstreak egg surveys*

123

124 Egg surveys are a recognized, standardized monitoring method for Brown hairstreak butterflies,
125 as the adults are rather difficult to locate due to their lifestyle and behaviour (they are
126 predominantly canopy dwelling; Butterfly Conservation, 2013; Fartmann & Timmerman, 2006).
127 Egg surveys were carried out in February or early March each year, after the ‘late winter plots’
128 had been cut but before the blackthorn was in flower or leaf, when the pale white eggs contrasted
129 with the dark blackthorn bark (Figure 1b). All blackthorn stems and shoots in the central 10m of
130 each experimental plot were searched intensively for Brown hairstreak eggs for 20 minutes on
131 each side of each plot (40 minutes per plot). Once an egg was located it was checked with a
132 magnifying glass, to verify species identity. During the verification and recording the stopwatch
133 was paused, to ensure a consistent 40 minutes of search time per plot regardless of how many
134 eggs were found. Trials of survey techniques during the first year of monitoring showed that
135 extending the search time beyond 40 minutes per plot did not result in more eggs being spotted,
136 and that a second fieldworker did not record any extra eggs in addition to those detected by one
137 person surveying a plot over the 40 minute period. As the woody species composition of each
138 experimental plot varied, the percentage cover of blackthorn was assessed on each side of each
139 plot was assessed July 2013.

140

141 *Statistical analyses*

142

143 The effects of cutting frequency, timing, intensity and the interactions between them on
144 cumulative Brown hairstreak egg counts over four years was tested using Generalised Linear
145 Models (GLMs) with a Poisson distribution. Percentage cover of blackthorn was included as a
146 covariate in all analyses. Interactions and factors that did not contribute significantly to GLMs
147 were removed one at a time, and changes in the explanatory power of the models were tested
148 using likelihood ratio tests in a backwards selection procedure (LRT, Faraway, 2015). All
149 analyses were carried out in R version 3.0.3 (R Core Development Team, 2014) using package
150 lme4 (Pinheiro et al., 2014). Final model output is given in the Electronic Supplementary
151 Information.

152

153 **Results and Discussion**

154

155 Two hundred and thirty eggs were recorded during this study, with the number varying between
156 years (total located across experimental plots, minimum = 25 in 2013, maximum = 91 in 2015),
157 in line with numbers of eggs found at a single site in previous studies on Brown hairstreak eggs
158 (e.g. Thomas, 1974). Assessing cumulative egg abundance per plot over all four years of the
159 survey, hedges cut to allow incremental growth had on average 2.3 times more Brown hairstreak
160 eggs on average than those cut back to a standard height and width (LRT $\chi^2_2=35.0$, $P<0.001$;
161 Figure 2). There was an interaction between the frequency and timing of hedgerow cutting on
162 cumulative Brown hairstreak egg abundance (LRT $\chi^2_2=16.7$, $P<0.05$); on hedgerow plots cut in
163 autumn, there were on average 1.3 times more Brown hairstreak eggs if plots were cut once
164 every three years compared to those cut every year (for GLM final model output see electronic
165 supplementary material)..Hedges cut in autumn also had nearly twice as many eggs as those cut

166 in winter, for those plots cut less frequently than every year (1.96 times more eggs when cut in
167 autumn vs. winter for plots cut once every three years; 1.83 times if cut once every two years).
168 More eggs were found on block 3 of the experiment than blocks 1 and 2.

169

170

171 Reductions in Brown hairstreak butterfly distributions have previously been linked to changes in
172 hedgerow management regimes, largely based on anecdotal observations and autecology. The
173 current multi-year study provides the first empirical evidence demonstrating that Brown
174 hairstreak egg abundance in late winter varies with the frequency, timing and intensity of
175 hedgerow cutting. These differences in abundance could be due to egg mortality as a direct result
176 of trimming blackthorn branches containing eggs, resulting in eggs being damaged and not
177 hatching, or hatching at the base of hedges in the mulch that is left by trimming with mechanical
178 flails, without open buds and leaves to provide a food source. Alternatively, the reduced
179 abundance could be caused by female oviposition choice, or by a combination of egg mortality
180 and oviposition choice. Previous studies on Brown hairstreak oviposition preference and the
181 position of eggs suggest that the differences in abundance are largely due to egg mortality. The
182 tendency for female Brown hairstreak butterflies to lay the majority of eggs on recent blackthorn
183 growth (de Vries et al., 2011; Merckx & Berwaerts, 2010; Fartmann & Timmerman, 2006)
184 probably results in high egg mortality as a result of hedgerow trimming, as young growth often
185 protrudes the furthest and is hence most likely to be trimmed. In a study where eggs were marked
186 and counted before and after a single hedgerow cutting event, egg mortality was found to be
187 around 80% (Thomas, 1974). Moreover, Thomas (1974) showed no effect of past management

188 on Brown hairstreak oviposition preference between blackthorn branches and stems in hedges
189 that had last been cut one vs. between one and two vs. more than two years previously.
190

191 Cutting at a standard intensity every year in autumn is the most common hedge management
192 outside AES (Sparks & Croxton, 2007; Staley et al., 2012). Current AES in England (ES and the
193 recently launched Countryside Stewardship) include guidance/prescriptions to cut hedgerows
194 either once every two years between September and February, or for a higher payment either
195 once every two years in January/February or once every three years between September –
196 February (Natural England, 2013, 2015). Within the ES AES the most common prescription is
197 cutting once every two years in autumn, to the same height and width on each occasion (Natural
198 England, 2009). The lower payments associated with autumn cutting once every two years
199 within ES are based on evidence that this cutting regime provides less benefit for wildlife, for
200 example in provision of berries for overwintering birds and mammals (Staley et al., 2012) or in
201 increasing Lepidoptera abundance and diversity (Facey et al., 2014; Staley et al., 2016). In
202 contrast to the benefits of winter cutting for other taxa, here we demonstrate that cutting in
203 September results in nearly twice the number of Brown hairstreak eggs compared with cutting
204 hedgerows in February. Brown hairstreak adults often persist until late September and in some
205 years early October, and therefore may have laid eggs after the autumn plots were cut in
206 September. Over the four years of this study, autumn plots were cut between 17th and 29th
207 September. Cutting dates were compared with the last date on which Brown hairstreak adults
208 were recorded in the three counties surrounding the Yarcombe experimental site (Devon, Dorset
209 and Somerset) for each of the four years, using data from the UK Butterfly Monitoring Scheme
210 (<http://www.ukbms.org/>). Brown hairstreak adults were recorded on the wing at nearby UKBMS

211 sites on or after the autumn cutting date in three of the four years of egg monitoring. The Brown
212 hairstreak species action plan states that cutting hedges in August may be beneficial for Brown
213 hairstreak egg survival (Bourn & Warren, 1998), but here we show that cutting hedges within
214 AES in mid to late September every two to three years can also nearly double the number of
215 eggs, compared with late winter. Clearly, there is a need to balance the benefits of earlier cutting
216 in relation to Brown hairstreak against the conservation of other taxa which are likely to do better
217 under late winter cutting regimes, including the broader Lepidoptera community in hedgerows
218 (Facey et al., 2014; Staley et al., 2016).

219

220 Cutting to allow incremental growth forms part of the discretionary management advice for
221 hedgerows within the new Countryside Stewardship AES in England (Natural England, 2015).
222 The prevalence of this form of management within AES cannot be estimated as it is not part of
223 the prescribed management for any AES option. The increase in Brown hairstreak egg
224 abundance under trimming for incremental growth shown here, together with the increased
225 Lepidoptera diversity and abundance found previously (Staley et al., 2016), demonstrate that this
226 form of reduced cutting intensity has benefits across the broader Lepidoptera community that
227 utilizes hedges, and could hence be considered as part of prescribed management under future
228 AES. Butterflies are sometimes considered indicators for invertebrate diversity more broadly,
229 leading to the adoption of butterfly population data as biodiversity indicators (e.g. Fox et al.,
230 2015; Merckx et al., 2013). Trimming for incremental growth thus has the potential to benefit
231 broader invertebrate assemblages.

232

233 Over the long term, cutting to allow incremental growth would result in hedges that are taller and
234 wider. However, if landowners do not want hedgerows that are larger eventually, they have the
235 option of cutting hedgerows back to their original height and width periodically, or of
236 rejuvenating hedgerows to encourage regrowth from the base using techniques such as coppicing
237 or hedge-laying (Amy et al., 2015; Staley et al., 2015), following a period of incremental cutting
238 intensity. As well as providing food resources for Lepidoptera larvae and other invertebrates,
239 hedges benefit invertebrates in other ways, such as through the provision of shelter against
240 convective cooling for (ectothermic) species in otherwise often exposed agricultural landscapes
241 (Merckx et al., 2008). The use of a range of hedgerow management regimes to create a
242 heterogenous landscape, including supporting a range of microclimates, may also balance the
243 conservation requirements of a range of taxa (Merckx et al., 2008; Merckx et al. 2010; Oliver et
244 al., 2010).

245

246 In conclusion, here we show using a four-year study that the abundance of Brown hairstreak
247 butterfly eggs is greater on hedgerow plots that are cut less frequently, in mid to late September,
248 or at a reduced cutting intensity, compared to the standard practice of cutting hedgerows back to
249 a standard height and width every year. Hedgerow management under AES has clear potential to
250 contribute to Brown hairstreak conservation, but the potentially conflicting needs of other taxa in
251 relation to the timing of hedgerow cutting need to be considered. One possible solution for this
252 would be the inclusion of reduced intensity cutting, shown to benefit both Brown hairstreak
253 butterflies and the wider Lepidoptera community.

254 **Acknowledgements**

255 Many thanks to Nigel Adams who helped to apply the experimental treatments at Yarcombe, to
256 Lucy Hulmes, Roselle Chapman, Jo Savage, Sarah Facey and Pete Nuttal for help with Brown
257 hairstreak egg surveys, to Hannah Dean for database work and to John Pidgeon for access to his
258 hedgerows. The hedgerow management treatments were funded under Defra grant BD2114.
259 Three reviewers helped to improve the manuscript.

260

261 **References**

262 Amy, S.R., Heard, M.S., Hartley, S.E., George, C.T., Pywell, R.F., & Staley, J.T. (2015)

263 Hedgerow rejuvenation management affects invertebrate communities through changes to
264 habitat structure. *Basic and Applied Ecology*, 16, 443-451.

265 Bourn, N.A.D. & Warren, M.S. (1998) Species Action Plan. Brown hairstreak *Thecla betula*.

266 Butterfly Conservation, Wareham, Dorset, U.K.

267 Butterfly Conservation. (2013) Brown hairstreak (*Thecla betulae*) factsheet. [http://butterfly-](http://butterfly-conservation.org/files/1.brown-hairstreak-species-factsheet.pdf)
268 [conservation.org/files/1.brown-hairstreak-species-factsheet.pdf](http://butterfly-conservation.org/files/1.brown-hairstreak-species-factsheet.pdf), accessed 03/05/2016

269 de Vries, H.H., Ens, S.H., de Graaf, G., Teunissen, L., van der Velde, R., Vogelaar, L.,

270 Winterink, A., & Visser, M.E. (2011) Synchronisation of egg hatching of brown hairstreak
271 (*Thecla betulae*) and budburst of blackthorn (*Prunus spinosa*) in a warmer future. *Journal*
272 *of Insect Conservation*, 15, 311-319.

273 Facey, S.L., Botham, M.S., Heard, M.S., Pywell, R.F., & Staley, J.T. (2014) Lepidoptera

274 communities and agri-environment schemes; examining the effects of hedgerow cutting
275 regime on Lepidoptera diversity, abundance and parasitism. *Insect Conservation and*
276 *Diversity*, 7, 543-552.

277 Faraway, J.J. (2015) Extending the Linear Model with R: Generalised Linear, Mixed Effects and
278 Nonparametric Regression Models. Second edition edn. CRC Press, Taylor & Francis
279 Group, Boca Raton, Florida, USA.

280 Fartmann, T. & Timmerman, K. (2006) Where to find the eggs and how to manage the breeding
281 sites of the Brown Hairstreak (*Thecla betulae* (Linnaeus, 1758)) in Central Europe? Nota
282 Lepidoptera, 29 117-126.

283 Fox, R., Brereton, T., Asher, J., August, T.A., Botham, M.S., Bourn, N.A.D., Cruickshanks,
284 K.L., Bulman, C.R., Ellis, S., Harrower, C.A., Middlebrook, I., Noble, D.G., Powney,
285 G.D., Randle, Z., Warren, M.S., & Roy, D.B. (2015) The State of the UK's Butterflies
286 2015. Butterfly Conservation and Centre for Ecology and Hydrology, Butterfly
287 Conservation, Wareham, Dorset, UK.

288 Merckx, T., Huertas, B., Basset, Y., & Thomas, J.A. (2013) A Global Perspective on Conserving
289 Butterflies and Moths and their Habitats. In: Key Topics in Conservation Biology 2 (edited
290 by D. Macdonald & K.J. Willis), pp. 237-257. Wiley, Oxford, UK.

291 Merckx, T. & Berwaerts, K. (2010) What type of hedgerows do Brown hairstreak (*Thecla*
292 *betulae* L.) butterflies prefer? Implications for European agricultural landscape
293 conservation. Insect Conservation and Diversity, 3, 194-204.

294 Merckx, T., Feber, R., Parsons, M., Bourn, N., Townsend, M., Riordan, P., & Macdonald, D.
295 (2010) Habitat preference and mobility of *Polia bombycina*: are non-tailored agri-
296 environment schemes any good for a rare and localised species? Journal of Insect
297 Conservation, 14, 499-510.

298 Merckx, T., Van Dongen, S., Matthysen, E., & Van Dyck, H. (2008) Thermal flight budget of a
299 woodland butterfly in woodland versus agricultural landscapes: an experimental
300 assessment. *Basic and Applied Ecology*, 9, 433-442.

301 Natural England (2009) Agri-environment schemes in England 2009: a review of results and
302 effectiveness. <http://publications.naturalengland.org.uk/publication/46002> Report NE 194

303 Natural England (2013) Entry Level Stewardship - Environmental Stewardship Handbook.
304 Fourth edition. <http://publications.naturalengland.org.uk/publication/2798159>

305 Natural England (2015) Countryside Stewardship - Options and
306 Supplements. https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/447651/cs-options-supplements.pdf

307

308 Oliver, T., Roy, D.B., Hill, J.K., Brereton, T., & Thomas, C.D. (2010) Heterogeneous landscapes
309 promote population stability. *Ecology Letters*, 13, 473-484.

310 Pinheiro, J., Bates, D., DebRoy, S., Sarkar, S., & R Core Development Team (2014) nlme:
311 Linear and Nonlinear Mixed Effects Models R package version 3.1-117. [http://CRAN.R-](http://CRAN.R-project.org/package=nlme)
312 [project.org/package=nlme](http://CRAN.R-project.org/package=nlme)>. R package version 3.1-117.

313 R Core Development Team (2014) R: A Language and Environment for Statistical Computing.
314 <http://www.R-project.org/>. R Foundation for Statistical Computing, Vienna, Austria.

315 Sparks, T.H. & Croxton, P.J. (2007) The influence of timing and frequency of hedgerow cutting
316 on hawthorn flowering and berry yields: preliminary results. *Aspects of Applied Biology*,
317 82, 103-106.

318 Staley, J.T., Botham, M.S., Chapman, R.E., Amy, S.R., Heard, M.S., Hulmes, L., Savage, J., &
319 Pywell, R.F. (2016) Little and late: How reduced hedgerow cutting can benefit
320 Lepidoptera. *Agriculture, Ecosystems & Environment*, 224, 22-28.

- 321 Staley, J.T., Amy, S.R., Adams, N.P., Chapman, R.E., Peyton, J.M., & Pywell, R.F. (2015) Re-
322 structuring hedges: rejuvenation management can improve the long term quality of
323 hedgerow habitats for wildlife. *Biological Conservation*, 186, 187-196.
- 324 Staley, J.T., Sparks, T.H., Croxton, P.J., Baldock, K.C.R., Heard, M.S., Hulmes, S., Hulmes, L.,
325 Peyton, J., Amy, S.R., & Pywell, R.F. (2012) Long-term effects of hedgerow management
326 policies on resource provision for wildlife. *Biological Conservation*, 145, 24-29.
- 327 Thomas, J.A. (1974) Factors influencing the numbers and distribution of the Brown Hairstreak,
328 *Thecla betulae* L. (Lepidoptera, Lycaenidae) and the Black Hairstreak, *Strymonidia pruni*
329 L. (Lepidoptera, Lycaenidae). PhD thesis, University of Leicester, Leicester, UK.
- 330 Thomas, J. & Lewington, R. (2010) *The Butterflies of Britain and Ireland*. British Wildlife
331 Publishing, Dorset, UK.

332 **Figure legends**

333

334 Figure 1

335 a) Layout of experimental hedgerow blocks and factorial combinations of treatments
336 manipulating the frequency (once every 1 vs. 2 vs. 3 years), timing (A = autumn, September vs.
337 W = winter, January or February) and intensity (S = cut back to standard height and width vs. I =
338 incremental growth, cut to allow 10 cm of recent growth to remain on sides and top) of hedgerow
339 cutting, and a control treatment that was not cut for the duration of the experiment. B) Brown
340 hairstreak butterfly eggs at the experimental plots at Yarcombe, Devon. © Lucy Hulmes, CEH

341


342 Figure 2

343 Cumulative abundance (mean \pm SE) of Brown hairstreak eggs over four years (2012 – 2015) on
344 blackthorn in 15m long hedgerow plots subject to cutting frequency (every 1 vs. 2 vs. 3 years),
345 timing (autumn (unfilled) vs. winter (striped)) and intensity (standard (a) vs. incremental growth
346 (b)) experimental treatments since 2010. A control treatment was not cut for the duration of the
347 experiment.

348

349

Figure 1



Plot number	Treatment number	Hedgerow cutting treatment frequency, timing, intensity
1	4	1, A, I
2	2	2, A, S
3	3	3, A, S
4	12	3, W, I
5	7	1, W, S
6	5	2, A, I
7	11	2, W, I
8	1	1, A, S
9	9	3, W, S
10	10	1, W, I
11	8	2, W, S
12	6	3, A, I
13	13	Control





Figure 2



Electronic Supplementary Information ICDIV-16-0232 R1: Statistical output for Generalised Linear Model analysis of cumulative number of Brown hairstreak eggs

```
> summary(m10a)
```

Call:

```
glm(formula = Lep_Thecla_betulae_eggs ~ blockBHcumNC +
  cutfreqBHcumNC +
  cuttimeBHcumNC + cutintBHcumNC +
  cutfreqBHcumNC:cuttimeBHcumNC,
  family = quasipoisson)
```

Deviance Residuals:

	Min	1Q	Median	3Q	Max
	-2.561	-1.428	-0.055	0.607	3.048

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	1.317	0.380	3.468	0.002	**
blockBHcumNC2	0.058	0.304	0.192	0.849	
blockBHcumNC3	0.806	0.262	3.080	0.005	**
cutfreqBHcumNC2	0.649	0.396	1.638	0.113	
cutfreqBHcumNC3	0.872	0.382	2.280	0.031	*
cuttimeBHcumNCw	0.578	0.401	1.441	0.161	
cutintBHcumNCs	-0.836	0.229	-3.658	0.001	**
cutfreqBHcumNC2:cuttimeBHcumNCw	-1.184	0.560	-2.115	0.044	*
cutfreqBHcumNC3:cuttimeBHcumNCw	-1.253	0.537	-2.333	0.027	*

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

(Dispersion parameter for quasipoisson family taken to be 2.370382)

Null deviance: 155.504 on 35 degrees of freedom
 Residual deviance: 66.145 on 27 degrees of freedom
 AIC: NA