



ELSEVIER

Available online at www.sciencedirect.com

ScienceDirect

Current Opinion in
Plant Biology

Stomatal development: focusing on the grasses

Christopher Hepworth¹, Robert S Caine², Emily L Harrison²,
Jennifer Sloan^{1,2} and Julie E Gray²

The development and patterning of stomata in the plant epidermis has emerged as an ideal system for studying fundamental plant developmental processes. Over the past twenty years most studies of stomata have used the model dicotyledonous plant *Arabidopsis thaliana*. However, cultivated monocotyledonous grass (or Gramineae) varieties provide the majority of human nutrition, and future research into grass stomata could be of critical importance for improving food security. Recent studies using *Brachypodium distachyon*, *Hordeum vulgare* (barley) and *Oryza sativa* (rice) have led to the identification of the core transcriptional regulators essential for stomatal initiation and progression in grasses, and begun to unravel the role of secretory signaling peptides in controlling stomatal developmental. This review revisits how stomatal developmental unfolds in grasses, and identifies key ontogenetic steps for which knowledge of the underpinning molecular mechanisms remains outstanding.

Addresses

¹ Department of Animal and Plant Sciences, University of Sheffield, S10 2TN, UK² Department of Molecular Biology and Biotechnology, University of Sheffield, S10 2TN, UKCorresponding author: Hepworth, Christopher
(c.hepworth@sheffield.ac.uk)**Current Opinion in Plant Biology** 2018, **41**:1–7This review comes from a themed issue on **Growth and development**Edited by **Gwyneth Ingram** and **Ari Pekka Mähönen**<http://dx.doi.org/10.1016/j.pbi.2017.07.009>1369-5266/© 2017 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Introduction

Stomata function as the interface between plants and atmosphere, exerting control over gaseous diffusion and balancing the uptake of carbon dioxide with the loss of water vapour [1]. Regulation of stomatal development is of critical importance in allowing plants to adjust their gaseous exchange to suit the prevailing environmental conditions [2–4]. Stomatal development has been extensively studied, and has emerged as an excellent system for investigating cell-fate specification and cellular differentiation [5,6**]. The distribution of stomata on the leaf


surface is a highly regulated process with a level of plasticity, and components regulating stomatal development continue to be identified [7,8]. Much of our current understanding stems from work conducted on the model dicot *Arabidopsis thaliana* and many comprehensive reviews are available [9,10].

Although cereal grasses provide the majority of human nutrition we still know surprisingly little about their stomata. As scientific focus moves towards the engineering of ‘climate ready crops’ that will be better suited to predicted warmer, drier, higher carbon dioxide environments, understanding the regulatory mechanisms of grass stomatal development and patterning could prove key to future success. In this review, we outline recent advances emerging from studies of grasses and discuss the outstanding questions.

The grass stomatal lineage

The development of stomatal complexes in grasses differs to that of the dicots in a number of ways. Most notably, grass stomata are formed from dumbbell-shaped guard cells (GCs) that are flanked by subsidiary cells (SC) which develop in parallel rows within defined and specific epidermal cell files. In contrast, the GCs of dicots are kidney-shaped and form stomata that are scattered throughout the epidermis in a less orderly pattern. In nascent leaves, grass stomatal development occurs along a spatiotemporal gradient with the earliest stages occurring basally, and proceeding as cells move upwards as the leaf expands [11,12]. This developmental pathway can be broken down into 6 stages and is illustrated using barley (*Hordeum vulgare*) in Figure 1. Initially, close to the leaf base, prior to stomatal-lineage cell specification, potential precursor cells proliferate in particular files (Stage 1). As undifferentiated cells are pushed further up the leaf blade alternate cells enter the stomatal development pathway via an asymmetric ‘entry’ division leading to a smaller guard mother cell (GMC) and a larger sister cell (Stage 2). Cells from files on either side of a newly formed GMC then also divide asymmetrically to form subsidiary mother cells (SMCs) (Stage 3). After the cells have increased in size, mature GMCs are flanked by two nascent SCs (Stage 4), a final symmetric division of the GMC leads to the formation of two immature GCs (Stage 5). The stomatal complex matures and expands to form a pair of dumbbell-shaped GCs, which separate to form the stomatal pore (Stage 6). Thus, each mature grass stomatal complex includes a central pore, a dumbbell-shaped GC pair and two flanking SCs. Each complex overlies an airspace,

Figure 1



Six stages of grass stomatal development. **(1)** Selection of stomatal lineage cells within defined rows. **(2)** Asymmetric entry divisions generate smaller guard mother cells (GMCs), depicted by white arrows, and larger epidermal cells. **(3)** GMCs then expand and laterally induce subsidiary mother cell (SMC) formation (see white arrows) via asymmetric divisions. **(4)** Subsidiary cell maturation. **(5)** GMCs divide symmetrically. **(6)** GMC elongation and maturation to form the guard cell (GC) complex. All confocal images were taken from the base of leaf 2 of 6-day-old barley seedlings (cv. Golden Promise) stained with propidium iodide. Scale bar = 5 μm .

or ‘sub-stomatal cavity’, which forms between the mesophyll cells of the underlying layer, to facilitate efficient gaseous diffusion in and out of the leaf. Several recent studies provide insights into the transcriptional and regulatory mechanisms underpinning grass stomatal development. These make use of grass genome sequences and build on knowledge gained from Arabidopsis.

Brachypodium: A model for recent discovery

Despite differences in morphology and patterning, the basic helix-loop-helix transcription factors underpinning stomatal fate in Arabidopsis, SPEECHLESS (SPCH), MUTE and FAMA together with heterodimeric partners

INDUCER OF CBF EXPRESSION1 (ICE1) and SCREAM2 (SCRM2) are highly conserved, with origins which predate the divergence of the mosses and hornworts from ancestral land plants [5,6^{**},13,14^{**}]. The discovery of functionally orthologous grass genes [15–18] has shed light on the mechanisms responsible for stomatal development and patterning in grasses. Liu *et al.* [12] investigated putative orthologues of SPCH, MUTE and FAMA in both rice and maize (*Zea mays*) and revealed at least one SPCH and a FAMA gene that are required for stomatal development in rice. More recently, Raissig *et al.* [14^{**}] used the wheat relative *Brachypodium distachyon* (Brachypodium), to dissect the roles of grass SPCH and

ICE/SCRM orthologues. They found that although Brachypodium uses *SPCH* and *ICE/SCRM* gene products to regulate stomatal formation, the grass pathway is 'alternatively wired' to achieve correctly patterned stomata. Specifically, a *SPCH* duplication event has occurred in grasses leading to two functional but partially redundant paralogs: *BdSPCH1* and *BdSPCH2* which both act early during stomatal development. For *ICE/SCRM* family members, a divergence of function has occurred in comparison to Arabidopsis orthologues. Rather than being functionally redundant, *BdICE1* and *BdSCRM2* control overlapping stages of stomatal development; *BdICE1* primarily functions during the initial asymmetric entry division, *BdSCRM2* acts later during the differentiation of GMCs prior to the formation of SMCs [14**]. The observation that the expression of the *BdSPCH1/2* and *BdICE1/SCRM2* genes is limited to stomatal cell files suggests that the regulation of these genes or proteins across the leaf blade is critical for the correct patterning of stomata across the leaf. How such spatial regulation is achieved is a key next line of enquiry.

The presence of flanking SCs is common to all grass stomatal complexes and these cells have long been believed to assist in altering aperture size in a timely and energy efficient manner [19]. However, despite their important role, little has been known about how SCs are developmentally programmed. Again, recent studies in Brachypodium are beginning to shed light on the area. The discovery that *BdMUTE* moves from GMCs, via plasmodesmata, into neighboring SMCs where it acts to establish SMC identity has advanced our understanding of monocot stomatal development considerably [6**]. Mutants lacking *BdMUTE* function known as *subsidiary cell identity defective (sid)* plants, produce GCs without flanking SCs. These plants have allowed researchers to test the importance of SCs in grass stomatal behaviour, for the first time. The finding that *sid* plants have reduced stomatal gas exchange and impaired growth, confirm the important role of SCs and suggests opportunities for the enhancement of stomatal aperture control and plant productivity via the targeted manipulation of SC development.

Signaling peptides regulate grass stomatal development


In concert with the bHLH transcription factors, a family of cysteine-rich cell-to-cell signaling peptides regulates the cellular divisions and cell fate transitions required for stomatal development. These epidermal patterning factors (EPFs) and their associated receptor components are well-characterised in Arabidopsis with EPF2 primarily regulating asymmetric entry divisions and EPF1 primarily overseeing the differentiation of GMCs and stomatal spacing. The EPF-like peptide known as EPFL9 or STOMAGEN positively regulates stomatal development by competing with EPF2 during early stomatal development to promote stomatal lineage cell fate [20–25].

Recently, Hughes *et al.* [26*] characterised the role of *HvEPF1*; a barley orthologue of AtEPF1/2 (Figure 2), which when ectopically expressed inhibits stomatal development. Analysis of *HvEPF1* over-expressing barley leaf epidermis revealed that many GMCs do not progress to form stomatal complexes. Moreover, high levels of expression of *HvEPF1* inhibit the asymmetric 'entry' division that produces GMCs, the maturation of GMCs, the production of SMCs and sub-stomatal cavity formation. Thus a grass signaling peptide similar in sequence to Arabidopsis EPF1 and 2 is able to prevent GMC formation and cause the arrest of GMC development prior to SMC generation but how *HvEPF1* functions at normal endogenous levels remains to be investigated. One potential function given the large number of arrested GMCs devoid of SCs, is that *HvEPF1* primarily downregulates *HvSPCH* protein levels thereby preventing GMCs from proceeding further through the stomatal lineage. Whether *HvEPF1* activity directly or indirectly regulates the *HvMUTE* gene or protein or other targets downstream of *HvSPCH* is intriguing area for future study.

The severe reductions in stomatal frequency and gas exchange brought about by increasing *HvEPF1* levels led to improved barley drought tolerance and water use efficiency. Any reduced capacity for photosynthesis did not impact on grain production under either well-watered or drought conditions. These results suggest promising routes for cereal crop improvement through stomatal density manipulation. Our knowledge of grass EPF/L function is further extended by a study describing the use of gene editing techniques to knock-out a rice orthologue of Arabidopsis *EPFL9*, *OsEPFL9a* (Figure 2) causing up to 8-fold reductions in stomatal density [27*]. These barley and rice *EPF/L* studies confirm that, as in Arabidopsis, both positive and negative stomatal development regulators are active in grasses.

Phylogenetic and functional analyses suggest that in addition to the stomatal bHLH transcription factors and the epidermal patterning factors, their cognate receptor components TMM and ERRECTA family, are almost certainly also conserved throughout land plants [28,29]. This provides a strong indication that a conserved functional stomatal development module exists in the grasses. However, whilst the evidence is clear that a number of EPF/L peptides are conserved between dicots and monocots [13,26*], the specifics relating to how each function in grasses is not clear. As several grass genome sequences are now accessible, we revisit the EPF/L story in grasses (Figure 2). Almost all of the grass genomes that we surveyed encode two peptides which cluster closely with Arabidopsis EPF1 and EPF2 stomatal regulators. The exception is wheat (*Triticum aestivum*) which being hexaploid has multiple orthologues of both *EPF1* and *EPF2*. Strikingly, our analysis reveals that for EPFL9, a

Figure 2



Phylogenetic analysis of EPIDERMAL PATTERNING FACTOR (EPF) and EPF-like (EPFL) peptides in grass species. Sequences were obtained via BLAST searches of peptides encoded by the *Hordeum vulgare* (barley), *Triticum aestivum* (wheat), *Brachypodium distachyon* and *Oryza sativa* (rice) genome sequences using Phytozome v12. Additional sequences from BLAST searches of the genomes of *Arabidopsis thaliana*, *Selaginella moellendorffii* and *Physcomitrella patens* are included to provide evolutionary context. All amino acid sequences with a BLAST score of at least 60 against *Arabidopsis* AtEPF1 (AT2G20875.1) and all sequences derived from *Arabidopsis* EPFL9 (AT4G12970.1 or STOMAGEN) BLAST searches were used for the subsequent alignment of retrieved sequences. EPFL6 is included to illustrate relatedness of EPF and EPFL peptides. SmEPF1-1 and SmEPF1-2 sequence information was taken from [28]. Three other sequences virtually identical to HvEPF1 (HORVU2Hr1G116030.1, HORVU2Hr1G116040.1 and HORVU2Hr1G116070.1) were omitted as they are assumed to be annotation errors and are not present when other barley genome browsers are interrogated. The evolutionary history was inferred using the Neighbor-Joining method [30]. The optimal tree with the sum of branch length = 4.77465428 is shown. The percentage of replicate trees in which the associated taxa clustered together in the bootstrap test (1000 replicates) are shown next to the branches [31]. The tree is drawn to scale, with branch lengths in the same units as those of the evolutionary distances used to infer the phylogenetic tree. The evolutionary distances were computed using the Poisson correction method [32] and are in the units of the number of amino acid substitutions per site. The analysis involved 33 amino acid sequences. All positions containing gaps and missing data were eliminated. There were a total of 38 positions in the final dataset. Evolutionary analyses were conducted in MEGA7 [33].

gene duplication event has occurred in the grasses leading to at least two distinct *EPFL9*-like genes in all species surveyed.


Having identified at least two *EPF1/EPF2* genes and two *EPFL9* equivalents (Figure 2), the next question is ‘Do all of the identified EPF/L genes encode peptides that regulate stomatal development, and if so, how do they facilitate communication between developing stomatal lineage cells?’ It is clear that grass stomatal complexes are formed by two distinct types of asymmetric divisions (which form the GMC and SMCs) and that the bHLH transcription factors regulating these divisions have to

some degree functionally diversified from *Arabidopsis* [6^{••},14^{••}]. It remains unknown whether EPF signaling peptides evolved in parallel to bHLH transcription factors to regulate SC development in grasses. Clearly, further functional studies of the potential regulators of grass stomatal formation identified here (Figure 2) and elsewhere [28,34,35] are required to further decipher stomatal development and patterning in grasses.

Complexity of stomatal patterning in grasses

Whilst in *Arabidopsis* the development of stomata is possible in most parts of nascent leaves this is not the case in grasses. As grass leaves grow and increase in width

Figure 3



The complexities of inter-file and intra-file patterning of grass stomatal precursors. **(a)** Rice stomatal lineage cells developing in neighboring stomatal files. Guard mother cells (GMCs) formed from asymmetric entry divisions can be seen (white arrows) developing in close proximity to GMCs flanked by subsidiary cells (SCs). **(b)** Barley stomatal lineage cells developing in adjacent rows. GMCs without SCs can be seen (white arrows) developing in close proximity to more advanced stomatal lineage complexes where subsidiary mother cells asymmetric divisions or symmetric GMC divisions are occurring. **(c)** Rice stomatal file with different stage stomatal lineage cells forming in a non-linear order from leaf base to tip. White arrows highlight the more mature developmental stage. **(d)** Barley stomatal file highlighting the non-linearity of stomatal development from leaf base to tip. White arrows highlight the more mature developmental stage. All images were generated using confocal microscopy from 6-day-old rice (cv. IR64) and barley (cv. Golden Promise) seedlings stained with propidium iodide. Scale bar = 10 μm .

more stomatal and non-stomatal files must form. How these are specified remains unknown. Our observations of developing leaves in rice and barley seedlings suggest that in the earlier forming leaves this process is dynamic with files containing stomatal lineage cells at differing developmental stages occurring in close proximity (Figure 3a and b). It is not uncommon to concurrently observe GMCs flanked by nascent SCs in one stomatal file and more recently formed GMCs without SCs in an adjacent file (white arrows, Figure 3a), or symmetrically dividing GMCs with flanking in close proximity to nascent GMCs where SCs have yet to form (white arrows, Figure 3b). With EPF/L peptides known to be important in regulating stomatal lineage cell placement in *Arabidopsis* [25], it will be interesting to learn to what extent their control extends both within and between stomatal files in grasses.

Our observations also revealed that the linearity of grass stomatal development within files of stomata is not always continuous and that earlier staged cells can occasionally form further from the leaf base than more advanced stomatal structures (Figure 3c and d). This suggests that the stomatal development module in grasses must not only be fluid between cell files but also within a file. We are yet to determine the importance of EPF/L peptide function in enabling such patterning.

Next steps

We have begun to gain insights into how grasses regulate the production of stomata. However, a number of fundamental questions remain unanswered. Most notably, ‘What are the regulatory switch(es) that specify which epidermal files will produce stomata during early leaf development?’ and ‘How is the regular spacing of stomata

within files achieved whilst also maintaining the development and spacing of SCs?'. Learning the answers to such questions could facilitate the generation of more refined cereal crop cultivars that are better suited to the predicted future climate, or increased frequency of severe weather events. For example, by increasing the number of stomata in grasses and or altering stomatal performance, we may be able to increase the photosynthetic potential of plants [35]. Moreover, increases in stomatal number could lead to transpirational water flux that may be beneficial in aiding root development, and nutrient uptake [36–39]. Conversely, by reducing stomatal number we should be able to improve soil water retention, drought tolerance and water use efficiency [26*].

Acknowledgements

We apologize to authors not cited due to space limitations, and thank the BBSRC Newton Rice Research Fund for funding.

References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Lawson T, Blatt M: **Stomatal size, speed and responsiveness impact on photosynthesis and water use efficiency.** *Plant Physiol* 2014, **164**:1556-1570.
 2. Casson S, Gray JE: **Influence of environmental factors on stomatal development.** *New Phytol* 2008, **178**:9-23.
 3. Hamanishi ET, Thomas BR, Campbell MM: **Drought induces alterations in the stomatal development program in *Populus*.** *J Exp Bot* 2012, **63**:4959-4971.
 4. Tricker PJ, Gibbins JG, Rodríguez López CM, Hadley P, Wilkinson MJ: **Low relative humidity triggers RNA-directed de novo DNA methylation and suppression of genes controlling stomatal development.** *J Exp Bot* 2012, **63**:3799-3813.
 5. Chater C, Caine RS, Fleming AJ, Gray JE: **Origins and evolution of stomatal development.** *Plant Physiol* 2017, **174**:624-638.
 6. Raissig MT, Matos JL, Gil MX, Kornfeld A, Bettadapur A, Abrash E, Allison HR, Badgley G, Vogel JP, Berry JA *et al.*: **Mobile MUTE specifies subsidiary cells to build physiologically improved grass stomata.** *Science* 2017, **355**:1215-1218.
- Transcription factor *MUTE* (an ortholog of Arabidopsis *MUTE*) was identified as being essential for the development of subsidiary cells in grasses. Interestingly, *MUTE* was able to move from guard mother cells, via the plasmodesmata, to flanking cells before specifying subsidiary cell formation.
7. Engineer CB, Ghassemian M, Anderson JC, Peck SC, Hu H, Schroeder JI: **Carbonic anhydrases, EPF2 and a novel protease mediate CO₂ control of stomatal development.** *Nature* 2014, **513**:246-250.
 8. Klermund C, Ranftl QL, Diener J, Bastakis E, Richter R, Schwachheimer C: **LLM-domain B-GATA transcription factors promote stomata development downstream from light signaling in *Arabidopsis thaliana* hypocotyls.** *Plant Cell* 2016, **28**:646-660.
 9. Han S-K, Torii KU: **Lineage-specific stem cells, signals and asymmetries during stomatal development.** *Development* 2016, **143**:1259-1270.
 10. Simmons AR, Bergmann DC: **Transcriptional control of cell fate in the stomatal lineage.** *Curr Opin Plant Biol* 2016, **29**:1-8.
 11. Stebbins GL, Jain SK: **Developmental studies of cell differentiation in the epidermis of monocotyledons: I. Allium, rhoeo, and commelina.** *Dev Biol* 1960, **2**:409-426.
 12. Liu T, Ohashi-Ito K, Bergmann DC: **Orthologs of *Arabidopsis thaliana* stomatal bHLH genes and regulation of stomatal development in grasses.** *Development* 2009, **136**:2265-2276.
 13. Liu F, Jensen CR, Shahanzari A, Andersen MN, Jacobsen S-E: **ABA regulated stomatal control and photosynthetic water use efficiency of potato (*Solanum tuberosum* L.) during progressive soil drying.** *Plant Sci* 2005, **168**:831-836.
 14. Raissig MT, Abrash E, Bettadapur A, Vogel JP, Bergmann DC: **Grasses use an alternatively wired bHLH transcription factor network to establish stomatal identity.** *Proc Natl Acad Sci U S A* 2016, **113**:8326-8331.
- Experiments demonstrating that the same core transcriptional regulators control stomatal initiation in both Arabidopsis and grasses. However, the specific structure, function and regulation of the gene families have diverged.
15. Ohashi-Ito K, Bergmann DC: **Arabidopsis FAMA controls the final proliferation/differentiation switch during stomatal development.** *Plant Cell* 2006, **18**:2493-2505.
 16. MacAlister CA, Ohashi-Ito K, Bergmann DC: **Transcription factor control of asymmetric cell divisions that establish the stomatal lineage.** *Nature* 2007, **445**:537-540.
 17. Pillitteri LJ, Torii KU: **Breaking the silence: three bHLH proteins direct cell-fate decisions during stomatal development.** *BioEssays* 2007, **29**:861-870.
 18. Kanaoka MM, Pillitteri LJ, Fujii H, Yoshida Y, Bogenschutz NL, Takabayashi J, Zhu JK, Torii KU: **SCREAM/ICE1 and SCREAM2 specify three cell-state transitional steps leading to arabidopsis stomatal differentiation.** *Plant Cell* 2008, **20**:1775-1785.
 19. Chen Z-H, Chen G, Dai F, Wang Y, Hills A, Ruan Y-L, Zhang G, Franks PJ, Nevo E, Blatt MR: **Molecular evolution of grass stomata.** *Trends Plant Sci* 2017, **22**:124-139.
 20. Hara K, Yokoo T, Kajita R, Onishi T, Yahata S, Peterson KM, Torii KU, Kakimoto T: **Epidermal cell density is autoregulated via a secretory peptide, EPIDERMAL PATTERNING FACTOR 2 in Arabidopsis leaves.** *Plant Cell Physiol* 2009, **50**:1019-1031.
 21. Hunt L, Gray JE: **The signaling peptide EPF2 controls asymmetric cell divisions during stomatal development.** *Curr Biol* 2009, **19**:864-869.
 22. Hunt L, Bailey KJ, Gray JE: **The signalling peptide EPFL9 is a positive regulator of stomatal development.** *New Phytol* 2010, **186**:609-614.
 23. Sugano SS, Shimada T, Imai Y, Okawa K, Tamai A, Mori M, Hara-Nishimura I: **Stomagen positively regulates stomatal density in Arabidopsis.** *Nature* 2010, **463**:241-244.
 24. Le J, Zou J, Yang K, Wang M: **Signaling to stomatal initiation and cell division.** *Front Plant Sci* 2014, **5**:297.
 25. Qi X, Han SK, Dang JH, Garrick JM, Ito M, Hofstetter AK, Torii KU: **Autocrine regulation of stomatal differentiation potential by EPF1 and ERECTA-LIKE1 ligand-receptor signaling.** *Elife* 2017 <http://dx.doi.org/10.7554/elife.24102>.
 26. Hughes J, Hepworth C, Dutton C, Dunn JA, Hunt L, Stephens J, Cameron D, Waugh R, Gray JE: **Reducing stomatal density in barley improves drought tolerance without impacting on yield.** *Plant Physiol* 2017, **174**:776-787.
- A functional barley orthologue of Arabidopsis EPF1 (*HvEPF1*) is described. Overexpression of *HvEPF1* led to reduced stomatal formation, predominantly by the prevention of guard mother cell maturation. Subsequent physiological analysis revealed reducing stomatal density could improve drought tolerance without a deleterious effect on yield.
27. Yin X, Biswal AK, Dionora J, Perdigon KM, Balahadia CP, Mazumdar S, Chater C, Lin HC, Coe RA, Kretzschmar T *et al.*: **CRISPR-Cas9 and CRISPR-Cpf1 mediated targeting of a stomatal developmental gene EPFL9 in rice.** *Plant Cell Rep* 2017, **36**:745-757.
- Through the use of CRISPR/Cas9 technology the authors mutated the stomatal regulatory gene *OseEPFL9* which is related to Arabidopsis *EPFL9*/*STOMAGEN*. This led to an 8-fold reduction in stomatal density. This finding suggests a role for positive signaling peptides in grass stomatal development.
28. Caine RS, Chater CC, Kamisugi Y, Cuming AC, Beerling DJ, Gray JE, Fleming AJ: **An ancestral stomatal patterning module**

- revealed in the non-vascular land plant *Physcomitrella patens*. *Development* 2016, **143**:3306-3314.
29. Chater CC, Caine RS, Tomek M, Wallace S, Kamisugi Y, Cuming AC, Lang D, MacAlister CA, Casson S, Bergmann DC et al.: **Origin and function of stomata in the moss *Physcomitrella patens***. *Nat Plants* 2016, **2**:16179.
 30. Saitou N, Nei M: **The neighbor-joining method: a new method for reconstructing phylogenetic trees**. *Mol Biol Evol* 1987, **4**:406-425.
 31. Felsenstein J: **Phylogenies and the comparative method**. *Am Nat* 1985, **125**:1-15.
 32. Zuckerkandl E, Pauling L: **Molecules as documents of evolutionary history**. *J Theor Biol* 1965, **8**:357-366.
 33. Kumar S, Stecher G, Tamura K: **MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets**. *Mol Biol Evol* 2016, **33**:1870-1874.
 34. Peterson KM, Rychel AL, Torii KU: **Out of the mouths of plants: the molecular basis of the evolution and diversity of stomatal development**. *Plant Cell* 2010, **22**:296-306.
 35. Tanaka N, Kato M, Tomioka R, Kurata R, Fukao Y, Aoyama T, Maeshima M: **Characteristics of a root hair-less line of *Arabidopsis thaliana* under physiological stresses**. *J Exp Bot* 2014, **65**:1497-1512.
 36. Matimati I, Verboom GA, Cramer MD: **Nitrogen regulation of transpiration controls mass-flow acquisition of nutrients**. *J Exp Bot* 2014, **65**:159-168.
 37. Franks PJ, Doheny-Adams TW, Britton-Harper ZJ, Gray JE: **Increasing water-use efficiency directly through genetic manipulation of stomatal density**. *New Phytol* 2015, **207**:188-195.
 38. Hepworth C, Turner C, Landim MG, Cameron D, Gray JE: **Balancing water uptake and loss through the coordinated regulation of stomatal and root development**. *PLoS One* 2016, **11**:e0156930.
 39. Hepworth C, Doheny-Adams T, Hunt L, Cameron DD, Gray JE: **Manipulating stomatal density enhances drought tolerance without deleterious effect on nutrient uptake**. *New Phytol* 2015, **208**:336-341.