



van Dijk, L., Hayton, A., Main, D., Booth, A., King, A., Barrett, D., ...
Reyher, K. (2016). Participatory Policy Making by Dairy Producers to
Reduce Anti-Microbial use on Farms. *Zoonoses and Public Health*.
<https://doi.org/10.1111/zph.12329>

Peer reviewed version

Link to published version (if available):
[10.1111/zph.12329](https://doi.org/10.1111/zph.12329)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the accepted author manuscript (AAM). The final published version (version of record) is available online via Wiley at DOI 10.1111/zph.12329. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/pure/about/ebr-terms>



**PARTICIPATORY POLICY MAKING BY DAIRY PRODUCERS TO
REDUCE ANTIMICROBIAL USE ON FARMS**

Journal:	<i>Zoonoses and Public Health</i>
Manuscript ID	ZPH-Jun-16-181.R2
Manuscript Type:	Original Article
Date Submitted by the Author:	n/a
Complete List of Authors:	van Dijk , Lisa ; University of Bristol, School of Veterinary Sciences, University of Bristol, Langford House, Langford, Bristol, Reyher, Kristen ; University of Bristol, School of Veterinary Sciences, University of Bristol, Langford House, Langford, Bristol, Main , David ; University of Bristol, School of Veterinary Sciences, University of Bristol, Langford House, Langford, Bristol, Barrett, David ; University of Bristol, School of Veterinary Sciences, University of Bristol, Langford House, Langford, Bristol, Buller, Henry ; University of Exeter College of Life and Environmental Sciences Hayton, Alastair; RAFT Solutions Ltd. King, Andrew; Coombe Farm Booth, Andrew; WDFarmers Ltd.
Subject Area:	Antimicrobial resistance, Dairy cattle, Veterinary medicine

SCHOLARONE™
Manuscripts



1
2
3 **Lisa van Dijk, University of Bristol, United Kingdom.**

4
5 **PARTICIPATORY POLICY MAKING BY DAIRY PRODUCERS TO REDUCE**
6 **ANTIMICROBIAL USE ON FARMS**

7
8
9 **L. van Dijk***, **A. Hayton[†]**, **D. C. J. Main***, **A. Booth[§]**, **A. King[#]**, **D. C. Barrett***, **H. J. Buller[‡]**, and
10 **K. K. Reyher***

11
12 **University of Bristol, School of Veterinary Sciences, Langford House, Langford,*
13 *Bristol, BS40 5DU, United Kingdom.*

14
15 *[†]RAFT Solutions Ltd, West Hill Barns, Evershot,*
16 *DT2 0LD, United Kingdom*

17
18 *[‡]College of Life and Environmental Sciences, University of Exeter, Amory Building, Rennes Drive,*
19 *Exeter, EX4 4RJ, United Kingdom.*

20
21 *[§]WD Farmers Ltd, Mary Street House, Mary Street, Taunton, TA1 3NW, United Kingdom*

22
23 *[#]Coombe Farm, Roundham, Crewkerne, TA18 8RR, United Kingdom*

24 *lisa.vandijk@bristol.ac.uk*

25
26 **Impacts**

- 27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
- Antimicrobials are widely used in food-producing animals, posing a potential risk of spread of antimicrobial resistant bacteria from animals to humans. Frequent use of certain antimicrobials in food animals may reduce their effectiveness as treatments for both animals and humans.
 - Intensive collaboration and dialogue between dairy producers, veterinarians, industry partners and researchers led to the development of an antimicrobial stewardship policy including practical measures designed to deliver real, on-farm changes in antimicrobial use, while maintaining or improving dairy herd health and welfare.
 - Although many producers were already stringent in their antimicrobial use (many were organic dairy farmers), almost 60% envisaged making changes to their use of antimicrobial medicines on farm as a result of their participation in the policy development process.

45
46 **Summary**

47
48 Pressures for more responsible use of antimicrobial (AM) medicines in food animals are likely to
49 increase from policy makers and the food industry, including retailers. To address this challenge,
50 participatory approaches to welfare interventions and disease prevention may also be necessary
51 alongside more conventional regulatory measures.
52
53
54
55
56
57
58
59
60

1
2
3 This paper describes the process of enabling groups of dairy producers to use a participatory policy-
4 making (PPM) approach to develop an AM stewardship policy. The policy includes measures agreed
5 to by all producers for more responsible use of AMs, while maintaining or improving dairy herd health
6 and welfare.
7
8
9

10
11 This process provided a unique opportunity for collaboration and dialogue between producers,
12 veterinarians, industry and researchers. Its participatory nature encouraged comprehensive learning for
13 all involved. This integration of science with producers' knowledge and experience led to credible and
14 practical recommendations designed to deliver real and lasting change in AM use.
15
16
17

18
19 The multidisciplinary nature of this research marks a significant contribution to embedding social
20 science skills and approaches into the veterinary sphere. As an initial step in creating better
21 understanding of how participatory approaches with farmers can be applied in a UK context and more
22 widely, this work serves as a pilot for promoting more responsible use of veterinary medicines in other
23 livestock species.
24
25
26
27
28
29

30 31 **Key words**

32 Antimicrobial resistance; public health; participation; antimicrobial stewardship policy; dairy
33 production; veterinary medicine use.
34

35 36 **Introduction**

37
38 The increase of antimicrobial resistance (AMR) is a growing problem for human health and an
39 increasing economic burden for society (Laxminarayan *et al.*, 2013). Antimicrobials (AMs) are widely
40 used in food-producing animals (O'Neill, 2015) and this has led to growing concern. There is a risk of
41 spread of AMR bacteria from animals to humans, and the frequent use of certain AMs in food animals
42 may reduce their effectiveness for treatment in humans (EFSA, 2015). Increasing resistance also has
43 an adverse effect on the livestock sector, leading to more recalcitrant bacterial infections in food-
44 producing animals, increased costs of treatment, productivity losses and animal welfare concerns
45 (McEwen & Fedorka-Cray, 2002).
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 High use of AMs in food-producing animals is often associated with intensive pig and poultry
4 production, and use of AMs in dairy cattle may not be seen as an immediate concern (Raymond *et al.*,
5 2006; Brunton *et al.*, 2012). However, the use of AMs, including those of major importance to treat
6 serious infections in humans (fluoroquinolones, 3rd and 4th generation cephalosporins and long-acting
7 macrolides; WHO, 2012), are commonplace in dairy systems (Raymond *et al.*, 2006; Brunton *et al.*,
8 2012). The majority of AMs in dairy cattle are administered for the treatment of mastitis during
9 lactation and at cessation of lactation, dry cow therapy (Swinkels *et al.*, 2015). A survey of AM usage
10 on dairy farms in the United Kingdom (UK) revealed that antibiotic tubes used for treating mastitis
11 frequently contain 4th generation cephalosporins, and that a large number of dairy producers fed their
12 calves waste milk from cows with mastitis that were potentially undergoing treatment with these AM
13 families (Brunton *et al.*, 2012).
14
15
16
17
18
19
20
21
22
23
24
25
26

27 The UK currently has no governmental regulations to reduce veterinary AM use similar to those
28 introduced in Netherlands in 2010, which required farms to reduce veterinary antibiotic use by 50% by
29 2013 and 70% by 2015 (as compared with the amounts that had been used in 2009; Speksnijder *et al.*,
30 2015b). The only UK regulatory measures are voluntary measures to ensure best practice and
31 responsible use through farm assurance schemes and compliance to retailer standards for meat and
32 animal products, as described in the UK Five Year AM Strategy 2013 to 2018 (Davies & Gibbens,
33 2013). Several major retailers and milk buyers in the UK have or are in the process of setting their
34 own guidelines regarding the use of AMs. For example, one milk buyer recently introduced a
35 requirement for their suppliers to use selective rather than universal dry cow therapy, and some
36 retailers are already monitoring the use of 3rd and 4th generation cephalosporins and fluoroquinolones
37 (Alderton, 2015). Recently, the Responsible Use of Medicines in Agriculture (RUMA) Alliance -
38 consisting of 24 organizations along the value chain, including farmer representative organizations,
39 retailers, veterinary practices, farm assurance organizations and animal welfare organizations -
40 published guidelines for more responsible use of AMs by cattle farmers (RUMA, 2015).
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Pressures to reduce AM use and to use AM medicines more responsibly in food-producing animals are
4 likely to increase from policy makers and the food industry, including retailers. To address this
5 challenge, participatory approaches to welfare interventions and disease prevention may also be
6 necessary to achieve sustainable farm practice change in addition to more conventional regulatory
7 methods (Buller *et al.*, 2015). Participatory approaches - in particular participatory policy-making
8 (**PPM**) - offers innovative solutions to the AMR challenge and fills the void in public policy making
9 regarding this challenge. These approaches shift the view of farmers as merely policy implementers to
10 viewing farmers as active partners in collaborative decision making and joint learning processes with
11 other stakeholders to identify and develop solutions together (Schut *et al.*, 2014; Sutton, 1999). Karl
12 (2002) states “*PPM implies empowerment of stakeholders to take part in the whole cycle of the policy*
13 *process: formulation, implementation, monitoring and evaluation of policy*”. Solutions developed by
14 farmers through these approaches often lead to on-farm practice change as they are more practical and
15 implementable than solutions developed by outsiders (Pretty, 1995; Chambers *et al.*, 1989). Moreover,
16 they allow farmers to maintain control and a sense of ownership over the solutions (Escobar & Buller,
17 2014). Evidence of successful PPM can be found in a number of areas, from citizen participation in
18 local governance to environment management (Van den Hove, 2000).
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36

37 This paper describes a policy development process in which two groups of dairy producers and their
38 veterinarians were actively encouraged and enabled to work together to develop their own AM
39 stewardship policy leading to changes in farming practice and reduced AM use.
40
41
42
43
44

45 **Materials and methods**

46
47
48 The policy development process was initiated by two dairy producer groups, a group of 25 organic
49 producers and a group of 48 conventional producers. These two groups exclusively supply one major
50 retailer in the UK. From the outset, the process was a focused initiative driven by producers as a
51 collective, and all producers in the two groups participated in the policy development process to
52 ensure ownership of the policy. The producer groups aimed to develop an enactable on-farm AM
53
54
55
56
57
58
59
60

1
2
3 stewardship policy that showed their commitment and responsibility to the milk buyer and retailer to
4 reduce and rationalise AM use without compromising animal health and welfare.
5
6
7

8
9 The leaders of the producer groups and their lead veterinary advisor initiated the formation of a project
10 team which also included a representative from the retailer and milk buyer and researchers. A
11 facilitator was contracted to guide the policy development process (Figure 1). A series of meetings and
12 workshops were held over the period May to August, 2015 to conceive and develop the policy. The
13 first team meeting was held in May, 2015 to identify policy objectives and policy areas, to clarify
14 specific terms (e.g. What do we mean by responsible use?) and to reach agreement on how to engage
15 all producers in the policy making process (Figure 1, Step 3). Taking workshop costs, group size as
16 well as travel time for producers to the workshop into consideration, it was agreed to organize four
17 regional workshop, with each workshop comprising 20 to 25 producers and their veterinarians. The
18 project team developed an outline for the regional workshops and structured the main discussions
19 around on-farm measures for more responsible use of AMs within four categories (policy principles),
20 based on a conceptual model derived from Speksnijder and colleagues (2015a): 1. disease
21 management strategies, 2. ensuring correct use of medicines, 3. avoiding prophylactic use and 4.
22 ensuring quality data recording and use. These categories provided an initial framework for policy
23 development.
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40

41 The regional workshops (Figure 1, Step 4) were key in the development of the policy, and were held in
42 June, 2015. Each regional workshop began with a presentation by the lead veterinary advisors to
43 provide insights into AMR and why it mattered to farmers, the dairy industry and the world. As part of
44 this presentation, a survey was conducted amongst producers and veterinarians using an interactive
45 audience response system (Turning Point 5, Turning Technologies, Belfast, UK) to pose multiple-
46 choice questions to which the attendees could respond. The answers to these questions were
47 amalgamated in order to better understand the perception of producers and their veterinarians on the
48 use of AMs and AMR as well as to raise their awareness on the issue and compare and contrast the
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3 different levels of awareness represented. These question sessions were followed by short focus group
4
5 discussions to create a better understanding of the benefits and challenges of current AM use on farms.
6
7 The main part of the workshop consisted of group discussions to identify practical measures that could
8
9 ensure responsible (and often reduced) use of AMs on farm in each category. Workshop participants
10
11 were divided into smaller groups, with producers and veterinarians kept in separate groups to ensure
12
13 producers felt comfortable to actively participate in the discussion (Morgan, 1996) as well as to
14
15 manage veterinary-client communication dynamics (Shaw, 2004) and thus reduce bias in responses.
16
17 Participants were requested to discuss and write down practical measures on sticky note paper (one
18
19 strategy per sticky note). After exhaustion of new ideas, each group was asked to rank the measures
20
21 they had identified according to ease of implementation on farms. Each group presented their
22
23 measures in the four categories, and all sticky notes (per category) were then combined. General
24
25 plenary discussion followed with both producers and veterinarians in order to further develop and
26
27 agree on the measures presented. The facilitator consolidated the outcomes of the four workshops, and
28
29 the project team met in July, 2015 to draft an initial policy document. Further feedback on this initial
30
31 document was achieved through a follow-up meeting at the beginning of August, 2015 with a
32
33 representation of producers from each region to refine the policy (Figure 1, Steps 5 and 6).
34
35
36

37 Figure 1. Outline of the process followed to develop an antimicrobial stewardship policy with dairy
38
39 farmers in the UK
40
41
42

43 After final review by the lead veterinarian and producer group leaders at the end of August, 2015
44
45 (Figure 1, Step 7), the policy was communicated by email to all producers in the two producer groups
46
47 (Figure 1, Step 8), and policy implementation tools were developed (Figure 1, Step 9). The two groups
48
49 then began the policy implementation (Figure 1, Step 10).
50
51
52

53 Results

54
55
56
57
58
59
60

1
2
3 The turnout at all regional workshops was high: 97 participants (70 producers out of a total of 73 in
4 the two producers groups) and 27 veterinarians, from all practices servicing the farms participated.
5
6
7

8 **Perception of producers and their veterinarians on the use of AMs and AMR**

9
10 Producers and their veterinarian were responsive and aware of the problem of AMR. When asked
11 whether they agreed or disagreed with the statement that “*AMR is just a fashionable topic and a*
12 *passing fad*”, 80% (76/95) disagreed with this statement.
13
14
15
16

17
18
19 In terms of antibiotic use, 79% (75/95) agreed with the statement that antibiotic use in the UK
20 livestock sector is too high. Asking specifically about whether they perceived that there were AMR
21 bacteria on dairy farms, 46% (44/97) of the respondents estimated the percentage of UK dairy farms
22 with AMR bacteria as 25 % or less (Figure 2), whilst 23% (22/97) of the respondents estimated the
23 percentage of UK farms with AMR bacteria as 100%. Narrowing down whether producers thought
24 there was AMR on their own farm, 41% (36/87) of the producers and their veterinarians thought there
25 was mild AMR on their farms, and 36% (31/87) thought there was moderate AMR on their farms
26 (none 1%, very limited 18%, and severe 3%).
27
28
29
30
31
32
33
34
35
36

37 Figure 2. Participants’ response to the question to estimate the presence of AMR bacteria on UK dairy
38 farms
39
40
41
42
43

44 In the fight against AMR, producers and veterinarians indicated that they see the role of veterinarians
45 as being somewhat more important than the role of producers (Figure 3). In response to a question
46 about the importance of the role of the producer in the fight against AMR, 70% (68/97) of respondents
47 indicated that the role of farmer was very important. This is compared with the response to a similar
48 question about the importance of the role of the veterinarian in the fight against AMR, where 87%
49 (84/97) of the respondents indicated the role of the veterinarian was very important
50
51
52
53
54
55
56
57
58
59
60

1
2
3 Figure 3. Participants' response to the question about the importance of the role of the producer and
4 the veterinarian in the fight against antimicrobial resistance
5
6

7
8 An important question for the future success of the policy development process was whether the
9 producers felt able to reduce the amount of AMs they used on their farm whilst also maintaining the
10 same production levels and health of their herds. 73% (71/97) of respondents indicated they felt they
11 were able to reduce the amount of AMs on their farms.
12
13
14
15

16
17
18 In the third and fourth regional workshops, a specific question was added. Producers and the
19 veterinarians were asked whether or not they felt they could specifically reduce the use of AMs of
20 major importance to treat serious infections in humans; 53% (27/51) indicated they could.
21
22
23

24 25 26 **Antimicrobial Stewardship Policy**

27
28 A conceptual model originally developed by Speksnijder and colleagues (2015a) was adapted by the
29 project team to provide a structure (core principles) around which the final stewardship policy was
30 developed:
31
32

33
34 Principle 1: Disease reduction strategies

35
36 Principle 2: Ensuring correct use of medicines

37
38 Principle 3: Avoiding prophylactic use

39
40 Principle 4: Encouraging quality data recording and use
41

42
43 Based on the strategies discussed during the workshops, policy-specific and practical measures were
44 developed (Table 1). Practical measures were organised under the following policy statements, which
45 were developed during subsequent meetings with the producer representatives after the regional
46 workshops. These policy measures adhered to the above-mentioned principles, and were stated in
47 active forms.
48
49
50

- 51
52 • All producers use animal production practices that reduce, and, where possible, eliminate the
53 need for AM therapies.
54
- 55
56 • All producers work with their veterinarians to assess and address disease risk regularly.
57
58
59
60

- On all farms, all staff engaged in using antibiotics use them responsibly or are adequately supervised by staff who are able to do so.
- On all farms, the use of antibiotics to prevent disease is minimalised.
- On all farms, unified data is collected and used to benchmark and compare medicine use within and between farms in order to work towards further reductions in AM use.

The term ‘responsible use’ was clarified to reflect an individual as being able to identify the correct animal to treat (only treating infections), as well as identifying the correct product to use at the correct dose (amount) and correct duration (including initiating treatment at the right time).

Table 1. Policy measures - Antimicrobial Stewardship Policy, Year 1

Whilst developing the policy, it became clear that, within the producer groups there was only limited information available as to the levels of AM use on farms. Hence, some of the policy measures could only be defined if more detailed information on current use was available. Thus it was decided that the measures agreed on would initially only apply to the first year of implementation, and that the policy would be updated and reviewed on a yearly basis as the understanding of use on farms develops and scientific evidence on the subject grows.

Participation in the policy development process

Evaluations at the end of the workshops indicated a consistently positive response from participants (Figure 4). Comments provided on the *benefits* of participating in the workshop included:

“made me think about antibiotic use more”

“knowing the opinions of my fellow producers”

“ideas for moving our practice forward, using drug usage as part of the Herd Health Plan”

“helped develop policies to advantage all parties”

“helped make policy that works for the farmer”

Figure 4. Participant’s response to regional workshop evaluation questions

1
2
3 (note: Question 4 was left blank by some respondents as this question was not relevant for
4 veterinarians.)
5
6
7

8
9 Although generally very positive about the workshops, participants also felt that the development of
10 such a policy was difficult and mentally taxing. Scoring their experience from 1 (not at all
11 challenging) to 10 (very challenging), over 42% (33/78) of participants gave a score of 8 and higher as
12 to whether they felt participation in this process was challenging. More than half of producers (58%;
13 45/78) gave a score of 8 or more as to whether they envisaged making any changes to their use of AM
14 medicines on farm as a result of their participation in the workshops. Changes mentioned by producers
15 and veterinarians included:
16
17

18
19
20
21
22
23 *“more targeted approaches”*

24 *“further reduction in use of cephalosporin, more selective dry cow therapy”*

25
26
27 *“selective dry cow therapy, bacteria testing on farm”*

28
29 *“better storage and disposal of partly used AMs”*

30
31 *“reduce use of Naxcel (ceftiofur, a 3rd generation cephalosporin)*

32
33 *“more prevention”*

34
35
36 *“stop using Marbocyl (marbofloxacin, a fluoroquinolone) and Cobactan (cefquinome, a 4th*
37 *generation cephalosporin) injectables”*

38
39 *“changing dry cow therapy, reducing 3rd and 4th generation AMs ”*
40
41
42

43 Discussion

44
45
46 This is the first known application of PPM to livestock health policy. The policy development process
47 provided a unique opportunity for intensive collaboration and dialogue between producers,
48 veterinarians, industry partners and researchers. The participatory process provided comprehensive
49 learning for all involved, and allowed for the integration of science and the producers' own knowledge
50 and experience. The process led to the development of credible and practical recommendations
51 designed to deliver real on-farm changes in antibiotic use. Achieving genuine and effective change in
52
53
54
55
56
57
58
59
60

1
2
3 AM use depends upon the collaborative engagement of farmers, veterinarians, and other actors along
4
5 the value chain, including milk buyers and retailers (Buller et al., 2015).
6
7

8
9 Although evidence of substantive increases in AMR bacteria in the dairy sector is currently limited
10
11 (EFSA, 2015; Landers *et al.*, 2012), three quarters of producers (and their veterinarians) in this survey
12
13 thought that they had mild to moderate levels of AMR on their farm. This result brings up a number of
14
15 other questions: On what evidence is this claim being made? Are farmers and veterinarians
16
17 experiencing reduced effectiveness of AMs on their farms, or is their perception of this issue driven by
18
19 media or by the concern of the retailer they supply?
20
21

22
23 In developing the practical policy, there was a tension between the producers' commitment to set
24
25 common and consistent measures as a group and their individual desire to retain personal autonomy of
26
27 action. Producers were willing to commit to the policy measures; however, how each measure was to
28
29 be implemented and achieved on farm needed to be farm-specific and in the hands of the individual
30
31 producer. As a result of this tension, the measures adopted were less prescriptive in nature than, for
32
33 example, the RUMA guidelines. Being less prescriptive should not necessarily mean less effective,
34
35 however, as Biggs and colleagues (2016) have shown less prescriptive and more bespoke criteria to
36
37 select cows for dry cow therapy on dairy farms can lead to better results in terms of reduced AM use.
38
39 Comparing the adopted policy measures in the current study with the RUMA guidelines revealed that
40
41 most aspects indicated in the guidelines had been included. Those specific issues which were not
42
43 included were nevertheless discussed within the workshops. For example, keeping closed herds (where
44
45 no introductions of new animals are made in order to ensure biosecurity) was tabled, but was not
46
47 considered possible for all producers, particularly those with organic herds.
48
49

50
51
52 Discussions over what measures to include and how best to achieve these measures was challenging at
53
54 times. One particular area of difficulty for farmers was maintaining a coherent and correct policy of
55
56 medicine use when staffing was often inconsistent and changing (particularly with the use of relief
57
58 milkers or casual workers). Contrary to human health where those who administer medicines need to
59
60

1
2
3 have a licence or specific training, in the farming sectors of the UK, AMs prescribed by the
4 veterinarian are largely administered by producers and farm staff. Producers indicated they were able
5 to ensure their permanent staff had the appropriate qualification and training relevant to administering
6 AMs, however this was seen as much more of a challenge when using temporary relief workers. Thus,
7 making certain that the correct animal is treated, with the right dose, at the right time and right
8 duration becomes more difficult. The producers agreed to overcome this by defining clear
9 responsibilities regarding AM use and having clear AM decision trees with protocols for applicable
10 diseases in the relevant languages for the nationalities of the relief workers.
11
12
13
14
15
16
17
18
19

20
21 Although recording AM use on farms is required by law (DEFRA, 2012), producers in the UK have
22 few methods to aggregate these records, therefore little comprehensive data is available which relates
23 to AM use on individual or group levels. For this reason, there is at present no clear understanding of
24 AM use per individual animal on UK farms. Medicine data recording is a mandatory requirement in
25 the Royal Society for the Prevention of Cruelty to Animals and Soil Association standards as part of
26 farm assurance in the Herd Health Plan (Hemsworth *et al.*, 2009). However, differences in recording
27 practice, little agreement on choosing specific numerators or denominators, and the absence of a single
28 universal recording system used by all group members was a limitation. During the policy
29 development process, producers realised that to be able to set realistic reduction targets (Key
30 Performance Indicators) and to benchmark AM use well, they would have to gain a better
31 understanding of how medicines were being used on both individual and group levels. Also, currently
32 in the UK, there are few methods of systematically integrating farm management and veterinary data.
33 The producer groups therefore agreed that in the first year of policy implementation, the focus would
34 first be to collate and use all on-farm data for analysis.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50

51
52 Through the four regional workshops, all producers in the two groups participated in the process. It is
53 anticipated, however, that engaging all producers of a particular group in the PPM process might not
54 always be practical. Initial discussions have now been held with sheep and beef producer groups
55 supplying the same retailer to implement a similar policy development process. Working with these
56
57
58
59
60

1
2
3 groups will bring a new set of challenges, as the number of producers are much larger and more
4
5 geographically diverse. Other challenges in terms of transferability of the PPM approach relate to the
6
7 efficacy of producers. The belief of producers in their own ability to make a change is essential for the
8
9 success of this process. The participating dairy producers felt they had the power and autonomy to
10
11 make changes on their farms; this, however, might not be the case in all livestock sectors nor in all
12
13 countries. Particularly in the poultry and pig sectors, producers are less autonomous and might not feel
14
15 empowered enough to make this process a success. Further work using the PPM approach in different
16
17 contexts and settings is necessary to explore transferability of the approach.
18
19

20 21 **Concluding Remarks**

22
23 This research provides insight on how to enable groups of dairy producers to conceive, design and
24
25 implement AM stewardship policies to address the challenge of more responsible use of medicines on
26
27 farms, while maintaining or improving dairy herd health and welfare. As an initial step in creating
28
29 better understanding of how participatory approaches with producers can be applied in a UK context
30
31 and more widely, this work serves as a pilot for promoting more responsible use of AMs in other
32
33 livestock species.
34
35
36
37

38 **Acknowledgements**

39
40 We would like to thank all producers who are members of WD Farmers and producers in the Coombe
41
42 Farm organic milk producer groups, along with their veterinarians, who so actively participated in the
43
44 policy workshops. We also would like to thank Paula Baker, Gwen Rees and Esther Jones for their
45
46 assistance with the workshops.
47
48
49
50

51 **References**

52
53
54 Alderton, S., 2015: News standards for Arla producers focuses on antibiotics use. *Farmers Weekly*,
55 Available at: <http://www.fwi.co.uk/livestock/new-standards-for-arla-producers-focuses-on-antibiotics-use.htm>.
56
57
58
59
60

- 1
2
3 Biggs, A., D. Barrett, A. Bradley, M. Green, K. Reyher, and R. Zadoks, 2016: Antibiotic dry cow
4 therapy: where next? *Veterinary Record*, **178**, 93-94.
- 5 Brunton, L., D. Duncan, N. Coldham, L. Snow, and J. Jones, 2012: A survey of antimicrobial usage on
6 dairy farms and waste milk feeding practices in England and Wales. *The Veterinary record*,
7 **171**, 296-296.
- 8 Buller, H., S. Hinchliffe, J. Hockenhull, D. Barrett, K. Reyher, A. Butterworth, and C. Heath, 2015:
9 Systematic review and social research to further understanding of current practice in the
10 context of using antimicrobials in livestock farming and to inform appropriate interventions
11 to reduce antimicrobial resistance within the livestock sector. Department of Environment,
12 Food and Rural Affairs. Report DO0558.
- 13 Chambers, R., A. Pacey, and L.A. Thrupp, 1989: Farmer First: Farmer Innovation and Agricultural
14 Research. Intermediate Technology Publications, London.
- 15 Davies, S., and N. Gibbens, 2013: UK Five Year Antimicrobial Resistance Strategy 2013 to 2018.
16 Department of Health, London.
- 17 DEFRA, 2012: Beef cattle and dairy cows: health regulations. Guidance on the main diseases that
18 affect cattle, disease prevention and legal controls in place to protect cattle health: Record-
19 keeping. Available at: <https://www.gov.uk/guidance/cattle-health#record-keeping>.
- 20 EFSA, 2015: ECDC/EFSA/EMA first joint report on the integrated analysis of the consumption of
21 antimicrobial agents and occurrence of antimicrobial resistance in bacteria from humans and
22 food-producing animals. *EFSA Journal*, **13**, 4006.
- 23 Escobar, M.P., and H. Buller, 2014: Projecting Social Science into Defra's Animal Welfare Evidence
24 Base: A review of current research and evidence gaps on the issue of farmer behaviour with
25 respect to animal welfare. Department for Environment, Food and Rural Affairs, United
26 Kingdom.
- 27 Hemsworth, P., J. Barnett, and G. Coleman, 2009: The integration of human-animal relations into
28 animal welfare monitoring schemes. *Animal Welfare*, **18**, 335-345.
- 29 Karl, M., 2002: Participatory Policy Reform from a Sustainable Livelihoods Perspective. Food and
30 Agriculture Organisation, Rome.
- 31 Landers, T.F., B. Cohen, T.E. Wittum, and E.L. Larson, 2012: A review of antibiotic use in food animals:
32 perspective, policy, and potential. *Public health reports*, 4-22.
- 33 Laxminarayan, R., A. Duse, C. Wattal, A.K. Zaidi, H.F. Wertheim, N. Sumpradit, E. Vlieghe, G.L. Hara,
34 I.M. Gould, and H. Goossens, 2013: Antibiotic resistance—the need for global solutions. *The
35 Lancet infectious diseases*, **13**, 1057-1098.
- 36 McEwen, S.A., and P.J. Fedorka-Cray, 2002: Antimicrobial use and resistance in animals. *Clinical
37 Infectious Diseases*, **34**, S93-S106.
- 38 Morgan, D.L., 1996: Focus groups as qualitative research (Vol. 16). *Sage publications*
- 39 O'Neill, J., 2015: Antimicrobials in agriculture and the environment: reducing unnecessary use and
40 waste. The Review on Antimicrobial Resistance
- 41 Pretty, J.N., 1995: Participatory learning for sustainable agriculture. *World Development*, **23**, 1247-
42 1263.
- 43 Raymond, M., R. Wohrle, and D. Call, 2006: Assessment and promotion of judicious antibiotic use on
44 dairy farms in Washington State. *Journal of dairy science*, **89**, 3228-3240.
- 45 Responsible Use of Medicines in Agriculture Alliance (RUMA), 2015: Responsible use of
46 antimicrobials in cattle production. Available at: [http://www.ruma.org.uk/cattle/responsible-
47 use-antimicrobials-dairy-beef-cattle-production/](http://www.ruma.org.uk/cattle/responsible-use-antimicrobials-dairy-beef-cattle-production/)
- 48 Schut, M., J. Rodenburg, L. Klerkx, A. van Ast, and L. Bastiaans, 2014: Systems approaches to
49 innovation in crop protection. A systematic literature review. *Crop Protection*, **56**, 98-108.
- 50 Shaw, J.R., C.L. Adams, and B.N. Bonnett, 2004: What can veterinarians learn from studies of
51 physician-patient communication about veterinarian-client-patient communication?. *Journal
52 of the American Veterinary Medical Association*, **224**(5), 676-684.
- 53
54
55
56
57
58
59
60

- 1
2
3 Speksnijder, D.C., A. Jaarsma, A. Gugten, T. Verheij, and J. Wagenaar, 2015a: Determinants
4 Associated with Veterinary Antimicrobial Prescribing in Farm Animals in the Netherlands: A
5 Qualitative Study. *Zoonoses and public health*, **62**, 39-51.
- 6 Speksnijder, D.C., D.J. Mevius, C.J.M. Bruschke, and J.A. Wagenaar, 2015b: Reduction of Veterinary
7 Antimicrobial Use in the Netherlands. The Dutch Success Model. *Zoonoses and Public Health*,
8 **62**, 79-87.
- 9 Sutton, R., 1999: The policy process: an overview. Overseas Development Institute, London.
- 10 Swinkels, J., A. Hilkens, V. Zoche-Golob, V. Krömker, M. Buddiger, J. Jansen, and T. Lam, 2015: Social
11 influences on the duration of antibiotic treatment of clinical mastitis in dairy cows. *Journal of*
12 *dairy science*, **98**, 2369-2380.
- 13 Van den Hove, S., 2000: Participatory approaches to environmental policy-making: the European
14 Commission Climate Policy Process as a case study. *Ecological Economics*, **33**, 457-472.
- 15 World Health Organization (WHO), 2012: Critically important antimicrobials for human medicine.
16 WHO Advisory Group on Integrated Surveillance of Antimicrobial Resistance, Geneva,
17 Zwitserland.
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Outline of the process followed to develop an antimicrobial stewardship policy with dairy farmers in the UK

Figure 1.

164x266mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2. Participants' response to the question to estimate the presence of AMR bacteria on UK dairy farms
Figure 2.
258x169mm (300 x 300 DPI)

Manuscript Only




Figure 3. Participants' response to the question about the importance of the role of the producer and the veterinarian in the fight against antimicrobial resistance

Figure 3

258x168mm (300 x 300 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 4. Participant’s response to regional workshop evaluation questions (note: Question 4 was left blank by some respondents as this question was not relevant for veterinarians.)

Figure 4
249x99mm (300 x 300 DPI)

Review Only

<i>Policy measures AM Stewardship Policy, Year 1</i>	
Aim	Policy measures
<i>Principle 1: Disease reduction strategies</i>	
On all farms, animal production practices are used that reduce and, where possible, eliminate the need for AM therapies.	All farms are required to engage in at least two events annually to share best practice amongst the producers groups concerning environmental and livestock management to reduce disease. At the end of the first year of implementation of this policy, Key Performance Indicators (KPI's) will be set based on data collected in Year 1 to review and benchmark AM use on individual farms. Support will be provided to farms to improve farm environments and/or husbandry practice if they do not meet the threshold, in order to further reduce AM use.
All producers work with their veterinarians to assess and address disease risk regularly.	All farms make proactive use of Herd Health Plans (HHP); these are regularly reviewed with their veterinarian (minimum of twice a year) and updated, specifically containing measures to reduce the need for AMs. These measures must be reported to the management offices.
<i>Principle 2: Ensuring correct use of medicines</i>	
On all farms, all staff engaged in using antibiotics use them responsibly or are adequately supervised by staff who are able to do so.	On all farms, clear responsibilities for staff (based on their experience, qualifications and training) are defined regarding AM use. Records (including training records) are kept for all staff to provide evidence of their ability. Each farm has an AM medicine use decision tree along with protocols for applicable diseases in relevant languages (dependent on nationality of its staff); these are displayed and discussed with their veterinarian. The decision tree includes aspects such as how to handle recurrent infections and culling, how to follow withdrawal periods, and use of non-AM treatments that help farm workers to reserve antibiotics for use as a last resort. Both producers and veterinarians work to reduce, if not avoid, the use of antibiotics classified as "critically important" for human health. Based on an initial understanding of the use of these antibiotics within the producers groups, KPI's will be set for reduction of these antibiotics in Year 2.
<i>Principle 3: Avoiding prophylactic use</i>	
On all farms, the use of antibiotics to prevent disease is minimized.	All farm use selected/targeted dry cow therapy, following consultation with and direct advice from their veterinarian. All herds perform at least six-weekly somatic cell counting on all milking cows and utilise these data in decision-making about lactating and dry cow therapy. Milk under antibiotic withdrawal is not fed to calves. Each farm's HHP includes a "temp and treat" plan for calves with pneumonia.
<i>Principle 4: Encouraging quality data recording and use</i>	
On all farms, unified data is collected and used to benchmark and compare medicine use within and between farms in order to work towards further reductions in AM use.	On all farms, simple and clear records of AM use (self-reporting of use) are kept and reviewed by the farmer and veterinarian regularly (at least twice yearly, preferably at the same time as the HHP review). All veterinarians prescribing or administering AMs provide clear reasons and records to justify use. The veterinary practice responsible for supply of medicines to a farm that is part of the producers groups must make full and transparent data of all AMs supplied available to that farm at least twice yearly. Across the two producer groups, data is compiled on AM use in order to clearly understand its use on farms within the producer groups and for further implementation of measures to reduce use in Year 2.

Table 1. Policy measures - Antimicrobial Stewardship Policy, Year 1
Table 1
265x157mm (300 x 300 DPI)