



Silvera, A., Knowles, T., Butterworth, A., Berckmans, D., Vranken, E., & Blokhuis, HJ. (2017). Lameness assessment with automatic monitoring of activity in commercial broiler flocks. *Poultry Science*, 96(7), 2013–2017. [pex083]. <https://doi.org/10.3382/ps/pex023>

Peer reviewed version

Link to published version (if available):
[10.3382/ps/pex023](https://doi.org/10.3382/ps/pex023)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Oxford University Press at <https://academic.oup.com/ps/article-lookup/doi/10.3382/ps/pex023>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/pure/about/ebr-terms>

LAMENESS ASSESSMENT WITH AUTOMATIC MONITORING OF ACTIVITY IN COMMERCIAL BROILER FLOCKS

A.M. Silvera*, T.G. Knowles‡, A. Butterworth‡, D. Berckmans‡, E. Vranken‡§, and H.J.

Blokhuis*

**Swedish University of Agricultural Sciences, Animal Environment and Health PO Box 7068 Uppsala, SE-750 07, †University of Bristol, School of Veterinary Sciences, Langford, UK BS40 5DU. ‡KU Leuven, Department of Biosystems, Division M3-BIORES: Measure, Model & Manage of Bioresponses, Kasteelpark Arenberg 30, 3001 Heverlee, Belgium. §FANCOM B.V., Research Department, P.O. Box 7131, 5980 AC Panningen, The Netherlands.*

CORRESPONDING AUTHOR:

Anna Silvera, Mail: anna@silvera.nu, Phone: +4670-6616614

Swedish University of Agricultural Sciences, Animal Environment and Health PO Box 7068
Uppsala, SE SE-750 07, SWEDEN

1 **ABSTRACT** The possibility of using automatic recordings of broiler chicken activity in
2 commercial flocks to assess the birds' walking ability (lameness) was investigated. Data were
3 collected from five commercial broiler farms in four European countries, using 16 flocks and
4 33 assessment occasions. Lameness was assessed using established gait scoring (GS)
5 methods (Kestin et al. 1992, Welfare Quality 2009) and took place at 3, 4 and 5 weeks of age.
6 Gait score (GS) was used to assess the birds walking ability and automatic recordings of bird
7 activity were collected using the eYeNamic™-camera system before, during and after an
8 assessor walked through the house. The variables used to predict the level of GS extracted
9 from the camera system were: baseline activity, time from assessor leaving the house to
10 resumption of baseline activity, average activity over that period, and Δ Amplitude
11 (difference between highest activity peak after assessor left the house and baseline level).
12 Age (< 0.001) and Δ Amplitude ($p = 0.0002$) were significantly related to GS, with the gait
13 getting poorer with increased age and Δ Amplitude decreasing with declining walking ability.
14 Both measures are thus included in a predictive equation. The results demonstrate a potential
15 method using image analysis techniques to realize an automated assessment of the level of
16 lameness in commercial broiler flocks. This could be of use in future animal welfare
17 assessment schemes.

18 **Key words:** gait score, precision livestock farming, image analysis, welfare

19

20

INTRODUCTION

21 Broiler chickens, reared for meat production, have for decades been genetically selected for
22 rapid growth and high meat yield and are reared intensively in large flocks with high stocking
23 densities (Knowles et al. 2008). One consequence of this approach is an increased risk of
24 lameness and impaired walking ability in the flock (Bradshaw et al. 2002). The breeding
25 companies have noted the problem and 25 years of selection to improve the leg health have

26 been successful (Kapell et al. 2012), but still it is important to monitor the leg health in
27 broiler flocks. Leg problems may have metabolic, developmental or infectious causes
28 (Butterworth and Haslam, 2009) and can impair the welfare of affected broilers (Bradshaw et
29 al. 2002). Welfare is compromised because lame birds have problems accessing feed and
30 water (Bessei, 2006) and they can experience pain (McGeown et al., 1999; Danbury et al.
31 2000; Caplen et al., 2013). Lameness is also negatively related to final weight at slaughter
32 (Butterworth and Haslam 2009) and may be associated with high flock mortality (Wideman
33 et al., 2012). Several methods have been developed to assess lameness in broiler flocks. To
34 determine the exact pathological cause of lameness, a post-mortem examination of the bird is
35 needed (Mench, 2004). Non-invasive methods include the Latency to Lie (Weeks et al. 2002)
36 and the Gait Scoring-method (Welfare Quality®, 2009, Kestin et al. 1992). In the former
37 method, the bird's latency (delay) before it lies down is measured when birds are placed in a
38 water-proof test pen with the floor covered with tepid water to depth of 30 mm; severe
39 lameness results in short latencies, whereas birds with good leg health will remain standing
40 for longer. This method provides an objective measurement of lameness but has time and
41 resource constraints that limit the number of birds which can be tested. The 'manual' Gait
42 Score-method is probably the most widely adopted; in this method, the bird's walking ability
43 is graded between 0 (perfect walking) to 5 (not able to move). This observational method
44 enables a large number of birds to be assessed during the same inspection, but it has been
45 criticized for being subjective, having poor reliability between observers (Mench, 2004) and
46 being costly in terms of the time, and hence the man hours required. Moreover the risk for
47 disease transfers when observers visit livestock houses should be reduced to a minimum.
48 Developing assessment methods that can be conducted on-farm by the caretaker during daily
49 routines can be one solution to above mentioned issues, but can also reduce the stressful
50 event of a un-known human enclosing and handling the birds (Marchewka et al, 2013).

51

52 Another approach that is a time-saving, continuous and objective way to assess lameness
53 could be provided by the use of currently available sensing technology to automatically
54 evaluate and monitor walking ability in the broiler flock. Several experimental methods have
55 been developed. Examples include; use of force plates (Corr et al. 2007) or image analysis
56 techniques, including the use of optical flow patterns (Dawkins et al. 2009). The eYeNamic
57 TM system (Fancom BV, The Netherlands) uses video cameras and image processing methods
58 (Kashiha et al. 2013) to monitor a relatively uncomplicated variable; the activity of a broiler
59 flock. The eYeNamic TM system was used to determine whether there were correlations
60 between broiler activity and the level of lameness, using the Gait Score-method as a reference
61 or ‘gold standard’ (Aydin et al. 2010, 2013). A non-linear relationship was found (Aydin et
62 al. 2010, 2013) where birds with gait score 3 showed the highest levels of activity. These
63 results suggest that general activity as such cannot be used to predict gait score in a flock
64 (Aydin et al. 2010, 2013, Dawkins et al. 2009 and Dawkins et al. 2012). Therefore, in the
65 current experiment, we decided to create a challenge, and induce movement of the birds by
66 the presence of a human. An assessment was made of the activity measures before and after a
67 human had walked through the flock, and it was then determined whether activity patterns
68 observed around this human activity were related to Gait Score.

69

70

MATERIALS AND METHODS

Flocks and farms

72 This experiment was conducted at five commercial broiler farms in four European countries
73 (Italy, Spain, the Netherlands and two farms in the U.K). All regulations regarding ethical
74 care and use of animals were strictly followed. In total, Gait Score data were collected from
75 16 flocks and 33 assessments occasions, at the ages of 3, 4 and/or 5 weeks (Table 1) whilst

76 automated image based calculation of activity was carried out continuously over the entire
77 rearing period. The farmers walked through the flock a minimum of twice per day, walking
78 up and down each feeder and drinker row. The human-animal contact was mainly visual and
79 the farmer examined the birds visually and removed sick or dead birds during the walk.

80

81 Table 1. The distribution of data collection

No. of farms	5
No. of flocks visited	16
No. of assessments	33
No. of assessments at age 3 weeks	12
No. of assessments at age 4 weeks	10
No. of assessments at age 5 weeks	11

82

83

84 *Measures*


85 The birds' walking ability was assessed using the Gait Scoring-method presented in the
86 Welfare Quality® protocols for poultry (Welfare Quality®, 2009; Kestin et al. 1992), where
87 the bird's gait is graded between 0 (perfect walking) to 5 (unable to move). At each
88 assessment, 100-200 birds were randomly selected and a portable fenced arena was placed
89 around the group of birds, using the procedures described in the Welfare Quality® protocol.
90 Up to 5 groups were fenced at different locations in the house, which were distributed in a
91 randomized fashion. All birds within a sampling pen were scored. The assessment scores
92 were collected by trained assessors, one in each country, except in the U.K. where two
93 trained assessors visited the farms.

94 Automatic activity recordings from the eYeNamic™ system were collected with overhead
95 cameras mounted on the ceiling. Only one camera was used to collect data in this experiment,
96 it was the one positioned close to the entrance. Recording began when the assessor entered
97 the poultry house. The system enabled three measurements per minute. Activity was
98 calculated as an index between zero (no activity) and 100 (all bird pixels have moved
99 between two consecutive frames), based on image analysis technology (Kashiha et al. 2013).

100

101 *Experimental procedure*

102 The experimental procedure started with the collection of baseline activity for 10 minutes. No
103 disturbance was allowed in the broiler flock during this time. After 10 minutes, the assessor
104 entered the animal house, walked along the side of the whole house, turned at the end and
105 walked back in a straight line through the middle of the house below the camera (see Figure
106 1). The aim was to mimic the way a farmer would move through the flock during the daily
107 check of the animals, in a usual and more or less standardized way, and with a uniform
108 moderate walking pace. The period when the human is present cannot be used within
109 calculations, since the used eYeNamic™-system does not differentiate between ‘human-
110 pixels’ and ‘bird-pixels’. Hence the activity recordings when the human is in the house are a
111 mix of bird and human movement. After the walking procedure, the flock was left alone
112 without any disturbance for 15 minutes of activity recording. After this re-settling period, the
113 assessor re-entered the house and gait scoring was carried out, by fencing a groups of birds (a
114 total of 100-200 birds) and scoring the birds while they walked out of the arena.



Entrance

115

116 Figure 1: The standardized procedure of walking through the broiler flock.

117

118 **Statistical analysis**

119 The variables calculated from the activity recordings are presented in table 2.

120 Table 2. Variables derived from the activity recordings.

121

Variable	Definition
Baseline activity	Average activity index during 10 minutes before the assessor entered
Time to return to baseline (s)	Time from when the assessor left the house to when the animals resumed baseline levels of activity
Average activity after	Average activity during the time to return to baseline activity
Δ Amplitude	The difference between highest activity peak after assessor left the flock and baseline level

122

123

124 The data sets were modelled using the multilevel statistical software package MLwiN
125 (Rasbash et al. 2009). The multilevel structure of the data was specified within MLwiN as
126 measurement occasion, within flock, within farm, whereas ‘Gait Score’ was modelled using a
127 GLM approach. Age, Δ Amplitude, Baseline activity, Average activity after and Time to
128 return to baseline were all tested as covariates within the model. Those significant at $p \leq 0.05$
129 were retained in the final model (see Table 3) which can be used to predict the degree of
130 lameness (gait score) in the flock. The statistical analysis resulted in the following predictive
131 equation for GS:

132

$$133 \text{ Gait Score}_{ijk} = \beta_0_{ijk} \text{ const} + \beta_1 \Delta \text{ Amplitude}_{ijk} + \beta_2 \text{ Age}_{ijk}$$

134

135

RESULTS

136 *Activity recordings*

137 Figure 2 describes the characteristics of the activity recordings. The period with the walking
138 human is visible in the graph due to the increase and irregularity of the registered activity
139 index. The reaction of the birds (Δ Amplitude) is visible as a peak in activity and a
140 characteristic slope is shown when the birds recover from the disturbance and return to
141 baseline activity levels.



142

143 Figure 2. A representative example of activity pattern of the broilers during the experimental
144 procedure. The striped lines indicate when the assessor entered and left the stable.

145

146 ***Gait Score***

147 In general, gait score means showed little variation over time and all the flocks had low GS
148 scores (3 weeks =1.4±0.6, 4 weeks = 1.5±0.6 and 5 weeks = 1.9±0.6, (mean ± SD)), thereby
149 showing good leg health status. The observed trend towards increased gait scores with
150 increasing age was noted, and was expected (see Discussion).

151

152 ***Prediction of Gait Score using activity index recordings***

153 An overview of the gait score - and activity measures used in the statistical analysis are
154 presented as mean values in table 3.

155

156 Table 3. Mean values of gait score and activity measures presented by age.

Age	Gait Score	Baseline activity	Time to return to baseline	Average activity after	Δ Amplitude
3	1.4	6.29	388.25	9.67	17.22
4	1.5	6.28	368.70	9.01	13.24
5	1.9	5.82	391.91	7.28	7.57

157

158

159 Δ Amplitude and Age were significantly related to Gait Score and are thus included in the
160 equation. The levels of an effect (β), the standard errors and p-values are presented in table 4.

161 The indices ‘ijk’ refer to the hierarchical levels; assessment occasion, flock then farm,
162 respectively.

163

164

165

166

167

168 Table 4. The parameter estimates from the model predicting GS from the automated activity
 169 measurements of the flocks (the parameter estimates and p values shown for the significant
 170 variables are estimates when the non-significant variables have been excluded from the
 171 model).

Variable	β	se	p-value
Constant	1.045	0.269	< 0.001
Δ Amplitude	- 0.011	0.003	<0.0002
Age	0.230	0.031	< 0.001
Baseline activity	0.009	0.011	0.416
Time to return to baseline (s)	-0.00011	0.0000085	0.185
Average activity after	-0.001	0.011	0.927

172

173

DISCUSSION

174 The present study used a repeated, simulated, animal inspection routine where a person
 175 walked through the flock to challenge the birds and elicit changes in their locomotor activity
 176 that could be automatically measured using the eYeNamic™ system. The statistical analysis
 177 then determined if such activity changes were related to walking ability (lameness).

178 The variable Δ Amplitude, (which is a measure of the birds' direct response to an
 179 approaching human (moving away reveals their walking ability)), was significantly related to
 180 walking ability and might thus be used to predict the Gait Score level in a flock. Similarly, an
 181 earlier study found that Δ Amplitude and age were the strongest variables for predicting the
 182 human-animal relationship in a broiler flock (Silvera et al, unpublished data). Using the same
 183 methods as in this experiment the results from an Avoidance Distance Touch Test, where the

184 distance from an approaching human is assessed as a measure of fear of humans (Graml et
185 al. 2008), could be predicted from activity recordings by the eYeNamic™ system. It can
186 also be considered that the Δ Amplitude can be related to birds moving away from the human
187 as a response of fear of humans (Jones and Waddington, 1992). Since the method used in
188 this experiment was designed as an imitation of the daily routine conducted by the farmer, the
189 level of fearfulness was considered to be low due to habituation and thus not affecting the
190 outcome.

191 The relationships presented in this study show promise for the future development of a fully
192 automatic continuous assessment system. Since the model is directly fitted to the current data
193 and experimental settings, it cannot be used to predict Gait Score as accurately as in this
194 study in commercial broiler flocks as such. Its performance needs to be tested in more flocks
195 where the background variability may be greater. Both lameness and activity levels may vary
196 between individual flocks and farms, which makes it desirable to test the method on a data set
197 with a larger span of gait scores and activity levels. As in several automated tools in livestock
198 monitoring a self-adapting algorithm can solve these problems. Nevertheless, the results
199 from this experiment show that there is a relationship worth further detailed study and that
200 could be of potential for commercial use. The kurtosis and scw of the distribution of the
201 activity data in this study would be interesting to add to future development of the method. In
202 the study of Dawkins et al. (2012) positive correlations was found between the skew and
203 kurtosis in optical flow data and gait score. The results showed that it could predict the gait
204 score at 28 days already several days on beforehand. The experiment had like the present
205 study a narrow range of gait scores (average score: 1.92 ± 0.23) which further advocates the
206 investigation of the correlations in future research. Further investigation of the scw and
207 kurtosis of the data is also supported by the results by Roberts et al. (2012), where Bayesian

208 regression on continuous optical flow data predicted future Gait Score results in broiler
209 flocks.

210 The birds' age also had a significant and positive effect in predicting Gait Score. This is
211 consistent with previous reports (Sørensen et al. 2000, Weeks et al. 2000, Kestin et al. 2001)
212 and is likely a side effect of the very rapid increase in live weight with age.

213 Our finding that baseline activity measures had no significant effect probably reflected the
214 fact that broiler chickens show low activity in general with 76-86% of the flock lying down
215 (Weeks et al. 2000). However, measurement of baseline activity is necessary because it is
216 included in the equation used to calculate Δ Amplitude. Furthermore, the age-related increase
217 in lameness (Weeks et al. 2000) suggests that a more detailed on-farm study of general
218 activity could be worthwhile.

219 ***Conclusion***

220 The present results demonstrate the potential value of using image analysis techniques for
221 automated assessment of lameness in commercial broiler flocks. The fact that the prediction
222 of Gait Score was possible, even when the general leg health in the studied flocks was very
223 good and had a narrow range (GS: 1.4-1.9) suggests future research to develop an automatic
224 continuous on-farm assessment method.

225

226

227

ACKNOWLEDGEMENTS

228 The authors gratefully acknowledge the European Community for financial support in
229 Collaborative Project EU-PLF KBBE.2012.1.1-02-311825 under the Seventh Framework
230 Programme. A special thanks to the participating farms and to Déborah Temple, Ilaria
231 Fontana, Gemma Richards, Henk Gunnink and Steven Brown for carrying out the
232 assessments. Thanks also to Luc Rooijackers, Eric Koenders and Tom van Hertem at
233 Fancom BV for technical support and for coordinating the video recordings. We also thank
234 Bryan Jones for editing the manuscript and helpful comments.

235

REFERENCES

- 236
237 Aydin, A., Cangar, O., Ozcan, S.E., Bahr, C. and Berckmans, D. 2010. Application of a fully
238 automatic analysis tool to assess the activity of broiler chickens with different gait scores.
239 *Comput. Electron. Agric.* 73:194-199.
- 240 Aydin, A., Pluk, A., Leroy, T., Berckmans, D. and Bahr, C. 2013. Automatic identification of
241 activity and spatial use of broiler chickens with different gait scores. *T ASABE* 56:1123-
242 1132.
- 243 Bessei, W. (2006) Welfare of broilers: a review. *World Poultry Sci J*, 62:455-466.
- 244 Bradshaw, R. H., R. D. Kirkden, and D. M. Broom. 2002. A review of the aetiology and
245 pathology of leg weakness in broilers in relation to welfare. *Avian Poult Biol Rev* 13:45-
246 103).
- 247 Butterworth, A. and Haslam, S.M. (2009) A lameness control strategy for broiler fowls,
248 Welfare Quality reports No. 13, Welfare Quality reports (Cardiff, UK, Cardiff School of City
249 and Regional Planning). 56 pp.
- 250 Caplen, G., Colbrone, G.R., Hothersall, B., Nicol, C.J., Waterman-Pearson, A.E., Weeks,
251 C.A. and Murrell, J.C. 2013. Lameness broiler chickens respond to non-steroidal anti-
252 inflammatory drugs with objective changes in gait function: A controlled clinical trial. *The*
253 *Vet J* 196:477-482.
- 254 Corr, S.A., McCorquodale, C., McDonald, J., Gentle, M. and McGovern, R. 2007. A force
255 plate study of avian gait. *J Biomech* 40: 2037-2043.
- 256 Danbury, T.C., Weeks, C.A., Waterman-Pearson, A.E., Kestin, S.C. and Chambers, J.P.
257 (2000) Self-selection of the analgesic drug carprofen by lame broiler chickens. *Vet. Rec.*
258 146:307-311.
- 259 Dawkins, M.S., Lee, H-J., Waitt, C.D. and Roberts, S.J. 2009. Optical flow patterns in broiler
260 chicken flocks as automated measures of behaviour and gait. *Appl. Anim. Behav. Sci.* 119:
261 203-209.

262 Dawkins, M.S., Cain, R. and Roberts, S.J. 2012. Optical flow, flock behaviour and chicken
263 welfare. *Anim Behav* 84 :219-223

264 Graml, C., Waiblinger, S. and Niebuhr, K. 2008. Validation of tests for on-farm assessment
265 of the hen-human relationship in non-cage systems. *Appl. Anim. Behav. Sci* 111, 301-310.

266 Jones, R.B. and Waddington, D. 1992. Modification of fear in domestic chicks, *Gallus gallus*
267 *domesticus*, via regular handling an early environmental enrichment. *Anim. Behav.* 43:1021-
268 1033.

269 Kashiha, M., Pluk , A., Bahr, C., Vranken, E. and Berckmans, D. 2013. Development of an
270 early warning system for a broiler house using computer vision. *Biosyst Eng* 116: 36-45.

271 Kapell, D.N.R.G., Hill, W.G., Neeteson, A-M., McAdam, J., Koerhuis, A.N.M. and
272 Avendaño, S. 2012. Twenty-five years of selection for improved leg health in purebred
273 broiler lines and underlying genetic parameters. *Poult Sci.* 91:3032-3043.

274 Kestin, S.C., Knowles, T.G., Tinch, A.E. and Gregory, N.G. 1992. Prevalence of leg
275 weakness in broiler chickens and its relationship with genotype. *Vet. Rec.* 131: 190-194.

276 Kestin, S.C., Gordon, S., Su, G. and Sørensen, P. 2001. Relationships in broiler chickens
277 between lameness, liveweight, growth rate and age. *Vet. Rec.* 148:195-197.

278 Knowles, T. G., S. C. Kestin, S. M. Haslam, S. N. Brown, L. E. Green, A. Butterworth, S. J.
279 Pope, D. Pfeiffer, and C. J. Nicol. 2008. Leg Disorders in Broiler Chickens: Prevalence, Risk
280 Factors and Prevention. *PLoS One* 3: e1545.

281 Marchewka ,J., Watanabe, T. T. N., Ferrante V. and Estevez , I. 2013. Welfare assessment in
282 broiler farms: Transect walks versus individual scoring. *Poult Sci.* 92:2588-2599.

283 Mench, J. 2004. Lameness. in: Weeks, C. and Butterworth, A. (Eds.) *Measuring and*
284 *Auditing Broiler Welfare.* CAB International. Oxfordshire. U.K. 3-17 pp.

285 McGeown, D., Danbury, T.C., Waterman-Pearson, A.E. and Kestin, S.C. 1999. Effect of
286 carprofen on lameness in broiler chickens. *Vet. Rec.* 144:668-671.

287 Rasbash, J., Charlton, C., Browne, W.J., Healy, M. and Cameron, B. 2009. MLwiN Version
288 2.1. Centre for Multilevel Modelling, University of Bristol.

289 Roberts, S.J., Cain, R. and Dawkins. M.S. 2012. Prediction of welfare outcomes for broiler
290 chickens using Bayesian regression on continuous optical flow data. . J. R. Soc. Interface
291 9:3436–3443.

292 Sørensen, P., Su, G. and Kestin, S.C. 2000. Effects of age and stocking density on leg
293 weakness in broiler chickens. *Poult Sci.* 79:864-870.

294 Weeks, C.A., Danbury, T.D., Davies, H.C., Hunt, P. and Kestin, S.C. 2000. The behaviour of
295 broiler chickens and its modification by lameness. *Appl. Anim. Behav. Sci.* 67:111-125.

296 Weeks, C.A., Knowles, T.G., Gordon, R.G., Kerr, A.E., Peyton, S.T. and Tillbrook, N.T.
297 2002. New method for objectively assessing lameness in broiler chickens. *Vet. Rec.* 151:762-
298 764.

299 Welfare Quality® (2009) Welfare Quality® assessment protocol for poultry (broilers, laying
300 hens), (Lelystad, The Netherlands, Welfare Quality® Consortium). 114 pp.

301 Wideman, R.F., Hamal, K.R., Stark, J.M., Blankenship, J., Lester, H., Mitchell, K.N.,
302 Lorenzoni, G. and Pevzner, I. 2012. A wireflooring model for inducing lameness in broilers:
303 Evaluation of probiotics as a prophylactic treatment. *Poult Sci.* 91:870-883.